

**On the Complete Elliptic Integrals and Babylonian Identity II:
An Approximation for the Complete Elliptic Integral of the first kind**

Edigles Guedes

edigles.guedes@gmail.com

December 15, 2013

He who did not spare his own Son but gave him up for us all, how will he not also with him graciously give us all things? — Romans 8:32.

Abstract. Using the Theorem 4 of previous paper, I evaluate the complete elliptic integral of the first kind in approximate analytical closed form, by means of Bessel functions.

1. Introduction

By means of the theorem 4 of previous paper, I demonstrated the following approximations:

$$K(k) \approx \left(\frac{\pi + \sqrt{\pi}}{2}\right) \left[I_0\left(\frac{k}{2}\right)\right]^2 + \frac{\sqrt{\pi}}{2} k I_0\left(\frac{k}{2}\right) I_1\left(\frac{k}{2}\right) - \frac{\sqrt{\pi}(4 + k^2)}{8}$$

and

$$K(k) \approx \left(\frac{\pi + \sqrt{\pi}}{2}\right) \left[I_0\left(\frac{k}{2}\right)\right]^2 + \pi \left[I_1\left(\frac{k}{2}\right)\right]^2 + \frac{7\sqrt{\pi}}{8} k I_0\left(\frac{k}{2}\right) I_1\left(\frac{k}{2}\right) - \left(\frac{2\pi k^2 + 7\sqrt{\pi} k^2 + 16\sqrt{\pi}}{32}\right),$$

for $0 < k \leq 1/2$.

2. THEOREMS

Theorem 1. For $0 < k \leq 1/2$, then

$$K(k) \approx \left(\frac{\pi + \sqrt{\pi}}{2}\right) \left[I_0\left(\frac{k}{2}\right)\right]^2 + \frac{\sqrt{\pi}}{2} k I_0\left(\frac{k}{2}\right) I_1\left(\frac{k}{2}\right) - \frac{\sqrt{\pi}(4 + k^2)}{8},$$

where $K(k)$ denotes the complete elliptic integral of first kind and $I_\nu(z)$ denotes the modified Bessel function of the first kind.

Proof. I will prove the equality above, developing the left hand side of the equation below

$$(1) \quad K(k) - \frac{\pi}{2} \left[I_0\left(\frac{k}{2}\right)\right]^2 = I_k.$$

First, I encounter

$$(2) \quad \begin{aligned} \frac{\pi}{2} \left[I_0\left(\frac{k}{2}\right)\right]^2 &= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \Gamma\left(m + \frac{1}{2}\right) \frac{k^{2m}}{2^{2m}(m!)^3} = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \frac{(2m)!}{2^{2m}m!} \cdot \frac{k^{2m}}{2^{2m}(m!)^3} = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \frac{(2m)! k^{2m}}{2^{4m}(m!)^4} \\ &= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \binom{2m}{m} \frac{k^{2m}}{2^{4m}(m!)^2}. \end{aligned}$$

On the other hand, in [1], I proved the identity

$$\begin{aligned}
(3) \quad K(k) &= \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)! \Gamma\left(m + \frac{3}{2}\right)}{(2m+1)} \frac{k^{2m}}{2^{2m}(m!)^3} = \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)!(2m+2)!}{(2m+1)2^{2m+2}(m+1)!} \frac{k^{2m}}{2^{2m}(m!)^3} \\
&= \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)!(2m+2)(2m+1)(2m)!}{(2m+1)2^{2m+2}(m+1)m!} \cdot \frac{k^{2m}}{2^{2m}(m!)^3} \\
&= \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)!2(m+1)(2m+1)(2m)!}{(2m+1)2^{2m+2}(m+1)m!} \cdot \frac{k^{2m}}{2^{2m}(m!)^3} \\
&= \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)!(2m)!}{2^{2m+1}m!} \cdot \frac{k^{2m}}{2^{2m}(m!)^3} = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \frac{(2m)!}{2^{2m}m!} \cdot \frac{(2m)!k^{2m}}{2^{2m}(m!)^3} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \frac{[(2m)!]^2 k^{2m}}{2^{4m}(m!)^4} = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \binom{2m}{m}^2 \frac{k^{2m}}{2^{4m}}.
\end{aligned}$$

provided that $0 < k < 1$. I subtract (3) of (2), and obtain

$$\begin{aligned}
(4) \quad K(k) - \frac{\pi}{2} \left[I_0\left(\frac{k}{2}\right) \right]^2 &= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \binom{2m}{m}^2 \frac{k^{2m}}{2^{4m}} - \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \binom{2m}{m} \frac{k^{2m}}{2^{4m}(m!)^2} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \left[\binom{2m}{m} - \frac{1}{(m!)^2} \right] \binom{2m}{m} \frac{k^{2m}}{2^{4m}} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \left[\frac{(2m)!}{(m!)^2} - \frac{1}{(m!)^2} \right] \frac{(2m)!k^{2m}}{(m!)^2 2^{4m}} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} [(2m)! - 1] \frac{(2m)!k^{2m}}{(m!)^4 2^{4m}} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=1}^{\infty} [(2m)! - 1] \frac{(2m)!k^{2m}}{(m!)^4 2^{4m}} = \frac{\sqrt{\pi}k^2}{2^4} + \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} [(2m)! - 1] \frac{(2m)!k^{2m}}{(m!)^4 2^{4m}}.
\end{aligned}$$

I note that, if $m \geq 2$, then

$$(5) \quad (2m)! - 1 \geq 5m + 1.$$

I take (5) into (4)

$$\begin{aligned}
(6) \quad K(k) - \frac{\pi}{2} \left[I_0\left(\frac{k}{2}\right) \right]^2 &> \frac{\sqrt{\pi}k^2}{2^4} + \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} (5m+1) \frac{(2m)!k^{2m}}{(m!)^4 2^{4m}} \\
&= \frac{\sqrt{\pi}k^2}{2^4} + \frac{5\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \frac{(2m)!mk^{2m}}{(m!)^4 2^{4m}} + \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \frac{(2m)!k^{2m}}{(m!)^4 2^{4m}} \\
&= \frac{\sqrt{\pi}k^2}{2^4} + \frac{5\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m-1)!m! 2^{4m}} + \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m!)^2 2^{4m}}.
\end{aligned}$$

I calculate

$$(7) \frac{5\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m-1)! m! 2^{4m}} = \frac{5\sqrt{\pi}k^2}{16} {}_1F_2\left(\frac{3}{2}; 2, 2; \frac{k^2}{4}\right) - \frac{5\sqrt{\pi}k^2}{16} = \frac{5\sqrt{\pi}}{4} k I_0\left(\frac{k}{2}\right) I_1\left(\frac{k}{2}\right) - \frac{5\sqrt{\pi}k^2}{16}$$

and

$$(8) \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m!)^2 2^{4m}} = \frac{\sqrt{\pi}}{2} {}_1F_2\left(\frac{1}{2}; 1, 1; \frac{k^2}{4}\right) = -\frac{\sqrt{\pi}k^2}{16} + \frac{\sqrt{\pi}}{2} \left[I_0\left(\frac{k}{2}\right)\right]^2 - \frac{\sqrt{\pi}}{2}.$$

From (6), (7) and (8), I obtain

$$K(k) - \frac{\pi}{2} \left[I_0\left(\frac{k}{2}\right)\right]^2 > \frac{\sqrt{\pi}k^2}{16} + \frac{5\sqrt{\pi}}{4} k I_0\left(\frac{k}{2}\right) I_1\left(\frac{k}{2}\right) - \frac{5\sqrt{\pi}k^2}{16} - \frac{\sqrt{\pi}k^2}{16} + \frac{\sqrt{\pi}}{2} \left[I_0\left(\frac{k}{2}\right)\right]^2 - \frac{\sqrt{\pi}}{2},$$

hence,

$$K(k) > \left(\frac{\pi + \sqrt{\pi}}{2}\right) \left[I_0\left(\frac{k}{2}\right)\right]^2 + \frac{5\sqrt{\pi}}{4} k I_0\left(\frac{k}{2}\right) I_1\left(\frac{k}{2}\right) - \sqrt{\pi} \left(\frac{5k^2 + 8}{16}\right),$$

since $0 < k < 1/2$, I conclude easily that

$$K(k) \approx \left(\frac{\pi + \sqrt{\pi}}{2}\right) \left[I_0\left(\frac{k}{2}\right)\right]^2 + \frac{5\sqrt{\pi}}{4} k I_0\left(\frac{k}{2}\right) I_1\left(\frac{k}{2}\right) - \sqrt{\pi} \left(\frac{5k^2 + 8}{16}\right).$$

This completed the proof. The reader can see the Table 1. \square

Theorem 2. For $0 < k \leq 1/2$, then

$$K(k) \approx \left(\frac{\pi + \sqrt{\pi}}{2}\right) \left[I_0\left(\frac{k}{2}\right)\right]^2 + \pi \left[I_1\left(\frac{k}{2}\right)\right]^2 + \frac{7\sqrt{\pi}}{8} k I_0\left(\frac{k}{2}\right) I_1\left(\frac{k}{2}\right) - \left(\frac{2\pi k^2 + 7\sqrt{\pi}k^2 + 16\sqrt{\pi}}{32}\right),$$

where $K(k)$ denotes the complete elliptic integral of first kind and $I_\nu(z)$ denotes the modified Bessel function of the first kind.

Proof. I will prove the equality above, developing the left hand side of the equation below

$$(9) \quad K(k) - \left\{ \frac{\pi}{2} \left[I_0\left(\frac{k}{2}\right)\right]^2 + \pi \left[I_1\left(\frac{k}{2}\right)\right]^2 \right\} = I_k.$$

First, I encounter

$$(10) \quad \frac{\pi}{2} \left[I_0\left(\frac{k}{2}\right)\right]^2 = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \Gamma\left(m + \frac{1}{2}\right) \frac{k^{2m}}{2^{2m} (m!)^3} = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \frac{(2m)!}{2^{2m} m!} \cdot \frac{k^{2m}}{2^{2m} (m!)^3} = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \frac{(2m)! k^{2m}}{2^{4m} (m!)^4} \\ = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \binom{2m}{m} \frac{k^{2m}}{2^{4m} (m!)^2}$$

and

$$(11) \quad \pi \left[I_1\left(\frac{k}{2}\right)\right]^2 = \sqrt{\pi} \sum_{m=0}^{\infty} \frac{\Gamma\left(\frac{2m+1}{2}\right)}{\Gamma(m+1)\Gamma(m+2)\Gamma(m)} \frac{k^{2m}}{2^{2m}} = \sqrt{\pi} \sum_{m=0}^{\infty} \frac{\Gamma\left(m + \frac{1}{2}\right)}{(m-1)! m! (m+1)!} \frac{k^{2m}}{2^{2m}} \\ = \pi \sum_{m=0}^{\infty} \frac{(2m)!}{(m-1)! (m!)^2 (m+1)!} \frac{k^{2m}}{2^{4m}} = \pi \sum_{m=0}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m-1)! (m+1)! 2^{4m}}.$$

On the other hand, in [1], I proved the identity

$$\begin{aligned}
(12) \quad K(k) &= \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)! \Gamma\left(m + \frac{3}{2}\right)}{(2m+1)} \frac{k^{2m}}{2^{2m}(m!)^3} = \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)! (2m+2)!}{(2m+1)2^{2m+2}(m+1)!} \frac{k^{2m}}{2^{2m}(m!)^3} \\
&= \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)! (2m+2)(2m+1)(2m)!}{(2m+1)2^{2m+2}(m+1)m!} \cdot \frac{k^{2m}}{2^{2m}(m!)^3} \\
&= \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)! 2(m+1)(2m+1)(2m)!}{(2m+1)2^{2m+2}(m+1)m!} \cdot \frac{k^{2m}}{2^{2m}(m!)^3} \\
&= \sqrt{\pi} \sum_{m=0}^{\infty} \frac{(2m)! (2m)!}{2^{2m+1}m!} \cdot \frac{k^{2m}}{2^{2m}(m!)^3} = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \frac{(2m)!}{2^{2m}m!} \cdot \frac{(2m)! k^{2m}}{2^{2m}(m!)^3} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \frac{[(2m)!]^2 k^{2m}}{2^{4m}(m!)^4} = \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \binom{2m}{m}^2 \frac{k^{2m}}{2^{4m}}.
\end{aligned}$$

provided that $0 < k < 1$. I subtract (12) of (11) and (10), to obtain

$$\begin{aligned}
(13) \quad K(k) &- \left\{ \frac{\pi}{2} \left[I_0\left(\frac{k}{2}\right) \right]^2 + \pi \left[I_1\left(\frac{k}{2}\right) \right]^2 \right\} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \binom{2m}{m}^2 \frac{k^{2m}}{2^{4m}} - \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \binom{2m}{m} \frac{k^{2m}}{2^{4m}(m!)^2} - \pi \sum_{m=0}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m-1)!(m+1)! 2^{4m}} \\
&= \sqrt{\pi} \sum_{m=0}^{\infty} \left[\frac{1}{2} \binom{2m}{m} - \frac{1}{2(m!)^2} - \frac{\sqrt{\pi}}{(m-1)!(m+1)!} \right] \binom{2m}{m} \frac{k^{2m}}{2^{4m}} \\
&= \sqrt{\pi} \sum_{m=0}^{\infty} \left[\frac{1}{2} \frac{(2m)!}{(m!)^2} - \frac{1}{2(m!)^2} - \frac{\sqrt{\pi}}{(m-1)!(m+1)!} \right] \frac{(2m)! k^{2m}}{(m!)^2 2^{4m}} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \left[\frac{(2m)!}{(m!)^2} - \frac{1}{(m!)^2} - \frac{2m\sqrt{\pi}}{m(m-1)!(m+1)m!} \right] \frac{(2m)! k^{2m}}{(m!)^2 2^{4m}} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \left[\frac{(2m)!}{(m!)^2} - \frac{1}{(m!)^2} - \frac{2m\sqrt{\pi}}{(m!)^2(m+1)} \right] \frac{(2m)! k^{2m}}{(m!)^2 2^{4m}} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=0}^{\infty} \left[(2m)! - 1 - \frac{2m\sqrt{\pi}}{m+1} \right] \frac{(2m)! k^{2m}}{(m!)^4 2^{4m}} \\
&= \frac{\sqrt{\pi}}{2} \sum_{m=1}^{\infty} \left[(2m)! - 1 - \frac{2m\sqrt{\pi}}{m+1} \right] \frac{(2m)! k^{2m}}{(m!)^4 2^{4m}} \\
&= -\left(\frac{\pi - \sqrt{\pi}}{16}\right) k^2 + \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \left[(2m)! - 1 - \frac{2m\sqrt{\pi}}{m+1} \right] \frac{(2m)! k^{2m}}{(m!)^4 2^{4m}}.
\end{aligned}$$

I note that, if $m \geq 2$, then

$$(14) \quad (2m)! - 1 - \frac{2m\sqrt{\pi}}{m+1} > \frac{7m}{2} + 1.$$

I take (14) into (13)

$$(15) \quad \begin{aligned} K(k) - \left\{ \frac{\pi}{2} \left[I_0 \left(\frac{k}{2} \right) \right]^2 + \pi \left[I_1 \left(\frac{k}{2} \right) \right]^2 \right\} &> \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \left(\frac{7m}{2} + 1 \right) \frac{(2m)! k^{2m}}{(m!)^4 2^{4m}} - \left(\frac{\pi - \sqrt{\pi}}{16} \right) k^2 \\ &= \frac{7\sqrt{\pi}}{4} \sum_{m=2}^{\infty} \frac{(2m)! m k^{2m}}{(m!)^4 2^{4m}} + \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \frac{(2m)! k^{2m}}{(m!)^4 2^{4m}} - \left(\frac{\pi - \sqrt{\pi}}{16} \right) k^2 \\ &= \frac{7\sqrt{\pi}}{4} \sum_{m=2}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m-1)! m! 2^{4m}} + \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m!)^2 2^{4m}} - \left(\frac{\pi - \sqrt{\pi}}{16} \right) k^2. \end{aligned}$$

I calculate

$$(16) \quad \begin{aligned} \frac{7\sqrt{\pi}}{4} \sum_{m=2}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m-1)! m! 2^{4m}} &= \frac{7\sqrt{\pi} k^2}{32} {}_1F_2 \left(\frac{3}{2}; 2, 2; \frac{k^2}{4} \right) - \frac{7\sqrt{\pi} k^2}{32} \\ &= \frac{7\sqrt{\pi}}{8} k I_0 \left(\frac{k}{2} \right) I_1 \left(\frac{k}{2} \right) - \frac{7\sqrt{\pi} k^2}{32} \end{aligned}$$

and

$$(17) \quad \frac{\sqrt{\pi}}{2} \sum_{m=2}^{\infty} \binom{2m}{m} \frac{k^{2m}}{(m!)^2 2^{4m}} = \frac{\sqrt{\pi}}{2} {}_1F_2 \left(\frac{1}{2}; 1, 1; \frac{k^2}{4} \right) = -\frac{\sqrt{\pi} k^2}{16} + \frac{\sqrt{\pi}}{2} \left[I_0 \left(\frac{k}{2} \right) \right]^2 - \frac{\sqrt{\pi}}{2}.$$

From (15), (16) and (17), I obtain

$$K(k) > \left(\frac{\pi + \sqrt{\pi}}{2} \right) \left[I_0 \left(\frac{k}{2} \right) \right]^2 + \pi \left[I_1 \left(\frac{k}{2} \right) \right]^2 + \frac{7\sqrt{\pi}}{8} k I_0 \left(\frac{k}{2} \right) I_1 \left(\frac{k}{2} \right) - \left(\frac{2\pi k^2 + 7\sqrt{\pi} k^2 + 16\sqrt{\pi}}{32} \right),$$

since $0 < k < 1/2$, I conclude easily that

$$K(k) \approx \left(\frac{\pi + \sqrt{\pi}}{2} \right) \left[I_0 \left(\frac{k}{2} \right) \right]^2 + \pi \left[I_1 \left(\frac{k}{2} \right) \right]^2 + \frac{7\sqrt{\pi}}{8} k I_0 \left(\frac{k}{2} \right) I_1 \left(\frac{k}{2} \right) - \left(\frac{2\pi k^2 + 7\sqrt{\pi} k^2 + 16\sqrt{\pi}}{32} \right).$$

This completed the proof. The reader can see the Table 2. \square

REFERENCES

- [1] Guedes, Edigles, *On the Complete Elliptic Integrals and Babylonian Identity I: The $1/\pi$ formulaes Involving Gamma functions and Summations*, 2013.

Table 1

In this table, I have: first column: m ; second column: $k = 1/m$; third column: $\left(\frac{\pi+\sqrt{\pi}}{2}\right)\left[I_0\left(\frac{1}{2m}\right)\right]^2 + \frac{5\sqrt{\pi}}{4m}I_0\left(\frac{1}{2m}\right)I_1\left(\frac{1}{2m}\right) - \frac{\sqrt{\pi}}{16}\left[5\left(\frac{1}{m}\right)^2 + 8\right]$; fourth column: $K\left(\frac{1}{m}\right)$; fifth column: $\frac{K\left(\frac{1}{m}\right)}{\left(\frac{\pi+\sqrt{\pi}}{2}\right)\left[I_0\left(\frac{1}{2m}\right)\right]^2 + \frac{5\sqrt{\pi}}{4m}I_0\left(\frac{1}{2m}\right)I_1\left(\frac{1}{2m}\right) - \frac{\sqrt{\pi}}{16}\left[5\left(\frac{1}{m}\right)^2 + 8\right]}$.

2	$\frac{1}{2}$	1.651757078	1.685750354	1.02058007
3	$\frac{1}{3}$	1.6057434	1.617386735	1.007251055
4	$\frac{1}{4}$	1.590251421	1.596242222	1.003767203
5	$\frac{1}{5}$	1.583187698	1.586867847	1.002324518
6	$\frac{1}{6}$	1.579378879	1.581878436	1.00158262
7	$\frac{1}{7}$	1.577091885	1.578903918	1.001148971
8	$\frac{1}{8}$	1.5756114	1.576986771	1.000872912
9	$\frac{1}{9}$	1.574598142	1.575678422	1.000686067
10	$\frac{1}{10}$	1.57387424	1.574745561	1.000553615
11	$\frac{1}{11}$	1.573339105	1.574056948	1.000456254
12	$\frac{1}{12}$	1.572932358	1.573534108	1.000382565
13	$\frac{1}{13}$	1.572615974	1.573127756	1.000325433
14	$\frac{1}{14}$	1.572365032	1.572805664	1.000280234
15	$\frac{1}{15}$	1.57216265	1.572546032	1.000243857
16	$\frac{1}{16}$	1.571997057	1.572333687	1.000214141
17	$\frac{1}{17}$	1.571859847	1.572157798	1.000189553
18	$\frac{1}{18}$	1.571744884	1.572010469	1.000168974
19	$\frac{1}{19}$	1.571647605	1.571885834	1.000151578
20	$\frac{1}{20}$	1.571564561	1.571779457	1.00013674
21	$\frac{1}{21}$	1.571493103	1.571687938	1.000123981
22	$\frac{1}{22}$	1.571431171	1.571608632	1.000112929
23	$\frac{1}{23}$	1.571377145	1.571539459	1.000103293
24	$\frac{1}{24}$	1.571329734	1.571478762	1.000094841
25	$\frac{1}{25}$	1.571287901	1.571425211	1.000087387
26	$\frac{1}{26}$	1.571250803	1.571377726	1.000080778
27	$\frac{1}{27}$	1.571217751	1.571335424	1.000074892
28	$\frac{1}{28}$	1.571188179	1.571297578	1.000069628
29	$\frac{1}{29}$	1.571161614	1.571263582	1.000064899
30	$\frac{1}{30}$	1.571137662	1.571232932	1.000060637
31	$\frac{1}{31}$	1.57111599	1.571205202	1.000056782
32	$\frac{1}{32}$	1.571096319	1.571180032	1.000053283

Table 2

In this table, I have: first column: m ; second column: $k = 1/m$; third column: $\left(\frac{\pi+\sqrt{\pi}}{2}\right)\left[I_0\left(\frac{1}{2m}\right)\right]^2 + \pi\left[I_1\left(\frac{1}{2m}\right)\right]^2 + \frac{7\sqrt{\pi}}{8m}I_0\left(\frac{1}{2m}\right)I_1\left(\frac{1}{2m}\right) - \left(\frac{2\pi(1/m)^2+7\sqrt{\pi}(1/m)^2+16\sqrt{\pi}}{32}\right)$; fourth column: $K\left(\frac{1}{m}\right)$; fifth column:

m	$k = 1/m$	$\left(\frac{\pi+\sqrt{\pi}}{2}\right)\left[I_0\left(\frac{1}{2m}\right)\right]^2 + \pi\left[I_1\left(\frac{1}{2m}\right)\right]^2 + \frac{7\sqrt{\pi}}{8m}I_0\left(\frac{1}{2m}\right)I_1\left(\frac{1}{2m}\right) - \left(\frac{2\pi(1/m)^2+7\sqrt{\pi}(1/m)^2+16\sqrt{\pi}}{32}\right)$	$K\left(\frac{1}{m}\right)$	$\frac{\left(\frac{\pi+\sqrt{\pi}}{2}\right)\left[I_0\left(\frac{1}{2m}\right)\right]^2 + \pi\left[I_1\left(\frac{1}{2m}\right)\right]^2 + \frac{7\sqrt{\pi}}{8m}I_0\left(\frac{1}{2m}\right)I_1\left(\frac{1}{2m}\right) - \left(\frac{2\pi(1/m)^2+7\sqrt{\pi}(1/m)^2+16\sqrt{\pi}}{32}\right)}{K\left(\frac{1}{m}\right)}$
2	$\frac{1}{2}$	1.651546952	1.685750354	1.020709918
3	$\frac{1}{3}$	1.605702276	1.617386735	1.007276852
4	$\frac{1}{4}$	1.590238451	1.596242222	1.00377539
5	$\frac{1}{5}$	1.583182393	1.586867847	1.002327876
6	$\frac{1}{6}$	1.579376323	1.581878436	1.001584241
7	$\frac{1}{7}$	1.577090506	1.578903918	1.001149847
8	$\frac{1}{8}$	1.575610592	1.576986771	1.000873425
9	$\frac{1}{9}$	1.574597637	1.575678422	1.000686387
10	$\frac{1}{10}$	1.573873909	1.574745561	1.000553825
11	$\frac{1}{11}$	1.573338879	1.574056948	1.000456398
12	$\frac{1}{12}$	1.572932199	1.573534108	1.000382666
13	$\frac{1}{13}$	1.572615858	1.573127756	1.000325507
14	$\frac{1}{14}$	1.572364946	1.572805664	1.000280289
15	$\frac{1}{15}$	1.572162584	1.572546032	1.000243898
16	$\frac{1}{16}$	1.571997006	1.572333687	1.000214173
17	$\frac{1}{17}$	1.571859807	1.572157798	1.000189578
18	$\frac{1}{18}$	1.571744852	1.572010469	1.000168995
19	$\frac{1}{19}$	1.57164758	1.571885834	1.000151595
20	$\frac{1}{20}$	1.57156454	1.571779457	1.000136753
21	$\frac{1}{21}$	1.571493086	1.571687938	1.000123992
22	$\frac{1}{22}$	1.571431157	1.571608632	1.000112938
23	$\frac{1}{23}$	1.571377134	1.571539459	1.000103301
24	$\frac{1}{24}$	1.571329725	1.571478762	1.000094848
25	$\frac{1}{25}$	1.571287892	1.571425211	1.000087392
26	$\frac{1}{26}$	1.571250796	1.571377726	1.000080783
27	$\frac{1}{27}$	1.571217745	1.571335424	1.000074896
28	$\frac{1}{28}$	1.571188174	1.571297578	1.000069631
29	$\frac{1}{29}$	1.571161609	1.571263582	1.000064902
30	$\frac{1}{30}$	1.571137657	1.571232932	1.00006064
31	$\frac{1}{31}$	1.571115987	1.571205202	1.000056784
32	$\frac{1}{32}$	1.571096316	1.571180032	1.000053285