

Time Runs Backwards Inside Black Holes?

Author: George Rajna

Abstract	1
Considering the chess game as a model of physics	1
Using this model in physics	4
Quantum Information Science	4
Quantum Computing Research	4
The General Weak Interaction	5
Black Holes revisited	6
Conclusions	6
References	7

Abstract

A new research paper published in Physical Review Letters has brought forward a significant new understanding of general relativity laws, and has found some strange physics taking place inside black holes. Specifically, that the direction of time could be reversed within them. [9]

Considering the positive logarithmic values as the measure of entropy and the negative logarithmic values as the measure of information we get the Information – Entropy Theory of Physics, used first as the model of the computer chess program built in the Hungarian Academy of Sciences.

Applying this model to physics we have an understanding of the perturbation theory of the QED and QCD as the Information measure of Physics. We have an insight to the current research of Quantum Information Science. The generalization of the Weak Interaction shows the arrow of time in the associate research fields of the biophysics and others. We discuss also the event horizon of the Black Holes, closing the information inside.

Considering the chess game as a model of physics

In the chess game there is also the same question, if the information or the material is more important factor of the game? There is also the time factor acting as the Second Law of Thermodynamics, and the arrow of time gives a growing disorder from the starting position.

When I was student of physics at the Lorand Eotvos University of Sciences, I succeeded to earn the master degree in chess, before the master degree in physics. I used my physics knowledge to see the chess game on the basis of Information – Entropy Theory and giving a presentation in the Hungarian Academy of Sciences, proposed a research of chess programming. Accepting my idea there has built the first Hungarian Chess Program "PAPA" which is participated on the 1st World Computer Chess Championship in Stockholm 1974. [1]

The basic theory on which one chess program can be constructed is that there exists a general characteristic of the game of chess, namely the concept of entropy.

This concept has been employed in physics for a long time. In the case of a gas, it is the logarithm of the number of those microscopic states compatible with the macroscopic parameters of the gas.


What does this mean in terms of chess? A common characteristic of every piece is that it could move to certain squares, including by capture. In any given position, therefore, the pieces by the rules of the game possess certain states, only one of which will be realized on the next move. The difference of the logarithm of the numbers of such states for Black and White respectively is the "entropy of the position". The task of the computer is then to increase this value for its own benefit.

Every chess player knows that the more mobility his pieces have and the more constrained are his opponent's, the better his position. For example, checkmate is the best possible state for the attacker, and the chess program playing according to the above principle without the prior notion of checkmate will automatically attempt it if possible.

Entropy is a principle of statistical physics and therefore is only applicable in statistical contexts. The number of microstates of a confined gas is very large and therefore the statistical approach is valid. In chess, however, the number of pieces, a macroscopic parameter, is very small and therefore in this context the "value" of a position cannot be an exact function of entropy. For example, it is possible to checkmate with a total force of a single pawn despite the fact that the opponent has many pieces and various positions available.

Examples of sacrificial combinations further demonstrate this consideration. Therefore we also need specific information about any given position. For example, entropy could be maximized by White giving check, but if the checking piece is then taken, the move was a bad one. The logarithm of the number of variations which have been examined in this way gives the amount of information. In the endgame it is rather inaccurate. Because of the small number of pieces the above noted inadequacy of the statistical principle becomes evident and we need to compute much more information to fill the gap.

We can think about the positive logarithmic values as the measure of entropy and the negative logarithmic values as the measure of information.



Shortly speaking:

- The evaluation of any position is based on the entropy + information.
- The entropy is the logarithm of the possible legal moves of the position.
- The information is simply the depth of the search, since it is the logarithm of the exponential growing number of possible positions, $\log e^x = x$.

E = entropy

I = information

D = depth of search

M = legal moves in any position, M_w for white moves and M_b for black moves

$$E = \log M_w - \log M_b = \log M$$

And since $\log e^x = x$, $I = D$

We get information + entropy, the value V of any position in the search tree of the current chess position:

$$V(D, M) = I + E = D + \log M$$

This naturally gives better values for a deeper search with greater mobility. [2]

Using this model in physics

Viewing the confined gas where the statistical entropy not needs the information addition is not the only physical system. There are for example quantum mechanical systems where the information is a very important qualification. The perturbation theory needs higher order calculations in QED or QCD giving more information on the system as in the chess games happens, where the entropy is not enough to describe the state of the matter. The variation calculation of chess is the same as the perturbation calculation of physics to gain information, where the numbers of particles are small for statistical entropy to describe the system. The role of the Feynman graphs are the same as the chess variations of a given position that is the depth of the variations tree, the Information is the same as the order of the Feynman graphs giving the Information of the micro system.

Quantum Information Science

Quantum information science is an area of study based on the idea that information science depends on quantum effects in physics. It includes theoretical issues in computational models as well as more experimental topics in quantum physics including what can and cannot be done with quantum information.

Quantum Computing Research

Quantum computing has been an intense research field since Richard Feynman in 1981 challenged the scientific community to build computers based on quantum mechanics. For decades, the pursuit remained firmly in the theoretical realm.

To understand the quantum world, researchers have developed lab-scale tools to manipulate microscopic objects without disturbing them. The 2012 Nobel Prize in Physics recognizes two of these quantum researchers: David Wineland, of the National Institute of Standards and Technology and the University of Colorado in Boulder, and Serge Haroche, of the Collège de France and the Ecole Normale Supérieure in Paris. Two of their papers, published in 1995 and '96 in Physical Review Letters, exemplify their contributions. The one by Wineland and collaborators showed how to use atomic states to make a quantum logic gate, the first step toward a superfast quantum computer. The other, by Haroche and his colleagues, demonstrated one of the strange predictions of quantum mechanics—that measuring a quantum system can pull the measuring device into a weird quantum state which then dissipates over time.

IBM scientists believe they're on the cusp of building systems that will take computing to a whole new level. On Feb 28, 2012 the IBM team presented major advances in quantum computing device performance at the annual American Physical Society meeting. Using a variety of techniques in the

IBM laboratories, scientists have established three new records for retaining the integrity of quantum mechanical properties in quantum bits, or qubits, and reducing errors in elementary computations. These breakthrough results are very close to the minimum requirements for a full-scale quantum computing system as determined by the world-wide research community. [3]

Quantum computing in neural networks is one of the most interesting research fields today. [4] The biological constructions of the brain are capable to memorize, associate and logically thinking by changing their quantum states. The machine learning of Artificial Intelligence will be one of the mainstreams of the Quantum Computing, when it will be available. Probably the main challenge will be to simulate the brain biologic capability to create new quantum states for logical reasoning, since we don't know nowadays how it is work exactly in the brain. [8]

The General Weak Interaction

The Weak Interactions T-asymmetry is in conjunction with the T-asymmetry of the Second Law of Thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes for example the Hydrogen fusion. The arrow of time by the Second Law of Thermodynamics shows the increasing entropy and decreasing information by the Weak Interaction, changing the temperature dependent diffraction patterns. A good example of this is the neutron decay, creating more particles with less known information about them. [5]

The neutrino oscillation of the Weak Interaction shows that it is a general electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. There should be a new research space of the Quantum Information Science the 'general neutrino oscillation' for the greater than subatomic matter structures as an electric dipole change.

There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also. [6]

The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction.

Black Holes revisited

The Black Holes are the counter example, where the matter is so highly concentrated that the entropy is very low and the information is high but closed inside the event horizon.

The problem is with the Black hole that it is not a logical physical state of the matter by the diffraction theory, because we cannot find a temperature where this kind of diffraction patterns could exist. [5]

Also the accelerating charges of the electric current say that the charge distribution maintains the accelerating force and this viewpoint of the relativity does not make possible an acceleration that can cause a Black Hole. The ever growing acceleration simply resolved in the spin. [7]

The spin is one of the most generic properties of the Universe, not only the elementary particles are spinning, but also the Sun, Earth, etc. We can say that the spin is the resolution of the constantly accelerating matter solving the problem of the relativity and the accelerating Universe. The gravity is the magnetic effect of the accelerating matter, the attracting force between the same charges; working by the electromagnetic oscillations, because of this is their universal force. Since this effect is relatively weak, there is no way for the gravitation force to compress the matter to a Black Hole.

New Research Suggests That Time Runs Backwards Inside Black Holes

A new research paper published in Physical Review Letters has brought forward a significant new understanding of general relativity laws, and has found some strange physics taking place inside black holes. Specifically, that the direction of time could be reversed within them. Several physical procedures are perfectly symmetric in time. Take a pendulum for instance. If someone shows you a video of a pendulum swinging, you cannot differentiate if the video is actually moving forward or backward. But some processes are not symmetric at all. We can tell that a pendulum will ultimately slow because of friction and we know that it was triggered at some point, so we can give a temporal direction to physics. The directionality of time and our view of it was called the "Arrow of Time" by British astronomer Arthur Eddington, and it has been connected to the entropy of the cosmos.

Event horizons ascend from the mathematics of general relativity, but the simplest mathematical explanation of an event horizon leads to a paradox. Event horizons must contain all the history of the cosmos, from the Big Bang to its death. This generates numerous complications as it assumes that the universe must be deterministic and that past and future are inscribed on the "skin" of black holes. While this paradox doesn't affect explanations and forecasts of the physics of black holes, it is evidently a limiting factor in correctly understanding how black holes work.

The new research tries to fill this important gap in black hole physics. The elementary notion of this paper is grounded on the holographic principle. According to the author of the paper the event horizon is essentially a holographic screen, a hypersurface with a precise entropy. You can have two

kinds of holographic screens, past holographic screens and future holographic screens, liable on whether the entropy inside the surface is increasing or decreasing.

Study author Netta Engelhardt told Phys.org "Holographic screens are in a sense a local boundary to regions of strong gravitational fields. Future holographic screens correspond to gravitational fields which pull matter together... whereas past holographic screens correspond to regions which spread matter out..."

If this principle is applied to the cosmos as a whole, the entropy arrow is steady with the second law of thermodynamics. Entropy increases, time moves forward. But the use of the law onto black holes yields an interesting outcome. Within a black hole, entropy decreases (things become more structured) and thus thermodynamically time runs backward.

Although this is an exciting consequence, the paper is important because it verifies the first area law in general relativity and it might have solved a long-standing problem in black holes physics. [9]

Conclusions

My opinion is that information and matter are two sides of the same thing in physics, because the matter is the diffraction pattern of the electromagnetic waves, giving the temperature dependent different structures of the matter, the information about them arrives by the electromagnetic waves and also the entropy or uncertainty as the measure of disorder. [7]

The Fluctuation Theory gives a probability for Information grow and Entropy decrease seemingly proportionally with the gravitational effect of the accelerating Universe, against the arrow of time by the Second Law of Thermodynamics. The information and entropy are the negative and positive sides of the logarithmic curve, describing together the state of the matter.

References

[1] The world computer chess championship, Stockholm 1974

Author: Jean E. Hayes, David Levy

Publisher: University Press (Edinburgh) ISBN 0852242859

<http://chessprogramming.wikispaces.com/Papa>

<http://chessprogramming.wikispaces.com/George+Rajna>

[2] Information – Entropy Theory of Artificial Intelligence

Author: George Rajna

Publisher: Vixra.org <http://vixra.org/abs/1201.0063>

[3] IBM Quantum Computing Press Release - Tues, Feb 28, 2012

Author: IBM Research Team

Publisher: IBM <http://ibmquantumcomputing.tumblr.com/>

[4] Quantum computing in neural networks

Author: P. GRALEWICZ

Publisher: arXiv.org <http://arxiv.org/abs/quant-ph/0401127>

[5] 3 Dimensional String Theory

Author: George Rajna

Publisher: Vixra.org <http://vixra.org/abs/1112.0064>

[6] The application of statistical physics to evolutionary biology

Author: Guy Sella and Aaron E. Hirsh

Publisher: Hebrew University of Jerusalem Department of Biological Sciences, Stanford University, Stanford, CA 94305 <http://www.pnas.org/content/102/27/9541.full.pdf+html>

[7] The Magnetic Field of the Electric Current and the Magnetic Induction

Author: George Rajna

Publisher: Vixra.org <http://vixra.org/abs/1112.0058>

[8] Automated Quantum Reasoning: Non-Logic _Semi-Logic _Hyper-Logic

Author: Bob Coecke

Publisher: Oxford University Computing Laboratory <http://www.researchgate.net>

[9] New Research Suggests That Time Runs Backwards Inside Black Holes

<http://www.physics-astronomy.com/2015/11/new-research-suggests-that-time-runs.html>