

A NEW DARK MATTER DENSITY PROFILE AS POWER OF GRAVITATIONAL FIELD FOR COMA CLUSTER

Author: Manuel Abarca Hernández mabarcaher1@gmail.com

INDEX

1. ABSTRACT..... 2

2. INTRODUCTION..... 2

3. COMA CLUSTER DATA..... 5

4. FITTING PARAMETERS OF NFW PROFILE 6

5. FITING PARAMETER OF DARK MATTER DENSITY AS POWER OF GRAVITATIONAL FIELD 8

 5.1 DARK MATTER DENSITY THROUGH DM DENSITY AS POWER OF E FOR GALAXIES..... 9

6. BERNOULLI DIFFERENTIAL EQUATION FOR GRAVITATIONAL FIELD IN COMA CLUSTER HALO 9

7. BERNOULLI PROFILE OF DARK MATTER DENSITY FOR COMA CLUSTER HALO..... 11

8. CALCULUS OF DARK MATTER IN COMA CLUSTER HALO UP TO 14 MPC WITH NFW PROFILE 15

9. CALCULUS OF DARK MATTER IN COMA CLUSTER HALO UP TO 14 MPC WITH BERNOULLI PROFILE 15

10. GRAVITATIONAL MASS IN COMA CLUSTER UP TO 14 MPC 16

11. CONCLUSION 16

12. BIBLIOGRAPHIC REFERENCES 17

1. ABSTRACT

For last two years I have published eight papers about DM on galaxies. Data have been taken from recent papers written by prestigious researches. The main target of papers have been to calculate DM density through an original profile called Bernoulli profile and check its results with other standard DM profiles such as NFW, Burkert or Pseudo-Isothermal. Results on every galaxy checked have been successful.

The importance these works is that Bernoulli profile is supported by an original hypothesis about Dark matter nature. This hypothesis stated that dark matter is generated locally by gravitational field according an unknown quantum mechanism.

Now, in this paper Dark matter density in Coma cluster will be studied with the same method.

Briefly is going to be explained the structure of this paper.

Firstly are presented data Coma Cluster, which come from recent papers published in 2016. They are two couples of radius and mass enclosed by its sphere. Radius considered are $R_{500} = 1.314$ Mpc and Virial radius = 3.1 Mpc

With only two couple of data is fitted NFW profile for Dark matter density in Coma cluster within an interval radius 1.314 Mpc to 3.1 Mpc. As DM through NFW profile gives an excessive amount of DM within halo up to 14 Mpc it has been introduced Bernoulli profile which is an original DM density profile developed by author, used successfully in several galaxies to study DM.

According Chernin, A. 2013. Firstly are introduced concepts of gravitating mass, dark energy or anti gravitating mass and matter mass apart from its formula $M_G(R) = M_M(R) + M_{DE}(R)$. Because at cluster halo scale dark energy are not negligible, so it has been considered Dark energy effect to calculate gravitating mass within Coma cluster halo up to 14 Mpc.

Result got of gravitating mass within cluster halo up to 14 Mpc through DM calculated with Bernoulli profile differ only 8% regarding gravitating mass got by Diferio A. & Geller M.J. 1997 whereas gravitating mass got with NFW differ more than 600%.

2. INTRODUCTION

For last two years I have published eight papers about DM on galaxies. Data have been taken from recent papers written by prestigious researches. The main target of papers have been to calculate DM density through an original profile called Bernoulli profile and check its results with other standard DM profiles such as NFW, Burkert or Pseudo-Isothermal. Results on every galaxy checked have been successful.

The importance these works is that Bernoulli profile is supported by an original hypothesis about Dark matter nature.

In my first paper [1] Abarca, M. 2014. called *Dark matter model by quantum vacuum*, were introduced the main hypothesis the model which is that DM is generated locally by gravitational field according an unknown quantum mechanism.

Now, in this paper Dark matter density in Coma cluster will be studied with the same method.

Briefly is going to be explained the structure of this paper.

Firstly are presented data Coma Cluster, which come from recent paper published in 2016. They are two couples of radius and mass enclosed by its sphere. $R_1 = R_{500}$ and $R_2 =$ Virial radius.

1.314 Mpc = R1	$6 \cdot 10^{14} M_{\odot} = M1$
3.1 ± 0.7 Mpc = R2	$2.9 \cdot 10^{15} M_{\odot} = M2$

Secondly has been fitted a NFW profile to get DM density at R1 and R2.

R1 = 1.314 Mpc	DM 1 = $2.67 \cdot 10^{-24} \text{ kg/m}^3$
R2 = 3.1 Mpc	DM 2 = $9.2 \cdot 10^{-25} \text{ kg/m}^3$

In galaxies are used dozens of data to get its NFW profile. However we have got NFW profile for Coma cluster with the minimum, two couples of data. In addition error data are important because of huge distance to Coma cluster, 95 Mpc.

Therefore it is clear that calculus have errors not very well quantified.

The following step is to fit data of NFW DM density profile as power of gravitational field, E. Particularly formula found is $\varphi_{DM}(E) = A \cdot E^B$ Where A and B are parameter to calculate with previous data.

Before calculate A& B is needed to know gravitational field at R1 & R2.

Data of Dark matter density and gravitational field at R1 and R2 radius.		
R1=1.314 Mpc	$E1 = 4.83 \cdot 10^{-11} \text{ m/s}^2$	$D1 = 2.67 \cdot 10^{-24} \text{ kg/m}^3$
R2= 3.1 Mpc	$E2 = 4.19 \cdot 10^{-11} \text{ m/s}^2$	$D2 = 9.2 \cdot 10^{-25} \text{ kg/m}^3$

Taking second and third column it is calculated easily parameters A& B of $\varphi_{DM}(E) = A \cdot E^B$

DM Density as power of gravitational field $D = AE^B$	
A International System	$5.563 \cdot 10^{53}$
B A dimensional	7.495

In chapter 6 it is got a Bernoulli differential equation for E depending on radius within halo dominion from 3.1 Mpc to 14 Mpc and thanks its solution it is possible to calculate Bernoulli profile which is

DM Density Bernoulli profile for Coma Cluster halo $3.1 < \text{radius} < 14 \text{ Mpc}$	
$E_{BER}(r) = (Cr^{\alpha} + Dr)^{\beta}$ $C = 3.5155 \cdot 10^{-233}$ $D = 2,52698 \cdot 10^{44}$ $\alpha = 12.99$ $\beta = -0,15396459$	
Density D.M. BERNOULLI (r) = $D_{DMB}(r) = A \cdot E^B$ Where $A = 5.563 \cdot 10^{53}$ and $B = 7.495$ unit kg/m^3	

In chapter seven is tabulated Bernoulli and NFW profiles at 0,1 Mpc step. It is clearly showed that NFW is a great deal bigger density that Bernoulli profile. In chapter eight and nine are calculated total mass enclosed within halo cluster up to 14 Mpc.

Quoting to [12] Chernin, A.D. 2013. In chapter ten are introduced concepts of Gravitating mass, Anti gravitating mass or Dark energy and Matter mass. According this author at scale halo cluster it is needed to consider dark energy to calculate gravitating mass, which is kind of matter measured by astrophysics.

Therefore $M_G(R) = M_M(R) + M_{DE}(R)$ where $M_{DE}(R)$ is a negative quantity.

According [12] Chernin, A.D. 2013 $M_{DE}(14 \text{ Mpc}) = -2.4 \cdot 10^{15} M_\odot$.

According numerical calculus of total dark matter through Bernoulli profile $M_M(14 \text{ Mpc}) = 4.6 \cdot 10^{15} M_\odot$ and therefore $M_G = 2.2 \cdot 10^{15} M_\odot$ which match perfectly with gravitating mass found by [13] Diaferio & Geller 1997 with a value $M_G = 2.4 \cdot 10^{15} M_\odot$. However with NFW profile $M_M(14 \text{ Mpc}) = 3.05 \cdot 10^{16} M_\odot$ which is a quantity 6 times bigger than $4.6 \cdot 10^{15} M_\odot$ therefore it is clear that NFW profile does not work at halo cluster scale.

The agreement between $M_G = 2.4 \cdot 10^{15} M_\odot$ got by Diaferio & Geller, 1997 and $M_G = 2.2 \cdot 10^{15} M_\odot$ got through Bernoulli profile is the main result this paper.

3. COMA CLUSTER DATA

According [10] De Martino, Plank Collaboration project [12] determined that the angular size subtended by R_{500} is $\theta_{500} = 48 \pm 1 \text{ arcmin}$ corresponding to $R_{500} = 1.314 \text{ Mpc}$ and the mass associated is $M_{500} \approx 6 \cdot 10^{14} M_{\odot}$.

As it is known R_{500} is the radius a sphere where the average density is 500 times bigger than critical density of

Universe which is $\rho_{CRITICAL} = \frac{3 \cdot H_0^2}{8\pi G}$ where H_0 is Hubble constant. Currently $H_0 = 67.8 \text{ Km/s/Mpc}$. so

$$\rho_{CRITICAL} = \frac{3 \cdot H_0^2}{8\pi G} = 8.64 \cdot 10^{-27} \text{ kg/m}^3 \text{ and } M_{500} \approx 6 \cdot 10^{14} M_{\odot}$$

According [9] Fuller, C.2016. Virial Radius has been estimated as the mean Virial radius published by four authors Kubo et al. 2007. Geller et al. 1999. Hughes, 1989. The & White 1986. $R_{VIRIAL} = 3.1 \pm 0.7 \text{ Mpc}$

[9] Fuller, C.2016 has published this graphs and data which allows to calculate velocity dispersion $\sigma = 905 \text{ Km/s}$




Figure 1. The velocity distribution of the CCC galaxies, showing a clearly relaxed system. The vertical dashed lines represent the 3σ velocity dispersion of the fitted Gaussian function (solid line).

Table 1. Derived parameters of the Coma cluster.

Virial Radius	3.1 Mpc
Velocity Dispersion, σ	905 km s ⁻¹
Mean Cluster velocity, μ	6984 km s ⁻¹
Minimum velocity limit, $\mu - 3\sigma$	4268 km s ⁻¹
Maximum velocity limit, $\mu + 3\sigma$	9700 km s ⁻¹

[9] Fuller, C.2016 has published table below where is shown baryonic mass and Virial Mass or dynamical mass at 3.1 Mpc radius.

It is remarkable the quotient $M_{\text{VIRIAL}} / M_{\text{BARYONIC}} = 104$. As reader knows the same proportion of masses, in average, 10 times lower in galaxies.

Table 11. Mass parameters for the Coma Cluster derived from optically selected galaxies

Component	Mass (M_{\odot})
Total Stellar Mass	2.8×10^{13}
Total Gas Mass	1.6×10^{11}
Total Dust Mass	2.5×10^9
Total Baryonic (Excluding X-ray gas)	2.8×10^{13}
Virial Mass	2.9×10^{15}

the cluster, we can measure the Virial mass of the cluster using:

$$M_{\text{Virial}} = \frac{5R_{\text{virial}}\sigma^2}{G} \quad (6)$$

Below are summarised Coma Cluster data needed in this paper.

DISTANCES VS MASSES IN	COMA CLUSTER
1.314 Mpc	$6 \cdot 10^{14} M_{\odot}$
3.1 ± 0.7 Mpc	$2.9 \cdot 10^{15} M_{\odot}$

Another important data used to contrast calculus got in this paper are the following one: According [12] Chernin, A.D. 2013 and [13] Diaferio, A., Geller M.J., 1997. Gravitating mass of Coma cluster halo up to 14 Mpc is $2.4 \cdot 10^{15} M_{\odot}$. In chapter 10 will be explained gravitating mass concept.

4. FITTING PARAMETERS OF NFW PROFILE

As Baryonic matter is under 1% of total mass. This kind of matter is neglected, so it can be considered previous data as Dark matter entirely.

Through previous data it is possible to fit a NFW profile for DM density.

As it is known $D_{\text{NFW}}(R) = \frac{4D_s}{x \cdot (1+x)^2}$ where $x = R/R_s$. R_s is called radius scale and D_s is density at $x = 1$ or $R = R_s$

As reader knows, Total DM enclosed by a sphere of radius R is $M(< R) = 16\pi R^3 \cdot D_s \cdot \left[\ln(1+x) - \frac{x}{1+x} \right]$

Calling $f(x) = \left[\ln(1+x) - \frac{x}{1+x} \right] = \ln(1+r/R_s) - \frac{r}{r+R_s}$ and $K = 16\pi R_s^3 \cdot D_s$ then $M(R) = K \cdot f(x)$

Having NFW profile two parameters R_s and D_s , it is needed a couple of data to calculate these parameters.

Below is explained method followed.

Naming these data, it is right to explain better calculus

DISTANCES VS MASSES IN	COMA CLUSTER
1.314 Mpc = R_1	$6 \cdot 10^{14} M_\odot = M_1$
3.1 Mpc = R_2	$2.9 \cdot 10^{15} M_\odot = M_2$

Using total mass formula at two radius we have.

$M(R_2) = K \cdot f(X_2)$ where $X_2 = R_2/R_s$ and $M(R_1) = K \cdot f(X_1)$ where $X_1 = R_1/R_s$

$M(R_2)/M(R_1) = f(X_2) / f(X_1) = 4.83333$

Are defined these auxiliary functions $g(r) = \left[\ln(1+3.1/r) - \frac{3.1}{3.1+r} \right]$ and $j(r) = \left[\ln(1+1.314/r) - \frac{1.314}{1.314+r} \right]$

Tabulating both functions by a computer it is right to get that for $r = 14.9$ quotient $g(14.9)/j(14.9) = 4.834$

Therefore parameter radius scale $R_s = 14.9$ Mpc.

Knowing $R_s = 14.9$ Mpc it is right to calculate $f(X_2) = 0.016788322$ y $f(X_1) = 0.00347278$

From formula $M(R_2) = K \cdot f(X_2)$ or formula $M(R_1) = K \cdot f(X_1)$ it is got $D_s = 6.98 \cdot 10^{-26} \text{ kg/m}^3$ thanks data masses.

PARAMETER NFW PROFILE FOR	COMA CLUSTER
R_s	14.9 Mpc
D_s	$6.98 \cdot 10^{-26} \text{ kg/m}^3$

Finally it is possible to calculate DM density at radius R_1 and R_2

Dark matter density for NFW profile at R_1 and R_2 radius.		
$R_1 = 1.314$ Mpc	$X_1 = 0.0882$	$D_1 = 2.67 \cdot 10^{-24} \text{ kg/m}^3$
$R_2 = 3.1$ Mpc	$X_2 = 0.208$	$D_2 = 9.2 \cdot 10^{-25} \text{ kg/m}^3$

5. FITING PARAMETER OF DARK MATTER DENSITY AS POWER OF GRAVITATIONAL FIELD

Density $DM = A \cdot E^B$. Having this function two parameters, are needed two couple of data to calculate it.

Dark matter density for NFW profile at R1 and R2 radius and total mass enclosed by its spheres		
R1=1.314 Mpc	M1= $6 \cdot 10^{14} M_{\odot}$	D1 = $2.67 \cdot 10^{-24} \text{ kg/m}^3$
R2= 3.1 Mpc	M2= $2.9 \cdot 10^{15} M_{\odot}$	D2 = $9.2 \cdot 10^{-25} \text{ kg/m}^3$

Through these data it is possible to calculate firstly gravitational field at radius R1 and radius R2.

$$E = \frac{GM}{R^2}$$

Data of Dark matter density and gravitational field at R1 and R2 radius.		
R1=1.314 Mpc	E1= $4.83 \cdot 10^{-11} \text{ m/s}^2$	D1 = $2.67 \cdot 10^{-24} \text{ kg/m}^3$
R2= 3.1 Mpc	E2= $4.19 \cdot 10^{-11} \text{ m/s}^2$	D2 = $9.2 \cdot 10^{-25} \text{ kg/m}^3$

Summarising data

E1= $4.83 \cdot 10^{-11} \text{ m/s}^2$	D1 = $2.67 \cdot 10^{-24} \text{ kg/m}^3$
E2= $4.19 \cdot 10^{-11} \text{ m/s}^2$	D2 = $9.2 \cdot 10^{-25} \text{ kg/m}^3$

Writing symbolically previous data $D1 = A \cdot E1^B$ and $D2 = A \cdot E2^B$ it is right to get that $B = \frac{\ln 2.902}{\ln 1.151} = 7.495$

because $D1/D2 = 2.902$ and $E1/E2 = 1.153$

Calculus of A may be made using (E1,D1) and (E2,D2) data. Both of them led to $A = 5.563 \cdot 10^{53}$, probably the bigger parameter never got ;

Calculus of A and B are a great deal easier than calculus made for NFW parameters.

DM Density as power of gravitational field $D = AE^B$	
A International System	$5.563 \cdot 10^{53}$
B Adimensional	7.495

In the following chapter it will be show that both DM profiles (NFW and power of E) give essentially the same results when radius is inside the interval 1.314Mpc – 3.1 Mpc. However DM as power of E is a great deal lower than NFW profile when radius is $R > 3.1 \text{ Mpc}$.

5.1 DARK MATTER DENSITY THROUGH DM DENSITY AS POWER OF E FOR GALAXIES

According [2] Abarca, M.2015 and [8] Abarca,M.2016. Formula below is a good approximation for dark matter density whatever giant galaxy.

<i>Dark matter density function as Universal law for big galaxies</i>	
Density _{DM} = 2,526 · 10 ⁻⁵ · E ^B where B= 1,74	
Where Unit for Density _{D.M.} is Kg / m ³ and Unit for E is m/s ²	

Now it will be checked with data we have for Coma cluster.

Data within Coma cluster for gravit. Field and DM density	
E1= 4.83·10 ⁻¹¹ m/s ²	D1 = 2.67·10 ⁻²⁴ kg/m ³
E2= 4.19·10 ⁻¹¹ m/s ²	D2 = 9.2·10 ⁻²⁵ kg/m ³

DM density through DM density as power of E for giant galaxies	
E1= 4.83·10 ⁻¹¹ m/s ²	D1 _{GALAXIES} = 2.8·10 ⁻²³ kg/m ³
E2= 4.19·10 ⁻¹¹ m/s ²	D2 _{GALAXIES} = 2.2 ·10 ⁻²³ kg/m ³

Reader can check that DM density got with formula for galaxies is more than ten times bigger than DM density formula fitted to Coma cluster data.

6. BERNOULLI DIFFERENTIAL EQUATION FOR GRAVITATIONAL FIELD IN COMA CLUSTER HALO

It will be considered the region 3.1 < Radius < 14 Mpc where density of baryonic matter is negligible versus baryonic density. So for radius bigger than 3.1 Mpc it will be considered that derivative of M(r) depend on dark matter density only.

As it is known in this formula $E = G \frac{M(r)}{r^2}$, M(r) represents mass enclosed by a sphere with radius r. If it is considered radius > 3.1 Mpc then the derivative of M(r) depend on dark matter density essentially and therefore $M'(r) = 4\pi r^2 \varphi_{DM}(r)$.

As $\varphi_{DM}(r) = A \cdot E^B(r)$ Where A= 5.563·10⁵³ and B= 7.495 then $M'(r) = 4\pi r^2 \cdot A \cdot E^B$

If $E = G \frac{M(r)}{r^2}$ is differentiated it is got $E'(r) = G \frac{M'(r) \cdot r^2 - 2rM(r)}{r^4}$

If $M'(r) = 4\pi r^2 \varphi_{DM}(r)$ is replaced above it is got $E'(r) = 4\pi G \varphi_{DM}(r) - 2G \frac{M(r)}{r^3}$ As $\varphi_{DM}(r) = A \cdot E^B(r)$ it

is right to get $E'(r) = 4\pi \cdot G \cdot A \cdot E^B(r) - 2 \frac{E(r)}{r}$ which is a Bernoulli differential equation.

$E'(r) = K \cdot E^B(r) - 2 \frac{E(r)}{r}$ being $K = 4\pi \cdot G \cdot A$ then $K = 4,66488631594 \cdot 10^{44}$ into I.S. of units.

Calling y to E , the differential equation is written this way $y' = K \cdot y^B - \frac{2 \cdot y}{r}$

Bernoulli family equations $y' = K \cdot y^B - \frac{2 \cdot y}{r}$ may be converted into a differential linear equation with this

variable change $u = y^{1-B}$.

General solution is $E(r) = \left(Cr^{2B-2} + \frac{Kr(1-B)}{3-2B} \right)^{\frac{1}{1-B}}$ with $B \neq 1$ and $B \neq 3/2$ where C is the parameter of initial condition of gravitational field at a specific radius.

Calling $\alpha = 2B - 2$ $\beta = \frac{1}{1-B}$ and $D = \left(\frac{K(1-B)}{3-2B} \right)$ formula may be written as

$E(r) = \left(Cr^\alpha + Dr \right)^\beta$ Where specifically values for these parameters are the following ones:

$$\alpha = 2B - 2 = 12,99$$

$$\beta = \frac{1}{1-B} = -0,15396459$$

$$D = \left(\frac{K(1-B)}{3-2B} \right) = 2,52698 \cdot 10^{44}$$

Initial condition for parameter C calculus

Suppose R_0 and E_0 are specific initial conditions for radius and gravitational field then $C = \frac{E_0^{1/\beta} - D \cdot R_0}{R_0^\alpha}$

As dominion radius begin at $R = 3.1 \text{ Mpc} = 9.57 \cdot 10^{22} \text{ m}$. it is right to consider as initial condition $R_0 = 3.1 \text{ Mpc} = 9.57 \cdot 10^{22} \text{ m}$. and $E_0 = 4.19 \cdot 10^{-11} \text{ m/s}^2$ being $C = 3.5155 \cdot 10^{-233}$

Probably the lower parameter never calculated ;

Initial condition values R_0 & E_0	
$R_0 =$	3.1 Mpc
$E_0 =$	$4.19 \cdot 10^{-11} \text{ m/s}^2$
$C =$	$3.5155 \cdot 10^{-233}$ units I.S.

Finally it is possible to write formula for DM density profile got through Bernoulli method.

Bernoulli Solution for Gravitational field inside halo $3.1 \text{ Mpc} < \text{Radius} < 14 \text{ Mpc}$
$E_{BER}(r) = (Cr^\alpha + Dr)^\beta$ $C = 3.5155 \cdot 10^{-233}$ $D = 2,52698 \cdot 10^{44}$ $\alpha = 12.99$ $\beta = -0,15396459$

7. BERNOULLI PROFILE OF DARK MATTER DENSITY FOR COMA CLUSTER HALO

Thanks Bernoulli solution for gravitational field is right to get DM density through power of E formula.

DM Density Bernoulli profile for Coma Cluster halo $3.1 < \text{radius} < 14 \text{ Mpc}$
$E_{BER}(r) = (Cr^\alpha + Dr)^\beta$ $C = 3.5155 \cdot 10^{-233}$ $D = 2,52698 \cdot 10^{44}$ $\alpha = 12.99$ $\beta = -0,15396459$
Density D.M. BERNOULLI (r) = $D_{DMB}(r) = A \cdot E^B$ Where $A = 5.563 \cdot 10^{53}$ and $B = 7.495$ unit kg/m^3

Thanks these formulas it is right to tabulate gravitational field and DM as power of E. The last column shows DM density of NFW profile. Both density are essentially the same at 3.1 Mpc. However DM density Bernoulli decrease more quickly than density NFW. Therefore total mass of DM in halo will be different for both densities profiles.

In next chapter will be calculated total mass, and surprisingly mass provided through Bernoulli density matches with DM calculated by observational methods whereas total mass through NFW profiles is far from it.

Radius	Gravt. field	DM Bernoulli	DM NFW profile
Mpc	m/s ²	Kg/m ³	kg/m ³
3,1	4,19E-11	9,20E-25	9,19E-25
3,2	4,15608E-11	8,66E-25	8,81E-25
3,3	4,11806E-11	8,08E-25	8,45E-25
3,4	4,07464E-11	7,46E-25	8,11E-25
3,5	4,02448E-11	6,80E-25	7,79E-25
3,6	3,9663E-11	6,10E-25	7,50E-25
3,7	3,8991E-11	5,37E-25	7,21E-25
3,8	3,82234E-11	4,62E-25	6,95E-25
3,9	3,73609E-11	3,90E-25	6,70E-25
4	3,64111E-11	3,21E-25	6,46E-25
4,1	3,53873E-11	2,59E-25	6,24E-25
4,2	3,43073E-11	2,06E-25	6,03E-25
4,3	3,31905E-11	1,60E-25	5,83E-25
4,4	3,20562E-11	1,24E-25	5,63E-25
4,5	3,09214E-11	9,44E-26	5,45E-25
4,6	2,98001E-11	7,15E-26	5,28E-25
4,7	2,87034E-11	5,40E-26	5,11E-25
4,8	2,7639E-11	4,07E-26	4,96E-25
4,9	2,66121E-11	3,06E-26	4,81E-25
5	2,56259E-11	2,31E-26	4,66E-25
5,1	2,46818E-11	1,74E-26	4,53E-25
5,2	2,37801E-11	1,32E-26	4,40E-25
5,3	2,29205E-11	1,00E-26	4,27E-25
5,4	2,21016E-11	7,62E-27	4,15E-25
5,5	2,13222E-11	5,82E-27	4,03E-25
5,6	2,05805E-11	4,46E-27	3,92E-25
5,7	1,98746E-11	3,44E-27	3,82E-25
5,8	1,92029E-11	2,66E-27	3,72E-25
5,9	1,85634E-11	2,06E-27	3,62E-25
6	1,79545E-11	1,60E-27	3,52E-25
6,1	1,73742E-11	1,25E-27	3,43E-25
6,2	1,68211E-11	9,84E-28	3,35E-25
6,3	1,62936E-11	7,75E-28	3,26E-25
6,4	1,57901E-11	6,13E-28	3,18E-25
6,5	1,53094E-11	4,86E-28	3,10E-25
6,6	1,48501E-11	3,87E-28	3,03E-25
6,7	1,4411E-11	3,09E-28	2,95E-25
6,8	1,3991E-11	2,47E-28	2,88E-25
6,9	1,3589E-11	1,99E-28	2,82E-25

7	1,32039E-11	1,60E-28	2,75E-25
7,1	1,2835E-11	1,30E-28	2,69E-25
7,2	1,24812E-11	1,05E-28	2,63E-25
7,3	1,21419E-11	8,55E-29	2,57E-25
7,4	1,18161E-11	6,97E-29	2,51E-25
7,5	1,15033E-11	5,70E-29	2,45E-25
7,6	1,12027E-11	4,68E-29	2,40E-25
7,7	1,09137E-11	3,85E-29	2,35E-25
7,8	1,06358E-11	3,17E-29	2,30E-25
7,9	1,03683E-11	2,62E-29	2,25E-25
8	1,01107E-11	2,17E-29	2,20E-25
8,1	9,86269E-12	1,80E-29	2,16E-25
8,2	9,62364E-12	1,50E-29	2,11E-25
8,3	9,39318E-12	1,25E-29	2,07E-25
8,4	9,17089E-12	1,04E-29	2,03E-25
8,5	8,9564E-12	8,74E-30	1,98E-25
8,6	8,74935E-12	7,34E-30	1,94E-25
8,7	8,54938E-12	6,17E-30	1,91E-25
8,8	8,3562E-12	5,20E-30	1,87E-25
8,9	8,16949E-12	4,39E-30	1,83E-25
9	7,98896E-12	3,71E-30	1,80E-25
9,1	7,81435E-12	3,14E-30	1,76E-25
9,2	7,64541E-12	2,67E-30	1,73E-25
9,3	7,48188E-12	2,27E-30	1,70E-25
9,4	7,32354E-12	1,93E-30	1,66E-25
9,5	7,17018E-12	1,65E-30	1,63E-25
9,6	7,02158E-12	1,41E-30	1,60E-25
9,7	6,87756E-12	1,21E-30	1,57E-25
9,8	6,73792E-12	1,04E-30	1,54E-25
9,9	6,60249E-12	8,89E-31	1,52E-25
10	6,4711E-12	7,65E-31	1,49E-25
10,1	6,3436E-12	6,59E-31	1,46E-25
10,2	6,21982E-12	5,68E-31	1,44E-25
10,3	6,09964E-12	4,91E-31	1,41E-25
10,4	5,9829E-12	4,25E-31	1,39E-25
10,5	5,86949E-12	3,68E-31	1,36E-25
10,6	5,75926E-12	3,19E-31	1,34E-25
10,7	5,65212E-12	2,77E-31	1,32E-25
10,8	5,54793E-12	2,41E-31	1,29E-25
10,9	5,4466E-12	2,10E-31	1,27E-25
11	5,34803E-12	1,83E-31	1,25E-25
11,1	5,2521E-12	1,60E-31	1,23E-25
11,2	5,15873E-12	1,40E-31	1,21E-25
11,3	5,06783E-12	1,22E-31	1,19E-25
11,4	4,97931E-12	1,07E-31	1,17E-25
11,5	4,89309E-12	9,41E-32	1,15E-25

11,6	4,80909E-12	8,27E-32	1,13E-25
11,7	4,72724E-12	7,27E-32	1,12E-25
11,8	4,64745E-12	6,40E-32	1,10E-25
11,9	4,56967E-12	5,64E-32	1,08E-25
12	4,49383E-12	4,97E-32	1,06E-25
12,1	4,41986E-12	4,39E-32	1,05E-25
12,2	4,3477E-12	3,88E-32	1,03E-25
12,3	4,27729E-12	3,43E-32	1,01E-25
12,4	4,20858E-12	3,04E-32	9,99E-26
12,5	4,14151E-12	2,70E-32	9,84E-26
12,6	4,07604E-12	2,39E-32	9,69E-26
12,7	4,0121E-12	2,13E-32	9,55E-26
12,8	3,94966E-12	1,89E-32	9,40E-26
12,9	3,88866E-12	1,68E-32	9,26E-26
13	3,82906E-12	1,50E-32	9,13E-26
13,1	3,77083E-12	1,34E-32	8,99E-26
13,2	3,71391E-12	1,19E-32	8,86E-26
13,3	3,65827E-12	1,06E-32	8,73E-26
13,4	3,60387E-12	9,51E-33	8,60E-26
13,5	3,55068E-12	8,51E-33	8,48E-26
13,6	3,49866E-12	7,62E-33	8,36E-26
13,7	3,44777E-12	6,83E-33	8,24E-26
13,8	3,39798E-12	6,12E-33	8,12E-26
13,9	3,34927E-12	5,49E-33	8,01E-26
14	3,30159E-12	4,93E-33	7,90E-26

8. CALCULUS OF DARK MATTER IN COMA CLUSTER HALO UP TO 14 MPC WITH NFW PROFILE

Total DM for NFW profiles is $M(< R) = 16\pi R^3 s \cdot D_s \left[\ln(1+x) - \frac{x}{1+x} \right]$.

In chapter four were calculated NFW parameters.

PARAMETER NFW PROFILE FOR	COMA CLUSTER
Rs	14.9 Mpc
Ds	$6.98 \cdot 10^{-26} \text{ kg/m}^3$

For $R = 14 \text{ Mpc}$ $x = 0.9396$ $M(14 \text{ Mpc}) = K \cdot f(0.9396) = 0.17805 \cdot K$ and $K = 1.71 \cdot 10^{17} M_{\odot}$

So $M(14 \text{ Mpc}) = 3.05 \cdot 10^{16} M_{\odot}$ which is more than 6 times bigger than dark matter calculated by other methods which includes observational techniques.

9. CALCULUS OF DARK MATTER IN COMA CLUSTER HALO UP TO 14 MPC WITH BERNOULLI PROFILE

Dark matter enclosed inside halo is calculated through a numerical integration using data table written in chapter six. In this table dominion radius extend from 3.1 Mpc up to 14 Mpc.

Upper bound for mass is $M_{\text{UPPER BOUND}} = 1.83 \cdot 10^{15} M_{\odot}$ and lower bound for mass is $M_{\text{LOWER BOUND}} = 1.58 \cdot 10^{15} M_{\odot}$

So is a good approximation consider its average value as total dark matter mass $M_{\text{DARK MATTER HALO}} = 1.7 \cdot 10^{15} M_{\odot}$

Adding this value to $2.9 \cdot 10^{15} M_{\odot}$, which is mass within radius 3.1 Mpc, results a total mass for Coma cluster

$$M_M(14 \text{ Mpc}) = 4.6 \cdot 10^{15} M_{\odot}$$

Author [12] Chernin, A.D. 2013, mentioned that [13] Diaferio & Geller 1997 and [14] Diaferio 1999 using the caustic technique got a mass up to 14 Mpc of $2.4 \cdot 10^{15} M_{\odot}$.

Apparently results got do not agree at all with observational results got by Diaferio. However if it is considered Dark Energy enclosed within halo, results match perfectly as it will be seen at following chapter.

10. GRAVITATIONAL MASS IN COMA CLUSTER UP TO 14 MPC

According [12] Chernin, A.D. 2013. At cluster scale is necessary consider three class of mass, because at this scale dark energy effect may be considerate as anti gravitating mass or negative mass. Therefore effective gravitating mass is calculated adding to total mass, dark energy as a negative quantity.

$$M_G(R) = M_M(R) + M_{DE}(R)$$

According [12] Chernin, A.D. 2013. Mass of dark energy within a given clustrocentric radius R is

$$M_{DE} = \frac{-8\pi \cdot \rho_{DE} \cdot R^3}{3} \text{ Where } \rho_{DE} = 7.1 \cdot 10^{-27} \text{ kg/m}^3 \text{ is density of dark energy, which is Universal. It is remarkable}$$

that DE density should be multiplied by two times volume of sphere to calculate mass anti gravitating effect.

$$\text{So } M_{DE}(14 \text{ Mpc}) = -2.4 \cdot 10^{15} M_{\odot}.$$

Therefore $M_G(14 \text{ Mpc}) = M_M(14 \text{ Mpc}) + M_{DE}(14 \text{ Mpc}) = 4.6 \cdot 10^{15} M_{\odot} - 2.4 \cdot 10^{15} M_{\odot} = 2.2 \cdot 10^{15} M_{\odot}$ which is a very close value to total gravitating mass found by [13] Diaferio & Geller 1997 with a value $M_G = 2.4 \cdot 10^{15} M_{\odot}$

11. CONCLUSION

It is known that NFW profile works perfectly at galactic scale. However at cluster halo scale has given a very high value for total dark matter which is not supported by observational method. More exactly DM calculated by NFW profile within cluster halo up to 14 Mpc is 6 times bigger over the expected value estimated with other methods.

The original view this work has been to calculate a new DM profile called Bernoulli taking as data NFW profile at radius 1.314 Mpc and 3.1 Mpc. and extend successfully its dominion up to 14 Mpc.

Values of DM density of Bernoulli profile tabulated at 0.1 Mpc from 3.1 to 14 Mpc has been used to calculate numerically total dark matter in cluster halo. So total mass at 14 Mpc sphere is $4.6 \cdot 10^{15} M_{\odot}$ which added to negative dark energy give a Gravitatory mass at 14 Mpc. $M_G = 2.2 \cdot 10^{15} M_{\odot}$ which match perfectly with gravitatory mass found by [13] Diaferio & Geller 1997 with a value $M_G = 2.4 \cdot 10^{15} M_{\odot}$.

During last two year I have published several papers about DM in galaxies through Bernoulli profile with successful results which always are very similar to got with NFW profile.

The importance Bernoulli profile is that is supported by hypothesis of dark matter generated by gravitational field.

In fact in chapter five was got $DM \text{ density} = AE^B$ Where A & B are parameters and E is gravitational field.

I think is not casual that total matter calculated, under this hypothesis, within Coma cluster halo agree perfectly with total matter calculated by observational methods by prestigious researchers.

In my opinion results got in this paper support strongly that DM is generated locally by gravitational field according a universal mechanism that works at galaxy halo scale as well.

In addition, as I published in [1] Abarca, M.2014. the ultimate theory of DM will be a quantum gravity theory.

12. BIBLIOGRAPHIC REFERENCES

[1] Abarca, M.2014,viXra:1410.0200

Dark matter model by quantum vacuum

[2] Abarca, M.2015,viXra:1510.0324

Dark matter density function depending on gravitational field as Universal law

[3] Abarca, M.2015. viXra.org/abs/1512.0309

A new dark matter density profile for NGC 3198 galaxy to demonstrate that dark matter is generated by gravitational field.

[4] Abarca, M.2016.viXra. 1601.0014

A New Dark Matter Density Profile for M33 Galaxy to Demonstrate that Dark Matter is Generated by Gravitational Field

[5] Abarca, M.2016.viXra. 1601.0242v1

A New Dark Matter Density Profile for M31 Galaxy to Demonstrate that Dark Matter is Generated by Gravitational Field.

[6] Abarca, M.2016. viXra.1602.0047

A new Dark matter density profile for Milky Way to demonstrate that dark matter is generated by gravitational field.

[7] Abarca, M.2016. viXra:1606.0007

A new Dark matter density profile for Milky Way which depend on gravitational field.

[8] Abarca,M.2016. viXra:1606.0279v1

Dark matter density profile on big galaxies depend on gravitational field as Universal law

[9] Fuller, C.2016.arXiv:1603.02970v1

H-Atlas: The far-infrared properties of galaxies in and around the Coma Cluster

[10] De Martino.2016.arXiv:1605.08223v1

A $f(r)$ - gravity model of the Sunyaev - Zeldovich profile of the Coma Cluster compatible with Plank data

[11] Plank collaboration. Plank intermediate results X: Physics of the hot gas in the Coma Cluster.

A&A,554,140 (2013).

[12] Chernin, A.D. 2013.arXiv:1303.3800v1.

Dark Energy and the structure of the Coma Cluster of galaxies.

[13] Diaferio,A., Geller M.J.,1997,ApJ,481,633

[14] Diaferio, A.1999, MNRAS,309,610