

The structure of the Pivot Universe

Arieh Sher

email: pivot.universe.theory@gmail.com

Abstract

The Pivot theory presents a new theory regarding the structure of the Universe. It postulates that the Universe is composed of a huge spinning black hole, the Pivot, and a ring of a finite visible Universe that is orbiting it. I claim that the missing dark matter, resides in the Pivot, rather than being spread out through the Universe, as is postulated by the Big Bang (BB) theory.

The Pivot theory, described in this article, consists of two parts. The first relates to the questions of how did it all begin and how did the Universe has evolved into the Pivot structure. This part is not fully addresses. However, the Pivot theory does not come out of the blue. It is based on the theory of the primeval hadron. Using the hadron's theory, the structure of the Universe is explained and its size is calculated. It is shown that the primeval hadron theory can answer questions relating to time, namely, what happened at the exact moment of the Universe creation i.e., $t=0$ and, more importantly, what happened before $t=0$.

The second part relates to the current structure of the Universe. Although the exact way the Universe evolved is not fully addressed, the structure of the Pivot Universe, quite strikingly, is in accord with many known cosmological observations such as: The flattened rotation curve in spiral Galaxies, the Spiral shape of Galaxies, Redshift of Galaxies, Blue shift of Galaxies, Cosmic Microwave Background, Universe rotation observed by Birch, Handedness of Galaxies, Origin of spinning and rotation of all celestial objects, Olbers' paradox, Hubble's observations, Mach's principle.

Key words: Universe structure, Pivot, dark matter, CMB, hadron, gravitational z shift

Contents

- 1. The Pivot theory- Overview**
- 2. Known constants and accepted parameters in cosmology**
- 3. The Primeval hadron theory**
- 4. Dimensions of the Universe**
- 5. The visible Universe is an inertial frame**
- 6. Frame dragging of the space around the Pivot**
- 7. The Origin of spinning and rotation of celestial bodies**
- 8. Rotation curve and spiral shape of a Galaxy**
- 9. The shape of a spiral Galaxy**
- 10. Gravitational z shift**
- 11. Blue shifted Galaxies**
- 12. Andromeda Galaxy**
- 13. Time dilation in the visible Universe**
- 14. Doppler shift Vs. gravitational shift**
- 15. Hubble's observations**
- 16. Cosmic Microwave Background (CMB)**
- 17. Olbers' paradox**
- 18. Handedness of Galaxies**
- 19. Formation of Galaxies, Clusters of Galaxies and Solar systems**
- 20. Mach's principle Vs. Newton's bucket**
- 21. Summary**
- 22. References**

1. The Pivot theory- Overview

The Pivot theory consists of two parts. The first part relates to the questions, how it all began and how has the Universe evolved into the Pivot structure. This part is not fully addressed. However, it has not out of the thin air. It is based on the theory that the Universe began as a primeval hadron. Muradian [1], showed that there is a universal relationship between mass and intrinsic angular momentum of cosmic objects. He showed, by comparing calculations and observations, that this relation is valid for superclusters, Galaxies, stars and planets. However, when he tried to calculate the mass of the entire Universe, he got a large discrepancy. In order to calculate the mass of the Universe, he assumed that the hadron can be modeled as a Kerr black hole. By doing so he limited the mass that was accumulated by the hadron to gravitational mass. I suggest re-modeling the hadron as a Kerr-Newman black hole (KNBH), thus enabling inclusion of the electromagnetic mass. This model answers questions relating to time: what happened at the exact moment of the Universe creation i.e., $t=0$ and, more importantly, what happened before $t=0$. These questions cannot be answered by the Big Bang (BB) theory because there is a singularity problem at $t=0$, indicating that the laws of physics collapse and cannot be applied at the moment of creation.

The second part describes the current structure of the Universe; a huge rotating black hole, the Pivot, with a ring of a finite visible Universe orbiting it in the opposite direction. The Pivot theory solves the dark matter issue by postulating that the dark matter is concentrated in the Pivot rather than being spread throughout the Universe, as claimed by the BB. Being located in this huge black hole is the reason why the dark matter cannot be observed, even though it influences any celestial body, starting from dust and ending in the largest structures in the Universe. The Pivot structure, answers, quite strikingly, observed and unsolved issues in cosmology: Origin of spinning and rotation of all celestial objects, Flattened rotation curve in spiral Galaxies, Spiral shape of Galaxies, Redshift of Galaxies, Blue shift of Galaxies, Cosmic Microwave Background, Universe rotation observed by Birch, Handedness of Galaxies, Olbers' paradox, Hubble's observations (but not Hubble's law!), Mach's principle. The Pivot theory answers two most mind-boggling theories in modern cosmology: the dark matter and the dark energy. The solution for the first is described here and the second is simply not needed in the Pivot theory.

2. Known constants and accepted parameters in cosmology

$Gly = 9.454 \cdot 10^{24} \text{ m}$...Billion light years.

$Mly = 9.454 \cdot 10^{21} \text{ m}$...Million light years.

$Kly = 9.454 \cdot 10^{18} \text{ m}$...Thousand light years.

$G = 6.67 \cdot 10^{-11} \text{ m}^3 / \text{kg} / \text{sec}^2$...Gravitational constant.

$C = 2.99 \cdot 10^8 \text{ m / sec}$...Light velocity.
$\rho = 10^{-26} \text{ kg / m}^3$...Density of ordinary matter in Universe.
$\hbar = 1.054571 \cdot 10^{-34} \text{ J} \cdot \text{sec}$...Reduced Planck's constant.
$m_{\text{proton}} = 1.6726219 \cdot 10^{-27} \text{ kg}$...Mass of proton.
$\omega = 10^{-13} \text{ rad / yr}$...Angular velocity of the Universe- Birch [6].

Birch [6] calculated from the study of position angles and polarization of high luminosity classical double radio sources a rotational velocity for the Universe of the order of $\omega = 10^{-13} \text{ rad / yr}$. Although his work has not been convincingly refuted, it was criticized for using improper statistics, see e.g., Nodland & Ralston [13]. It is interesting to note that Sistero [9] tried to calculate the angular momentum and mass of a rotating Universe based on Birch's finding and Muradian estimations. However, he assumed that the Universe is a homogeneous spherical body with Hubble's radius. The results he got are in great discrepancy from accepted values. He himself acknowledges that the uncertainty in the values he got is very large. Birch's number of $\omega = 10^{-13} \text{ rad / yr}$ plays an important role in the calculations of the Pivot theory, but while viewing the criticism, it should be taken into account that it is possible that this number is not accurate.

$M_{vu} = 1.5 \cdot 10^{53} \text{ kg}$...Mass of visible Universe, Wikipedia [7].
$R = \frac{26.8\%}{4.9\%} = 5.47$... Ratio of dark matter mass to visible Universe mass, Wikipedia [8].
$M_p = R \cdot M_{vu} = 8.2 \cdot 10^{53} \text{ kg}$...Calculated mass of Pivot.
$M_{total} = M_p + M_{vu} = 9.7 \cdot 10^{53} \text{ kg}$...Calculated total mass of the Universe.
$M_{bh} = 10^{36} \text{ kg}$...Assumed average mass of a black hole in the center of a Galaxy.
$M_{gal} = 10^{39} \text{ kg}$...Assumed average mass of a Galaxy.

3. The Primeval hadron theory

The Pivot theory uses the hadron theory described by Muradian [1] and others suggesting that the Universe began as a super dense primeval hadron. He uses the generalized Regge's law general formula which connects the maximal spin J and the mass m of a celestial object:

$$J = \hbar \cdot \left[\frac{m}{m_{proton}} \right]^{1+1/n} \quad \text{where for Galaxies, clusters of Galaxies and globular Galaxies the two-}$$

dimensional disk like hadron $n=2$. For stars, planets and asteroids $n=3$. He showed that the calculated values of J and m of all celestial bodies, namely superclusters, Galaxies, stars and planets, are in good agreement with observations. Muradian [1] is aware that magnetic fields play an important role in astrophysics and that these fields can be a remnant of the magnetic field of the primeval hadron. However, the magnetic fields are not included in his calculations, as he claims that the charge cannot be calculated theoretically. For calculating the mass of the Universe Muradian [2] assumes that the primeval hadron can be modeled as a Kerr black hole.

He equates the maximal angular momentum of the Universe $J = \frac{G \cdot m^2}{C}$ to the angular

momentum of the primeval hadron $J = \hbar \cdot \left(\frac{m}{m_{proton}} \right)^{\frac{3}{2}}$. By doing so he gets that the Universe

mass m is:

$$m = \frac{\hbar^2 \cdot C^2}{G^2 \cdot m_{proton}^3} = 4.77 \cdot 10^{49} \text{ kg} \quad (3.1)$$

The calculated mass from (3.1) is significantly smaller than the accepted mass of the of the Universe that is $m = 1.5 \cdot 10^{53} \text{ kg}$. My claim is that the electromagnetic mass must be added to the hadron mass. I suggest that this discrepancy can be resolved by modeling the hadron as a Kerr-Newman black hole (KNBH). The mass of the KNBH is composed of gravitational mass and electromagnetic mass. The mass of the Universe calculated in equation (3.1) is merely the gravitational mass of the hadron and therefore it is designated hereafter as M_g . The electromagnetic mass Q is a part of M_{total} (the total mass of the Universe that was calculated before).

Let assume $Q = \gamma \cdot M_{total}$, γ can be calculated in two ways:

1) Calculating the total mass of the hadron:

$$M_{total} = M_g + Q = M_g + \gamma \cdot M_{total} \Rightarrow \gamma = 1 - \frac{M_g}{M_{total}} = 0.999951$$

2) Using the KNBH solution that claims that in order not to destroy the horizon, KNBH must also satisfy equation (8)

$$Q^2 + a^2 \leq M_{total}^2 \quad (3.2)$$

The spin parameter a is defined as $a = \frac{J_{proton} \cdot C}{G \cdot M_{total}}$ (3.3) and $J_{proton} = \hbar \cdot \left[\frac{M_{total}}{m_{proton}} \right]^{3/2}$ (3.4)

Substituting (3.4) in (3.3) $\Rightarrow a^2 = \frac{\hbar^2 \cdot C^2}{G^2 \cdot m_{proton}^3} \cdot M_{total} = M_g \cdot M_{total}$

Further substituting Q and a^2 in equation (3.2) gives:

$$\gamma^2 \cdot M_{total} + M_g \leq M_{total} \quad \text{and finally, } \gamma \leq \left(1 - \frac{M_g}{M_{total}} \right)^{0.5} = 0.999975$$

The calculated value of γ shows that practically the entire mass of the primeval hadron was contributed by electromagnetic charge. It should be noted that γ was found here based on the accepted estimation of the total mass of the Universe. From the two calculated values of γ , it is shown the primeval KNBH exploded before the M_{total} reached its maximum allowed value. An open question remains: Why did the primeval hadron explode at this specific γ ?

According to quantum physics, it is possible that the primeval hadron or KNBH accumulated energy from the vacuum. As a result of accumulating the vacuum energy, the mass of the KNBH, its angular momentum and its charge grew. Thus, the total mass of the hadron originated from two sources; the gravitational mass and the electromagnetic charge mass. The KNBH growth continued until it reached an instability state and exploded. KNBHs are not observed in the Universe. Thus, according to the Pivot theory, the existence of KNBH was a unique event in the evolution of the Universe. Two questions are not addressed here: First, how much time did it take for the hadron to accumulate the energy from the vacuum. Second, why did the KNBH's explosion occur at the specific total mass of the current Universe. Explosion of KNBH is an ongoing research. Hod [4], e.g., describes the super radiant instabilities in the composed Kerr-Newman-black-hole-charged massive-scalar-field bomb. The mechanism in which the explosion of KNBH has evolved into the structure of the Pivot Universe can be explained as follows: The explosion produced, according to Newton's 3rd law, an equal inward force. This resulted in particles moving towards the center. Such a mechanism is known as implosion and is practically used in nuclear bombs. When the mass of the inward rushing particles was big enough, a new black hole, the Pivot, was created. A dynamic process occurred involving particles that spread out and particles that were swallowed by the Pivot. Consequently, the Pivot grew bigger until it reached its current size. The Pivot theory postulates that due to the fact that the disk-like primeval hadron had a spin, sub atomic particles, during the explosion, flung off tangentially from it. This tangential motion of the particles accounts for the ring shape of the current visible Universe. It should also be noted that particles that were swallowed by the Pivot impinged it tangentially, causing it to rotate in the opposite direction of the particles that

flew away from it. Eventually, a state of dynamic equilibrium was reached. The particles that escaped the Pivot's event horizon began to orbit around it, creating the visible Universe.

Elementary particles can be presented as black holes, see e.g., Oldershaw [3]. Oldershaw notes that if the primeval hadron, or proton, is modeled as a Kerr black hole, one can retrodict its mass and its radius. However, more accurate mass and radius values for the proton are retrodicted using the full KNBH solution. The conclusion is, that black holes can present the elementary particles as well as the largest cosmic structure in the Universe, including the entire Universe.

4. Dimensions of the Universe

The size of the Universe is calculated based on the equity of total angular momentum of the primeval hadron and its mass to that of the current Universe.

The inner radius of Universe ring must satisfy $R_{in} \geq R_s$ (Schwarzschild radius), for simplicity it is assumed that $R_{in} = R_s$:

$$R_{in} = R_s = \frac{2 \cdot G \cdot M_p}{C^2} = 129.48 \cdot Gly \quad (4.1)$$

The Pivot theory relies on two conservation laws: the mass conservation law and the total angular momentum conservation law. Based on the mass conservation law it is postulated that the total mass M_{total} of the current Universe is equal to the mass of the primeval hadron. The conservation law of the total angular momentum states that the sum of the orbital angular momentum plus the intrinsic spin of a system is constant. The Universe began as a hadron that had an intrinsic angular spin. Given that at the time of creation, the spinning primeval hadron was the only existing body, the Universe orbital angular momentum was zero. Following the hadron explosion, two distinct bodies were created: the Pivot, having only a spin angular momentum and the visible Universe that has orbital angular momentum. (Note: there is also the spin of celestial bodies, Galaxies, stars and planets. But it will be shown that the sum of spin angular momentum of the celestial bodies is negligible in comparison to the orbital angular momentum of the visible Universe ring). Since the orbital angular momentum at the time of the hadron explosion was zero, the sum of the total angular momentum in the current Universe must be also zero. Thus, the Pivot spin and the orbital angular momentum of the visible Universe must be equal, albeit, with opposite signs.

The spinning angular momentum of a disk-like hadron (proton) is given by Muradian [1]:

$$J_{proton} = \hbar \cdot \left[\frac{M_{total}}{m_{proton}} \right]^{3/2} = 1.474 \cdot 10^{94} \text{ erg} \cdot \text{sec} \quad (4.2)$$

The outer radius of the visible Universe, R_{out} , can be calculated from (4.3). The left side of equation (4.3) is the angular momentum of the ring shaped visible Universe J_{vu} . It is equal to half of the angular momentum of the proton:

$$\frac{M_{vu}}{2} \cdot (R_{in}^2 + R_{out}^2) \cdot \omega = J_{vu} = \frac{J_{proton}}{2} \Rightarrow R_{out} = 133.87 Gly \quad (4.3)$$

All celestial bodies i.e., Galaxies, stars, planets, interstellar planets, also have a spinning angular momentum. However, it is shown now, that the sum of the spin angular momentum of all these celestial bodies is negligible in comparison to the orbital angular momentum of the visible Universe disk. Based on Muradian [1], an estimated total angular momentum of all celestial objects is:

$$J_{obj} = J_{gal} \cdot N_{gal} + J_{star} \cdot N_{star} + J_{planet} \cdot N_{planet} = 2 \cdot 10^{85} erg \cdot sec \quad (4.4)$$

Where:

Average angular momentum of a Galaxy $J_{gal} = 10^{74} erg \cdot sec$, Number of Galaxies $N_{gal} = 2 \cdot 10^{11}$,

Angular momentum of an average star $J_{star} = 10^{49} erg \cdot sec$, Number solar systems $N_{star} = 10^{22}$,

Angular momentum of an average planet $J_{planet} = 10^{40} erg \cdot sec$, Number planets $N_{planet} = 10^{24}$.

From (4.4) $J_{obj} = 2 \cdot 10^{85} erg \cdot sec \ll J_{vu} = 7 \cdot 10^{93} erg \cdot sec$ and therefore J_{obj} can be neglected in (4.3).

The width of the visible Universe ring, W , can be calculated from equation (4.5):

$$M_{vu} = W \cdot \rho \cdot \pi \cdot (R_{out}^2 - R_{in}^2) \Rightarrow W = 4.9 Gly \quad (4.5)$$

The Pivot's spin ω_p can be calculated from equation (4.6). The left side of (4.6) is the angular momentum of the Pivot J_p and is equal to half the angular momentum of the proton. It is assumed in (4.6), that the Pivot is a sphere with an equally distributed density. The minus sign indicates that the Pivot spins in opposite direction to the visible Universe.

$$-\frac{2}{5} \cdot M_p \cdot R_s^2 \cdot \omega_p = J_p = \frac{J_{proton}}{2} \Rightarrow \omega_p = -4.73 \cdot 10^{-14} rad / yr \quad (4.6)$$

Fig. 4.1 shows the dimensions of the Pivot Universe according to the above calculations.




Fig. 4.1 – Structure and dimensions of the Pivot Universe

5. The visible Universe is an inertial frame

The celestial bodies in the visible Universe, although located around such a huge black hole, can be considered as inertial frames. An inertial frame must fulfill two requirements:

- 1) The Galaxies are not accelerated (or the net force acting on them is zero).
- 2) The Galaxies move at a constant velocity in a straight line.

The first requirement is fulfilled by calculating the gravity of the Pivot acting in the visible Universe. The gravity is calculated at R_{in} (5.1) and at R_{out} (5.2):

$$g_{p_in} = \frac{G \cdot M_p}{R_{in}^2} = 3.65 \cdot 10^{-11} \frac{m}{\text{sec}^2} \quad (5.1)$$

$$g_{p_out} = \frac{G \cdot M_p}{R_{out}^2} = 3.41 \cdot 10^{-11} \frac{m}{\text{sec}^2} \quad (5.2)$$

This gravity of the Pivot is miniscule and is almost the same everywhere in the visible Universe.

The second requirement is fulfilled by the fact that the visible Universe radius is so enormous that the observable Galaxies can be considered as moving in straight lines. In addition, each Galaxy is orbiting the Pivot at a constant velocity.

This gives rise of using Newton's law of universal gravitation and the laws of SR. This assumption is used later for other calculation e.g., the flattened rotation curve in spiral Galaxies and the spiral shape of Galaxies.

6. Frame dragging of the space around the Pivot

The average velocity V_{avg} of the visible Universe at $R_{avg} = \frac{(R_{in} + R_{out})}{2}$ can be calculated in two ways:

1) Using Newton's law of universal gravitation (6.1):

$$V_{avg} = \left(\frac{G \cdot M_p}{R_{avg}} \right)^{0.5} \cong 0.7C \quad (6.1)$$

2) Using angular velocity formula (6.2):

$$V_{avg} = \omega \cdot R_{avg} \cong 0.0132C \quad (6.2)$$

There is a big discrepancy between the results of (6.1) and (6.2). This discrepancy is resolved by taking into account the frame dragging effect the Pivot has on the visible Universe. The twist of the space is such that the visible Universe that rotates in the opposite direction to the Pivot's spin will move slower in reference to the Pivot axis.

The frame dragging angular speed Ω is calculated by (6.3):

$$\Omega = \frac{R_{in} \cdot \alpha \cdot C}{R_{in}^3 + \alpha^2 \cdot R_{avg} + R_{in} \cdot \alpha^2} = 1.798 \cdot 10^{-14} \frac{rad}{yr} \quad (6.3)$$

Where $\alpha = \frac{J_p}{M_p \cdot C} = 3 \cdot 10^{24} m$ and the angular momentum of the Pivot: $J_p = \frac{J_{proton}}{2}$

The velocity of the Universe around the Pivot axis is

$$V_{avg} = (\omega - \Omega) \cdot R_{avg} = 0.0108C \quad (6.4)$$

The result of (6.4) is in reasonable agreement with (6.2), when taking into account that some of the parameters used in this calculation are estimated

7. The Origin of spinning and rotation of celestial bodies

Following the explosion of the primeval hadron, it took the visible Universe some time (how long?) to arrange itself in an orbit around the Pivot. The temperature of the visible Universe, after the explosion, was extremely high and contained only a soup of sub atomic particles. Later on, as the Universe cooled down, ordinary atoms were created and this was the time that gravity forces between atoms appeared. The local density of the visible Universe was the cause of the variety of celestial bodies, which enabled several scenarios. If the density of atoms at a certain region in the visible Universe was too low to enable attraction between them, they remained as a cloud of gas that orbits the Pivot. If the density of atoms was sufficient for interaction between them, then a nucleus of a star was created. The nucleus of the star encountered atoms from various orbits having different velocities. Stars that are farther from the Pivot than the Galaxy's center, move faster than the Galaxy's center. Stars located closer to the Pivot move slower. This created a torque on the star nucleus, consequently causing the star to spin while orbiting the Pivot, simultaneously. Fig 7.1 shows trajectories of stars (or any other celestial bodies) around the center of the Galaxy. If the mass of the star became big enough, it collapsed and created a black hole. The black hole attracted other stars that began orbiting it. The stars already orbiting the Galaxy's black hole attracted additional stars and thus giving birth to a full-sized Galaxy.




Fig. 7.1 – Trajectories of stars around a Galaxy black hole

8. Rotation curve of a Galaxy

The rotation curve of a Galaxy is a plot of the orbital speeds of visible stars or gas in that Galaxy versus their radial distance from that Galaxy's center. (See Fig. 8.2). The rotation curves of Galaxies were measured accurately by Vera Rubin and her team. They verified that the rotation curve was flattened, or in other words, that the velocities of stars orbit at roughly the same speed, not depending on their distance from the Galaxy's center. This observation is contrary to Newton mechanics that claims that stars that are located further from the Galaxy center will move slower. The rotation curves can be explained by hypothesizing the existence of a substantial amount of matter influencing the Galaxy. Cosmologists have calculated the amount of this matter to be 5.5 times the amount of the mass of the visible matter. The material responsible for the extra mass was dubbed, "dark matter" because it has not been found by observations and its nature is not known.

The Pivot theory accepts the hypothesis that the curve flattening is caused by the dark matter, but claims that the dark matter resided in the Pivot rather than being spread in and around Galaxies as postulated by current cosmology. The fact that the dark matter is located in the Pivot black hole is the reason why it cannot be detected. The Pivot theory explains in this paragraph the flattened curve of Galaxies by calculating the velocities of stars in Galaxies. The mass of the Pivot used in the calculations is assumed to be equal to the amount of the dark matter calculated by cosmologists to explain the curve flattening.

In the previous paragraph, the origin of spinning and rotation of celestial bodies was explained. It has been also calculated in paragraph 6 that any celestial body orbits the Pivot at an average speed of $\sim 0.7C$. A celestial body that is part of a Galaxy performs simultaneously two motions. The first is orbiting the Pivot at a speed of $\sim 0.7C$. The second is orbiting the center of the Galaxy at a much lower speed of \sim tens to hundreds km/sec. The superposition of the two velocities results in the flattened curve of stars in a Galaxy. The following calculation is done on an arbitrary Galaxy located 150Mly from the Milky Way. The distance of 150Mly between the Galaxies is assumed in order to assure that there is no gravitational pull between Galaxies. (See paragraph 19, where the gravitational pull is calculated for Galaxies that are much closer to each other). It is shown that the exact shape of the curve is dependent on several parameters. This is the reason why, when curves of many Galaxies are depicted on the same graph, the motion of stars seems somewhat chaotic (See Fig. 8.2).




Fig. 8.1– Velocities of a star

The gravitational forces exerted on a star in a Galaxy are caused by:

- Distributed mass of the Galaxy.
- The Black Hole in the center of a Galaxy.
- The Pivot.

Note: The calculations are done with MathCad.

a) Velocity of a star due to distributed mass of the Galaxy

$$r := 0.0001\text{Kly}, 0.01\text{Kly} \dots 100\text{Kly}$$

Distance range of star to its Galaxy center

$$r_0 := 30\text{Kly}$$

Assumed characteristic radius.

$$V_{\text{gal}}(r) := \begin{cases} \left(\frac{G \cdot M_{\text{gal}}}{r_0}\right)^{0.5} \cdot \frac{r}{r_0} & \text{if } 0\text{Kly} \leq r < r_0 \\ \left(\frac{G \cdot M_{\text{gal}}}{r}\right)^{0.5} & \text{if } r \geq r_0 \end{cases} \quad \text{Rotation velocity due to distributed mass of a Galaxy}$$

b) The velocity of a star around the Galaxy's Black Hole:

$$V_{\text{bh}}(r) := \left(\frac{G \cdot M_{\text{bh}}}{r}\right)^{0.5}$$

c) The rotation velocity of a star in a Galaxy orbiting the Pivot:

$$d := 150 \cdot \text{Mly}$$

Arbitrary assumed distance

$$R_{\text{gal}} := R_{\text{mw}} + d$$

$$V_{\text{p_star}}(r, \alpha) := \left(\frac{G \cdot M_{\text{p}}}{R_{\text{gal}} - r \cdot \cos(\alpha)}\right)^{0.5}$$

Summation of the three velocities on the star:

$$V_{\text{sum}}(r, \alpha) := V_{\text{p_star}}(r, \alpha) + (V_{\text{gal}}(r) + V_{\text{bh}}(r)) \cdot \cos(\alpha)$$

Only as a reference, the magnitudes of the three velocities are calculated for a star located 10Kly from the Galaxy's black hole. It is clear that the velocity of a star around the Pivot is dominant.

$$V_{\text{p_star}}(10\text{Kly}, 0\text{deg}) = 210864 \frac{\text{km}}{\text{s}}$$

$$V_{\text{gal}}(10\text{Kly}) = 5.1 \cdot \frac{\text{km}}{\text{s}}$$

$$V_{\text{bh}}(10\text{Kly}) = 0.8 \cdot \frac{\text{km}}{\text{s}}$$

Orbital velocity of Milky Way around the Pivot:

$$V_{p_mw} := \left(\frac{G \cdot M_p}{R_{mw}} \right)^{0.5}$$

The velocities of Galaxies in the Universe, as shown earlier in this paper, are $\sim 0.7C$. The relativistic velocity of a star in a Galaxy as seen by a Milky Way observer is given by (8.1):

$$V_{star}(r, \alpha) := \frac{V_{p_mw} - V_{sum}(r, \alpha)}{1 - \frac{V_{p_mw} \cdot V_{sum}(r, \alpha)}{c^2}} \quad (8.1)$$

The rotation curve as observed for Galaxies is somewhat chaotic and is shown in Fig. 8.2. The common characteristic for all observations is that as the distance of a star to the center of its Galaxy increases the curve flattens out.




Fig. 8.2 -Rotation curves of Galaxies (from: Sofue [10])

The calculated velocity of a star in a Galaxy is given by (8.1) and is shown Fig. 8.3. It is seen that the observed velocities of stars in a Galaxy will be confined between the two extreme curves of the graph (solid red and dashed blue). The exact shape of the graph is dependent mainly on α , d and the mass of Galaxy. It is seen that Fig. 8.3 resembles Fig. 8.2.




Fig. 8.3 – Rotation curve for a Galaxy at a distance of 150Mly from Milky Way

9. The shape of a spiral Galaxy

The spiral shape of Galaxies is based on the equations used in paragraph 7 and 8. Most spiral Galaxies contain a central bulge surrounded by a flat rotating disk of stars. The bulge in the Galaxy center contains a massive black hole. The bulge has an ellipsoidal shape, because stars that were attracted by the central black hole in the Galaxy, originated not only from stars that were located on the equatorial plane of the Galaxy but also from planes that are “above” or “below” the equatorial plane. Orbiting the bulge, the rotating disk separates itself into arms that circle the Galaxy center. The spiral arms are the result of the superposition of the velocity of $\sim 0.7C$ of a star around the Pivot and its velocity around the center of the Galaxy (several to hundreds km/sec). An additional observation explained by the Pivot theory is why spiral Galaxies may have more than two arms. A Galaxy has more than two arms if during its rotation around the Pivot the Galaxy encountered several dense regions at various times. Each time the Galaxy crosses a dense region new arms are added to the Galaxy. The width of the arm depends on the size of the dense region. Some Galaxies may encounter so many dense regions, that its arms may appear as an elliptical Galaxy.

The angular displacement of a star orbiting the Galaxy’s black hole (θ) (See Fig. 8.1) is the integral $\int \frac{V_{star}}{r} dt$, where V_{star} is given in (8.1) This integration is done only for stars that are located on the main axis (i.e., the axis that connects the Pivot and the Galaxy’s black hole, i.e.,

angles of 0deg and 180deg – (See Fig. 7.1 and Fig. 8.1). Only at these angles, stars begin to orbit around the Galaxy’s center. Fig. 9.1 shows the shape of a spiral shape Galaxy 13.3 Billion years after the Galaxy creation. It can be shown that the spiral shape is changing at a very slow pace over Billions of years.

Note: In the following equations, the “mod” operator of MathCad- returns the remainder on dividing x by y (x modulo y). In the current case y=360deg. The reason for using this function is that stars in Galaxies have completed many full rotations around the Galaxy’s black hole during 13.3 Billion years.

$$\theta_1(r, \alpha) := \text{mod} \left[\int_{0\text{yr}}^{13.3 \cdot 10^9 \text{yr}} \frac{V_{\text{star}}[r, (\alpha) \cdot \text{deg}]}{r} dt \cdot \text{deg}, 360\text{deg} \right]$$

$$\theta_2(r, \alpha) := \text{mod} \left[\int_{0\text{yr}}^{13.3 \cdot 10^9 \text{yr}} \frac{V_{\text{star}}[r, (\alpha) \cdot \text{deg}]}{r} dt \cdot \text{deg}, 360\text{deg} \right] + 180\text{deg}$$




Fig. 9.1 – Shape of spiral Galaxy 13.3 Billion years after its creation.

10. Gravitational z shift

The Pivot theory postulates that the z shift of Galaxies is caused by the gravitational field. It states that electromagnetic radiation originating from a Galaxy orbiting the Pivot is reduced in frequency, or redshifted, when observed from a Galaxy that is closer to the Pivot. This is contrary to Hubble’s assumption that the z shift of Galaxies is caused by the Doppler effect. The

Doppler shift is caused by the difference in the velocities of the observed object and the Milky Way. However, since the Pivot theory claims that Galaxies are orbiting the Pivot at different velocities, the Doppler shift cannot be excluded. It will be shown later that its contribution to the measured z shift is small.




Fig. 10.1 – The Pivot Universe structure shows Galaxies orbiting the Pivot

Fig. 10.1 is a schematic structure of the Pivot Universe. It includes the Pivot, containing the entire dark matter in the Universe, with a ring of the visible Universe orbiting it. Clearly, the inner radius of the ring must be greater than the event horizon of the Pivot. Calculations of the gravitational z shift of a Galaxy Z_{gal_pivot} orbiting the Pivot at a radius R_{gal} , in the gravitational field of the Pivot, is given in equation (10.1) and shown in Fig. 10.2:

$$Z_{gal_pivot} = \frac{1}{\left(1 - \frac{2 \cdot G \cdot M_p}{R_{gal} \cdot C^2}\right)^{0.5}} - 1 \quad (10.1)$$




Fig. 10.2 – z shift in the Pivot’s gravitational field.

The next step is finding the z shift of Galaxies as seen from the Milky Way (Z_{gal}). This is done by using the following assumption: Galaxy GN-z11 was found to have the highest z shift ever measured $Z_{gal} = 11.09$. It is **assumed** that GN-z11 is located on the outside radius of the Universe R_{out} . On the other hand, from equation (10.1), the z shift of GN-z11, as seen from the Pivot, is 4.527. Thus, the z shift of the Milky Way Z_{mw} , as seen from the Pivot, can be calculated by equation (10.2). Note: Should Galaxies with higher z shift, will be found in the future, Z_{mw} be corrected accordingly.

$$Z_{mw} = 11.09 + 4.527 = 15.617 \quad (10.2)$$

Having Z_{mw} the orbiting radius of the Milky Way R_{mw} is calculated by (10.3)

$$R_{mw} = \frac{2 \cdot G \cdot M_p}{\left(1 - \frac{1}{(Z_{mw} + 1)^2}\right) \cdot C^2} = 129.954703 Gly \quad (10.3)$$

Now, the z shift of any Galaxy, as seen from the Milky Way Z_{gal} is calculated from (10.4).

$$Z_{gal} = Z_{mw} - Z_{gal_pivot} \quad (10.4)$$

Having Z_{gal} , the orbiting radius of any Galaxy R_{gal} can be calculated by (10.5) and shown in Fig. 10.3.

$$R_{gal} = \frac{2 \cdot G \cdot M_p}{\left(1 - \frac{1}{(Z_{gal} + 1)^2}\right) \cdot C^2} \quad (10.5)$$




Fig. 10.3 – z shift of Galaxies as seen from Milky Way. The dashed lines relate to the Milky Way: $Z_{mw}=0$ and $R_{mw} = 129.95Gly$

11. Blue shifted Galaxies

Fig. 10.1 shows that Galaxies observed from the Milky Way can be either red shifted or blue shifted. For Galaxies that are closer to the Pivot than the Milky Way, $Z_{gal} < 0$. Galaxies that are orbiting the Pivot at a radius larger than the Milky Way radius are red shifted $Z_{gal} > 0$. Had the Milky Way been located on the outer radius of the Universe ring, all Galaxies would have been blue shifted. Referring to Fig. 10.1, GAL-A and GAL-B are shown to have different distances from the Milky Way. Nevertheless, they have the same z shift, because both are located on the same orbit. In order to determine whether the observed Galaxy is approaching or receding the Milky Way the change in the measured distance should be found.

It is clear that the Milky Way orbiting radius is between R_{in} and R_{out} . The distance of the Milky Way from R_{in} is: $R_{mw} - R_{in} = 129.954703 Gly - 129.4840 Gly = 0.471Gly = 471Mly$. The distance of the Milky Way from R_{out} is $R_{out} - R_{mw} = 133.868Gly - 129.954703 Gly = 3.91Gly$. This result explains why there are more observed red-shifted Galaxies than blue-shifted Galaxies.

12. Andromeda Galaxy

Andromeda Galaxy is blue shifted. For Andromeda, it was measured: $Z_{gal} = -0.001001$, distance - 2.5Mly and it is predicted, by others, that the Milky Way and Andromeda will collide in 4Gly.

See Wikipedia [14]. It is interesting to examine the prediction of a collision between Andromeda and the Milky Way, based on the Pivot theory. The orbiting radius of Andromeda R_{andro} is calculated by (12.1):

$$R_{andro} = \frac{2 \cdot G \cdot M_p}{\left(1 - \frac{1}{(Z_{mw} + 0.001001 + 1)^2}\right)} \cdot C^2 = 129.95465 Gly \quad (12.1)$$

The difference in orbiting radiuses of the Milky Way and Andromeda is:

$R_{mw} - R_{andro} = 129.954703 Gly - 129.95465 Gly = 56.914 Kly$. The disk diameter of Andromeda is 220Kly and the disk diameter of the Milky Way is 180Kly. The result is that there is a chance of collision, because $(220Kly + 180Kly)/2 = 150Kly > 56.914Kly$. A collision will occur if Andromeda and the Milky Way are located on the same plane. However, if the two Galaxies are located on different planes that are separated more than 150Kly, no collision is expected to occur.

To find the time to collision, first the velocity of each Galaxy is found (12.2) and (12.3), then the relativistic velocity between the two Galaxies is calculated (12.4) and finally, the time is calculated (12.5).

$$V_{mw} = \left(\frac{G \cdot M_p}{R_{mw}}\right)^{0.5} \quad (12.2) \quad V_{andro} = \left(\frac{G \cdot M_p}{R_{andro}}\right)^{0.5} \quad (12.3)$$

$$V = \frac{V_{mw} - V_{andro}}{1 - \frac{V_{andro} \cdot V_{mw}}{C^2}} = 0.092 \frac{km}{sec} \quad (12.4) \quad t = \frac{2.5 Mly}{V} = 8 \cdot 10^{12} yr \quad (12.5)$$

The time for a collision between Andromeda and the Milky Way calculated by the Pivot theory differs significantly from the prediction of the BB cosmology: $4 \cdot 10^9 yr$

13. Time dilation in the visible Universe

The Pivot is a Kerr black hole. The time dilation equation of Kerr metric takes the form of:

$$t_f = \frac{t_0}{\left(1 - \frac{2 \cdot G \cdot M_p \cdot R_{gal}}{C^2 \cdot \rho^2}\right)^{0.5}} \quad \text{where} \quad \rho^2 = R_{gal}^2 + \frac{J_p}{M_p \cdot C^2} \cdot \cos^2 \theta \quad (13.1)$$

The parameters of the Pivot are such that it can be considered as a Schwarzschild black hole. Assuming $(\cos \theta = 1, R_{gal} = R_s)$.

Calculating (13.1) gives: $\frac{J_p}{M_p \cdot C^2} = 9 \cdot 10^{48} m^2 \ll R_{gal}^2 = 1.5 \cdot 10^{54} m^2$ therefore (13.1) becomes:

$$\rho^2 = R_{gal}^2$$

and Schwarzschild time dilation equation (13.2) can be used:

$$t_f = \frac{t_0}{\left(1 - \frac{2 \cdot G \cdot M_p}{C^2 \cdot R_{gal}}\right)^{0.5}} \quad (13.2)$$

Assuming the time elapsed at the Milky Way is 1 sec, then the time of other Galaxies in the visible Universe is calculated by (13.3) and is shown in Fig. (13.1):

$$t_f = \frac{1 \text{ sec}}{\left(1 - \frac{2 \cdot G \cdot M_p}{C^2 \cdot R_{mw}}\right)^{0.5}} \cdot \left(1 - \frac{2 \cdot G \cdot M_p}{C^2 \cdot R_{gal}}\right)^{0.5} \quad (13.3)$$




Fig 13.1 – Time dilation in the visible Universe

Assuming the elapsed time in the Milky Way is 1 sec, the elapsed time for a Galaxy located at R_{out} is 3 sec. A Galaxy near R_s the elapsed time approaches 0 sec.

14. Doppler shift Vs. gravitational shift

This paragraph calculates the contribution of the Doppler effect to the observed z shift of a Galaxy. For finding the Doppler shift of an observed Galaxy, first, the orbiting velocities around the Pivot of the Milky Way (14.1) and the observed Galaxy (14.2) should be found. Then, the relativistic velocity of the two Galaxies is calculated (14.3). Subsequently, the Doppler shift is calculated (14.4). Finally, the ratio of the doppler shift to the total z shift is calculated (14.5) and shown in Fig. 14.1. The Doppler effect contribution to the z shift is less than 0.2%.

$$V_{mw} = \left(\frac{G \cdot M_P}{R_{mw}}\right)^{0.5} = 0.706C \quad (14.1)$$

$$V_{gal} = \left(\frac{G \cdot M_P}{R_{gal}}\right)^{0.5} \quad (14.2)$$

$$V = \frac{V_{mw} - V_{gal}}{1 - \frac{V_{gal} \cdot V_{mw}}{C^2}} \quad (14.3)$$

$$Z_{doppler} = \frac{V}{C} \quad (14.4)$$

$$Ratio = \frac{Z_{doppler}}{Z_{gal} + Z_{doppler}} \quad (14.5)$$




Fig. 14.1 - The ratio of Doppler shift to the total shift

15. Hubble's observations

This paragraph's aim is to show that Hubble's law is wrong. The cosmological principle is supported by Hubble's law. This is in contrast to the Pivot theory that claims the Universe has a rotation axis. I am not disputing the correctness of the observations made by Hubble and others. Nevertheless, I claim that the laws he derived from these observations are wrong.

Hubble's law is a cornerstone in the Big Bang (BB) theory. Hubble's law consists actually of two laws. The first states that Galaxies are receding from each other at a velocity that is linear to their relative distances $V = H_0 \cdot d$. An important conclusion that can be derived from the BB theory is that the Universe "on large-scale" is isotropic and homogenous. The second law claimed by Hubble is $V = z \cdot C$, meaning that there is a linear relation between the receding velocity and the redshift of a Galaxy. Hubble's laws seemed to be correct at his time, when measurements were done on Galaxies near the Milky Way. However, when Hubble's law was used for observations of distant Galaxies with high redshifted the graph curved up. This finding was a basis of an additional paradigm: The Universe is expanding forever in all directions, at an ever-accelerating speed. There are observed Galaxies with $z > 1.5$ that are receding from the Milky Way at speeds greater than the speed of light, Davis & Lineweaver [11]. The accelerated expanding Universe gives rise to yet another theory claiming that in order to accelerate the Universe expansion at the observed rates, there must be an additional force. This force is dubbed "dark energy". The mass of this dark energy was calculated to be ~70% of all the mass in the Universe.

The confusion, among the scientific community, is great. Kirshner [12] relates to Hubble's diagram (Fig. 15.1): "Staring at his original Hubble diagram, you can see that there is a handful of nearby Galaxies with blue shifts, and a large scatter of velocities at any given distance. Hubble shrewdly used plausible methods to average the data for Galaxies that are at the same distance to make his result stand out more clearly from the noise. He was fortunate to have data that behaved so well."




Fig. 15.1 – Hubble's original graph (1929)

Hubble's graph (Fig. 15.1) shows that there is a significant scatter of the measurements. There are Galaxies that are located at the same distance from the Milky Way but have different velocities. The Pivot theory can explain Hubble's observations. Fig. 10.1 shows two Galaxies

GAL-A and GAL-C having the same distance (d) from the Milky Way, but are located on different radii. According to the Pivot theory, they have different z shifts. But according to Hubble, they should have the same redshift and the same velocities. Hubble's law is valid only in a special case, when the observed Galaxies are located on the main axis (the axis that connects the Milky Way and the Pivot), as is shown e.g., for GAL-B. In this case, $\Delta R=d$ and the relation velocity-distant is linear. To sum up: the linear relation in Hubble's law is velocity-distance, whereas in the Pivot theory the linear relation is between the velocity and the difference in the orbiting radii. The relation between the relativistic velocity of a Galaxy and its radius is calculated by (14.3) and the linear relation is shown in Fig. 15.2.




Fig. 15.2 – Linear relation between Galaxy's velocity and its orbiting radius.

From Fig. 15.2 - The linearity constant is 5.1 km/s/Mpc . (This is in comparison to Hubble's constant, that is 70 km/s/Mpc). The maximal and constant recessional velocity of a Galaxy located at R_{out} is $\sim 6200 \text{ km/s}$.

From the Pivot model, the relation between the recessional velocity and z can be found by combining (10.4) and (14.3) and the result is shown in Fig. 15.3. Clearly, this is not a linear relation as claimed by Hubble's law.




Fig. 15.3 – Velocity of a Galaxy as function of z shift

16. Cosmic Microwave Background (CMB)

A known paradigm in modern cosmology is the cosmological principle. It states that at a “large-scale” (hundreds of million light years), the spatial distribution of matter in the Universe is isotropic and homogeneous. The strongest observational evidence for the cosmological principle is the CMB. Therefore, people may claim that the Pivot Universe, that has a preferred axis, is improbable. I claim that even though the Pivot Universe has a preferred axis, it still agrees with the CMB observation. It is demonstrated in this article, that the Milky Way is located inside a huge observable Universe. The observable Universe is a sector of a ring having a cross section of $4.9\text{Gly} \times 4.4\text{Gly}$. According to the calculation based on z shift, shown in paragraph 10, the Milky Way is located 470Mly from R_m . This means that the Milky Way is located inside a “large-scale Universe”. A Milky Way observer sees a huge number of Galaxies in whatever direction he looks. He may conclude that the entire Universe is isotropic and homogeneous. But for an observer in a Galaxy located on the outer surface of the ring, the Universe is not isotropic and homogeneous. In one direction, he will be facing the vacuum which is totally dark and has a temperature of absolute zero. Likewise, an observer that is located on the inner surface of the ring, i.e. facing the Pivot, sees a total darkness and measures the temperature of the Pivot that is practically absolute zero ($1.5 \times 10^{-31}\text{K}$). Only an observer that will travel outside the ring of the visible Universe will be able to see that the Universe has an axis of rotation.

17. Olbers' paradox

The paradox is that a static, infinitely old Universe with an infinite number of stars distributed in an infinitely large space would be bright rather than dark. The BB theory explains the paradox by the fact that space is expanding and therefore the emitted light is reduced via redshift. The Pivot Universe, on the other hand, is a finite Universe with a finite number of stars. Outside the ring of the visible Universe there is the infinite darkness of the vacuum and the darkness of the Pivot. An observer located in the visible Universe sees a huge number of Galaxies in whatever direction he looks, but eventually, in the background, he sees the total darkness.

18. Handedness of Galaxies

The Pivot theory explains yet an additional observation related to the handedness of the Universe. Longo and others [5] found abundance of left-handed, or counter-clockwise (CCW) Galaxies. The excess is small, about 7 percent, but the chance that it could happen in an isotropic and homogeneous Universe is slim. Fig. 18.1 shows that the observable Universe is contained within a sector of the visible Universe ring. It shows the equatorial plane of the visible Universe and three Galaxies, the Milky Way, and two arbitrary Galaxies: Galaxy A and Galaxy B. The Milky Way is located "above" Galaxy B and "below" Galaxy A. The Pivot theory

postulates that all Galaxies rotate in the same direction, opposite to the Universe ring rotation. The location of the Galaxies in relation to the Milky Way will determine the direction of their rotation as seen from the Milky Way. Thus, Galaxy A will be seen as rotating clockwise (CW), while Galaxy B will be seen as rotating counter clockwise (CCW) when observed from the Milky Way. Had the Milky Way been located on the equatorial plane then the number of CCW and CW Galaxies would have been the same. However, if the Milky Way is offset by 7% from the equatorial plane then the number of CCW Galaxies and CW Galaxies will differ by 7%.




Fig. 18.1- Handedness of Galaxies

19. Formation of Galaxies, Clusters of Galaxies and Solar systems

In paragraph 8 and 9, the shape of a Galaxy was calculated. It was assumed there that the distance between the Milky Way and the Galaxy is 150Mly, so that the gravitational pull between the Galaxies, in comparison to the gravitational pull of the Pivot, can be neglected. In this paragraph, the distance between Galaxies was reduced to 2.5Mly.

The explanation of formations of galaxies, clusters of Galaxies and Solar systems is based on calculating the gravitational pull between celestial bodies. Newton's law of universal gravitation states that a particle attracts every other particle in the universe by a force that is directly proportional to the product of their masses and inversely proportional to the square of the distance between their centers. The force on each celestial body is the sum of all the forces exerted on it by all other celestial bodies. It has been claimed earlier that the Pivot influences every celestial body in the Visible Universe. However, celestial bodies are also influenced by other celestial bodies. In this paragraph the gravitational pull between celestial bodies is compared to the gravitational pull exerted on them by the Pivot. Three cases are calculated:

- 1) The gravitational pull on Galaxies
- 2) The gravitational pull on Galaxies in clusters and
- 3) The gravitational pull on planets and stars.

1) Gravitational pull between the Milky Way and Andromeda Vs. the Pivot pull:

(Note: Andromeda Galaxy is the closest Galaxy to the Milky Way).

$$M_{mw} = 2 \cdot 10^{42} \text{ kg} \quad \dots \text{ Mass of Milky Way}$$

$$M_{andro} = 3 \cdot 10^{42} \text{ kg} \quad \dots \text{ Mass of Andromeda}$$

$$R_{andro} \cong R_{mw} = 129.95 \cdot \text{Gly} \quad \dots \text{ Orbiting radius of Milky Way around the Pivot}$$

$$D_{mw_andro} = 2.5 \cdot \text{Mly} \quad \dots \text{ Distance of Andromeda from Milky Way}$$

$$F_{p_mw} = \frac{G \cdot M_p \cdot M_{mw}}{R_{mw}^2} \quad (19.1) \quad \dots \text{ The Pivot gravitational pull on the Milky Way}$$

$$F_{mw_andro} = \frac{G \cdot M_{andro} \cdot M_{mw}}{D_{mw_andro}^2} \quad (19.2) \quad \dots \text{ The force between the Milky Way and Andromeda}$$

The ratio between (19.1) and (19.2) gives: $R1 = \frac{F_{p_mw}}{F_{mw_andro}} = 101$. The meaning is that the Pivot

gravitational pull is 100 times higher than the gravitational pull between neighboring Galaxies.

Note: Andromeda is the closest Galaxy to the Milky Way. The average distant between Galaxies is ~6Mly. This means that the influence of the Pivot on Galaxies is even higher than the ratio R1.

2) Gravitational pull between Milky Way and Sagittarius Dwarf Spheroid Vs. the Pivot pull:

Note: Sagittarius Dwarf Spheroid is the closest Galaxy to the Milky Way in the Milky Way local group.

$$M_{mw} = 2 \cdot 10^{42} \text{ kg} \quad \dots \text{ Mass of Milky Way}$$

$$M_{sagit} = 1 \cdot 10^{33} \text{ kg} \quad \dots \text{ Mass of Sagittarius Dwarf Spheroid}$$

$$R_{sagit} = R_{mw} = 129.95 \cdot \text{Gly} \quad \dots \text{ Orbiting radius of Sagittarius around the Pivot}$$

$D_{mw_sagit} = 0.08 \cdot Mly$ Distance of Sagittarius from Milky Way

$$F_{p_sagit} = \frac{G \cdot M_p \cdot M_{sagit}}{R_{sagit}^2} \quad (19.3) \quad \dots \text{The Pivot gravitational pull on Sagittarius}$$

$$F_{mw_sagit} = \frac{G \cdot M_{sagit} \cdot M_{mw}}{D_{mw_sagit}^2} \quad (19.4) \quad \dots \text{The force between Milky Way and Sagittarius}$$

The ratio between (19.3) and (19.4) gives: $R2 = \frac{F_{p_sagit}}{F_{mw_sagit}} = 0.16$. The meaning is that the Pivot gravitational pull is smaller than the pull between local group Galaxies. Therefore, clusters of galaxies can be formed where Galaxies orbit the center of the cluster.

3) Gravitational pull between Earth and Sun Vs. the Pivot pull:

$M_{earth} = 6 \cdot 10^{24} kg$ Mass of Earth

$M_{sun} = 2 \cdot 10^{30} kg$ Mass of Sun

$R_{earth} = R_{mw} = 129.95 \cdot Gly$ Orbiting radius of Earth around the Pivot

$D_{earth_sun} = 150 \cdot 10^6 \cdot km$ Distance of Earth from Sun

$$F_{p_earth} = \frac{G \cdot M_p \cdot M_{earth}}{R_{mw}^2} \quad (19.5) \quad \dots \text{The Pivot gravitational pull on Earth}$$

$$F_{sun_earth} = \frac{G \cdot M_{sun} \cdot M_{earth}}{D_{earth_sun}^2} \quad (19.6) \quad \dots \text{Gravitational pull between the Sun and Earth}$$

The ratio between (19.5) and (19.6) gives: $R3 = \frac{F_{p_earth}}{F_{sun_earth}} = 6 \cdot 10^{-9}$. The meaning is that the Pivot influence in a solar system is negligible.

Conclusion: It was shown that the Pivot Universe structure can explain the formations of Galaxies, Clusters of Galaxies and solar systems. The formation of the various structures of celestial bodies depends on the masses and distances of the celestial bodies. It was shown in

paragraph 5 that the magnitude of the gravity of the Pivot in the visible Universe is very small. Nevertheless, for Galaxies, the main gravitational pull is caused by the Pivot. The Galaxies in a cluster may orbit around the center of the cluster, but the gravitational pull of the Pivot must also be taken into consideration. On the other hand, for a solar system the main gravitational pull is between the star and its planets and between the planets themselves. The influence of the Pivot in solar system is negligible. For example, the trajectory of a man-made satellite can be calculated quite accurately when considering only the gravitational pull on the satellite by the Sun and the planets.

20. Mach's principle Vs. Newton's bucket

There is an ongoing profound debate in the scientific community relating to the issue whether the Universe is absolute or relative.

Newton conducted an experiment with a bucket containing water. He filled a bucket with water and suspend it from a fixed point with a rope. He rotated the bucket, twisting the rope more and more. When the rope has taken all the twisting that it can take he released the bucket. What happened? The bucket started to rotate because of the twisted rope. At first, the water in the bucket did not rotate with the bucket but remained fairly stationary. Slowly, however, the water began to rotate with the bucket and as it did so the surface of the water become concave. At that stage, the water and vessel rotated together, and there was no relative motion between them. Yet somehow the water "knows" to create a concave surface. Newton concluded from the bucket experiment that the concavity must be due to the water's rotation with respect to something else—absolute space. In other words, the rotation is absolute, not relative. That answer stood largely unchallenged for two centuries until Mach flatly declared Newton to be wrong.

Mach never gave a precise formulation of what became known as Mach's Principle. He claimed that Inertia—that is the tendency of massive objects to move at constant velocity—must depend on other bodies because motion itself must be measured relative to other bodies. In other words, local inertial frames are determined by the large-scale distribution of matter, as exemplified by this anecdote: You are standing in a field looking at the stars. Your arms are resting freely at your side, and you see that the distant stars are not moving. Now start spinning. The stars are whirling around you and your arms are pulled away from your body. Why should your arms be pulled away when the stars are whirling? Why should they be dangling freely when the stars don't move?

The Pivot theory claims that Newton is correct. The absolute rotating frame of the Universe is the Pivot. On the other hand, the Pivot theory can explain the anecdote given above. You, as well all the celestial bodies in the Universe are orbiting the Pivot at approximately the same orbital speed. During the time you spin (even if you spin for millions of years) celestial bodies, because of the huge radius of the Universe, are hardly moving relative to each other. So, for you, the distant stars and Galaxies are not moving. When you spin around yourself the distant stars seem to whirl around you.

21. Summary

The Pivot theory describes a Universe that began as a primeval hadron. This hadron can be modeled as a KNBH. It ended in the current Universe that is composed of a spinning Kerr black hole, the Pivot, and a ring of visible Universe orbiting it in an opposite direction to the Pivot's spin. The Pivot theory describes a Universe that is in a state of dynamic equilibrium, indicating that it will exist forever. Everything in the visible Universe may change, stars will consume their energy, Galaxies orbiting at too close orbits will eventually collide etc. The Pivot will last forever because there is no other object in the Universe that can threaten its existence. The Pivot Universe structure explains many known cosmic observations. It addresses the issue of the dark matter and suggests that it resides in the Pivot and therefore, most probably, will never be observed. The dark energy theory is not required in this model.

The issues of the dynamic process that started with the primeval hadron and ended with the current Universe are not addressed here in a satisfactory manner, for example: 1) What is the mechanism that caused the hadron to explode when its mass reached the total mass of the current Universe? 2) How long did it take the primeval hadron to reach this mass?

Acknowledgment

I would like to thank my wife, Talia and my daughter, Liad, for reviewing this article.

22. References

1. R. M. Muradian (1980) – “The primeval hadron: origin of stars, Galaxies and astronomical Universe” http://articles.adsabs.harvard.edu/cgi-bin/nph-iarticle_query?bibcode=1980Ap%26SS..69..339M&db_key=AST&page_ind=0&plate_select=NO&data_type=GIF&type=SCREEN_GIF&classic=YES
2. R. M. Muradian (1987) - “The primeval hadron: origin of rotation and magnetic fields in the Universe” <http://adsabs.harvard.edu/full/1987IAUS..121..341M>
3. R. L. Oldershaw (2006) - “Hadrons as Kerr-Newman Black Holes” <https://arxiv.org/ftp/astro-ph/papers/0701/0701006.pdf>
4. S. Hod (2016) – “Analytic treatment of the system of a Kerr-Newman black hole and a charged massive scalar field” <https://arxiv.org/abs/1609.07146>
5. M. J. Longo (2007) - “Evidence for a Preferred Handedness of Spiral Galaxies” <https://arxiv.org/ftp/arxiv/papers/0707/0707.3793.pdf>
6. P. Birch (1982) – “Is the Universe rotating?” <http://buildengineer.com/www.paulbirch.net/IsTheUniverseRotating.pdf>
7. Wikipedia - https://en.wikipedia.org/wiki/Observable_Universe#cite_ref-15
8. Wikipedia – https://en.wikipedia.org/wiki/Dark_matter
9. R.F. Sistero (1982) “Roation of the Universe and the Universal Angular Momemtum-Mass Relationship” http://articles.adsabs.harvard.edu/cgi-bin/nph-iarticle_query?bibcode=1983ApL....23..235S&db_key=AST&page_ind=0&plate_select=NO&data_type=GIF&type=SCREEN_GIF&classic=YES

10. Y. Sofue (1999) – “Central Rotation Curves of Spiral Galaxies”
arXiv:astro-ph/9905056
11. T.M. Davis & C.H. Lineweaver (2004) “Expanding Confusion: Common Misconception of Cosmological Horizons and the Superluminal Expansion of the Universe”.
<http://www.publish.csiro.au/AS/pdf/AS03040>
12. R.P. Kirshner (2003) “Hubble's diagram and cosmic expansion”.
<http://www.pnas.org/content/101/1/8.long>
13. B. Nodland & J.P. Ralston (1997) “Indication of Anisotropy in Electromagnetic Propagation over Cosmological Distances” <https://arxiv.org/pdf/astro-ph/9704196.pdf>
14. Andromeda Galaxy https://en.wikipedia.org/wiki/Andromeda_Galaxy