

# The structure of the Pivot Universe

Arieh Sher

email: [pivot.universe.theory@gmail.com](mailto:pivot.universe.theory@gmail.com)

## Abstract

The Pivot theory describes the structure of the Universe. It postulates that the Universe is composed of a massive slowly spinning blackhole, the Pivot, and a ring-shaped visible Universe orbiting it. This combined structure, namely a blackhole and an orbiting ring, is an established phenomenon in general relativity. In general, our Universe is a finite island fixed in the infinite space. The Pivot can also be described from the quantum physics point of view. From this view, the Pivot is a Neutron star, composed of nucleons that are packed to the maximum density possible in the Universe.

The Pivot theory postulates that the Universe started as a primeval nucleus that was a spinning Neutron star. This Neutron star accumulated energy from the vacuum, until the velocity on its equator surface reached the speed of light and then exploded. It exploded into two distinct parts: The Pivot and the ring of the visible Universe. Using these assumptions, and relying on: known physical constants, cosmological observations, and conservation laws, the size of the Universe can be calculated.

Finally, the theory is verified against known cosmological observations, such as the flat rotation curve in Spiral Galaxies, the Spiral shape of Galaxies, Redshift of Galaxies, and other observations.

Keywords: Universe structure, frame dragging, Blackhole, Neutron star, QM, GR, gravitational constant, gravitational z shift

## **Contents**

- 1.** The Pivot theory- Overview
- 2.** Known constants and accepted parameters
- 3.** The Primeval Universe
- 4.** The Pivot Universe
- 5.** Gravitational z shift
- 6.** The Origin of spinning and rotation of celestial bodies
- 7.** Rotation curve and spiral shape of a Galaxy
- 8.** Hubble's observations
- 9.** Cosmic Microwave Background (CMB)
- 10.** Olbers' paradox
- 11.** Handedness of Galaxies
- 12.** Historical notes
- 13.** Summary
- 14.** References
  - Appendix A - The origin of gravity
  - Appendix B - Is a blackhole a neutron star?

## 1. The Pivot theory - Overview

The Pivot theory postulates that the current Universe is composed of a massive and slowly spinning Neutron star, the Pivot, located at its center. Orbiting the Pivot is a ring-shaped visible Universe. The Pivot can describe by two theories: on the one hand, quantum physics where it is described as a nucleus that is composed of an enormous number of nucleons that are packed in this nucleus to the maximum density possible. On the other hand, the Pivot can be described by GR as a huge slowly spinning Kerr blackhole, that due to its spin, it drags the space around it. The Pivot theory answers three questions:

The first question is: How did it all begin?

This part of the theory is speculative and is not adequately addressed; however, it is not out of the blue. The theory is based on quantum physics. Space, or vacuum, permeates infinitely in all directions. I claim that the origin of the matter Universe is the vacuum of the space. Quantum theory describes all elementary particles as vibrational modes in fundamental fields that exist at all points of space and time. In this vacuum, virtual particles are constantly popping in and out of existence. These virtual particles are coming in pairs, e.g., a quark and antiquark, exist for an extremely short time, and then mutually annihilate. Quantum physics teaches that it is possible to boost the pair apart using external energy (e.g., a very strong electric field – that pushes them in opposite directions because they have opposite electrical charges) so that they avoid annihilation and become actual (long-lived) particles. It all started with the creation of one hadron somewhere in space. From QCD it is known that if additional external energy (from the vacuum) was supplied to this primeval hadron, it did not dissolve to its constituents, but instead, caused the number of gluons and quarks inside the hadron grew considerably. Eventually, new hadrons were added to the first hadron. It is important to note that although the behavior of the hadron described above is not yet fully explained theoretically by quantum physics, it was verified by a large number of accelerator experiments over the last 50 years. Following that, it happened that quarks and gluons that were boosted by energy from the vacuum attracted each other to form a primeval nucleus. Speculative note: It is possible that according to the above description, additional Universes exist in the infinite space.

The model includes several assumptions:

- 1) The primeval nucleus started to spin from the beginning, and the new nucleons that accumulated around it continued to spin in the same direction.
- 2) The spinning of the nucleus that eventually became very massive, and according to GR, dragged space around it.
- 3) The nucleons were added to the primeval nucleus in such a way that they were packed to the maximum density possible in the Universe. I postulate that there is a maximum density possible in the Universe. It is approximately the density of the nucleus of an Atom or a Neutron star. At maximum density, the nucleons are so close to each other that they start to repel each other. QCD teaches that bringing the nucleons too close requires infinite force.

4) An additional assumption relates to the origin of gravity. During the epoch of the primeval Universe only the primeval nucleus existed. The only sources for holding the nucleus together were the nucleons themselves. Quantum physics teaches that there is a very strong attractive force between nucleons that are located at a sub-atomic distance. This strong force was the one that kept the nucleons together in the primeval nucleus. My claim is that this strong force between nucleons is the origin of gravity between celestial bodies. At cosmological distances the strong force (= gravity) decreases to extremely small values, however, as the number of nucleons in celestial bodies is enormous, there is still considerable gravity force between celestial bodies. Appendix A elaborates on this topic.

The second question is: Why did the primeval nucleus stop growing?

I postulate that the size of the primeval nucleus, or the Neutron star, grew until the tangential velocity on its equatorial surface reached the velocity of light and then it exploded. It is possible that at the moment of the explosion the Neutron star had an oblate spheroid shape rather than a sphere. Based on the Big Bang theory, this explosion occurred 13.7 billion years ago. What is not clear to me is how long did it take for the primeval nucleus to reach its maximum size. An additional note is that the explosion raised the temperature of the nucleus to very high temperatures.

The third question is: How has the Universe evolved into the Pivot structure?

The explosion shattered the outer layers of the primeval nucleus, specifically those regions that were near the equatorial plane. It caused the nucleons of these layers to flung off tangentially in the equatorial plane of the primeval nucleus and the same direction of the primeval nucleus spin. The final result was as follows: The more significant part of the primeval nucleus became the Pivot. The other nucleons that flung off tangentially from the primeval nucleus arranged around the Pivot in a ring-shaped visible Universe. This combined structure, namely a blackhole and an orbiting ring, is an established phenomenon in general relativity. One characteristic of the combined blackhole-ring structure is that there is a stable region around the blackhole where the orbital velocity of a particle equals the angular velocity of the dragged space. An observer in this stable region will orbit the blackhole but will be stationary in regard to the dragged space.

The center point of the Pivot is fixed in the space. In other words, the Universe has an absolute reference frame. From GR point of view, the Pivot is a Kerr Blackhole. Its existence is possible according to the Kerr solution because it is located inside the singularity ring. The inner radius of the ring-shaped visible Universe must be bigger than the Schwarzschild radius of the Pivot. Simultaneously, as is known from the Kerr solution, the space around the Pivot is dragged by the Pivot in the same direction of Pivot's spin. An important assumption in the calculations is that the angular speed of space dragged by the Pivot at the radius of the Milky Way is equal to the angular speed of the visible Universe measured by Birch. This fact explains the Michelson-Morley experiment. An observer located in the ring will not be able to measure his speed relative to the space that surrounds him because there is no relative motion between him and space. One

can sum up that Newton was correct by arguing that there is a fixed center of the Universe but to explain additional observations Einstein's GR must be taken into consideration.

After the explosion of the primeval nucleus, the visible Universe contained only very hot nucleons that were orbiting the Pivot. It took the visible Universe 380,000 years to cool down. When this happened, ordinary atoms were formed. The local density of the visible Universe was the cause of the variety of celestial bodies. If the density of atoms at a particular region in the visible Universe was too low to enable attraction between them, they remained as a cloud of gas that orbits the Pivot. If the density of atoms was sufficient for interaction between them, stars were created. The variety of celestial body was dependent on the mass of the born star. Some stars that had enough mass to collapse by gravity into Neutron stars, more massive stars collapsed into blackholes. The blackhole was powerful enough to attract additional stars that started to orbit it. Thus, stars performed two simultaneously trajectories. One around the Pivot and the other around the blackhole of a Galaxy- this was the way that Galaxies were created.

So far, the description of the Pivot Universe was qualitative. The next paragraphs are quantitative. In these paragraphs, I calculate the size of the primeval nucleus and the sizes of the Pivot and the visible Universe ring.

In order to test the theory, the Pivot structure is verified against known cosmological observations: Origin of spinning and rotation of all celestial objects, flattened rotation curve in Spiral Galaxies, Spiral shape of Galaxies, Redshift of Galaxies, Blue shift of Galaxies, Cosmic Microwave Background, Handedness of Galaxies, Olbers' paradox, Hubble's observations.

Finally, there are spin-offs of the Pivot theory. I relate to them in the Appendixes:

Appendix A- The origin of gravity.

It is shown how quantum physics, Newton's gravitational law, and GR can be unified.

Appendix B- Is a blackhole a neutron star?

I claim that there is one difference between the Pivot and the blackholes in the visible Universe. The accumulation of nucleons in the primeval Universe built the Pivot whereas a blackhole in the visible Universe ring is the result of the gravitational collapse of a massive star. However, the final result is the same. A nucleus in which nucleons are packed to the maximum allowed density in the Universe.

## 2. Known constants and accepted parameters

### Definitions:

$Gly = 9.454 \cdot 10^{24} \text{ m}$  ...is the distance of billion light years.

$Mly = 9.454 \cdot 10^{21} \text{ m}$  ...is the distance of million light years.

$Kly = 9.454 \cdot 10^{18} \text{ m}$  ...is the distance of thousand light years.

### Accepted parameters:

$G = 6.67 \cdot 10^{-11} \text{ m}^3 / \text{kg} / \text{sec}^2$  ...is the Gravitational constant.

$C = 2.9979 \cdot 10^8 \text{ m} / \text{sec}$  ...is light velocity.

$\rho_{ring} = 10^{-26} \text{ kg} / \text{m}^3$  ...is the density of ordinary matter in Universe.

$\hbar = 1.054571 \cdot 10^{-34} \text{ J} \cdot \text{sec}$  ...is the reduced Planck's constant.

$m_{neutron} = 1.674927471 \cdot 10^{-27} \text{ kg}$  ...is the mass of neutron (= ~ mass of proton).

$R_{neutron} = 0.8 \cdot 10^{-13} \text{ cm}$  ... is the radius of the neutron

### Evaluated and estimated parameters:

Note: Each of the following parameters is the best estimations I found in the literature. However, I adjusted some parameters to fit the model.

$\rho_{max} = 6.5 \cdot 10^{17} \cdot \text{kg} / \text{m}^3$  ... is the maximum density in Universe

Note: One of the main claims of the Pivot theory is that there is a maximum density of matter in the Universe. This density occurs when nucleons (neutrons/protons) of matter are packed so densely, that they cannot be squeezed any more. It is known that Neutron stars and atom nucleus have approximately the same maximum density.

$\omega_{Birch} \approx 10^{-13} \text{ rad} / \text{yr}$  ...is the observed angular velocity of the Universe- Birch [1], as measured from the Milky Way.

Note: Birch calculated from the study of position angles and polarization of high luminosity classical double radio sources a rotational velocity for the Universe of the order of  $\omega_{Birch} \approx 10^{-13} \text{ rad / yr}$ . Although his work has not been convincingly refuted, it was criticized for using improper statistics. In the following calculations, I assume  $\omega_{Birch} = 1.05 \cdot 10^{-13} \text{ rad / yr}$ .

$$M_{ring} = 1.45 \cdot 10^{53} \text{ kg} \quad \dots \text{is the mass of visible Universe ring, Wikipedia [2].}$$

Note: There are various methods for calculating the mass of the universe. It ranges anywhere from  $10^{53} \text{ kg}$  to  $10^{60} \text{ kg}$ .

$$R = \frac{26.8\%}{4.9\%} = 5.47 \dots \text{ is the ratio of "dark matter" mass to visible Universe mass, Wikipedia [3].}$$

Note: I concur with other theories that claim that adding "dark matter" to the Universe solves the issue of curve flattening of Galaxies. However, I claim that this "dark matter" resides in the Pivot rather than being spread in and around Galaxies as postulated by current cosmology. In the calculations, I am using the mass of the "dark matter" calculated by others, as the mass  $M_{pivot}$  of the Pivot.

$$M_{pivot} = R \cdot M_{ring} = 7.93 \cdot 10^{53} \text{ kg} \quad \dots \text{is the calculated mass of Pivot.}$$

$$M_{total} = M_{pivot} + M_{ring} = 9.38 \cdot 10^{53} \text{ kg} \quad \dots \text{is the calculated total mass of the Universe.}$$

### 3. The Primeval Universe

The primeval Universe consisted of a spinning Neutron star that dragged space around it. The Pivot theory uses the primeval hadron theory described by Muradian [4] and others suggesting that the Universe began as a super dense primeval hadron. He uses the generalized Regge's law general formula which connects the maximal spin  $J$  and the mass  $m$  of a celestial object:

$$J = \hbar \cdot \left[ \frac{m}{m_{proton}} \right]^{1+1/n} \quad \text{where for Galaxies, clusters of Galaxies and globular Galaxies the two-}$$

dimensional disk-like hadron  $n=2$ . For stars, planets and asteroids  $n=3$ . He showed that the calculated values of  $J$  and  $m$  of all celestial bodies, namely superclusters, Galaxies, stars, and planets, are in good agreement with cosmological observations. I am using this equation for calculating the angular momentum of the primeval Universe. Note: I am replacing the mass of the proton ( $m_{proton}$ ) in Muradian's equation by the mass of a neutron ( $m_{neutron}$ ), because I postulate that the primeval Universe was a Neutron star, composed mainly from Neutrons.

$$J_{total} = \hbar \cdot \left( \frac{M_{total}}{m_{neutron}} \right)^{\frac{3}{2}} \quad (3.1)$$

At the moment of explosion there were regions of the primeval Neutron star that were moving at velocities near the speed of light, therefore  $M_{total}$  must take into consideration relativistic effect.

In order to find  $J_{total}$  the mass  $M_{total}$  is replaced by the relativistic mass  $M_{rel\_total}$  of the Neutron star.

Finding  $J_{total}$  is done in the following steps:

1. Finding the radius of the primeval Neutron star. Assuming that the nucleons are densely packed in a solid sphere, the radius of this primeval nucleus ( $R_{nucleus}$ ) can be calculated:

$$N_{nucleons} = \frac{M_{total}}{m_{neutron}} = 5.6 \cdot 10^{80} \quad (3.2)$$

$$R_{nucleus} = R_{neutron} \cdot (N_{nucleons})^{\frac{1}{3}} = 6.594 \cdot 10^8 \text{ km} \quad (3.3)$$

2. Finding the total relativistic mass of the primeval Neutron star  $R_{rel\_total}$ . (Note: In this equation, it is assumed, for simplification, that the primeval Neutron star is a solid sphere. However, it is most likely that at the moment of the explosion the Neutron star had an oblate spheroid shape). The relativistic mass change is related to the radius of the region.



$$M_{rel\_total} = 4 \cdot \pi \cdot \rho_{max} \cdot \int_0^{R_{nucleus}} \frac{r^2}{[1 - (\frac{r}{R_{nucleus}})^2]^{0.5}} \cdot dr = 1.839 \cdot 10^{54} \cdot kg \quad (3.4)$$

3. The total angular momentum of the primeval Universe at the moment of explosion is:

$$J_{total} = \hbar \cdot \left( \frac{M_{rel\_total}}{m_{neutron}} \right)^{\frac{3}{2}} = 3.836 \cdot 10^{87} \cdot J \cdot s \quad (3.5)$$

This value of  $J_{total}$  will be used later when evaluating the shape and size of the current Pivot Universe.

In the following paragraph, the ratio of the primeval Universe angular momentum to the angular momentum of the Neutron star is calculated. It will be shown that the angular momentum of the Neutron star is negligible when compared to the angular momentum of the dragged space.

The primeval Universe angular momentum is the sum of the angular momentum of the primeval solid sphere Neutron star, plus the angular momentum of space that is dragged by this primeval Neutron star. Dragging of space by a spinning celestial body is a result of GR. The ratio of the angular momentum of the primeval Neutron star  $J_{nucleus}$  and the total angular momentum  $J_{total}$  is calculated as follows:

1. The explosion of the primeval Neutron star occurred when the tangential velocity at the equator of the Neutron star was equal to the speed of light  $C$ . The angular velocity of the nucleus was at that moment:

$$\Omega_{nucleons} = \frac{C}{R_{nucleus}} = 1.431 \cdot 10^4 \cdot rad / yr \quad (3.6)$$

2. The primeval nucleus is a solid sphere of Neutrons. The moment of inertia of this sphere, while taking into considerations relativistic effect, is:

$$I_{nucleus} = 4 \cdot \pi \cdot \rho_{max} \cdot \int_0^{R_{nucleus}} \frac{r^4}{[1 - (\frac{\Omega_{neucleus} \cdot r}{C^2})^2]^{0.5}} \cdot dr = 1.2 \cdot 10^{78} \cdot kg \cdot m^2 \quad (3.7)$$

And then the angular momentum of the nucleus is calculated:

$$J_{nucleus} = I_{nucleus} \cdot \Omega_{neucleus} = 5.455 \cdot 10^{74} \cdot J \cdot s \quad (3.8)$$

3. The ratio of  $\frac{J_{total}}{J_{nucleus}} = 7.03 \cdot 10^{12}$  shows that almost the entire angular momentum of the primeval Universe was due to the frame dragging of space. It can be claimed that the viscous forces are  $\gg$  than the inertial forces.

Speculative Note: The above description of a spinning Neutron sphere in an unbound space resembles a known problem from the field of fluid dynamics. In Stokes flow, when the inertial forces are negligible compared to the viscous forces, i.e., Reynolds number is very small, the solution yields a steady state rotation of the sphere together with the fluid around it. Therefore, Stokes flow solution on the surface of a sphere is also applicable to the Neutron star spinning in space.

Solving Stokes flow on the surface of the spinning primeval Neutron star gives the torque  $\tau$  :

$$\tau = 8 \cdot \pi \cdot \mu_{space} \cdot R_{nucleus}^3 \cdot \Omega_{nucleus} \dots \text{where } \mu \text{ is the viscosity of the space (3.9).}$$

On the other hand,  $\tau = J_{total} \cdot \Omega_{nucleus}$  where the dragged space on the Neutron star surface is also spinning at  $\Omega_{nucleus}$ . Therefore, the viscosity of space  $\mu_{space}$  on the surface of the primeval Neutron star can be calculated by:

$$\mu_{space} = \frac{J_{total}}{8 \cdot \mu \cdot R_{nucleus}^3} = 3.87 \cdot 10^{50} \cdot Pa \cdot s \quad (3.10)$$

The high viscosity of space on the surface of the Neutron star means that space clings strongly to this surface. I speculate that, unlike Stokes flow where the viscosity is constant everywhere in the fluid, the viscosity of space is dependent on gravity. It is maximal near a celestial body and drops substantially far away from this celestial body. I also speculate that the viscosity of space is the parameter that limits the speed of light in space. But, as space viscosity changes it is possible that the speed of light is not constant.

#### 4. The Pivot Universe

After the explosion of the primeval Universe, the Pivot Universe rearranged. An open issue is how long did it take the Pivot Universe to reorganize. In this paragraph, the sizes of the Pivot Universe are calculated. Calculating the dimensions of the Pivot Universe is based on conservation laws of angular momentum and mass. Namely, the angular momentum of the Pivot Universe must be equal to the angular momentum of the primeval Universe  $J_{total}$  that was calculated in (3.5). The angular momentum of the Pivot Universe is the sum of the angular momentum of the Pivot + the visible Universe + the dragged space. The mass of the Pivot + the mass of the visible Universe must be equal to the mass of the primeval nucleus. The calculation of the dimensions of the Pivot Universe is done in the following steps:

1. Schwarzschild radius of the Pivot is:

$$R_H = \frac{2 \cdot G \cdot M_{pivot}}{C} = 124.424 Gly \quad (4.1)$$

2. Main dimensions of the visible Universe ring are:

##### Note:

There are several assumptions and adjustment of data I did to find the dimensions of the Universe:

- 2.1 The Pivot Universe structure is similar to the known GR structure of a black hole surrounded by a thin orbiting ring. In GR there is a term called Innermost stable circular orbit (ISCO) which marks the inner edge of the ring orbiting the black hole. According to GR, a particle can stably stay in orbit around a massive object if its orbiting radius is greater than ISCO. The composed blackhole-ring system is an on-going research field in GR, and several papers relate to the calculation of the radius of ISCO. I tried to use Kerr ISCO calculator by Stein that is based on an article from 1972 by Bardeen & al. <https://duetosymmetry.com/tool/kerr-calculator-v2/> However, the results are not in agreement with my model. Therefore, I assumed a relation between the inner radius of the visible Universe ring and the Schwarzschild radius. The inner radius must be greater than the Schwarzschild radius.

$$R_m = 1.01 \cdot R_H = 125.668 \cdot Gly \quad (4.2)$$

- 2.2 The total angular momentum  $J_{total}$  of the primeval Universe (see 3.5) is divided equally between the Pivot and the ring. Namely, the angular momentum of the visible Universe

ring  $J_{ring}$  is equal to the angular momentum of the Pivot + the space dragged by the Pivot  
 $J_{pivot}$  :

$$J_{ring} = J_{pivot} = 0.5 \cdot J_{total} = 1.918 \cdot 10^{87} \cdot J \cdot s \quad (4.3)$$

2.3 The dimensions of the ring were adapted so that the radius of the Milky Way  $R_{mw}$  is in agreement with the cosmological observations of the z shift of galaxies in the Universe (see paragraph 5)

The outer radius of the visible universe  $R_{out}$  is found by using the formula of a ring's angular momentum:

$$\frac{M_{ring}}{2} \cdot (R_{in}^2 + R_{out}^2) \cdot \omega_{Birch} = J_{ring} \dots \Rightarrow R_{out} = 270.29 Gly \quad (4.4)$$

The width of the visible Universe  $W$  is found by:

$$M_{ring} = W \cdot \rho_{ring} \cdot \pi \cdot (R_{out}^2 - R_{in}^2) \dots \Rightarrow W = 95 Mly \quad (4.5)$$

### 3. Dimensions of the Pivot:

Radius of the Pivot is:

$$R_{pivot} = R_{neutron} \cdot (N_{pivot})^{\frac{1}{3}} \quad (4.6)$$

Where:

$R_{pivot}$  ... is the radius of the Pivot

$R_{neutron} = 0.8 \cdot 10^{-13} cm$  ... is the radius of the Neutron

$N_{pivot} = \frac{M_{pivot}}{m_{neutron}} = 4.734 \cdot 10^{80}$  ... is the number of Neutrons in the Pivot

$R_{pivot} = R_{neutron} \cdot (N_{pivot})^{\frac{1}{3}} = 6.235 \cdot 10^8 km$  ... is the radius of the Pivot

The spin and the tangential velocity of the Pivot:

This calculation is based on GR frame dragging of Kerr blackhole in the plane of the equator. See Wikipedia [5]. <https://en.wikipedia.org/wiki/Frame-dragging>).

$$\Omega(r) = \frac{R_H \cdot \alpha \cdot C}{r^3 + \alpha^2 \cdot r + R_H \cdot \alpha^2} \quad (4.7)$$

Where:  $\alpha = \frac{J_{total}}{M_{pivot} \cdot C} = 1.614 \cdot 10^{25} \cdot m = 1.7 Gly$

The graph of  $\Omega(r)$  according to (4.7) is shown in Fig. 4.1




Fig. 4.1 – Frame dragging caused by the Pivot

The angular velocity of the Pivot:

$$\Omega(R_{pivot}) = 5.86 \cdot 10^{-10} \text{ rad / yr}$$

The tangential velocity on the surface of the Pivot is:

$$V_{pivot} = \Omega(R_{pivot}) \cdot R_{pivot} = 0.0116 \cdot mm / sec \quad (4.8)$$

Conclusion: In GR terms the Pivot is a very slow Kerr blackhole.

#### 4. The radius of the Milky Way

In order to find the orbiting radius of the Milky Way  $R_{mw}$ , the measured  $\omega_{Birch}$  is equated to the angular velocity  $\Omega(r)$  of dragged space.

$$\omega_{Birch} = \frac{R_H \cdot \alpha \cdot C}{R_{mw}^3 + \alpha^2 \cdot R_{mw} + R_H \cdot \alpha^2} \dots \Rightarrow R_{mw} = 125.67 Gly \quad (4.9)$$

#### 5. Newton Vs. Birch

Having found the dimensions of the Pivot Universe, one can compare the velocity of the Milky Way orbiting the Pivot at a radius of  $R_{mw}$  according to Newton's gravitational law in comparison the velocity derived from Birch's observation:

$$V_{Newton} = \left( \frac{G \cdot M_{pivot}}{R_{mw}} \right)^{0.5} \cong 0.701C \quad (4.10)$$

The velocity of a celestial body orbiting the Pivot at a radius of  $R_{mw}$  can be also calculated by:

$$V_{Birch} = \omega_{Birch} \cdot R_{mw} = 0.013C \quad (4.11)$$

There is a discrepancy between the two calculations. The ratio is:

$$R_{ratio} = \frac{V_{Newton}}{V_{Birch}} = 52.84 \quad (4.12)$$

GR can explain this discrepancy. GR teaches that space is wrapped around a celestial body. Newton was not aware that all celestial bodies spin and while spinning drag and wrap the space around them. I postulate that the distance between two bodies that is a straight line between their centers according to Newton's gravitational law should be replaced by the length of the geodesic formed by the dragged space between the two bodies.

In order to equate the velocity of  $V_{Newton}$  to  $V_{Birch}$  the length of the curve  $L$  should be:

$$L = \frac{G \cdot M_p}{V_{Birch}^2} = 3.5 \cdot 10^5 \cdot Gly \quad \text{And the ratio } \frac{L}{R_{mw}} = 2.79 \cdot 10^3 \quad (4.13)$$

The angular displacement  $\theta(r)$  of the dragged space is calculated:

$$\theta(r) = \Omega(r) \cdot r \cdot t \quad (4.14)$$

Where:  $\Omega(r)$  is calculated according to Eq. (4.7) and  $t$  is the time required for the system to be in a steady state. The time  $t$  is unknown, and it will be calculated so Eq. (4.13) is satisfied.

The length of the geodesic in curved space is:

$$L = \int_{0-m}^{R_{out}} (1 + (\frac{d\theta}{dr})^2)^{0.5} dr = 2.79 \cdot 10^3 \cdot R_{mw} \quad (4.15)$$

Equating Eq. (4.14) to Eq. (4.15) results in:  $t = 8.12 \cdot 10^{13} \text{ yr}$ . Speculation: Does this result mean that the Pivot Universe has not yet reached a steady state?

Another interesting question arises from Fig. 4.2. It shows the velocity of the dragged space  $V(r) = \Omega(r) \cdot r$ . In the range of  $r = 1.7 \text{ Gly} \leftrightarrow 13.3 \text{ Gly}$  the velocity of the dragged space is higher than  $C$ , reaching a peak of  $2.1C$ . It is to be noted that the lower number of the range, i.e.,  $1.7 \text{ Gly}$ , is also the value of  $\alpha$  calculated by Eq. (4.7).

Speculation: This range where the velocity of dragged space is higher than  $C$  represent the singularity ring that is derived from the Kerr solution. Material that crosses the event horizon of the Pivot, will not reach the Pivot but instead will turn into energy in the singularity ring.




Fig. 4.2 – Velocity of dragged space around the Pivot

Notes:

- a) The calculated radius of the Pivot is  $6.23 \cdot 10^8 \text{ km}$ . For comparison only, the distance of Jupiter from the Sun is  $7.8 \cdot 10^8 \text{ km}$ . The size of the Pivot is impressive, but it is dwarfed when compared to the radius of the visible Universe ring. This ratio is:

$$Ratio = \frac{R_{pivot}}{R_{out}} = 2.4 \cdot 10^{-16}$$

b) It is interesting to calculate the surface gravity of the Pivot:

$$g_{pivot} = \frac{G \cdot M_{pivot}}{R_{pivot}^2} = 1.2 \cdot 10^{20} \cdot m / sec^2 \quad (4.10).$$

I claim that this calculated number is the maximal gravity in the Universe. I compare this result to the maximal gravity in the Universe estimated by Potzel [6]. In his article, he describes a theory and uses experimental data for calculating the maximum acceleration in the Universe. He predicts that the maximal acceleration in the universe is  $> 1.5 \cdot 10^{21} \cdot m / sec^2$ .

c) While calculating the angular momentum of the visible Universe ring, the spinning angular momentum of all celestial bodies, i.e., Galaxies, stars, planets, interstellar planets should have been included. However, it is shown now, that the sum of the spinning angular momentum of all these celestial bodies is negligible in comparison to the orbital angular momentum of the ring-shaped visible Universe. Based on Muradian [4], an estimated total angular momentum of all celestial objects is:

$$J_{objects} = J_{gal} \cdot N_{gal} + J_{star} \cdot N_{star} + J_{planet} \cdot N_{planet} = 2 \cdot 10^{85} \text{ erg} \cdot \text{sec}$$

Where:

The average angular momentum of a Galaxy:  $J_{gal} = 10^{74} \text{ erg} \cdot \text{sec}$ , Number of Galaxies:

$N_{gal} = 2 \cdot 10^{11}$ , Angular momentum of an average star:  $J_{star} = 10^{49} \text{ erg} \cdot \text{sec}$ , Number

solar systems:  $N_{star} = 10^{22}$ , Angular momentum of an average planet:  $J_{planet} = 10^{40} \text{ erg} \cdot \text{sec}$ ,

Number planets:  $N_{planet} = 10^{24}$ .  $J_{objects} = 2 \cdot 10^{85} \text{ erg} \cdot \text{sec} \ll J_{ring} = 1.91 \cdot 10^{94} \text{ erg} \cdot \text{sec}$  and therefore, can be neglected.

d) GR solution gives also rise to a photon sphere. The photon sphere prevents photons to leave the Pivot Universe. The radius of the photon sphere is an opened question and is assumed.

Based on the description and the calculations above, Figure 4.1 depicts the structure of Pivot Universe. The Pivot is a Neutron star is fixed in the infinite space but spins around this fixed point very slowly. The visible Universe ring is located outside the event horizon and rotates in the same direction of the Pivot. The Pivot cannot be observed by an observer from the visible Universe ring. In the figure, the dragging curves of space by the Pivot are schematically shown. The ring location around the Pivot is such that the orbital velocity of a celestial body in this ring is equal to the angular velocity of the space dragged by the Pivot. An observer from this celestial body cannot measure his velocity relative to the space. This phenomenon was proven in the Michelson-Morley experiment. Also, finally, from the GR solution, the Universe has a photon sphere.

To sum up, the Pivot and the Universe ring are in a dynamic equilibrium and can exist forever. On the other hand, celestial bodies located in visible Universe ring can change. Stars will consume their energy; galaxies may collide and so on.






Fig. 4.1- Structure and dimensions of the Pivot Universe

### 5. Gravitational z shift

The Pivot theory postulates that the gravitational field of the Pivot causes the z shift of Galaxies. GR states that electromagnetic radiation originating from a Galaxy orbiting the Pivot is reduced in frequency, or redshifted when observed from a Galaxy that is closer to the Pivot. But, since the Pivot theory claims that Galaxies are orbiting the Pivot at different velocities, the Doppler shift cannot be excluded. However, it will be shown later (see paragraph 8) that the Doppler redshift contribution to the measured z shift is small.




Fig. 5.1 – The Pivot Universe structure shows Galaxies orbiting the Pivot

Fig. 5.1 shows that Galaxies observed from the Milky Way can be either red shifted or blue shifted. For Galaxies that are closer to the Pivot than the Milky Way, e.g., Andromeda galaxy,  $Z_{gal} < 0$ . Galaxies that are orbiting the Pivot at a radius larger than the Milky Way radius are red shifted  $Z_{gal} > 0$ . Referring to Fig. 5.1, GAL-A and GAL-B are shown to have different distances from the Milky Way. Nevertheless, they have the same z shift, because both Galaxies are located on the same orbit. Had the Milky Way been located on the outer radius of the Universe ring, all Galaxies would have been blue shifted. It was shown in Eq. (4.9) that the Milky Way orbiting radius  $R_{mw}$  is closer to  $R_{in}$  than most of the Galaxies, therefore from the Milky Way most of the Galaxies will be redshifted.

The gravitational z shift of a Galaxy  $Z_{gal}$  orbiting the Pivot at a radius  $R_{gal}$ , is calculated by GR Eq. (5.1) and is shown in Fig. 5.2:

$$Z_{gal}(R_{gal}) = \frac{1}{\left(1 - \frac{2 \cdot G \cdot M_p}{R_{gal} \cdot C^2}\right)^{0.5}} - 1 \quad (5.1)$$




Fig 5.2 – Gravitational z shift caused by the Pivot

The z shift of the Milky Way is calculated by:

$$Z_{mw} = \frac{1}{\left(1 - \frac{2 \cdot G \cdot M_{pivot}}{R_{mw} \cdot C^2}\right)^{0.5}} - 1 = 6.95 \quad (5.2)$$

Now, the orbiting radius  $R_{gal}$  of any Galaxy can be found by the measurement of z shift as seen the Milky Way  $z_{gal}$ .

$$R_{gal} = \frac{2 \cdot G \cdot M_{pivot}}{\left(1 - \frac{1}{(z_{mw} - z_{gal} + 1)^2}\right) C^2} \quad (5.3)$$

Two examples of finding the orbiting radius of Galaxies using Eq. (5.3):

- 1) Andromeda Galaxy measured blue shift: -0.001001 gives that radius of Andromeda around the Pivot is:  $R_{andromeda} = 126.4218 \cdot Gly$ . The Milky Way radius is  $R_{mw} = 126.4223 \cdot Gly$ . This means that Andromeda is closer to Pivot by 510.8Kly than the Milky Way.
- 2) Galaxy GN-z11 is the Galaxy with the height redshift ever measured: 11.09. The radius of GN-z11 is 138.52Gly i.e., 12.1Gly further from the Pivot than the Milky Way.

## 6. The Origin of spinning and rotation of celestial bodies

After the explosion of the primeval nucleus, the visible Universe contained only very hot nucleons that were orbiting the Pivot. It took the visible Universe 380,000 years to cool down. When this happened, ordinary atoms were formed. All the atoms orbited the Pivot because they were dragged by the viscous space and simultaneously spin around their axis in the opposite direction to the Pivot spin. The explanation of this combined movement is not intuitive. The reader is referred to an experiment of Stokes flow that is shown in NSF [7] <https://www.youtube.com/watch?v=QcBpDVzBPMk>, start time: 3:38 min.

The celestial bodies created as a result of atoms attracting each other by gravity retained the spin direction of the atoms. The local density of the visible Universe was the cause of the variety of celestial bodies. If the density of atoms at a particular region in the visible Universe ring was too low to enable significant attraction between them, they remained as a cloud of gas that orbits the Pivot. If the density of atoms was sufficient for interaction between them, stars were created. The variety of celestial body was dependent on the mass of the born star. Some stars that had enough mass to collapse by gravity into neutron stars. Galaxies were formed in the following way: if the mass of the star was big enough it collapsed by gravitation to form a blackhole. Once a blackhole was created it started to accumulate matter and stars from the surrounding space. The blackhole swallowed some of the matter/stars, but other matter/stars began to orbit around it. Fig 6.1 shows trajectories of stars around the Galaxy's blackhole. Stars that are orbiting the Pivot at a bigger radius than the Galaxy's blackhole move slower than the Galaxy's blackhole. Stars that are orbiting the Pivot at a smaller radius than the Galaxy's blackhole move faster than the Galaxy's blackhole. This created a torque on the Galaxy, consequently causing it to

simultaneously spin around its blackhole and orbiting the Pivot in a counter direction of the Pivot's spin. The blackhole in the Galaxy's center played a crucial role in the first stages of the Galaxy evolution. Even though the mass of the blackhole in the center of the Galaxy is relatively huge, it can influence only stars that are orbiting near it. For example, the radius of the Milky Way is 100,000 light years Vs. the Schwarzschild radius of its blackhole that is  $1.3 \cdot 10^{-6}$  light years. But at this stage, additional stars were attracted by the distributed mass of the newborn Galaxy, rather than the gravity caused by the supermassive blackhole.




Fig. 6.1 – Trajectories of stars around a Galaxy blackhole

## 7. Rotation curve and spiral shape of a Galaxy

The rotation curve of a Galaxy is a plot of the orbital speeds of stars or gas in that Galaxy versus their radial distance from that Galaxy's center. (See Fig. 7.2). The rotation curves of Galaxies were measured accurately by Vera Rubin and her team. They verified that the rotation curve was flattened, or in other words, that the velocities of stars orbit at roughly the same speed not depending on their distance from the Galaxy's center. This observation is contrary to Newton mechanics that claims that stars that are located further from the Galaxy center will move slower. The Pivot structure can explain the flattening of the rotation curves.




Fig. 7.1 – Velocities of a star in a Galaxy

The gravitational forces exerted on a star in a Galaxy are caused mainly by:

- Distributed mass of the Galaxy.
- The Blackhole in the center of a Galaxy.
- The Pivot.

Data:

$M_{bh} = 10^{36} \text{ kg}$  ... is the average mass of a blackhole in the center of a Galaxy.

$d = 20 \cdot \text{Kly} = R_{gal} - R_{mw}$  ... is the distance between a Galaxy and the Milky way.

$r = 0 \dots 100 \text{ Kly}$  ... is the distance from the star to the blackhole of the Galaxy

$r_0 = 30 \text{ Kly}$  ... is the characteristic radius of distributed mass of the Galaxy.

$V_{sun\_bh} = 213 \cdot \text{km} / \text{s}$  ... is the velocity of the Sun around Milky Way blackhole

$V_{earth\_sun} = 30 \cdot \text{km} / \text{s}$  ... is the velocity of Earth around the Sun

a) Velocity of a star due to distributed mass of the Galaxy:

$$V_{gal}(r) := \begin{cases} \text{if } 0 \cdot \text{Kly} < r \leq r_0 \\ \left( \frac{G \cdot M_{gal}}{r_0} \right)^{0.5} \cdot \frac{r}{r_0} \\ \text{else} \\ \left( \frac{G \cdot M_{gal}}{r} \right)^{0.5} \end{cases} \quad (7.1)$$

b) The velocity of a star around the Galaxy's Blackhole:

$$V_{bh} = \left( \frac{G \cdot M_{bh}}{r} \right)^{0.5} \quad (7.2)$$

c) The velocity of a star in a Galaxy orbiting the Pivot:

$$V_{p\_star} = \omega_{Birch} \cdot (R_{gal} - r \cdot \cos(\alpha)) \quad (7.3)$$

Summation of the three velocities on the star gives:

$$V_{sum} = V_{p\_star} + (V_{gal} + V_{bh}) \cdot \cos(\alpha) \quad (7.4)$$

Orbital velocity of Milky Way around the Pivot:

$$V_{p\_mw} = \omega_{Birch} \cdot R_{mw} = 0.013C \quad (7.5)$$

The velocity of a star in a Galaxy as seen by an Earth observer is given by:

$$V_{star} = V_{p\_mw} - V_{sum} + (V_{sun\_bh} + V_{earth\_sun}) \quad (7.6)$$

Figure 7.2 shows the velocities of stars in an arbitrary Galaxy as seen from Earth. The red curve is a star velocity in a Galaxy and it is flattened out as  $r$  becomes bigger, according to Rubin's measurements. Most of the curve velocities of stars will be similar to the red curve. In any case, all the curves are confined between the two extreme curves of the graph (the black and the blue). The exact shape of the graph is dependent on  $\alpha$ ,  $d$ ,  $r$ , and the mass of the Galaxy's blackhole.




Fig. 7.2 – Rotation curve for a Galaxy at as seen by Earth observer

### The shape of a spiral Galaxy

Most spiral Galaxies contain a central bulge surrounded by a flat rotating disk of stars. The bulge in the Galaxy center includes a supermassive blackhole. The supermassive blackhole is created from the cloud of dust and stars that are orbiting the Pivot. This blackhole was the kernel of the Galaxy. The size of this blackhole varies, and it is also possible that there are Galaxies without blackholes at their center. The bulge has an ellipsoidal shape because stars that were attracted by the central blackhole in the Galaxy originated not only from stars that were located on the equatorial plane of the Galaxy but also from planes that are “above” or “below” the equatorial plane. A rotating disk that separates itself into arms circle the Galaxy center. The spiral arms are the result of the superposition of the motion of a star around the Pivot and its motion around the supermassive blackhole at the center of the Galaxy. Spiral Galaxies may have more than two arms. A Galaxy may have more than two arms if the Galaxy encountered several dense regions at various times. Each time the Galaxy crosses a dense region a new arm is added to the Galaxy. The width of the arm depends on the size of the dense region.

The angular displacement, during elapsed time  $t$ , of a star orbiting the Galaxy's blackhole is designated ( $\theta$ ) (See Fig. 7.1).  $\theta(r, \alpha) = \int_0^t \frac{V_{star}(r, \alpha)}{r} dt$ , where  $V_{star}(r, \alpha)$  is given in (7.6). This

integration is done only for stars that are located on the main axis (i.e., the axis that connects the Pivot and the Galaxy's blackhole, i.e., angles of 0deg and 180deg – (See Fig. 6.1 and Fig. 7.1). Only at these angles, stars begin to orbit around the Galaxy's center. Fig. 7.3 shows the shape of a spiral shape Galaxy 10 billion years after its creation. Note: The modulo operator “mod” before the integral enables the spiral shape of only the last rotation to be shown. The “mod” operator - returns the remainder on dividing x by y (x modulo y). In the current case y=360deg. The reason for using this operator is that stars in Galaxies have completed many full rotations around the Galaxy's blackhole during its life. For example, the Milky Way makes a full rotation around the blackhole located at its center every ~250 million years. Thus, the Milky Way has completed during its life ~54 full rotation.

It is interesting to show the prediction of the Pivot theory as to what will be the shape of the spiral Galaxy 50 Billion years after its creation. From Fig. 7.4 it can be concluded that the spiral shape of a spiral Galaxy is changing at a slow pace over billions of years.

$$\theta_1(r, \alpha) := \text{mod} \left[ \int_{0\text{yr}}^t \frac{V_{star}[r, (\alpha) \cdot \text{deg}]}{r} dt \cdot \text{deg} \cdot 360\text{deg} \right]$$

$$\theta_2(r, \alpha) := \text{mod} \left[ \int_{0\text{yr}}^t \frac{V_{star}[r, (\alpha) \cdot \text{deg}]}{r} dt \cdot \text{deg} \cdot 360\text{deg} \right] + 180\text{deg}$$




Fig. 7.3 – Shape of a spiral Galaxy at  $t = 10$  Billion years after its creation.




Fig. 7.4 – Shape of a spiral Galaxy at  $t = 50$  Billion years after its creation.



## 8. Hubble's observations

The aim of this paragraph is to show that Hubble's law is wrong. I am elaborating on this issue because Hubble's law is in clear contrast to the Pivot theory that claims the Universe has a rotation axis. I am not disputing the correctness of the observations made by Hubble and others. But, I argue that the laws he derived from these observations are wrong.

Hubble's law is a cornerstone in the Big Bang (BB) theory. Hubble's law consists actually of two laws. The first states that Galaxies are receding from each other at a velocity that is linear to their relative distances  $V = H_0 \cdot d$ . An important conclusion that can be derived from the BB theory is that the Universe "on large-scale" is isotropic and homogenous. The second law claimed by Hubble is  $V = z \cdot C$ , meaning that there is a linear relationship between the receding velocity and the redshift of a Galaxy. Hubble's laws seemed to be correct at his time when measurements were done on Galaxies near the Milky Way. However, when Hubble's law was used for observations of distant Galaxies with high redshifted the graph curved up. This finding was a basis of an additional paradigm: The Universe is expanding forever in all directions, at an ever-accelerating speed. According to Hubble's laws, there are observed Galaxies with  $z > 1.5$  that are receding from the Milky Way at speeds greater than the speed of light, Davis & Lineweaver [8]. The accelerated expanding Universe gives rise to yet another theory claiming that to accelerate the Universe expansion at the observed rates, there must be an additional force. This force is dubbed "dark energy". The mass of this dark energy was calculated to be  $\sim 70\%$  of all the mass in the Universe. The confusion, among the scientific community, is great. Kirshner [9] relates to Hubble's diagram (Fig. 7.1): "Staring at his original Hubble diagram, you can see that there is a handful of nearby Galaxies with blue shifts, and a large scatter of velocities at any given distance. Hubble shrewdly used plausible methods to average the data for Galaxies that are at the same distance to make his result stand out more clearly from the noise. He was fortunate to have data that behaved so well."




Fig. 7.1 – Hubble's original graph (1929)

Hubble's diagram shows that there is a significant scatter of the measurements. Some Galaxies are located at the same distance from the Milky Way but have different velocities. The Pivot theory can explain Hubble's observations. Fig. 6.1 shows two Galaxies GAL-A and GAL-C having the same distance ( $d$ ) from the Milky Way, but are located on different radiuses. According to the Pivot theory, they have different  $z$  shifts. But according to Hubble, they should

have the same redshift and the same velocities. Hubble's law is valid only in a particular case, when the observed Galaxies are located on the main axis (the axis that connects the Milky Way and the Pivot), as is shown, e.g., for GAL-B. In this case,  $\Delta R=d$  and the relation velocity-distance is linear. To sum up: the linear relation in Hubble's law is velocity-distance of the Galaxies, whereas in the Pivot theory the linear relation is between the velocity and the orbiting radiuses of the Galaxies.

### Doppler shift Vs. gravitational shift

Hubble claims that the redshift of Galaxies is due only to the Doppler effect. The Pivot theory claim that although there is a Doppler effect, it is small in comparison to the gravitational shift. (The gravitational shift is calculated in paragraph 6). For finding the Doppler shift of an observed Galaxy, first, the orbiting velocities around the Pivot of the Milky Way (8.1) and the observed Galaxy (8.2) should be found. Then, the velocity difference between the two Galaxies is calculated (8.3). Subsequently, the Doppler shift is calculated (8.4). Finally, the ratio of the Doppler shift to the total z shift is calculated (8.5) and shown in Fig. 8.2. (Note:  $Z_{gal}$  is the gravitational z shift). From this graph, the maximal Doppler effect contribution to the total z shift can reach 2.8% at  $R_{out}$ . However, the maximum redshift measured is of Galaxy GN-z11 located at radius 138.51Gly, the Doppler contribution is less than 0.2%.

$$V_{mw} = \omega_{Birch} \cdot R_{mw} = 0.0132C \quad (8.1)$$

$$V_{gal} = \Omega(R_{gal}) \cdot R_{gal} \quad (8.2)$$

$$\Delta V = V_{mw} - V_{gal} \quad (8.3)$$

$$Z_{doppler} = \frac{\Delta V}{C} \quad (8.4)$$

$$R_{percent} = \frac{Z_{doppler}}{Z_{gal} + Z_{doppler}} \cdot 100 \quad (8.5)$$




Fig. 8.2 - The ratio (%) of Doppler shift to the total z shift

## **9. Cosmic Microwave Background (CMB)**

A known paradigm in modern cosmology is the cosmological principle. It states that at a “large-scale” the spatial distribution of matter in the Universe is isotropic and homogeneous. The strongest observational evidence for the cosmological principle is the CMB. Therefore, people may claim that the Pivot Universe that has a preferred axis is improbable. I claim that even though the Pivot Universe has a preferred axis, it still agrees with the CMB observation. It was demonstrated in this article, that the Milky Way is located inside a huge observable Universe ring. A Milky Way observer sees a considerable number of Galaxies in whatever direction he looks. He may conclude that the entire Universe is isotropic and homogeneous. But for an observer in a Galaxy located on the outer surfaces of the ring, the Universe is not isotropic and homogenous. In one direction, he will be facing the vacuum which is totally dark and has a temperature of absolute zero. Likewise, an observer that is located on the inner surface of the ring, i.e., facing the Pivot, sees total darkness and measures the temperature of the vacuum. Only an observer that will travel outside the ring of the visible Universe will be able to see that the Universe has an axis of rotation.

## **10. Olbers' paradox**

Olber's paradox is that an infinitely old Universe with an infinite number of stars distributed in an infinitely large space would be bright rather than dark. The BB theory explains this paradox by claiming that space is expanding and therefore the emitted light from distance Galaxies is reduced via redshift so that the most distant Galaxies are not seen from the Milky Way. The Pivot Universe, on the other hand, is a finite Universe with a finite number of stars and Galaxies. Outside the ring of the visible Universe, there is the darkness of space. An observer located inside the visible Universe ring sees a considerable number of Galaxies in whatever direction he looks, but eventually, in the background, he sees the total darkness.

## 11. Handedness of Galaxies

The Pivot theory explains yet an additional observation related to the handedness of the Universe. Longo and others [10] found an abundance of left-handed, or counter-clockwise (CCW) Galaxies. The excess is small, about 7 percent, but the chance that it could happen in an isotropic and homogeneous Universe is slim. Fig. 11.1 shows a sector of the observable Universe ring. It shows the equatorial plane of the visible Universe ring and three Galaxies, the Milky Way, and two arbitrary Galaxies: Galaxy A and Galaxy B. The Milky Way is located "above" Galaxy B and "below" Galaxy A. The Pivot theory postulates that all Galaxies rotate in the same direction, opposite to the Universe ring rotation. The location of the Galaxies in relation to the Milky Way will determine the direction of their rotation as seen from the Milky Way. Thus, Galaxy A will be seen as rotating clockwise (CCW), while Galaxy B will be seen as rotating counterclockwise (CW) when observed from the Milky Way. Had the Milky Way been located on the equatorial plane then the number of CCW and CW Galaxies would have been the same. However, if the Milky Way is offset by 7% from the equatorial plane than the number of CCW Galaxies and CW Galaxies will differ by 7%.




Fig. 11.1- Handedness of Galaxies

## 12. Historical notes

There is an ongoing debate in the scientific community in regard as to what is space. It started as a philosophical issue around 1700. There are two opposing views regarding space. One was argued by Leibniz, who claimed that space is merely the spatial relation between real things in the Universe. This means that space would not exist undependably from the things. In other words, remove the real things from the Universe and space would not exist. This view is called relationism. Clarke argued the opposing view. He claimed that space is a sort of a substance that exists everywhere. Space contains all the things in the Universe: stars, planets, Galaxies, etc. The entire Universe is moving through space that is absolute. This view is called absolutism.

Newton took part in this debate. He argued that that absolutism was the correct view and that the Universe should have a center. At this center the group density of the stars is maximal, and as we

proceed outwards of this center, the group density of the stars decreases until finally at great distances, it is succeeded by an infinite region of emptiness. In other words, the Universe is a finite island in an infinite ocean of space. Newton, in 1718, suggested that the infinite space was filled with an aether that serves as a fixed frame in the Universe. People argued against this concept because it leads to the result that light emitted by the stars and also individual stars are perpetually passing into the infinite space, never to return. Such a finite material Universe should be destined to become impoverished. Modern cosmologists abandoned Newton's theory. Specifically, because of an experiment done, in 1887, by Michelson-Morley. They showed that light was traveling at the same velocity no matter in what direction it was moving. This fact proved that there is no reference frame in the Universe, so Newton was wrong.

Einstein related to this debate. In 1905 he published his Special Relativity that did not need the existence of aether. However, this debate has not been solved, because another experiment done by Sagnac in 1913, showed contrary to the M-M experiment that there is an aether. Einstein himself did not abandon the concept of the aether. Einstein, in 1920, said: "According to the general theory of relativity space without aether is unthinkable." As for Einstein's Universe; when he published GR, in 1917, he supposed the universe to be static and unchanging, but in 1930 he changed his mind after the discovery of Hubble that the Universe was expanding in all directions. Based on the theory of GR and observations the current prevailing theory in cosmology is the Big Bang theory. According to the BB the Universe started as a point of infinite density 13.8 Billion years ago and since then is expanding at an ever-increasing acceleration in all directions. Before the BB there was nothing, no matter, no space and no time.

It is important to mention a recent experiment that was published in 2011, named Gravity Probe B. In this experiment space dragging around Earth that was predicted by GR was proven to be correct. This means that space is a real substance that has intrinsic properties. After all, if space was nothing, there were nothing to drag.

Modern cosmology has no clear answer to the question: what is space? The Big Bang theory can explain various cosmological observations but on the other hand, has raised fundamental problems. For example, how did the Universe start? Known physics fail at the moment of creation. There are additional discrepancies in the BB theory. To solve them, cosmologists suggest new hypothesis such as the dark matter and dark energy. Currently, these theories have no observational evidence.

The Pivot theory describes in general, a Universe that resembles Newton's Universe, i.e., it is an isolated island in an infinite space and has a fixed center in space. However, to explain cosmological observations, it combines Newton's Universe and Einstein's GR. Such a structure could not be imagined by Newton, because his gravitational law did not take into consideration the fact that all bodies in the Universe spin and drag space around them.

### 13. Summary

The Pivot theory describes a Universe that began as a primeval nucleus. It originated from the vacuum as can be explained by QFT. This primeval nucleon was as a Neutron star that exploded when the velocity on its surface reached the speed of light. Following the explosion, the current Universe was created. The current Universe is composed of a Neutron star the Pivot, that can also be described as Kerr blackhole and a ring of visible Universe orbiting it in the direction of the Pivot's spin. This structure describes a Universe that is in a state of dynamic equilibrium. Although the Pivot is very massive, it occupies a tiny volume of the Universe. In this sense, the Universe resembles the structure of the atom. The Pivot can be compared to the nucleon of the atom. The Pivot, like the proton, may last forever. On the other hand, the visible Universe may change, stars will consume their energy; Galaxies orbiting in too close orbits will eventually collide, etc. The entire Universe is a finite isolated island in the infinite space, that may last forever.

The Pivot Universe explains fundamental questions in physics:

Does the Universe have a reference frame? What is the origin of gravity? What is the structure of a blackhole? How are QM and GR connected? Was there time before the Universe was created?

However, there are opened issues:

How long did it take to the primeval nucleus to reach its final size before it exploded? How long did it take the Pivot Universe to rearrange after the primeval nucleus exploded? Also, there are some issues I designated as speculative.

## 14. References

1. P. Birch (1982) – “Is the Universe rotating?”.  
<http://buildengineer.com/www.paulbirch.net/IsTheUniverseRotating.pdf>
2. Wikipedia. [https://en.wikipedia.org/wiki/Observable\\_Universe#cite\\_ref-15](https://en.wikipedia.org/wiki/Observable_Universe#cite_ref-15)
3. Wikipedia. [https://en.wikipedia.org/wiki/Dark\\_matter](https://en.wikipedia.org/wiki/Dark_matter)
4. R. M. Muradian (1980). “The primeval hadron: origin of stars, Galaxies and astronomical Universe”. <http://adsabs.harvard.edu/full/1980Ap%26SS..69..339M>
5. Wikipedia <https://en.wikipedia.org/wiki/Frame-dragging>
6. Potzel (2014) “Clock hypothesis of relativity theory, maximal acceleration, and Mossbauer spectroscopy”. <https://arxiv.org/abs/1403.2412>
7. NSF “Low Reynolds number flow” <https://www.youtube.com/watch?v=QcBpDVzBPMk>
8. T.M. Davis & C.H. Lineweaver (2004) “Expanding Confusion: Common Misconception of Cosmological Horizons and the Superluminal Expansion of the Universe”.  
<http://www.publish.csiro.au/AS/pdf/AS03040>
9. R.P. Kirshner (2003) “Hubble's diagram and cosmic expansion”.  
<http://www.pnas.org/content/101/1/8.long>
10. M. J. Longo (2009) - “Evidence for a Preferred Handedness of Spiral Galaxies”.  
<https://arxiv.org/abs/0904.2529>
11. Ma & Wang “Duality Theory of Strong Interaction”.  
<http://www.ejtp.com/articles/ejtpv11i31.pdf#page=113>
12. Wikipedia. [https://en.wikipedia.org/wiki/Nuclear\\_force](https://en.wikipedia.org/wiki/Nuclear_force)
13. Wikipedia. [https://en.wikipedia.org/wiki/Neutron\\_star](https://en.wikipedia.org/wiki/Neutron_star)
14. Wikipedia. [https://en.wikipedia.org/wiki/XTE\\_J1650-500](https://en.wikipedia.org/wiki/XTE_J1650-500)

## Appendix A - The origin of gravity

The gravitational constant  $G$  is of profound importance in physics as it is involved in the calculations of gravitational effects in Newton's gravitational law and Einstein's general relativity theory. However, no theory explains its origin.  $G$  is an empirical physical constant first measured by Cavendish in 1789 and since then was verified several times.

I claim that the strong force and gravity are the same force. The strongest attractor in the Universe is the Pivot, but it contains only nucleons. Nucleons attract each other by the strong force. The magnitude of this force changes considerably as a function of the distances between nucleons. In the quantum world, the strong force reigns, but in the Universe, the weak gravity force takes over.

The theory of the origin of gravity is based on the work done by Ma and Wang [11]. They developed formulas based on field theory. The formulas developed relate to the strong force interaction between quarks, nucleons, and atoms. They developed a modified Yukawa potential equation. Yukawa, around 1930, developed a formula for the strong force that indicates that the strong nucleon force between two nucleons is always attractive. However, experimentally, it is now known that the force is attractive and repulsive depending on the distance between nucleons.

The strong force between two nucleons  $F_n$  is described by formula 6.12 in [11].

$$F_n = g^2 \cdot \left( \frac{1}{4 \cdot e^2} \cdot \frac{1}{r^2} - \frac{2 \cdot r}{r_1^3} \cdot e^{-\frac{r}{r_1}} \right) \quad (\text{A.1})$$

Where:

$$g^2 = 10 \cdot \hbar \cdot C \quad \dots \text{Nucleon interaction constant.}$$

Note: Ma and Wang are using the above value of  $g^2$ , however this value depends on the energy of the interaction or the distance between particles (TBD).

$$r \quad \dots \text{is the distance between nucleons centers.}$$

$$r_1 = 10^{-13} \text{ cm} \quad \dots \text{is Yukawa radius}$$






Fig. A.1 – Force between two nucleons as a function of the distance between them.

Figure A.1 is in good agreement with measurements. From the graph, it is also clear that the strong force is reduced substantially as the distance between nucleons grows. For example, at a distance of one hundred centimeters between nucleons, the attractive force is  $4.8 \cdot 10^{-26} N$ , dropping down from the maximum force of  $2 \cdot 10^5 N$  at a distance of  $1.3 \cdot 10^{-13} cm$ . Here, I postulate that although the force between nucleons in the Universe is extremely small, there is nevertheless, a significant attraction force between celestial bodies in the Universe because each of them contains an enormous number of nucleons. This applies specifically to the Pivot that its attraction is felt on any celestial body even if the distance between the celestial body and the Pivot is over hundreds of billions light years. In the following paragraph, I will show that the gravitational constant  $G$  can be calculated by using other fundamental physical constants.

Examining equation (A.1) shows that  $\frac{1}{4 \cdot e^{0.5} \cdot r^2} \gg \frac{2 \cdot r}{r_1^3} \cdot e^{-\frac{r}{r_1}}$  when the distance between nucleons increases. The meaning of this is that at cosmic distances the second part of the equation (A.1) can be discarded and the attraction force can be written as:

$$F_n = g^2 \cdot \frac{1}{4 \cdot e^2} \cdot \frac{1}{r^2}$$

Now, consider the force between two celestial bodies one with mass  $M_1$  and the second  $M_2$ .

The number of the protons in the first body is:  $N_1 = \frac{M_1}{m_{proton}}$ . The number of protons in the

second body is  $N_2 = \frac{M_2}{m_{proton}}$

The force between the two bodies each of them containing many nucleons, according to Ma and Wang, is:

$$F_n = 3 \cdot g^2 \cdot N_1 \cdot N_2 \cdot \left(\frac{\rho_0}{\rho_n}\right)^6 \cdot \frac{1}{4 \cdot e^2} \cdot \frac{1}{r^2} = 3 \cdot g^2 \cdot \frac{1}{m_{proton}^2} \cdot \left(\frac{\rho_0}{\rho_n}\right)^6 \cdot \frac{1}{4 \cdot e^2} \cdot \frac{M_1 \cdot M_2}{r^2} \quad (A.2)$$

On the other hand, in Newton's gravitational theory, the force between two bodies is given by:

$$F_G = G \cdot \frac{M_1 \cdot M_2}{r^2} \quad (\text{A.3})$$

where  $G$  - the universal gravitational constant.

By equating  $F_n$  (A.2) and  $F_G$  (A.3) the value of  $G$  can be calculated:

$$G = 3 \cdot g^2 \cdot \frac{1}{m_{proton}^2} \cdot \left(\frac{\rho_0}{\rho_n}\right)^6 \cdot \frac{1}{4 \cdot e^2} \quad (\text{A.4})$$

Where:  $\rho_n = 0.85 \cdot 10^{-13} \text{ cm}$  - the radius of the proton.

Note: In equation (A.4) the effective radius of the quark  $\rho_0$  is not accurately known, therefore, its value is calculated. The result is:  $\rho_0 = 0.2809 \cdot 10^{-19} \text{ cm}$ . From experiments it is known that its radius is smaller than about  $\rho_0 = 20 \cdot 10^{-19} \text{ cm}$ .

Equation (A.4) unifies quantum physics, Newton's gravitational law, and GR.

## **Appendix B - Is a blackhole a neutron star?**

The Pivot was described in this article as a Neutron star from QM point of view, and as a Kerr blackhole from GR point of view.

Blackholes existence in the visible Universe were predicted by solving GR equations. There is one profound difference between the Pivot and the Blackholes in the visible Universe. The accumulation of nucleons built the Pivot whereas a Blackhole in the visible Universe ring is the result of the gravitational collapse of a massive star. However, the final result is the same. A nucleus in which nucleons are packed to the maximum allowed density in the Universe.

Karl Schwarzschild in 1915 found an exact solution to Einstein's field equations. This solution predicted that Blackholes exist in the Universe. In this solution there is an essential singularity at  $r = 0$ , meaning that the density at the center of the Blackhole is infinite. This singularity implies that the known laws of physics break down. Einstein rejected this singularity. In a paper from 1939, Einstein concluded that there was no way a Schwarzschild singularity could ever be possible and therefore the Schwarzschild singularity does not exist in physical reality.

This issue can be resolved by postulating that in the physical world all parameters have bounds. There is no mathematical reason why there is a maximum speed in the Universe; however, physics does not allow any speed to exceed the speed of light. The same implies to density. I postulate that the maximum density of matter in the Universe is the density of a proton (or a neutron), i.e.,  $\sim 6.5 \times 10^{17} \text{ kg/m}^3$ . This maximum density is found in the nucleus of an atom and a Neutron star. The density of a Neutron star is  $3.7 \times 10^{17}$  to  $5.9 \times 10^{17} \text{ kg/m}^3$ , which is comparable to the approximate density of an atomic nucleus of  $3 \times 10^{17} \text{ kg/m}^3$ . Wikipedia [13]

Some theories claim that if the mass of a Neutron star exceeds  $\sim 3$  sun-masses, it will collapse further to become a Blackhole. One hypothesis assumes that at this point the fermions of the neutron star turn into bosons. Other theories postulate the fermions break to their constituents namely quark gluon plasma. The common denominator to these theories is that the fermions are converted into other elementary particles.

**I postulate that also a Blackhole, precisely as a Neutron star and a nucleus of an atom, have the same structure. Namely, the nucleons are densely packed to the maximum density possible in the Universe. Specifically, the maximum possible density in the Universe is the density observed in the Universe  $\sim 6.5 \times 10^{17} \text{ kg/m}^3$ .**

While GR allows the density theoretically to become infinite, quantum theory and quantum experiments show that it is not possible. One of the theoretical reasons is Pauli's exclusion principle which forbids from two identical fermion particles to occupy the same place at the same time. In a Neutron star, the neutrons are packed so densely that they touch each other. To

turn into other elementary particles, first the neutrons must be squeezed together so that they overlap, but this is not possible according to Pauli's exclusion principle.

There are QCD experiments that show why squeezing nucleons in a nucleus more than the density of a proton is not possible.

- 1) Experiments that measure the force between two nucleons as a function of the distance between them show that the force between them can be described by the graph shown in [https://en.wikipedia.org/wiki/Nuclear\\_force](https://en.wikipedia.org/wiki/Nuclear_force) . This graph is based on Reid's potential formula. It shows that for a distance smaller than 0.8fm, the force becomes a sizeable repulsive force. Further analyzing Reid's equation shows that at  $r=0$  the potential as well the force between nucleons becomes infinite.
- 2) Physicists at Jefferson Lab did another experiment <https://www.jlab.org/news/releases/quarks-feel-pressure-proton> . They measured the distribution of pressure inside the proton. The findings show that the proton's building blocks, the quarks, are subjected to a pressure of 100 decillions Pascal ( $10^{35}$ ) near the center of a proton, which is about ten times greater than the pressure in the heart of a Neutron star. The meaning is that the outward-directed pressure from the center of the proton is higher than the inward-directed pressure near the proton's periphery and therefore a Neutron star cannot collapse.

Given the description above, the question now is how come that Blackholes are not directly observed in the Universe, while Neutron stars are seen. My answer is: **The visibility depends on the relation between the physical radius of the nucleus and its Schwarzschild radius.** A celestial body will be observed if its physical radius is bigger than its Schwarzschild radius. On the other hand, if a celestial body has a physical radius that is smaller than its Schwarzschild radius, it will be hidden.

The limiting mass and radius between a Neutron star and a Blackhole can be found in the following manner:

1. Given a celestial body with mass  $M$ .

2. The radius of a densely packed spherical celestial body is:

$$R_n = R_{neutron} \cdot \left(\frac{M}{m_{neutron}}\right)^{1/3} \quad (1)$$

where:

$$\text{Mass of Neutron:} \quad m_{neutron} = 1.6749275 \cdot 10^{-27} \text{ kg}$$

$$\text{Radius of Neutron:} \quad R_{neutron} = 0.8 \cdot 10^{-13} \text{ cm}$$

3. The Schwarzschild radius of a celestial body is:

$$R_H = \frac{2 \cdot G \cdot M}{C^2} \quad (2)$$

where:

$$\text{Gravitational constant:} \quad G = 6.67 \cdot 10^{-11} \frac{m^3}{kg \cdot sec^2}$$

$$\text{Light velocity:} \quad C = 2.99 \cdot 10^8 \frac{m}{sec}$$

4. Equating Schwarzschild radius of the celestial body to its physical radius; ( $R_H = R_n$ ):

$$\text{Gives:} \quad M_{limit} = \left(\frac{R_{neutron} \cdot C^2}{2 \cdot G \cdot m_{neutron}}\right)^{3/2} = 9.67 \cdot 10^{30} \text{ kg} \sim 4.86 \text{ Sun} - \text{masses}. \quad (3)$$

$$\text{and} \quad R_{limit} = 14.35 \text{ km}$$

From the above calculations, it is shown that the limit between a Neutron star and a Blackhole is 4.86 Sun-masses and a radius of 14.35km. A celestial body with a mass higher than 4.86 Sun-masses will become a Blackhole because its physical radius is smaller than its Schwarzschild radius.

**Observations:** This result is in good agreement with observations. The smallest Blackhole observed in the Universe is XTE\_J1650-500. Its mass is estimated to be ~5 to 10 Sun masses. On the other hand, the smallest measured Blackhole is GRO 1655-40 with a mass of 6.3 Sun masses. The conclusion is that the limit mass between a Blackhole and a Neutron star is between ~5 to 6.3 Sun masses. [https://en.wikipedia.org/wiki/XTE\\_J1650-500](https://en.wikipedia.org/wiki/XTE_J1650-500) Wikipedia [14]

So far, the development of the above equations is based only on classical mechanics. The Schwarzschild radius that was found from the solution of GR equations can be derived directly from the classical equation of the escape velocity from a celestial body. (The first who suggested this was John Michell in 1783). However, in Newton's equations, the spinning of celestial bodies is not taken into considerations. It is known now that all bodies in the Universe spin. A Blackhole that is formed by gravitational collapse of a star must retain the angular momentum of the star. In SBH (Schwarzschild blackhole), the mass collapses to the point that cannot have angular momentum. The conclusion is that SBH is a theoretical solution. But there is an additional solution to GR equations. This solution takes into consideration the spinning of bodies. It was suggested in 1963 by Roy Kerr and is called KBH (Kerr blackhole). Analyzing the Kerr solution shows:

- 1) KBH has a singularity ring at its center rather than the point singularity of SBH.
- 2) There is no singularity inside this ring.
- 3) The KBH has two event horizons the outer event horizon, and an inner event horizon.
- 4) There is a frame dragging of the space around the blackhole nucleus.

I postulate that:

- 1) The nucleus of the Blackhole resides inside the ring singularity.
- 2) The density of the nucleus inside the ring singularity is the maximum density possible in the Universe.
- 3)  $R_n$  of the solid nucleus sphere, is equal to the spin parameter  $\alpha$  defined by the Kerr solution.

$$\alpha = R_n \quad \dots \text{is an assumption}$$

where:

$$\alpha = \frac{J}{M \cdot C} \quad \dots \text{is the spin parameter defined by Kerr}$$

$J$  Angular momentum of the Black hole

$M$  Mass of Black hole

$$R_{outer} = R_H + \sqrt{R_H^2 - R_n^2} \quad (4) \quad \dots \text{is the outer event horizon}$$

Frame dragging in the equatorial plane around the nucleus is calculated by:

$$\Omega(r) = \frac{R_{outer} \cdot R_n \cdot C}{r^3 + R_n^2 \cdot r + R_{outer} \cdot R_n^2} \quad (5)$$

Velocity at r:

$$V(r) = \Omega(r) \cdot r \quad (6)$$

In the following examples it is shown that the tangential velocity on the surface of the nucleus  $V(R_n)$  is between  $0.35C$  to  $1C$ .

As an example, the biggest supermassive black hole found :  $M = 2.1 \cdot 10^{10} \cdot \text{Sunmasses} = 4.2 \cdot 10^{40} \text{ kg}$

$$R_n = 2.34 \cdot 10^4 \text{ km} \quad \text{from (1)}$$

$$R_H = 6.23 \cdot 10^{10} \text{ km} \quad \text{from (2)}$$

$$R_{outer} = 6.2 \cdot 10^{11} \text{ km} \quad \text{from (4)}$$

$$\Omega(R_n) = 12.8 \text{ Hz} \quad \text{from (5)}$$

$$V(R_n) = 1 \cdot C \quad \text{from (6)}$$

Another example, the smallest black hole (according to the current article):  $M = 4.86 \cdot \text{Sunmasses} = 9.67 \cdot 10^{30} \text{ kg}$

$$R_n = 14.35 \text{ km} \quad \text{from (1)}$$

$$R_H = 14.35 \text{ km} \quad \text{from (2)}$$

$$R_{outer} = 14.35 \text{ km} \quad \text{from (4)}$$

$$\Omega(R_n) = 6.96 \cdot 10^3 \text{ Hz} \quad \text{from (5)}$$

$$V(R_n) = 0.333 \cdot C \quad \text{from (6)}$$

#### Note:

At the beginning of this appendix the value of  $M_{limit}$  was calculated for a SBH eq. (3). It seems that this must be changed for KBH because now,  $R_n = R_{outer}$  rather than  $R_n = R_s$ .

However solving  $R_n = R_s + \sqrt{R_s^2 - R_n^2}$  shows that  $R_n = R_s$ .

Thus the value of  $M_{limit}$  applies for SBH as well as for KBH.

