

Electromagnetic Waves Within Small Dielectric Particles

Scientists from the Lomonosov Moscow State University together with their Russian and foreign colleagues have done for the first time direct measurements of giant electromagnetic fields, emerging in dielectric particles with the high refractive index at the scattering of electromagnetic waves. [17]

AMOLF researchers have developed nanoscale strings whose motion can be converted to light signals with unprecedented strength. [16]

Twisted PCFs show some amazing features, from circular birefringence to conservation of the angular momentum. [15]

Photonics applications rely greatly on what physicists call nonlinear optics - the different way in which materials behave depending on the intensity of light that passes through them. The greater the nonlinearity, the more promising the material for real-life applications. Now a team, led by Robert W. Boyd, Professor of Optics and Physics at the University of Rochester and the Canada Excellence Research Chair in Quantum Nonlinear Optics at the University of Ottawa, has demonstrated that the transparent, electrical conductor indium tin oxide can result in up to 100 times greater nonlinearity than other known materials. [14]

Harnessing the power of the sun and creating light-harvesting or light-sensing devices requires a material that both absorbs light efficiently and converts the energy to highly mobile electrical current. Finding the ideal mix of properties in a single material is a challenge, so scientists have been experimenting with ways to combine different materials to create "hybrids" with enhanced features. [13]

Condensed-matter physicists often turn to particle-like entities called quasiparticles—such as excitons, plasmons, magnons—to explain complex phenomena. Now Gil Refael from the California Institute of Technology in Pasadena and colleagues report the theoretical concept of the topological polariton, or “topolariton”: a hybrid half-light, half-matter quasiparticle that has special topological properties and might be used in devices to transport light in one direction. [12]

Solitons are localized wave disturbances that propagate without changing shape, a result of a nonlinear interaction that compensates for wave packet dispersion. Individual solitons may collide, but a defining feature is that they pass through one another and emerge from the collision unaltered in shape,

amplitude, or velocity, but with a new trajectory reflecting a discontinuous jump.

Working with colleagues at the Harvard-MIT Center for Ultracold Atoms, a group led by Harvard Professor of Physics Mikhail Lukin and MIT Professor of Physics Vladan Vuletic have managed to coax photons into binding together to form molecules – a state of matter that, until recently, had been purely theoretical. The work is described in a September 25 paper in Nature.

New ideas for interactions and particles: This paper examines the possibility to origin the Spontaneously Broken Symmetries from the Planck Distribution Law. This way we get a Unification of the Strong, Electromagnetic, and Weak Interactions from the interference occurrences of oscillators. Understanding that the relativistic mass change is the result of the magnetic induction we arrive to the conclusion that the Gravitational Force is also based on the electromagnetic forces, getting a Unified Relativistic Quantum Theory of all 4 Interactions.

Giant Enhancement of Electromagnetic Waves Revealed Within Small Dielectric Particles.....	3
Nanoscale motion sends light into overdrive	5
Tightly confined light.....	5
Distorted signals	5
Revealing quantum mechanical behavior.....	5
Helically twisted photonic crystal fibres	6
Topological effects.....	6
Helical Bloch waves.....	7
Guidance of light in twisted space	8
Conclusions.....	8
'Game-changer' for photonics applications: Researchers demonstrate record optical nonlinearity ...	8
Quantum dots enhance light-to-current conversion in layered semiconductors.....	10
Quasiparticles dubbed topological polaritons make their debut in the theoretical world	11
'Matter waves' move through one another but never share space	12
Photonic molecules	13
The Electromagnetic Interaction	13
Asymmetry in the interference occurrences of oscillators	13

Spontaneously broken symmetry in the Planck distribution law.....	15
The structure of the proton	17
The Strong Interaction	17
Confinement and Asymptotic Freedom	17
The weak interaction	18
The General Weak Interaction	19
Fermions and Bosons.....	19
The fermions' spin	19
The source of the Maxwell equations	20
The Special Relativity.....	21
The Heisenberg Uncertainty Principle.....	21
The Gravitational force	21
The Graviton.....	22
What is the Spin?	23
The Casimir effect	23
The Fine structure constant.....	23
Path integral formulation of Quantum Mechanics.....	24
Conclusions	24
References	25

Author: George Rajna

Giant Enhancement of Electromagnetic Waves Revealed Within Small Dielectric Particles

Scientists from the Lomonosov Moscow State University together with their Russian and foreign colleagues have done for the first time direct measurements of giant electromagnetic fields, emerging in dielectric particles with the high refractive index at the scattering of electromagnetic waves. The researchers have presented their project results in the Scientific Reports.

The problem of miniaturization of basic elements in electronics requires new approaches. In this connection, it has become very important to be able to create intensive electromagnetic fields

concentrated in the smallest possible volume. Scientists from the Lomonosov Moscow State University in collaboration with an international team for the first time conducted direct measurements of a giant resonant field excited inside a subwavelength dielectric particle at the scattering of a plane electromagnetic wave and provided the complete quantitatively theoretical explanation of the observed effect.

Professor Michael Tribelsky, Doctor of Physics and Mathematics and the leading article author, says: "In theory this effect has been known: in this case the scattering particle acts as a funnel, gathering incident radiation from a large area and concentrating it in a small volume within the particle. However, one faces many difficulties on the way of its practical realization. Metallic nanoparticles were the first candidates for such "field concentrators." Unfortunately, they have deceived the expectations. The point is that metals have high dissipative losses in the most interesting for applications area of visible light frequencies of the incident waves. The dissipation leads to significant energy losses, wasted for fruitless (and often harmful) heating of the nanoparticle, and diminishes the resonant enhancement of the electromagnetic field. In such a case, it would be natural to turn to dielectric particles. Unfortunately, it is not so simple to deal with them too."

If a particle does not have a high value of its refractive index, the resonant effects are weak. Regarding the high-index particles, whose size is smaller than the wavelength of the incident radiation, the common belief was that the electromagnetic field hardly penetrated in such a particle. However, it has turned out to be that at some frequencies of the incident radiation the case is just the opposite. Namely, the field not only penetrates into the particle, but its giant concentration may be observed. In a sense, the effect is analogous to a gradual swinging of a swing due to weak but made in a proper moment of time pushes.

"Our main result is that, to the best of our knowledge, we are the first who has obtained the direct experimental evidence of the effect and measured the profiles of the excited fields," -- Michael Tribelsky explains.

The difficulties of the corresponding measurements at optical frequencies are related to the fact that one has to measure fields within a nanoparticle, and the spatial resolution of the measurements should be of the order of nanometers! We have overcome the difficulties, modeling the scattering of light by a nanoparticle by means of the identical scattering of radio waves by a centimeter-size particle. To be able to move a probe inside the particle, liquid dielectric (usual distilled water held at a certain fixed temperature) poured into a transparent for the incident radio wave container has been employed.

The research area, to which the project refers to, lies on the very frontier of modern studies on subwavelength optics (namely, optics, dealing with objects whose scales are smaller than the wavelength of the incident radiation). Interest in these phenomena is caused by their extensive applications ranging from medicine (diagnosis and treatment of diseases, including oncological ones; targeted drug delivery and others), biology (various sensors and markers), telecommunications (nanoantennas) to systems for information recording and storage and other spheres. This phenomenon could also be used for the creation of revolutionary new optical computers where information is transferred not by electric pulses but by light packages.

The scientist concludes: "In a broad perspective our project may initiate the creation of a new landscape for design and fabrication of super-miniature nanodevices and metamaterials -- namely, materials, which are artificially formed and structured in a special way to possess unusual electromagnetic properties." [17]

Nanoscale motion sends light into overdrive

AMOLF researchers have developed nanoscale strings whose motion can be converted to light signals with unprecedented strength. This could allow for extremely precise sensors and comes with an important side effect. "Analogous to a guitar amplifier in overdrive producing distorted sound waves, our strong motion-to-light conversion leads to distorted light signals," says group leader Ewold Verhagen. "But these signals actually carry information about the motion that may lead to new ways of measuring quantum mechanical motion." The researchers published their results on 7 July 2017 in Nature Communications.

When a guitar player turns up the volume of the amplifier to the highest gain levels, the 'clean' harmonic vibrations of the guitar strings are converted to distorted sound waves. This 'dirty' sound is often desirable—the screaming of overdriven guitar amplifiers has defined the sound of rock music for decades.

Tightly confined light

The Photonic Forces group at AMOLF studies the interaction of motion and light using nanoscale silicon strings that vibrate like guitar strings at millions of times per second. The researchers use light to measure these vibrations with extreme precision. PhD student Rick Leijssen and his colleagues developed strings with a particular shape to squeeze light between the strings at a scale of tens of nanometers. "The tight confinement causes mechanical motion to be converted to light signals with unprecedented strength. This is a big step forward to creating motion sensors with extreme precision," Leijssen says. "Such sensors could detect string vibrations with amplitudes as small as the size of a proton and could be used to measure small forces and masses."

Distorted signals

The strong conversion of motion into light in the silicon strings has a side effect: The conversion is so strong that even for the tiny intrinsic fluctuations of the strings, the light is 'overdriven,' analogous to what happens in rock guitar amplifiers. Verhagen says, "Similar to the screaming sound of an overdriven amplifier, the light signals in our experiment contain many higher harmonics ('overtones') of the fundamental string resonance. This is because the conversion between motion and light is no longer linear."

Revealing quantum mechanical behavior

In a way, this nonlinear conversion forms a practical limit to the motion sensitivity. However, the distorted light signals could be put to new use. An important reason why the AMOLF physicists study tiny string vibrations is to reveal whether objects like strings behave according to the laws of quantum mechanics. Postdoctoral researcher Juha Muhonen says, "The higher harmonics of the produced light signals carry different kinds of information about the nanostring motion. For example, we demonstrated that they allow measuring the energy of the vibration with a high

accuracy. This could potentially lead to direct observations of quantized energy in the string, as we would expect from the theory of quantum mechanics." [16]

Helically twisted photonic crystal fibres

Photonic crystal fibres (PCF) are strands of glass, not much thicker than a human hair, with a lattice of hollow channels running along the fibre. If they are continuously twisted in their production, they resemble a multi-helix. Twisted PCFs show some amazing features, from circular birefringence to conservation of the angular momentum. The biggest surprise, however, is the robust light guidance itself, with no visible fibre core. The basis for this are forces which, like gravitation, are based on the curvature of space.

Chiral materials consist of many identical units (molecules or nanostructured elements) that are either randomly oriented in solution or arranged in an ordered fashion. They are ubiquitous in nature—for example, most biological molecules come in right and left-handed forms—and are finding an increasing number of applications in science and technology. Twisted photonic crystal fibre (t-PCF), in contrast, consists of a single uniaxial chiral unit that is infinitely extended in the third dimension—the direction of the twist. PCF itself typically consists of a hexagonal array of hollow microchannels running along the length of a glass fibre $\sim 100\ \mu\text{m}$ thick, so that when twisted it resembles a "multi-helix" of spiralling microchannels around a central axis (Fig. 1(a)).

Over the past several years we have been studying the behaviour of light in a range of different types of t-PCF, in the process uncovering some surprising phenomena and exploring potential applications.

We use two techniques for producing t-PCF. In the first, an untwisted PCF is post-processed under CO₂ laser heating, the fibre being mounted between a motorized rotation stage and a rigid support (Fig. 2(a)). As the motor rotates, the focused $10\ \mu\text{m}$ laser beam is scanned along the fibre using a steering mirror fixed to a precision motorized translation stage. Once the target twist period and sample length are set, the laser power and the scanning speed are chosen so as to heat the fibre to the glass-softening temperature. The writing process is computer-controlled and is capable of achieving twist periods as short as $300\ \mu\text{m}$. The second technique involves spinning the glass preform during fibre drawing, using a motor rotating at a few thousand rpm and a rotary joint with multiple inlets for controlling the pressure inside the hollow channels (Fig. 2(b)). It has the advantage that long lengths (100s of metres) of helical PCF with twist periods of a few millimetres can be readily fabricated.

Topological effects

The propagation of electromagnetic waves in helical structures began in earnest in the 1940s, with the invention of the travelling-wave tube amplifier. In this device a microwave signal is guided along a helical wire that spirals around an axially propagating electron beam. Since the physical distance over which the spiralling microwave signal travels is longer than the distance directly along the axis, its group and phase velocities are both effectively reduced. By appropriate design the velocity difference between the two waves can be adjusted, permitting the microwave signal to be amplified with power from the electron beam. In a similar manner, the geometrical stretching of the cladding structure in a t-PCF causes the effective optical path-length along the axis, and thus the effective

refractive index, to increase topologically with radius r following the relationship $n_{\text{eff}}(r) = n_0(1 + ar^2)^{1/2}$ where n_0 is the index in the untwisted case and a the twist rate in rad/m (see Fig. 1(b)).

Spectral dips in t-PCF with single core

This topological effect makes it possible for example to phase-match light guided in a central solid glass core (modal index n_c) to the fundamental space-filling mode in the cladding (phase index n_{SM} in the untwisted fibre) with the result that light can leak out into cladding modes at certain wavelengths. This results in a series of dips in the transmission spectrum, caused by anti-crossings between the core mode and leaky ring-shaped cladding modes (Fig. 1(c)) carrying orbital angular momentum (OAM), each dip corresponding to a different OAM order. Since the cladding light is diverted by the hollow channels into a spiral path, the azimuthal component of its wavevector must take values that yield a round-trip phase advance that is an integer multiple of 2π , where l is the OAM order. This leads to the condition:

$$l\lambda / (2\pi) = n_{az} r = n_{SM} r \sin\theta \sim n_{SM} a r^2 \quad (1)$$

where λ is the dip wavelength of the OAM order, n_{az} the azimuthal component of refractive index, and θ the local angle between the hollow channels and the fibre axis. Eq. (1) yields remarkably good agreement with experimental measurements, showing in particular that the dip wavelengths scale linearly with the twist rate. We have used the twist and strain sensitivity of these dips to construct an all-optical twist-strain transducer.

Helical Bloch waves

Understanding the physics of light propagation in t-PCF is quite challenging, because the natural coordinate system—helical—is non-orthogonal. This led us to introduce a new concept: helical Bloch waves. The optical Bloch waves of any untwisted periodic structure are described by the product of a periodic function $P(r)$ (with periodicities that match the structure) and a term representing the phase progression of the Bloch wave. A convenient physical picture for the modes guided in a t-PCF can be constructed by generalising Bloch's theorem so that the azimuthally periodic function follows the twist, taking the form where r is the radial coordinate and the azimuthal angle. At any given value of z , P will repeat at angular intervals $2\pi/N$, where N is the number of times the structure repeats over one revolution 2π . The Bloch waves can then be calculated analytically using an expansion in terms of azimuthal harmonics of OAM order l . Substituting this field Ansatz into Maxwell's equations allows the dispersion relation to be derived.

To explore the properties of helical Bloch waves, we fabricated a t-PCF with a ring of six solid glass "satellite" cores around its axis (Fig. 3(a)). The hollow channels had diameter $2\ \mu\text{m}$, spaced by $3\ \mu\text{m}$, and the twist rate was $2.9\ \text{rad/mm}$. This structure supports 6 non-degenerate helical Bloch modes with different values of orbital angular momentum, in both left and right circularly polarised states. To determine the OAM order of the modes guided through the t-PCF, the output was superimposed on to a divergent Gaussian beam and the resulting fringe pattern imaged using a CCD camera. The single- and double-spiral interference patterns in Fig. 3(b), which were recorded at a wavelength of $632.8\ \text{nm}$, confirm that the fibre generates optical vortices and preserves the magnitude and sign of the OAM for all four modes. Similar experiments carried out at multiple wavelengths and for fibres up to $50\ \text{m}$ long have confirmed that the t-PCFs preserve the magnitude and sign of the OAM.

Guidance of light in twisted space

We have discovered a new mechanism of light guidance, based on a t-PCF with no core. Cleaving the fibre and examining its cross-section reveals a complete absence of any structure at which light could be trapped (see Fig. 4(a)). Nevertheless it supports a guided mode: the helical twist creates a topological channel within which light is robustly trapped. This arises from the quadratic increase in optical path length with radius (mentioned above), which produces a radial gradient in axial refractive index, creating a potential well within which light is confined by photonic bandgap effects. Using mathematical tools from general relativity, we have shown that the geodesics of the light follow closed spiral paths within the topological channel, forming modes that carry OAM. The effective area of these modes decreases with twist rate a , so that by varying the twist rate along the fibre, it would be possible to create fibres whose mode-field diameter changes with position. Unlike in conventional index-guiding fibres, where the guided mode shifts towards the outside of the bend ("normal cornering"), this highly unusual mode shifts inwards towards the bend ("anomalous cornering"). Hamiltonian optics shows that the mode may be viewed as having negative effective mass (caused by the opposite sign of the dispersion surface curvature), so that it moves in the opposite direction when subject to bend forces.

Conclusions

The ability of t-PCF to generate and support OAM modes, as well as providing optical activity and circular dichroism, suggests that it may become useful in many applications. The series of transmission dips at twist-tunable wavelengths in solid-core PCF has applications in sensing and filtering. The transmission and preservation of circular polarisation states makes t-PCF very interesting for current sensing based on Faraday rotation. Its ability to robustly transmit pure OAM states over long distances may lead to applications in particle manipulation and telecommunications. It seems likely that many of these effects and phenomena will move into real-world applications in the near future. As yet unexplored is the use of t-PCF in nonlinear optics and fibre lasers, where the combination of circular and OAM birefringence with control of group velocity dispersion may offer opportunities for new kinds of mode-locked soliton lasers, wavelength conversion devices and supercontinuum sources. [15]

'Game-changer' for photonics applications: Researchers demonstrate record optical nonlinearity

Photonics applications rely greatly on what physicists call nonlinear optics - the different way in which materials behave depending on the intensity of light that passes through them. The greater the nonlinearity, the more promising the material for real-life applications. Now a team, led by Robert W. Boyd, Professor of Optics and Physics at the University of Rochester and the Canada Excellence Research Chair in Quantum Nonlinear Optics at the University of Ottawa, has demonstrated that the transparent, electrical conductor indium tin oxide can result in up to 100 times greater nonlinearity than other known materials.

"This result is a game-changer for photonics applications," said Boyd. "It rests on the core of what I've worked on for over 30 years at Rochester. I find it very rewarding that even after all this time there are still fundamental questions to be answered in the field of nonlinear optics."

The result will be published online by the journal Science on Thursday, April 28, 2016.

Photonics uses light to transmit information. Photonics also uses light to perform logical operations, just as electronics does with electrons. A key aspect to being able to exert control over light is to be able to control a specific property - the refractive index - of the material that is transmitting the light.

Tweaking the refractive index of a material, which leads to light travelling faster or slower, is the key way in which photonics applications control light. When the refractive index is different for different light intensities, the material is described as being optically nonlinear.

When a pulse of light is sent through the material, the refractive index changes according to that intensity. The refractive index of the material is changed for only a few femtoseconds - a few millionths of a billionth of a second. For some potential applications, it is possible to send a second pulse through the material before it has time to recover for the first pulse. This second pulse then "sees" the material as having the refractive index as modified by the first pulse.

In general though, it is the quickness with which the material recovers as well as the range of values that the refractive index can take - how strongly nonlinear the material is - that makes this system particularly attractive to photonics applications.

Boyd, his PhD student at Ottawa, M. Zahirul Alam, and then research associate Israel De Leon (currently a professor in Monterrey, Mexico) were able to improve on the previous record for optical nonlinearities by a factor of 100. This improvement took place because these researchers exploited the unusual optical properties of a material that occurs under certain conditions, what is known as the epsilon-near-zero region.

"It was surprising that showing such a strong optical nonlinearity in a known metal was this easy," said Boyd. "This material has been around for many years, but until now the community had overlooked the potential that the 'epsilon-near-zero' region of materials offered."

"The optical nonlinear response that we have observed introduces a new paradigm in nonlinear optics," said De Leon, now a professor at Tecnológico de Monterrey, Mexico. "The common knowledge had always been that nonlinear effects are tiny compared with the linear ones; but in our work we have measured a nonlinear response that is 170% larger than the linear response."

The result opens the door for more careful study of this region of materials, with a view to finding a material that can offer just the right properties for certain photonics applications.

The 'epsilon-near-zero' region for this material is linked to light of a specific frequency, roughly a wavelength of 1.2 micrometers. This wavelength is of interest because it is in between that of visible light and light of wavelength 1.5 micrometers. This wavelength is of particular interest to optical communications, which uses devices such as fiber optics to transmit information in light.

It is possible that changes in the chemical composition of the material could lead to a change in the frequency at which epsilon near zero occurs, thus bringing this frequency closer to that used by optical communications. [14]

Quantum dots enhance light-to-current conversion in layered semiconductors

Harnessing the power of the sun and creating light-harvesting or light-sensing devices requires a material that both absorbs light efficiently and converts the energy to highly mobile electrical current. Finding the ideal mix of properties in a single material is a challenge, so scientists have been experimenting with ways to combine different materials to create "hybrids" with enhanced features.

In two just-published papers, scientists from the U.S. Department of Energy's Brookhaven National Laboratory, Stony Brook University, and the University of Nebraska describe one such approach that combines the excellent light-harvesting properties of quantum dots with the tunable electrical conductivity of a layered tin disulfide semiconductor. The hybrid material exhibited enhanced light-harvesting properties through the absorption of light by the quantum dots and their energy transfer to tin disulfide, both in laboratory tests and when incorporated into electronic devices. The research paves the way for using these materials in optoelectronic applications such as energy-harvesting photovoltaics, light sensors, and light emitting diodes (LEDs).

According to Mircea Cotlet, the physical chemist who led this work at Brookhaven Lab's Center for Functional Nanomaterials (CFN), a DOE Office of Science User Facility, "Two-dimensional metal dichalcogenides like tin disulfide have some promising properties for solar energy conversion and photodetector applications, including a high surface-to-volume aspect ratio. But no semiconducting material has it all. These materials are very thin and they are poor light absorbers. So we were trying to mix them with other nanomaterials like light-absorbing quantum dots to improve their performance through energy transfer."

One paper, just published in the journal *ACS Nano*, describes a fundamental study of the hybrid quantum dot/tin disulfide material by itself. The work analyzes how light excites the quantum dots (made of a cadmium selenide core surrounded by a zinc sulfide shell), which then transfer the absorbed energy to layers of nearby tin disulfide.

"We have come up with an interesting approach to discriminate energy transfer from charge transfer, two common types of interactions promoted by light in such hybrids," said Prahlad Routh, a graduate student from Stony Brook University working with Cotlet and co-first author of the *ACS Nano* paper. "We do this using single nanocrystal spectroscopy to look at how individual quantum dots blink when interacting with sheet-like tin disulfide. This straightforward method can assess whether components in such semiconducting hybrids interact either by energy or by charge transfer."

The researchers found that the rate for non-radiative energy transfer from individual quantum dots to tin disulfide increases with an increasing number of tin disulfide layers. But performance in laboratory tests isn't enough to prove the merits of potential new materials. So the scientists incorporated the hybrid material into an electronic device, a photo-field-effect-transistor, a type of photon detector commonly used for light sensing applications.


As described in a paper published online March 24 in *Applied Physics Letters*, the hybrid material dramatically enhanced the performance of the photo-field-effect transistors—resulting in a

photocurrent response (conversion of light to electric current) that was 500 percent better than transistors made with the tin disulfide material alone.

"This kind of energy transfer is a key process that enables photosynthesis in nature," said Chang-Yong Nam, a materials scientist at Center for Functional Nanomaterials and co-corresponding author of the APL paper. "Researchers have been trying to emulate this principle in light-harvesting electrical devices, but it has been difficult particularly for new material systems such as the tin disulfide we studied. Our device demonstrates the performance benefits realized by using both energy transfer processes and new low-dimensional materials."

Cotlet concludes, "The idea of 'doping' two-dimensional layered materials with quantum dots to enhance their light absorbing properties shows promise for designing better solar cells and photodetectors." [13]

Quasiparticles dubbed topological polaritons make their debut in the theoretical world



Condensed-matter physicists often turn to particle-like entities called quasiparticles—such as excitons, plasmons, magnons—to explain complex phenomena. Now Gil Refael from the California Institute of Technology in Pasadena and colleagues report the theoretical concept of the topological polariton, or “topolariton”: a hybrid half-light, half-matter quasiparticle that has special topological properties and might be used in devices to transport light in one direction.

The proposed topolaritons arise from the strong coupling of a photon and an exciton, a bound state of an electron and a hole. Their topology can be thought of as knots in their gapped energy-band structure. At the edge of the systems in which topolaritons emerge, these knots unwind and allow the topolaritons to propagate in a single direction without back-reflection. In other words, the topolaritons cannot make U-turns. Back-reflection is a known source of detrimental feedback and loss in photonic devices. The topolaritons’ immunity to it may thus be exploited to build devices with increased performance.

The researchers describe a scheme to generate topolaritons that may be feasible to implement in common systems—such as semiconductor structures or atomically thin layers of compounds known as transition-metal dichalcogenides—embedded in photonic waveguides or microcavities. Previous approaches to make similar one-way photonic channels have mostly hinged on effects that are only applicable at microwave frequencies. Refael and co-workers’ proposal offers an avenue to make

such “one-way photonic roads” in the optical regime, which despite progress has remained a challenging pursuit. [12]

'Matter waves' move through one another but never share space

Physicist Randy Hulet and colleagues observed a strange disappearing act during collisions between forms of Bose Einstein condensates called solitons. In some cases, the colliding clumps of matter appear to keep their distance even as they pass through each other. How can two clumps of matter pass through each other without sharing space? Physicists have documented a strange disappearing act by colliding Bose Einstein condensates that appear to keep their distance even as they pass through one another.

BECs are clumps of a few hundred thousand lithium atoms that are cooled to within one-millionth of a degree above absolute zero, a temperature so cold that the atoms march in lockstep and act as a single "matter wave." Solitons are waves that do not diminish, flatten out or change shape as they move through space. To form solitons, Hulet's team coaxed the BECs into a configuration where the attractive forces between lithium atoms perfectly balance the quantum pressure that tends to spread them out.

The researchers expected to observe the property that a pair of colliding solitons would pass through one another without slowing down or changing shape. However, they found that in certain collisions, the solitons approached one another, maintained a minimum gap between themselves, and then appeared to bounce away from the collision.

Hulet's team specializes in experiments on BECs and other ultracold matter. They use lasers to both trap and cool clouds of lithium gas to temperatures that are so cold that the matter's behavior is dictated by fundamental forces of nature that aren't observable at higher temperatures.

To create solitons, Hulet and postdoctoral research associate Jason Nguyen, the study's lead author, balanced the forces of attraction and repulsion in the BECs.

Cameras captured images of the tiny BECs throughout the process. In the images, two solitons oscillate back and forth like pendulums swinging in opposite directions. Hulet's team, which also included graduate student De Luo and former postdoctoral researcher Paul Dyke, documented thousands of head-on collisions between soliton pairs and noticed a strange gap in some, but not all, of the experiments.

Many of the events that Hulet's team measures occur in one-thousandth of a second or less. To confirm that the "disappearing act" wasn't causing a miniscule interaction between the soliton pairs -- an interaction that might cause them to slowly dissipate over time -- Hulet's team tracked one of the experiments for almost a full second.

The data showed the solitons oscillating back and fourth, winking in and out of view each time they crossed, without any measurable effect.

"This is great example of a case where experiments on ultracold matter can yield a fundamental new insight," Hulet said. "The phase-dependent effects had been seen in optical experiments, but there has been a misunderstanding about the interpretation of those observations." [11]

Photonic molecules

Working with colleagues at the Harvard-MIT Center for Ultracold Atoms, a group led by Harvard Professor of Physics Mikhail Lukin and MIT Professor of Physics Vladan Vuletic have managed to coax photons into binding together to form molecules – a state of matter that, until recently, had been purely theoretical. The work is described in a September 25 paper in Nature.

The discovery, Lukin said, runs contrary to decades of accepted wisdom about the nature of light. Photons have long been described as massless particles which don't interact with each other – shine two laser beams at each other, he said, and they simply pass through one another.

"Photonic molecules," however, behave less like traditional lasers and more like something you might find in science fiction – the light saber.

"Most of the properties of light we know about originate from the fact that photons are massless, and that they do not interact with each other," Lukin said. "What we have done is create a special type of medium in which photons interact with each other so strongly that they begin to act as though they have mass, and they bind together to form molecules. This type of photonic bound state has been discussed theoretically for quite a while, but until now it hadn't been observed. [9]

The Electromagnetic Interaction

This paper explains the magnetic effect of the electric current from the observed effects of the accelerating electrons, causing naturally the experienced changes of the electric field potential along the electric wire. The accelerating electrons explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the electron's spin also, building the bridge between the Classical and Quantum Theories. [2]

Asymmetry in the interference occurrences of oscillators

The asymmetrical configurations are stable objects of the real physical world, because they cannot annihilate. One of the most obvious asymmetry is the proton – electron mass rate $M_p = 1840 M_e$ while they have equal charge. We explain this fact by the strong interaction of the proton, but how remember it his strong interaction ability for example in the H – atom where are only electromagnetic interactions among proton and electron.

This gives us the idea to origin the mass of proton from the electromagnetic interactions by the way interference occurrences of oscillators. The uncertainty relation of Heisenberg makes sure that the particles are oscillating.

The resultant intensity due to n equally spaced oscillators, all of equal amplitude but different from one another in phase, either because they are driven differently in phase or because we are looking at them an angle such that there is a difference in time delay:

$$(1) I = I_0 \sin^2 n \phi/2 / \sin^2 \phi/2$$

If ϕ is infinitesimal so that $\sin\phi = \phi$, then

$$(2) I = n^2 I_0$$

This gives us the idea of

$$(3) M_p = n^2 M_e$$




Fig. 30-3. A linear array of n equal oscillators, driven with phases $\alpha_s = s\alpha$.

Figure 1.) A linear array of n equal oscillators

There is an important feature about formula (1) which is that if the angle ϕ is increased by the multiple of 2π , it makes no difference to the formula.

So

$$(4) d \sin \theta = m \lambda$$

and we get m -order beam if λ less than d . [6]

If d less than λ we get only zero-order one centered at $\theta = 0$. Of course, there is also a beam in the opposite direction. The right choices of d and λ we can ensure the conservation of charge.

For example

$$(5) 2(m+1) = n$$

Where $2(m+1) = N_p$ number of protons and $n = N_e$ number of electrons.

In this way we can see the H_2 molecules so that $2n$ electrons of n radiate to $4(m+1)$ protons, because $d_e > \lambda_e$ for electrons, while the two protons of one H_2 molecule radiate to two electrons of them, because of $d_e < \lambda_e$ for this two protons.

To support this idea we can turn to the Planck distribution law, that is equal with the Bose – Einstein statistics.

Spontaneously broken symmetry in the Planck distribution law

The Planck distribution law is temperature dependent and it should be true locally and globally. I think that Einstein's energy-matter equivalence means some kind of existence of electromagnetic oscillations enabled by the temperature, creating the different matter formulas, atoms molecules, crystals, dark matter and energy.

Max Planck found for the black body radiation

As a function of wavelength (λ), Planck's law is written as:

$$B_{\lambda}(T) = \frac{2hc^2}{\lambda^5} \frac{1}{e^{\frac{hc}{\lambda T}} - 1}.$$




Figure 2. The distribution law for different T temperatures

We see there are two different λ_1 and λ_2 for each T and intensity, so we can find between them a d so that $\lambda_1 < d < \lambda_2$.

We have many possibilities for such asymmetrical reflections, so we have many stable oscillator configurations for any T temperature with equal exchange of intensity by radiation. All of these configurations can exist together. At the λ_{\max} is the annihilation point where the configurations are symmetrical. The λ_{\max} is changing by the Wien's displacement law in many textbooks.

$$(7) \quad \lambda_{\max} = \frac{b}{T}$$

where λ_{\max} is the peak wavelength, T is the absolute temperature of the black body, and b is a constant of proportionality called *Wien's displacement constant*, equal to $2.8977685(51) \times 10^{-3} \text{ m} \cdot \text{K}$ (2002 CODATA recommended value).

By the changing of T the asymmetrical configurations are changing too.

The structure of the proton

We must move to the higher T temperature if we want look into the nucleus or nucleon arrive to $d < 10^{-13}$ cm. If an electron with $\lambda_e < d$ move across the proton then by (5) $2(m+1) = n$ with $m = 0$ we get $n = 2$ so we need two particles with negative and two particles with positive charges. If the proton can fraction to three parts, two with positive and one with negative charges, then the reflection of oscillators are right. Because this very strange reflection where one part of the proton with the electron together on the same side of the reflection, the all parts of the proton must be quasi lepton so $d > \lambda_q$. One way dividing the proton to three parts is, dividing his oscillation by the three direction of the space. We can order $1/3$ e charge to each coordinates and $2/3$ e charge to one plane oscillation, because the charge is scalar. In this way the proton has two $+2/3$ e plane oscillation and one linear oscillation with $-1/3$ e charge. The colors of quarks are coming from the three directions of coordinates and the proton is colorless. The flavors of quarks are the possible oscillations differently by energy and if they are plane or linear oscillations. We know there is no possible reflecting two oscillations to each other which are completely orthogonal, so the quarks never can be free, however there is an asymptotic freedom while their energy are increasing to turn them to the orthogonally. If they will be completely orthogonal then they lose this reflection and take new partners from the vacuum. Keeping the symmetry of the vacuum the new oscillations are keeping all the conservation laws, like charge, number of baryons and leptons. The all features of gluons are coming from this model. The mathematics of reflecting oscillators show Fermi statistics.

Important to mention that in the Deuteron there are 3 quarks of $+2/3$ and $-1/3$ charge, that is three u and d quarks making the complete symmetry and because this its high stability.

The Pauli Exclusion Principle says that the diffraction points are exclusive!

The Strong Interaction

Confinement and Asymptotic Freedom

For any theory to provide a successful description of strong interactions it should simultaneously exhibit the phenomena of confinement at large distances and asymptotic freedom at short distances. Lattice calculations support the hypothesis that for non-abelian gauge theories the two domains are analytically connected, and confinement and asymptotic freedom coexist. Similarly, one way to show that QCD is the correct theory of strong interactions is that the coupling extracted at various scales (using experimental data or lattice simulations) is unique in the sense that its variation with scale is given by the renormalization group. [4]
Lattice QCD gives the same results as the diffraction theory of the electromagnetic oscillators, which is the explanation of the strong force and the quark confinement. [1]

The weak interaction

The weak interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and time reversal symmetry.

Another important issue of the quark model is when one quark changes its flavor such that a linear oscillation transforms into plane oscillation or vice versa, changing the charge value with 1 or -1. This kind of change in the oscillation mode requires not only parity change, but also charge and time changes (CPT symmetry) resulting a right handed anti-neutrino or a left handed neutrino.

The right handed anti-neutrino and the left handed neutrino exist only because changing back the quark flavor could happen only in reverse, because they are different geometrical constructions, the u is 2 dimensional and positively charged and the d is 1 dimensional and negatively charged. It needs also a time reversal, because anti particle (anti neutrino) is involved.

The neutrino is a 1/2 spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $\frac{1}{2}$ spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell–Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The limit of the velocity of particles as the speed of light appropriate only for electrical charged particles, since the accelerated charges are self maintaining locally the accelerating electric force. The neutrinos are CP symmetry breaking particles compensated by time in the CPT symmetry, that is the time coordinate not works as in the electromagnetic interactions, consequently the speed of neutrinos is not limited by the speed of light.

The weak interaction T-asymmetry is in conjunction with the T-asymmetry of the second law of thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes the weak interaction, for example the Hydrogen fusion.

Probably because it is a spin creating movement changing linear oscillation to 2 dimensional oscillation by changing d to u quark and creating anti neutrino going back in time relative to the proton and electron created from the neutron, it seems that the anti neutrino fastest then the velocity of the photons created also in this weak interaction?

A quark flavor changing shows that it is a reflection changes movement and the CP- and T- symmetry breaking. This flavor changing oscillation could prove that it could be also on higher level such as atoms, molecules, probably big biological significant molecules and responsible on the aging of the life.

Important to mention that the weak interaction is always contains particles and antiparticles, where the neutrinos (antineutrinos) present the opposite side. It means by Feynman's interpretation that these particles present the backward time and probably because this they seem to move faster than the speed of light in the reference frame of the other side.

Finally since the weak interaction is an electric dipole change with $\frac{1}{2}$ spin creating, it is limited by the velocity of the electromagnetic wave, so the neutrino's velocity cannot exceed the velocity of light.

The General Weak Interaction

The Weak Interactions T-asymmetry is in conjunction with the T-asymmetry of the Second Law of

Thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes for example the Hydrogen fusion. The arrow of time by the Second Law of Thermodynamics shows the increasing entropy and decreasing information by the Weak Interaction, changing the temperature dependent diffraction patterns. A good example of this is the neutron decay, creating more particles with less known information about them.

The neutrino oscillation of the Weak Interaction shows that it is a general electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. There should be a new research space of the Quantum Information Science the 'general neutrino oscillation' for the greater than subatomic matter structures as an electric dipole change.

There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction. [5]

Fermions and Bosons

The fermions are the diffraction patterns of the bosons such a way that they are both sides of the same thing.

The Higgs boson or Higgs particle is a proposed elementary particle in the Standard Model of particle physics. The Higgs boson's existence would have profound importance in particle physics because it would prove the existence of the hypothetical Higgs field - the simplest of several proposed explanations for the origin of the symmetry-breaking mechanism by which elementary particles gain mass. [3]

The fermions' spin

The moving charges are accelerating, since only this way can self maintain the electric field causing their acceleration. The electric charge is not point like! This constant acceleration possible if there is

a rotating movement changing the direction of the velocity. This way it can accelerate forever without increasing the absolute value of the velocity in the dimension of the time and not reaching the velocity of the light.

The Heisenberg uncertainty relation says that the minimum uncertainty is the value of the spin: $1/2 \hbar = \Delta x \Delta p$ or $1/2 \hbar = \Delta t \Delta E$, that is the value of the basic energy status.

What are the consequences of this in the weak interaction and how possible that the neutrinos' velocity greater than the speed of light?

The neutrino is the one and only particle doesn't participate in the electromagnetic interactions so we cannot expect that the velocity of the electromagnetic wave will give it any kind of limit.

The neutrino is a $1/2$ spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $1/2$ spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell–Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The source of the Maxwell equations

The electrons are accelerating also in a static electric current because of the electric force, caused by the potential difference. The magnetic field is the result of this acceleration, as you can see in [2].

The mysterious property of the matter that the electric potential difference is self maintained by the accelerating electrons in the electric current gives a clear explanation to the basic sentence of the relativity that is the velocity of the light is the maximum velocity of the matter. If the charge could move faster than the electromagnetic field than this self maintaining electromagnetic property of the electric current would be failed.

Also an interesting question, how the changing magnetic field creates a negative electric field? The answer also the accelerating electrons will give. When the magnetic field is increasing in time by increasing the electric current, then the acceleration of the electrons will increase, decreasing the charge density and creating a negative electric force. Decreasing the magnetic field by decreasing the electric current will decrease the acceleration of the electrons in the electric current and increases the charge density, creating an electric force also working against the change. In this way we have explanation to all interactions between the electric and magnetic forces described in the Maxwell equations.

The second mystery of the matter is the mass. We have seen that the acceleration change of the electrons in the flowing current causing a negative electrostatic force. This is the cause of the relativistic effect - built-in in the Maxwell equations - that is the mass of the electron growing with its acceleration and its velocity never can reach the velocity of light, because of this growing negative electrostatic force. The velocity of light is depending only on 2 parameters: the magnetic permeability and the electric permittivity.

There is a possibility of the polarization effect created by electromagnetic forces creates the negative and positive charges. In case of equal mass as in the electron-positron pair it is simply, but

on higher energies can be asymmetric as the electron-proton pair of neutron decay by weak interaction and can be understood by the Feynman graphs.

Anyway the mass can be electromagnetic energy exceptionally and since the inertial and gravitational mass are equal, the gravitational force is electromagnetic force and since only the magnetic force is attractive between the same charges, is very important for understanding the gravitational force.

The Uncertainty Relations of Heisenberg gives the answer, since only this way can be sure that the particles are oscillating in some way by the electromagnetic field with constant energies in the atom indefinitely. Also not by chance that the uncertainty measure is equal to the fermions spin, which is one of the most important feature of the particles. There are no singularities, because the moving electron in the atom accelerating in the electric field of the proton, causing a charge distribution on Δx position difference and with a Δp momentum difference such a way that their product is about the half Planck reduced constant. For the proton this Δx is much less in the nucleon, than in the orbit of the electron in the atom, the Δp is much higher because of the greatest proton mass.

The Special Relativity

The mysterious property of the matter that the electric potential difference is self maintained by the accelerating electrons in the electric current gives a clear explanation to the basic sentence of the relativity that is the velocity of the light is the maximum velocity of the matter. If the charge could move faster than the electromagnetic field than this self maintaining electromagnetic property of the electric current would be failed. [8]

The Heisenberg Uncertainty Principle

Moving faster needs stronger acceleration reducing the Δx and raising the Δp . It means also mass increasing since the negative effect of the magnetic induction, also a relativistic effect!

The Uncertainty Principle also explains the proton – electron mass ratio since the Δx is much less requiring bigger Δp in the case of the proton, which is partly the result of a bigger mass m_p because of the higher electromagnetic induction of the bigger frequency (impulse).

The Gravitational force

The changing magnetic field of the changing current causes electromagnetic mass change by the negative electric field caused by the changing acceleration of the electric charge.

The gravitational attractive force is basically a magnetic force.

The same electric charges can attract one another by the magnetic force if they are moving parallel in the same direction. Since the electrically neutral matter is composed of negative and positive charges they need 2 photons to mediate this attractive force, one per charges. The Big Bang caused parallel moving of the matter gives this magnetic force, experienced as gravitational force.

Since graviton is a tensor field, it has spin = 2, could be 2 photons with spin = 1 together.

You can think about photons as virtual electron – positron pairs, obtaining the necessary virtual mass for gravity.

The mass as seen before a result of the diffraction, for example the proton – electron mass ratio $M_p = 1840 M_e$. In order to move one of these diffraction maximum (electron or proton) we need to intervene into the diffraction pattern with a force appropriate to the intensity of this diffraction maximum, means its intensity or mass. [1]

The Big Bang caused acceleration created radial currents of the matter, and since the matter is composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles. The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces!?

The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

The Graviton

In physics, the graviton is a hypothetical elementary particle that mediates the force of gravitation in the framework of quantum field theory. If it exists, the graviton is expected to be massless (because the gravitational force appears to have unlimited range) and must be a spin-2 boson. The spin follows from the fact that the source of gravitation is the stress-energy tensor, a second-rank tensor (compared to electromagnetism's spin-1 photon, the source of which is the four-current, a first-rank tensor). Additionally, it can be shown that any massless spin-2 field would give rise to a force indistinguishable from gravitation, because a massless spin-2 field must couple to (interact with) the stress-energy tensor in the same way that the gravitational field does. This result suggests that, if a massless spin-2 particle is discovered, it must be the graviton, so that the only experimental verification needed for the graviton may simply be the discovery of a massless spin-2 particle. [3]

What is the Spin?

So we know already that the new particle has spin zero or spin two and we could tell which one if we could detect the polarizations of the photons produced. Unfortunately this is difficult and neither ATLAS nor CMS are able to measure polarizations. The only direct and sure way to confirm that the particle is indeed a scalar is to plot the angular distribution of the photons in the rest frame of the centre of mass. A spin zero particles like the Higgs carries no directional information away from the original collision so the distribution will be even in all directions. This test will be possible when a much larger number of events have been observed. In the mean time we can settle for less certain indirect indicators.

The Casimir effect

The Casimir effect is related to the Zero-point energy, which is fundamentally related to the Heisenberg uncertainty relation. The Heisenberg uncertainty relation says that the minimum uncertainty is the value of the spin: $1/2 h = dx dp$ or $1/2 h = dt dE$, that is the value of the basic energy status.

The moving charges are accelerating, since only this way can self maintain the electric field causing their acceleration. The electric charge is not point like! This constant acceleration possible if there is a rotating movement changing the direction of the velocity. This way it can accelerate forever without increasing the absolute value of the velocity in the dimension of the time and not reaching the velocity of the light. In the atomic scale the Heisenberg uncertainty relation gives the same result, since the moving electron in the atom accelerating in the electric field of the proton, causing a charge distribution on delta x position difference and with a delta p momentum difference such a way that they product is about the half Planck reduced constant. For the proton this delta x much less in the nucleon, than in the orbit of the electron in the atom, the delta p is much higher because of the greater proton mass. This means that the electron is not a point like particle, but has a real charge distribution.

Electric charge and electromagnetic waves are two sides of the same thing; the electric charge is the diffraction center of the electromagnetic waves, quantified by the Planck constant h.

The Fine structure constant

The Planck constant was first described as the proportionality constant between the energy (E) of a photon and the frequency (ν) of its associated electromagnetic wave. This relation between the energy and frequency is called the **Planck relation** or the **Planck–Einstein equation**:

$$E = h\nu .$$

Since the frequency ν , wavelength λ , and speed of light c are related by $\lambda\nu = c$, the Planck relation can also be expressed as

$$E = \frac{hc}{\lambda}.$$

Since this is the source of Planck constant, the electric charge countable from the Fine structure constant. This also related to the Heisenberg uncertainty relation, saying that the mass of the proton should be bigger than the electron mass because of the difference between their wavelengths.

The expression of the fine-structure constant becomes the abbreviated

$$\alpha = \frac{e^2}{\hbar c}$$

This is a dimensionless constant expression, 1/137 commonly appearing in physics literature.

This means that the electric charge is a result of the electromagnetic waves diffractions, consequently the proton – electron mass rate is the result of the equal intensity of the corresponding electromagnetic frequencies in the Planck distribution law, described in my diffraction theory.

Path integral formulation of Quantum Mechanics

The path integral formulation of quantum mechanics is a description of quantum theory which generalizes the action principle of classical mechanics. It replaces the classical notion of a single, unique trajectory for a system with a sum, or functional integral, over an infinity of possible trajectories to compute a quantum amplitude. [7]

It shows that the particles are diffraction patterns of the electromagnetic waves.

Conclusions

The proposed topolaritons arise from the strong coupling of a photon and an exciton, a bound state of an electron and a hole. Their topology can be thought of as knots in their gapped energy-band structure. At the edge of the systems in which topolaritons emerge, these knots unwind and allow the topolaritons to propagate in a single direction without back-reflection. In other words, the topolaritons cannot make U-turns. Back-reflection is a known source of detrimental feedback and loss in photonic devices. The topolaritons' immunity to it may thus be exploited to build devices with increased performance. [12]

Solitons are localized wave disturbances that propagate without changing shape, a result of a nonlinear interaction that compensates for wave packet dispersion. Individual solitons may collide, but a defining feature is that they pass through one another and emerge from the collision unaltered in shape, amplitude, or velocity, but with a new trajectory reflecting a discontinuous jump. This remarkable property is mathematically a consequence of the underlying integrability of the one-dimensional (1D) equations, such as the nonlinear Schrödinger equation, that describe solitons in a variety of wave contexts, including matter waves^{1, 2}. Here we explore the nature of soliton collisions using Bose–Einstein condensates of atoms with attractive interactions confined to a quasi-1D waveguide. Using real-time imaging, we show that a collision between solitons is a complex event

that differs markedly depending on the relative phase between the solitons. By controlling the strength of the nonlinearity we shed light on these fundamental features of soliton collisional dynamics, and explore the implications of collisions in the proximity of the crossover between one and three dimensions where the loss of integrability may precipitate catastrophic collapse. [10]

"It's a photonic interaction that's mediated by the atomic interaction," Lukin said. "That makes these two photons behave like a molecule, and when they exit the medium they're much more likely to do so together than as single photons." To build a quantum computer, he explained, researchers need to build a system that can preserve quantum information, and process it using quantum logic operations. The challenge, however, is that quantum logic requires interactions between individual quanta so that quantum systems can be switched to perform information processing. [9]

The magnetic induction creates a negative electric field, causing an electromagnetic inertia responsible for the relativistic mass change; it is the mysterious Higgs Field giving mass to the particles. The Planck Distribution Law of the electromagnetic oscillators explains the electron/proton mass ratio by the diffraction patterns. The accelerating charges explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the electron's spin also, building the bridge between the Classical and Relativistic Quantum Theories. The self-maintained electric potential of the accelerating charges equivalent with the General Relativity space-time curvature, and since it is true on the quantum level also, gives the base of the Quantum Gravity. The electric currents causing self-maintaining electric potential is the source of the special and general relativistic effects. The Higgs Field is the result of the electromagnetic induction. The Graviton is two photons together.

References

- [1] http://www.academia.edu/3834454/3_Dimensional_String_Theory
- [2] http://www.academia.edu/3833335/The_Magnetic_field_of_the_Electric_current
- [3] http://www.academia.edu/4158863/Higgs_Field_and_Quantum_Gravity
- [4] http://www.academia.edu/4196521/The_Electro-Strong_Interaction
- [5] http://www.academia.edu/4221717/General_Weak_Interaction
- [6] The Feynman Lectures on Physics p. 274 (30.6)
Author: Richard Phillips Feynman
Publisher: Addison Wesley Longman (January 1970)
ISBN-10: **0201021153** | ISBN-13: **978-0201021158**
- [7] Path Integral Formulation of Quantum Mechanics
http://en.wikipedia.org/wiki/Path_integral_formulation
- [8] https://www.academia.edu/4215078/Accelerated_Relativity
- [9] <http://phys.org/news/2013-09-scientists-never-before-seen.html>

- [10] <http://www.nature.com/nphys/journal/vaop/ncurrent/full/nphys3135.html>
- [11] <http://www.sciencedaily.com/releases/2014/11/141102160109.htm>
- [12] <http://physics.aps.org/synopsis-for/10.1103/PhysRevX.5.031001>
- [13] Quantum dots enhance light-to-current conversion in layered semiconductors
<http://phys.org/news/2016-04-quantum-dots-light-to-current-conversion-layered.html>
- [14] 'Game-changer' for photonics applications: Researchers demonstrate record optical nonlinearity
<http://phys.org/news/2016-04-game-changer-photonics-applications-optical-nonlinearity.html>
- [15] Helically twisted photonic crystal fibres
<https://phys.org/news/2017-07-helically-photonic-crystal-fibres.html>
- [16] Nanoscale motion sends light into overdrive
<https://phys.org/news/2017-07-nanoscale-motion-overdrive.html>
- [17] Giant Enhancement of Electromagnetic Waves Revealed Within Small Dielectric Particles
<https://www.rdmag.com/news/2017/07/giant-enhancement-electromagnetic-waves-revealed-within-small-dielectric-particles>