




Article

# Rigorous proof for Riemann Hypothesis obtained by adopting Algebra-Geometry Approach in Geometric Langlands Program

John Yuk Ching Ting <sup>1,2,†,‡</sup>  0000-0002-3083-5071

<sup>1</sup> University of Tasmania, Churchill Avenue, Hobart TAS 7005, Australia; jycting@utas.edu.au

<sup>2</sup> Dental and Medical Surgery, 729 Albany Creek Road, Albany Creek QLD 4035, Australia; jycting@hotmail.com or jycting1@gmail.com

\* Correspondence: jycting@utas.edu.au or jycting1@gmail.com; Tel.: +614 1775 1859

† The correctness of this paper has been certified by local Australian mathematicians. Derived antiderivatives and mathematical arguments that were also present in the author's 2020-dated published research paper have all been confirmed to be correct and complete using computer algebra system Maxima. Previous use of exact and inexact Dimensional analysis homogeneity are adapted onto this paper.


‡ This paper is dedicated to the author's daughter Jelena prematurely born 13 weeks early on May 14, 2012 and all front-line health workers globally fighting against the deadly COVID-19 Pandemic. Declaration of interests: This work was supported by the author's January 20, 2020 AUS \$5,000 private research grant from Mrs. Connie Hayes and Mr. Colin Webb. He also received AUS \$3,250 and AUS \$510 reimbursements from Q-Pharm for participating in EyeGene Shingles trial on March 10, 2020 and Biosimilar Study on February 10, 2021.

**Abstract:** The 1859 Riemann hypothesis conjectured all nontrivial zeros in Riemann zeta function are uniquely located on  $\sigma = 1/2$  critical line. Derived from Dirichlet eta function [proxy for Riemann zeta function] are, in chronological order, simplified Dirichlet eta function and Dirichlet Sigma-Power Law. Computed Zeroes from the former uniquely occur at  $\sigma = 1/2$  resulting in total summation of fractional exponent ( $-\sigma$ ) that is twice present in this function to be integer  $-1$ . Computed Pseudo-zeroes from the later uniquely occur at  $\sigma = 1/2$  resulting in total summation of fractional exponent ( $1 - \sigma$ ) that is twice present in this law to be integer  $1$ . All nontrivial zeros are, respectively, obtained directly and indirectly as the one specific type of Zeroes and Pseudo-zeroes only when  $\sigma = 1/2$ . Thus, it is proved that Riemann hypothesis is true whereby this function and law rigidly comply with Principle of Maximum Density for Integer Number Solutions. The geometrical-mathematical [unified] approach used in our proof is equivalent to the algebra-geometry [unified] approach of geometric Langlands program that was formalized by Professor Peter Scholze and Professor Laurent Fargues. A succinct treatise on proofs for Polignac's and Twin prime conjectures is also outlined in this research paper.

**Keywords:** Coherent sheaf; Dirichlet Sigma-Power Law; Etale sheaf; Fargues-Fontaine curve; Geometric Langlands program; Gram's Law; Polignac's and Twin prime conjectures; Pseudo-zeroes; Riemann hypothesis; Rosser Rule; Zeroes

## Contents

19	<b>Introduction</b>	2
20	General notations and Figures 1, 2, 3 and 4 . . . . .	3
21	Equivalence of our unified geometrical-mathematical approach and the ap-	
22	proach of geometric Langlands program including p-adic Riemann zeta	
23	function $\zeta_p(s)$ . . . . .	3
24	<b>Sketch of the Proof for Riemann hypothesis including the Modified Equations</b>	
25	<b>for simplified Dirichlet eta function and Dirichlet Sigma-Power Law that are</b>	
26	<b>expressed using trigonometric identities</b>	6



27	<b>The Completely Predictable and Incompletely Predictable entities</b>	15
28	<b>The exact and inexact Dimensional analysis homogeneity for Equations</b>	16
29	<b>Gauss Circle Problem and Primitive Circle Problem</b>	17
30	<b>Gauss Areas of Varying Loops and Principle of Maximum Density for Integer</b>	
31	<b>Number Solutions</b>	18
32	<b>Shift of Varying Loops in <math>\zeta(\sigma + it)</math> Polar Graph and Principle of Equidistant for</b>	
33	<b>Multiplicative Inverse with General Equations for simplified Dirichlet eta</b>	
34	<b>function and Dirichlet Sigma-Power Law</b>	20
35	<b>Riemann zeta function, Dirichlet eta function, simplified Dirichlet eta function</b>	
36	<b>and Dirichlet Sigma-Power Law</b>	23
37	<b>Conclusions</b>	24
38	<b>References</b>	26
39	<b>Gram's Law and Rosser Rule for Gram points</b>	27
40	<b>Miscellaneous Materials</b>	28


## 41 1. Introduction

42 Riemann hypothesis is an intractable open problem in Number theory that was  
 43 proposed in 1859 by famous German mathematician Bernhard Riemann (September  
 44 17, 1826 - July 20, 1866). This hypothesis conjectured all nontrivial zeros in Riemann  
 45 zeta function are uniquely located on  $\sigma = \frac{1}{2}$  critical line. By applying Euler formula to  
 46 Dirichlet eta function [*proxy* for Riemann zeta function], we obtain simplified Dirichlet  
 47 eta function whereby its computed Zeroes uniquely occur at  $\sigma = \frac{1}{2}$  resulting in total  
 48 summation of fractional exponent ( $-\sigma$ ) that is twice present in this function to be integer  
 49  $-1$ . Dirichlet Sigma-Power Law is the solution from performing integration on simplified  
 50 Dirichlet eta function whereby its computed Pseudo-zeroes uniquely occur at  $\sigma = \frac{1}{2}$   
 51 resulting in total summation of fractional exponent  $(1 - \sigma)$  that is twice present in this  
 52 law to be integer 1.

53 All nontrivial zeros are, respectively, obtained directly and indirectly as one specific  
 54 type [out of three different types] of Zeroes and Pseudo-zeroes only when  $\sigma = \frac{1}{2}$ . Then  
 55 [non-existent] virtual nontrivial zeros and [non-existent] virtual Pseudo-nontrivial zeros  
 56 cannot be obtained directly and indirectly as a type of virtual Zeroes and virtual Pseudo-  
 57 zeroes when  $\sigma \neq \frac{1}{2}$ . As per Lemma 1 on these three different types of entities, all (virtual)  
 58 Pseudo-zeroes can be precisely converted to (virtual) Zeroes.

59 From fully solving Theorem 1, Corollary 2 and Theorem 3 (that contains Proposition  
 60 1 and Proposition 2); we confirm the following *sine qua non* statement to be true: "Valid  
 61 only at unique  $\sigma = \frac{1}{2}$  critical line, **geometrical** Origin intercept points in Figure 2 are  
 62 precisely equivalent to **mathematical** nontrivial zeros in Eq. (1) [directly] as Zeroes  
 63 and Eq. (3) [indirectly] as Pseudo-zeroes when expressed with using trigonometric  
 64 identities, and in Eq. (9) [directly] as Zeroes and Eq. (10) [indirectly] as Pseudo-zeroes  
 65 when expressed without using trigonometric identities". Thus, it is proved that Riemann  
 66 hypothesis is true whereby this function and law rigidly obey Principle of Maximum  
 67 Density for Integer Number Solutions. They additionally manifest Principle of Equidistant  
 68 for Multiplicative Inverse and are [serendipitously] amendable to treatment with  
 69 trigonometric identities.

70 Together with number theory, geometry and analysis; algebra is one of the broad  
 71 areas of mathematics. In its most general form forming the unifying thread of all mathe-  
 72 matics, algebra is the study of mathematical symbols and rules for manipulating these



**Figure 1.** INPUT for  $\sigma = \frac{1}{2}, \frac{2}{5},$  and  $\frac{3}{5}$ .  $\zeta(s)$  has countable infinite set of Completely Predictable trivial zeros located at  $\sigma =$  all negative even numbers and [conjectured] countable infinite set of Incompletely Predictable nontrivial zeros located at  $\sigma = \frac{1}{2}$  given by various  $t$  values.

73 symbols. Geometry is concerned with properties of space that are related with distance,  
 74 shape, size, and relative position of figures. We arbitrarily use the term ‘mathematical’  
 75 instead of ‘algebra’, and explain in subsection 1.2 the unified geometrical-mathematical  
 76 approach used in our proof of Riemann hypothesis [that essentially unites mathematics  
 77 and geometry] is essentially equivalent to algebra-geometry approach used by geomet-  
 78 ric Langlands program [that essentially unites algebra and geometry]. We provide an  
 79 assortment of information on various important topics although these need not form  
 80 an essential part of our proof for Riemann hypothesis: brief synopsis regarding Gram’s  
 81 Law and Rosser Rule for Gram points in Appendix A, and Miscellaneous Materials such  
 82 as on cardinality, certain types of infinite series, Zeroes and Pseudo-zeroes in Appendix  
 83 B. A succinct treatise on rigorous proofs for Polignac’s and Twin prime conjectures is  
 84 also outlined in the Conclusions section.


85 **1.1. General notations and Figures 1, 2, 3 and 4**

86 *The following is a short list of abbreviations used by this paper.*


- 87 **CFS:** countable finite set
- 88 **CIS:** countable infinite set
- 89 **UIS:** uncountable infinite set
- 90 **CP:** Completely Predictable – see section 3 on CP entities
- 91 **IP:** Incompletely Predictable – see section 3 on IP entities
- 92 **DA:** Dimensional analysis – see section 4 on exact and inexact DA homogeneity
- 93 **NTZ:** nontrivial zeros (Gram[ $x=0,y=0$ ] points) = Origin intercept points when  $\sigma = \frac{1}{2}$
- 94  $\zeta(s): f(n)$  Riemann zeta function containing variable  $n$ , and parameters  $t$  and  $\sigma$
- 95  $\eta(s): f(n)$  Dirichlet eta function containing variable  $n$ , and parameters  $t$  and  $\sigma$
- 96 **sim- $\eta(s)$ :**  $f(n)$  simplified Dirichlet eta function containing variable  $n$ , and parameters  $t$   
 97 and  $\sigma$
- 98 **DSPL:**  $F(n)$  Dirichlet Sigma-Power Law= $\int sim - \eta(s)dn$  containing variable  $n$ , and  
 99 parameters  $t$  and  $\sigma$

100 **1.2. Equivalence of our unified geometrical-mathematical approach and the approach of geometric**  
 101 **Langlands program including  $p$ -adic Riemann zeta function  $\zeta_p(s)$**


102 An L-function consists of a Dirichlet series with a functional equation and an  
 103 Euler product. Examples of L-functions come from modular forms, elliptic curves,  
 104 number fields, and Dirichlet characters, as well as more generally from automor-  
 105 phic forms, algebraic varieties, and Artin representations. They form an integrated  
 106 component of ‘L-functions and Modular Forms Database’ (LMFDB, located at URL



**Figure 2.** OUTPUT for  $\sigma = \frac{1}{2}$  as Gram points. Schematically depicted polar graph of  $\zeta(\frac{1}{2} + it)$  plotted along critical line for real values of  $t$  running from 0 to 34, horizontal axis:  $Re\{\zeta(\frac{1}{2} + it)\}$ , and vertical axis:  $Im\{\zeta(\frac{1}{2} + it)\}$ . Total presence of all Origin intercept points.



**Figure 3.** OUTPUT for  $\sigma = \frac{2}{5}$  as virtual Gram points. Varying Loops are shifted to left of Origin with horizontal axis:  $Re\{\zeta(\frac{2}{5} + it)\}$ , and vertical axis:  $Im\{\zeta(\frac{2}{5} + it)\}$ . Total absence of Origin intercept points.



**Figure 4.** OUTPUT for  $\sigma = \frac{3}{5}$  as virtual Gram points with horizontal axis:  $Re\{\zeta(\frac{3}{5} + it)\}$ , and vertical axis:  $Im\{\zeta(\frac{3}{5} + it)\}$ . Varying Loops are shifted to right of Origin. Total absence of Origin intercept points.

107 <https://www.lmfdb.org/>) with far-reaching implications. In proper perspective,  $\zeta(s)$  is  
 108 then the simplest example of an L-function.

109 The unified geometrical-mathematical approach used in our proof on Riemann  
 110 hypothesis that specifically involve only the [isolated]  $\zeta(s)$  as one type of L-function  
 111 must then be equivalent to the unified algebra-geometry approach of geometric Lang-  
 112 lands program that generally involve all types of L-functions. Named after German  
 113 mathematician Adolf Hurwitz (March 26, 1859 - November 18, 1919), Hurwitz zeta  
 114 function is one of the many zeta functions. It is formally defined for complex arguments  
 115  $s$  with  $\text{Re}(s) > 1$  and  $q$  with  $\text{Re}(q) > 0$  by  $\zeta(s, q) = \sum_{n=0}^{\infty} \frac{1}{(n+q)^s}$ . This series is absolutely  
 116 convergent for given values of  $s$  and  $q$ , and can be extended to a meromorphic function  
 117 defined for all  $s \neq 1$ . With this scheme, our Riemann zeta function  $\zeta(s)$  is equivalently  
 118 given as  $\zeta(s, 1)$ .

119 Using the innovative research method of p-adic analysis popularized by renowned  
 120 German mathematician Professor Peter Scholze who won the 2018 Fields Medal; a p-adic  
 121 zeta function, or more generally a p-adic L-function, is a function analogous to Riemann  
 122 zeta function, or more general L-functions, but whose domain and target are p-adic  
 123 (where  $p$  is a prime number). In p-adic Riemann zeta function  $\zeta_p(s)$ , values at negative  
 124 odd integers are those of Riemann zeta function  $\zeta(s)$  at negative odd integers (up to  
 125 an explicit correction factor). The p-adic L-functions arising in this fashion as sourced  
 126 from p-adic interpolation[1] of special values of L-functions are typically referred to as  
 127 analytic p-adic L-functions. The other major source of p-adic L-functions is from the  
 128 arithmetic of cyclotomic fields, or more generally, certain Galois modules over towers of  
 129 cyclotomic fields or even more general towers.

130 There is no clear delineation between algebra and analysis: The involved mathe-  
 131 matics is considered more "algebraic" if it focuses more on structure and interaction of  
 132 operations that underlie the objects of study e.g. groups, rings, fields, etc. The involved  
 133 mathematics is considered more "analytic" if it focuses more on real numbers and mea-  
 134 surable quantities, and the approximation and computation thereof e.g. calculus, Taylor  
 135 series, derivatives, integrals, etc. Galois groups arise in the branch of mathematics called  
 136 algebra (reflecting the way we use algebra to solve equations), and automorphic forms  
 137 arise in the different branch of mathematics called analysis (which can be considered as  
 138 an enhanced form of calculus).

139 Formulated by renowned Canadian mathematician Robert Langlands in late 1960s,  
 140 Langlands correspondence classically refers to collection of results and conjectures  
 141 relating number theory and representation theory. Langlands conjecture for rational  
 142 numbers is further referred to as "global" Langlands correspondence [since rational  
 143 number system contain all prime numbers], and for p-adics as "local" Langlands corre-  
 144 spondence [since p-adic number systems deal with one prime number at a time]. The  
 145 coined *geometric Langlands program* is a reformulation of Langlands correspondence  
 146 obtained by replacing number fields appearing in original number theoretic version by  
 147 function fields and applying techniques from algebraic geometry, thus relating algebraic  
 148 geometry to representation theory. The aim is to find geometric objects with properties  
 149 that could stand in for Galois groups and automorphic forms in Langlands' conjectures.  
 150 The perfectoid spaces are adic spaces of special kind occurring in the study of problems  
 151 of "mixed characteristic" such as local fields of characteristic zero which have residue  
 152 fields of characteristic prime  $p$ . Based on p-adic geometry, Professor Scholze's 2012 PhD  
 153 thesis on perfectoid spaces[2] yields solution to a special case of weight-monodromy  
 154 conjecture.

155 Named after renowned French mathematicians Professor Laurent Fargues and  
 156 Professor Jean-Marc Fontaine (March 13, 1944 - January 29, 2019), Fargues-Fontaine curve  
 157 as a geometric object is a curve whose points each represented a version of an important  
 158 object called a p-adic ring. Professor Fargues and Professor Scholze subsequently came  
 159 up with two different kinds of more complicated geometric objects called sheaves:

160 coherent sheaves correspond to representations of p-adic groups, and étale sheaves to  
 161 representations of Galois groups. In their paper[3] with Fargues-Fontaine curve now  
 162 merging with Scholze's p-adic geometry, they develop the foundations of geometric  
 163 Langlands program whereby it is proved that there is always a way to match a coherent  
 164 sheaf with an étale sheaf, and as a result there is always a way to match a representation  
 165 of a p-adic group with a representation of a Galois group. In this ground-breaking way  
 166 of studying "local" Langlands correspondence based on these geometric objects called  
 167 sheaves, they finally proved the one direction of translation for this correspondence  
 168 although the other direction of translation remains an open question. This is the basic  
 169 premise of Langlands program which is a broad vision for investigating Galois groups –  
 170 essentially polynomials – through these types of translations.

171 Finally, since the infinitely many prime numbers  $\geq 2$  are a subset of the infinitely  
 172 many integers  $\geq 1$ ; we can derive the following alternative equally valid deductions  
 173 involving (positive) prime number system instead of the valid deductions as outlined in  
 174 Proposition 1 involving (positive) [non-prime number] integer number system: "Only  
 175 at  $\sigma = \frac{1}{2}$  critical line which involves applying  $f(n)$  as fractional exponent  $\frac{1}{2}$  or square  
 176 root on  $n =$  all perfect squares of prime numbers 4, 9, 25, 49, 121, 169, 289, 361, 529,  
 177 841... will we obtain maximum number of rational roots as consecutive prime number  
 178 solutions 2, 3, 5, 7, 11, 13, 17, 19, 23, 29... (viz, all prime numbers  $\geq 2$ ). This observation  
 179 uniquely comply with **Principle of Maximum Density for Prime Number Solutions** at  
 180  $\sigma = \frac{1}{2}$  critical line." We immediately recognize from above commentaries that using this  
 181 **Principle of Maximum Density for Prime Number Solutions** instead of **Principle of**  
 182 **Maximum Density for Integer Number Solutions** from Proposition 1 will also crucially  
 183 confer the proof for Theorem 3 to be fully complete.

## 184 **2. Sketch of the Proof for Riemann hypothesis including the Modified Equations for** 185 **simplified Dirichlet eta function and Dirichlet Sigma-Power Law that are expressed** 186 **using trigonometric identities**

187 Symbolically named after German mathematician Gustav Lejeune Dirichlet (Febru-  
 188 ary 13, 1805 - May 5, 1859), the word "Law" in DSPL represent a convenient terminology  
 189 to describe this function – viz, there is resemblance to Zipf's law via power law functions  
 190 in  $\sigma$  from  $s = \sigma + it$  being exponent of a power function as similar format to  $n^\sigma$ , logarithm  
 191 scale use, and  $\zeta(s)$  harmonic series connection. Respectively, we use Zeroes (as three  
 192 types of Gram points) and Pseudo-zeroes (as three types of Pseudo-Gram points) at  
 193  $\sigma = \frac{1}{2}$  to collectively refer to corresponding  $f(n)$ 's and  $F(n)$ 's x-axis intercept points,  
 194 y-axis intercept points and Origin intercept points. Respectively, we use virtual Zeroes  
 195 (as two types of virtual Gram points) and virtual Pseudo-zeroes (as two types of virtual  
 196 Pseudo-Gram points) at  $\sigma \neq \frac{1}{2}$  to collectively refer to corresponding  $f(n)$ 's and  $F(n)$ 's  
 197 x-axis intercept points and y-axis intercept points [with absent Origin intercept points].

198 *Geometrical and mathematical definitions for Gram points and virtual Gram points.* Figure  
 199 1 depicts complex variable  $s (= \sigma \pm it)$  as INPUT with x-axis denoting real part  $\text{Re}\{s\}$   
 200 associated with  $\sigma$ , and y-axis denoting imaginary part  $\text{Im}\{s\}$  associated with  $t$ . The  
 201 critical line:  $\sigma = \frac{1}{2}$ ; non-critical lines:  $\sigma \neq \frac{1}{2}$  viz,  $0 < \sigma < \frac{1}{2}$  and  $\frac{1}{2} < \sigma < 1$ ; and critical  
 202 strip:  $0 < \sigma < 1$ . Both the unique  $\sigma = \frac{1}{2}$  value and the non-unique  $\sigma \neq \frac{1}{2}$  values  $\in$  Set  
 203 **all  $\sigma$  values** whereby Set **all  $\sigma$  values** =  $\sigma \mid \sigma$  is a real number, and  $0 < \sigma < 1$ . With  
 204 including its complex conjugate,  $s = \sigma \pm it$  is present in our chosen  $f(n)$  and  $F(n)$  whereby  
 205 these are well-defined continuous [complex] functions that are always defined for any  
 206 arbitrarily chosen intervals  $[a,b]$ . With  $f(n) = 0$  and  $F(n) = 0$  giving rise to relevant derived  
 207 equations that are *dependently*-related [via Varying Loops], they generate corresponding  
 208 types of IP entities. These IP entities will inherently belong to the correctly assigned  
 209 *mutually exclusive CIS of Gram points and virtual Gram points* constituted by  $t$  values as  
 210 transcendental numbers except for first Gram $[y=0]$  point (and first virtual Gram $[y=0]$   
 211 point) given by  $t = 0$ . Origin intercept points, x-axis intercept points and y-axis intercept  
 212 points are geometrical definitions for IP entities of Gram $[x=0,y=0]$  points, Gram $[y=0]$

213 points and Gram[x=0] points at  $\sigma = \frac{1}{2}$ . These geometrical definitions are equivalent to  
 214 mathematical definitions as given by the equations below in this section.

215 Origin intercept points at  $\sigma = \frac{1}{2}$  consisting of Gram[x=0,y=0] points or NTZ are  
 216 computed directly from equations  $\eta(s) = 0$  and  $\text{sim-}\eta(s) = 0$ ; and indirectly from equation  
 217 DSPL = 0. x-axis intercept points at  $\sigma = \frac{1}{2}$  consisting of Gram[y=0] points or (traditional)  
 218 'usual' Gram points are computed directly from equation Gram[y=0] points- $\text{sim-}\eta(s) =$   
 219 0; and indirectly from equation Gram[y=0] points-DSPL = 0. y-axis intercept points at  
 220  $\sigma = \frac{1}{2}$  consisting of Gram[x=0] points are computed directly from equation Gram[x=0]  
 221 points- $\text{sim-}\eta(s) = 0$ ; and indirectly from equation Gram[x=0] points-DSPL = 0.

222 Relevant functions and equations are unique mathematical objects usefully clas-  
 223 sified as three types of infinite series: *Harmonic series*, *Alternating harmonic series* or  
 224 *Alternating series with trigonometric terms*. We perform crucial *de novo* analysis on these  
 225 functions and equations by noting their manifested intrinsic properties. Without loss  
 226 of validity in our correct and complete set of mathematical arguments, we adopt the  
 227 convention of providing focused analysis predominantly on appropriately chosen Alter-  
 228 nating series with trigonometric terms throughout our presentation. The complex  $f(n)$   
 229  $\zeta(s)$  is a Harmonic series that does not converge in critical strip. The complex  $f(n)$   $\eta(s)$  is  
 230 an Alternating harmonic series that converge in critical strip. Through analytic continua-  
 231 tion,  $\eta(s)$  must act as *proxy* function for  $\zeta(s)$  in this strip. [Caveat: the limit of an analytic  
 232 continuation is not the analytic continuation of the limit.] Derived as Euler formula  
 233 application to  $\eta(s)$  is the complex  $f(n)$   $\text{sim-}\eta(s)$ , and derived as  $\int \text{sim} - \eta(s) dn$  is the  
 234 complex  $F(n)$  DSPL. Both  $\text{sim-}\eta(s)$  and DSPL are Alternating series with trigonometric  
 235 terms that converge in critical strip.

236 The  $f(n)$   $\eta(s)$  will converge infinitely often to a zero value as  $\eta(s) = 0$  equation  
 237 giving rise to all NTZ or Gram[x=0,y=0] points. This event will only happen when  $\eta(s)$   
 238 is substituted with one unique  $\sigma$  value which is conjectured to be  $\sigma = \frac{1}{2}$  by Riemann  
 239 hypothesis. Being an Alternating harmonic series [without trigonometric terms that  
 240 graphically cater for all possible types of x-axis and y-axis intercept points], we inherently  
 241 cannot derive valid functions to obtain corresponding equations Gram[y=0] points-  
 242  $\eta(s) = 0$  and Gram[x=0] points- $\eta(s) = 0$  that will enable mathematical computations of  
 243 Gram[y=0] points as x-axis intercept points and Gram[x=0] points as y-axis intercept  
 244 points. Then, computed Zeroes are mathematically defined as  $\eta(s) = 0$  and  $\text{sim-}\eta(s) = 0$   
 245 when parameter  $\sigma = \frac{1}{2}$ ; computed virtual Zeroes are mathematically defined as  $\eta(s) \neq 0$   
 246 and  $\text{sim-}\eta(s) = 0$  when parameter  $\sigma \neq \frac{1}{2}$ ; computed Pseudo-Zeroes are mathematically  
 247 defined as DSPL = 0 when parameter  $\sigma = \frac{1}{2}$ ; and computed virtual Pseudo-zeroes are  
 248 mathematically defined as DSPL = 0 when parameter  $\sigma \neq \frac{1}{2}$ .

249 For  $0 \leq \delta \leq 1$ , let  $f(n) = \sin(n) \pm \delta$  and  $f(n) = \cos(n) \pm \delta$  represent two [simple]  
 250 trigonometric functions which are periodic transcendental-type functions. Both  $\sin(n) \pm$   
 251  $\delta = 0$  and  $\cos(n) \pm \delta = 0$  as equations will generate infinitely many CP x-axis intercept  
 252 points (Zeroes) for any given values of  $\delta$ . This will additionally include the solitary  
 253 Origin intercept point (Zero) obtained from  $\sin(n) \pm \delta = 0$  when  $\delta = 0$ . For both  
 254  $\sin(n) \pm \delta$  and  $\cos(n) \pm \delta$ , only when  $\delta = 0$  will their progressive / cumulative *Areas*  
 255 *Above the horizontal axis* be overall identical to *Areas Below the horizontal axis*. Otherwise,  
 256 these mentioned *Areas* will not be overall identical to each other when  $\delta \neq 0$ . We  
 257 now provide analogical reasoning for existence of infinitely many substituted  $\sigma$  values  
 258 (including  $\sigma = \frac{1}{2}$ ) that will all contribute to two conditions  $\text{sim-}\eta(s) = 0$  and DSPL = 0  
 259 being satisfied while simultaneously giving rise to (i) IP Zeroes and IP Pseudo-zeroes  
 260 [when  $\sigma = \frac{1}{2}$ ], and (ii) IP virtual Zeroes and IP virtual Pseudo-zeroes [when  $\sigma \neq \frac{1}{2}$ ]. With  
 261 (complex) sine and/or cosine terms present in  $f(n)$   $\text{sim-}\eta(s)$  and  $F(n)$  DSPL also being  
 262 periodic transcendental-type functions, we intuitively deduce  $\sigma = \frac{1}{2}$  and  $\sigma \neq \frac{1}{2}$  must  
 263 respectively act as the analogical equivalence of  $\delta = 0$  and  $\delta \neq 0$ . This deduction allows  
 264 intuitive and valid explanations for our two conditions to be satisfied by the infinitely  
 265 many substituted  $\sigma$  values. Consequently, we must rigorously prove additional property  
 266 of  $\text{sim-}\eta(s)$  and DSPL that they will characteristically, inevitably and uniquely comply

267 with Principle of Maximum Density for Integer Number Solutions only when  $\sigma = \frac{1}{2}$   
 268 with this Principle signifying complete presence of NTZ in  $\text{sim-}\eta(s)$  or Pseudo-NTZ in  
 269 DSPL as one unique type of Gram points or Pseudo-Gram points [which are otherwise  
 270 totally absent when  $\sigma \neq \frac{1}{2}$ ].

271 Figures 2, 3 and 4 are  $\zeta(\sigma + it)$  Polar Graphs [see Remark 10 on intimate relationship  
 272 between Cartesian Coordinates and Polar Coordinates] with x-axis denoting real part  
 273  $\text{Re}\{\zeta(s)\}$  and y-axis denoting imaginary part  $\text{Im}\{\zeta(s)\}$  generated by  $\zeta(s)$ 's output as real  
 274 values of  $t$  running from 0 to 34. There are infinite types-of-spirals (Varying Loops)  
 275 possibilities associated with each  $\sigma$  value arising from all infinite  $\sigma$  values in  $0 < \sigma < 1$   
 276 critical strip whereby the unique and solitary  $\sigma = \frac{1}{2}$  value that denote critical line  
 277 is located in this strip. We observe that Figure 3 [with  $\sigma = \frac{2}{5}$ ] and Figure 4 [with  
 278  $\sigma = \frac{3}{5}$ ] show associated shifts of Varying Loops that manifest Principle of Equidistant  
 279 for Multiplicative Inverse – see Proposition 2 from section 7. From observing Figure 2,  
 280 we can geometrically define NTZ (or Gram $[x=0,y=0]$  points) as Origin intercept points  
 281 occurring when  $\sigma = \frac{1}{2}$ . Then, two remaining types of Gram points as part of continuous  
 282 Varying Loops are consequently defined as x-axis intercept points and y-axis intercept  
 283 points occurring when  $\sigma = \frac{1}{2}$ .

284 Lemma 2 confirms the paired IP two types of Gram points [as Zeroes] situation,  
 285 paired IP two types of virtual Gram points [as virtual Zeroes] situation, paired IP two  
 286 types of Pseudo-Gram points [as Pseudo-zeroes] situation, and paired IP two types of  
 287 virtual Pseudo-Gram points [as virtual Pseudo-zeroes] situation are always  $\frac{1}{2}\pi$  out-of-  
 288 phase with each other in every one of these situations. Lemma 1 confirms IP Zeroes,  
 289 IP virtual Zeroes, IP Pseudo-zeroes and IP virtual Pseudo-zeroes are precisely related  
 290 as  $\frac{1}{2}\pi$  (for NTZ case) or  $\frac{3}{4}\pi$  (for Gram $[y=0]$  points and Gram $[x=0]$  points cases) out-  
 291 of-phase with each other. Thus from Lemma 1, corresponding three types of  $F(n)$ 's  
 292 Pseudo-zeroes or Pseudo-Gram points and two types of  $F(n)$ 's virtual Pseudo-zeroes or  
 293 virtual Pseudo-Gram points can be precisely converted to three types of  $f(n)$ 's Zeroes  
 294 or Gram points and two types of  $f(n)$ 's virtual Zeroes or virtual Gram points. Then,  
 295 Statement (I) – (IV) are valid whereby  $\sigma = \frac{1}{2}$ 's derived entities from Statement (III) can  
 296 be precisely converted to those from Statement (I), and  $\sigma \neq \frac{1}{2}$ 's derived virtual entities  
 297 from Statement (IV) can be precisely converted to those from Statement (II):

298 Statement (I) The  $f(n)$ 's Zeroes at  $\sigma = \frac{1}{2}$  [directly] equates to three types of Gram  
 299 points.

300 Statement (II) The  $f(n)$ 's virtual Zeroes at  $\sigma \neq \frac{1}{2}$  [directly] equates to two types of  
 301 virtual Gram points.

302 Statement (III) The  $F(n)$ 's Pseudo-zeroes at  $\sigma = \frac{1}{2}$  [indirectly] equates to three types  
 303 of Gram points.

304 Statement (IV) The  $F(n)$ 's virtual Pseudo-zeroes at  $\sigma \neq \frac{1}{2}$  [indirectly] equates to two  
 305 types of virtual Gram points.

306  
 307 **Remark 1.** Of particular relevance to Riemann hypothesis, we mathematically de-  
 308 duce from above materials that  $f(n)$ 's NTZ or Gram $[x=0,y=0]$  points as one type of Gram  
 309 points will conjecturally only exist at unique  $\sigma = \frac{1}{2}$  critical line [but not at non-unique  
 310  $\sigma \neq \frac{1}{2}$  non-critical lines]. This can be equivalently stated as:  $F(n)$ 's Pseudo-NTZ or  
 311 Pseudo-Gram $[x=0,y=0]$  points as one type of Pseudo-Gram points will conjecturally only  
 312 exist at unique  $\sigma = \frac{1}{2}$  critical line [but not at non-unique  $\sigma \neq \frac{1}{2}$  non-critical lines].

313  
 314 Useful analogy for Remark 2: A line consists of infinitely many points. Graphically,  
 315 the Origin is a zero-dimensional [single] point; x-axis or horizontal axis and y-axis or  
 316 vertical axis are one-dimensional lines [containing infinitely many points].

317  
 318 **Remark 2.** In Figure 3 and Figure 4, we note Origin intercept points as Gram $[x=0,y=0]$   
 319 points or NTZ cannot exist when  $\sigma \neq \frac{1}{2}$ . In Figure 2, we note Origin intercept points as



320 Gram[ $x=0,y=0$ ] points or NTZ only exist when  $\sigma = \frac{1}{2}$ . Of particular relevance to Riemann  
 321 hypothesis, we deduce  $\text{sim-}\eta(s)$  as periodic transcendental-type function only contain  
 322 one solitary  $\sigma$ -valued type of Origin intercept points (when  $\sigma = \frac{1}{2}$  for Gram[ $x=0,y=0$ ]  
 323 points or NTZ as conjectured by Riemann hypothesis) but infinitely many different  
 324  $\sigma$ -valued types of x-axis intercept points and y-axis intercept points (constituted by  
 325 solitary  $\sigma = \frac{1}{2}$  value for Gram[ $y=0$ ] points and Gram[ $x=0$ ] points as well as infinitely  
 326 many  $\sigma \neq \frac{1}{2}$  values for virtual Gram[ $y=0$ ] points and virtual Gram[ $x=0$ ] points). We can  
 327 conjure up an equivalent statement for DSPL as periodic transcendental-type function  
 328 whereby we replace NTZ and (virtual) Gram points with their counterparts Pseudo-NTZ  
 329 and (virtual) Pseudo-Gram points.

330

331 We can now propose Theorem 1 (with  $\sigma = \frac{1}{2}$  connoting exact DA homogeneity) and  
 332 Corollary 2 (with  $\sigma \neq \frac{1}{2}$  connoting inexact DA homogeneity) to fully represent Remark 1  
 333 and Remark 2. Their successful proofs will firstly, denote rigorous proof for Riemann  
 334 hypothesis that involves conjecture on location of NTZ as one type of Gram points  
 335 [viz, Origin intercept points] and secondly, provide precise explanations for remaining  
 336 two types of Gram points [viz, x-axis intercept points and y-axis intercept points]. In  
 337 addition, we incorporate Theorem 3 on rigid compliance by  $\text{sim-}\eta(s)$  and DSPL with  
 338 Principle of Maximum Density for Integer Number Solutions whereby its successful  
 339 proof will only eventuate when  $\sigma = \frac{1}{2}$ .

340

341 **Theorem 1.** Rigidly complying with exact DA homogeneity,  $f(n)$   $\text{sim-}\eta(s)$  and  $F(n)$   
 342 DSPL as relevant equations can incorporate three types of Gram points and Pseudo-  
 343 Gram points onto solitary  $\sigma = \frac{1}{2}$  critical line thus fully supporting Riemann hypothesis  
 344 to be true.

345

346 **Proof.** Using  $f(n)$   $\text{sim-}\eta(s)$  and  $F(n)$  DSPL, Riemann hypothesis propose all NTZ  
 347 are located on  $\sigma = \frac{1}{2}$  critical line in these functions. The three types of Gram points  
 348 and Pseudo-Gram points are each infinite in magnitude consisting of mutually exclu-  
 349 sive entities. Amounting to direct Proof by Positive, we show CIS of Gram[ $x=0,y=0$ ]  
 350 points or NTZ constitutes one type of Gram points only when  $\sigma = \frac{1}{2}$  thus fully sup-  
 351 porting Riemann hypothesis to be true. The preceding sentence is equally valid when  
 352 we replace Gram[ $x=0,y=0$ ] points, NTZ and Gram points with corresponding Pseudo-  
 353 Gram[ $x=0,y=0$ ] points, Pseudo-NTZ and Pseudo-Gram points. Respectively, the conven-  
 354 iently defined term of exact DA homogeneity denote [exact] integer  $-1$  and  $1$  derived  
 355 from  $\sum(\text{all fractional exponents}) = 2(-\sigma)$  and  $2(1 - \sigma)$ . These act as surrogate markers  
 356 in  $\text{sim-}\eta(s)$  and DSPL on [solitary]  $\sigma = \frac{1}{2}$  situation. Generated by relevant functions  
 357 and laws when  $\sigma = \frac{1}{2}$ , the three types of Gram points are mathematically defined as  
 358 equations  $\text{sim-}\eta(s) = 0$ , Gram[ $y=0$ ] points- $\text{sim-}\eta(s) = 0$  and Gram[ $x=0$ ] points- $\text{sim-}\eta(s)$   
 359  $= 0$ ; and the three types of Pseudo-Gram points are mathematically defined as equa-  
 360 tions DSPL = 0, Gram[ $y=0$ ] points-DSPL = 0 and Gram[ $x=0$ ] points-DSPL = 0. They all  
 361 correspond to relevant geometrically defined Origin intercept points, x-axis intercept  
 362 points and y-axis intercept points. Thus, three types of IP Gram points [IP Zeroes] and IP  
 363 Pseudo-Gram points [IP Pseudo-Zeroes] are mathematically and geometrically defined  
 364 to be located on  $\sigma = \frac{1}{2}$  critical line. Based solely on these definitive definitions, we can  
 365 uniquely incorporate three types of IP Gram points [IP Zeroes] and IP Pseudo-Gram  
 366 points [IP Pseudo-zeroes] onto  $\sigma = \frac{1}{2}$  critical line. *The proof is now complete for Theorem 1*  $\square$ .

366

367 **Corollary 2.** Rigidly complying with inexact DA homogeneity,  $f(n)$   $\text{sim-}\eta(s)$  and  
 368  $F(n)$  DSPL as relevant equations can incorporate two types of virtual Gram points and  
 369 virtual Pseudo-Gram points onto infinitely many  $\sigma \neq \frac{1}{2}$  non-critical lines thus also fully  
 370 supporting Riemann hypothesis to be true.

371

372 **Proof.** Using  $f(n)$   $\text{sim-}\eta(s)$  and  $F(n)$  DSPL, Riemann hypothesis equivalently pro-  
 373 pose all NTZ are not located on  $\sigma \neq \frac{1}{2}$  non-critical lines in these functions. The two types  
 of virtual Gram points and virtual Pseudo-Gram points are each infinite in magnitude

374 consisting of mutually exclusive entities. Amounting to indirect Proof by Contrapositive,  
 375 we show [non-existent] virtual Gram $[x=0,y=0]$  points or virtual NTZ will not constitute  
 376 one type of [non-existent] virtual Gram points when  $\sigma \neq \frac{1}{2}$  thus also fully supporting  
 377 Riemann hypothesis to be true. The preceding sentence is equally valid when we replace  
 378 virtual Gram $[x=0,y=0]$  points, virtual NTZ and virtual Gram points with corresponding  
 379 virtual Pseudo-Gram $[x=0,y=0]$  points, virtual Pseudo-NTZ and virtual Pseudo-Gram  
 380 points. Respectively, conveniently defined term of inexact DA homogeneity denote  
 381 [inexact] fractional (non-integer) number  $\neq -1$  and  $\neq 1$  derived from  $\sum$ (all fractional  
 382 exponents) =  $2(-\sigma)$  and  $2(1 - \sigma)$ . These act as surrogate markers in sim- $\eta(s)$  and DSPL  
 383 on [infinitely many]  $\sigma \neq \frac{1}{2}$  situations. Generated by relevant functions and laws when  
 384  $\sigma \neq \frac{1}{2}$ , the two types of virtual Gram points are mathematically defined as equations  
 385 virtual Gram $[y=0]$  points-sim- $\eta(s) = 0$  and virtual Gram $[x=0]$  points-sim- $\eta(s) = 0$ ; and  
 386 the two types of virtual Pseudo-Gram points are mathematically defined as equations  
 387 virtual Gram $[y=0]$  points-DSPL = 0 and virtual Gram $[x=0]$  points-DSPL = 0. They all  
 388 correspond to relevant geometrically defined x-axis intercept points and y-axis intercept  
 389 points. Thus, two types of IP virtual Gram points [IP virtual Zeroes] and IP virtual  
 390 Pseudo-Gram points [IP virtual Pseudo-Zeroes] are mathematically and geometrically  
 391 defined to be located on  $\sigma \neq \frac{1}{2}$  non-critical lines. Based solely on these definitive defini-  
 392 tions, we can uniquely incorporate two types of IP virtual Gram points [IP virtual Zeroes]  
 393 and IP virtual Pseudo-Gram points [IP virtual Pseudo-zeroes] onto  $\sigma \neq \frac{1}{2}$  non-critical  
 394 lines. *The proof is now complete for Corollary 2*□.

395  
 396 **Theorem 3.** Conforming to the solitary  $\sigma = \frac{1}{2}$  critical line [and not the infinitely  
 397 many  $\sigma \neq \frac{1}{2}$  non-critical lines e.g.  $\sigma = \frac{1}{3}$  or  $\frac{2}{3}$ ] whereby  $\sigma$  forms part of relevant fractional  
 398 exponents from base quantities  $(2n)$  and  $(2n-1)$  in sim- $\eta(s)$  [as Riemann sum  $\Delta n \rightarrow 1$   
 399 with variable n involving all integers  $\geq 1$ ] or DSPL [as definite integral  $\Delta n \rightarrow 0$  with  
 400 variable n involving all real numbers  $\geq 1$ ]; square roots of perfect squares [and not  
 401 e.g. cube roots of perfect cubes or squared cube roots of perfect cubes] when applied to  
 402 combined base quantities  $(2n)$  and  $(2n-1)$  in sim- $\eta(s)$  or DSPL will generate the maximum  
 403 number of integer solutions (constituted by all integers  $\geq 1$ ) that uniquely comply with  
 404 Principle of Maximum Density for Integer Number Solutions while also manifesting  
 405 Principle of Equidistant for Multiplicative Inverse.

406 **Proof.**  $\int \text{sim-}\eta(s)dn = \text{DSPL}$ . Whereas the two subsets of rational roots as integers  
 407 and irrational roots as irrational numbers can be generated by combined base quantities  
 408  $(2n)$  and  $(2n-1)$  from sim- $\eta(s)$  [as Riemann sum  $\Delta n \rightarrow 1$  with variable n involving all  
 409 integers  $\geq 1$ ], so must these two exact same subsets be generated by combined base  
 410 quantities  $(2n)$  and  $(2n-1)$  from DSPL [as definite integral  $\Delta n \rightarrow 0$  with variable n  
 411 involving all real numbers  $\geq 1$ ]. Thus in sim- $\eta(s)$  or DSPL, its computed CIS rational  
 412 roots (subset) as integers [rational numbers] + computed CIS irrational roots (subset) as  
 413 irrational numbers = computed CIS total roots. These two mutually exclusive subsets  
 414 belong to UIS real numbers. Using subset rational roots as integers at  $\sigma = \frac{1}{2}$  critical line,  
 415 and by comparing and contrasting this subset with [different] subset rational roots as  
 416 integers at  $\sigma = \frac{1}{3}$  or  $\frac{2}{3}$  non-critical lines corollary situation; we will show that square  
 417 roots of perfect squares [and not e.g. cube roots of perfect cubes or squared cube roots of  
 418 perfect cubes] when applied to combined base quantities  $(2n)$  and  $(2n-1)$  from sim- $\eta(s)$   
 419 or DSPL giving rise to maximum number of integer solutions (constituted by all integers  
 420  $\geq 1$ ) must uniquely comply with Principle of Maximum Density for Integer Number  
 421 Solutions (see Proposition 1 in section 6) while also manifesting Principle of Equidistant  
 422 for Multiplicative Inverse (see Proposition 2 in section 7). We apply concepts from  
 423 elegant Gauss Circle Problem and Primitive Circle Problem in section 5 onto materials  
 424 on aptly-named Gauss Areas of Varying Loops to justifiably obtain correct and complete  
 425 set of mathematical arguments that fully support Theorem 3. *The proof is now complete for*  
 426 *Theorem 3*□.

427

428 By conveniently employing only  $\text{sim-}\eta(s)$  for analysis here [with analysis using  
 429 DSPL being equally valid], Theorem 1 and Corollary 2 above can also be insightfully  
 430 combined as follows. Let Set  $\mathbf{G}$  = all Gram points =  $\text{Gram}[x=0,y=0]$  points +  $\text{Gram}[y=0]$   
 431 points +  $\text{Gram}[x=0]$  points and Set  $\mathbf{vG}$  = all virtual Gram points = virtual  $\text{Gram}[y=0]$   
 432 points + virtual  $\text{Gram}[x=0]$  points with virtual  $\text{Gram}[x=0,y=0]$  points = null set  $\emptyset$ . We can  
 433 apply **inclusion-exclusion principle**  $|\mathbf{G} \cup \mathbf{vG}| = |\mathbf{G}| + |\mathbf{vG}| - |\mathbf{G} \cap \mathbf{vG}| = |\mathbf{G}| + |\mathbf{vG}|$   
 434 because  $|\mathbf{G} \cap \mathbf{vG}| = 0$ . Since exclusive presence of Gram points and absence of virtual  
 435 Gram points on critical line denotes exclusive absence of Gram points and exclusive  
 436 presence of virtual Gram points on non-critical lines; then Gram points and virtual  
 437 Gram points as mutually exclusive entities must mathematically and geometrically be  
 438 incorporated, respectively, onto unique (solitary) critical line and non-unique (infinitely  
 439 many) non-critical lines of  $\text{sim-}\eta(s)$ .

440 **Derived  $f(n) = 0$  and  $F(n) = 0$  equations** – see  $\sigma = \frac{1}{2}$  (via Proposition 4.3 and Proposition  
 441 5.3) and  $\frac{2}{5}$  (via Corollary 4.4 and Corollary 5.4) representative examples given in [4], p. 27-28,  
 442 29-30 and section 4 below – comply with exact DA homogeneity at  $\sigma = \frac{1}{2}$  critical line and inexact  
 443 DA homogeneity at  $\sigma \neq \frac{1}{2}$  non-critical lines. NTZ are synonymous with  $\text{Gram}[x=0,y=0]$   
 444 points which is one type of Gram points. Whenever applicable, all modified equations  
 445 below are expressed using trigonometric identities. Together with  $\text{Gram}[y=0]$  points  
 446 and  $\text{Gram}[x=0]$  points as remaining two types of Gram points, these three types of Gram  
 447 points are fully **located** in allocated complex equations (akin to *Complex Containers*) as  
 448 IP entities whereby their overall location [but not actual positions] are **intrinsically**  
 449 **incorporated** in these complex equations – see section 3 for additional clarification. Eqs.  
 450 (1), (3), (5), (6), (7) and (8) that comply with exact DA homogeneity at  $\sigma = \frac{1}{2}$  all have  
 451 fractional exponents  $\frac{1}{2}$ . Eqs. (2) and (4) that comply with inexact DA homogeneity at  
 452  $\sigma = \frac{2}{5}$  have fractional exponents  $\frac{2}{5}$  in the former and  $\frac{3}{5}$  in the later that are mixed with  
 453 fractional exponents  $\frac{1}{2}$ .

$$\sum_{n=1}^{\infty} (2n)^{-\frac{1}{2}} 2^{\frac{1}{2}} \cos(t \ln(2n) + \frac{1}{4}\pi) - \sum_{n=1}^{\infty} (2n-1)^{-\frac{1}{2}} 2^{\frac{1}{2}} \cos(t \ln(2n-1) + \frac{1}{4}\pi) = 0 \quad (1)$$

With exact DA homogeneity, Eq. (1) is  $f(n)$   $\text{sim-}\eta(s)$  at  $\sigma = \frac{1}{2}$  that will incorporate all  
 NTZ [as Zeroes]. There is total absence of (non-existent) virtual NTZ [as virtual Zeroes].

$$\sum_{n=1}^{\infty} (2n)^{-\frac{2}{5}} 2^{\frac{1}{2}} \cos(t \ln(2n) + \frac{1}{4}\pi) - \sum_{n=1}^{\infty} (2n-1)^{-\frac{2}{5}} 2^{\frac{1}{2}} \cos(t \ln(2n-1) + \frac{1}{4}\pi) = 0 \quad (2)$$

454 With inexact DA homogeneity, Eq. (2) is  $f(n)$   $\text{sim-}\eta(s)$  at  $\sigma = \frac{2}{5}$  that will incorporate  
 455 all (non-existent) virtual NTZ [as virtual Zeroes]. There is total absence of NTZ [as  
 456 Zeroes].

$$\frac{1}{2^{\frac{1}{2}}} \left( t^2 + \frac{1}{4} \right)^{\frac{1}{2}} \cdot \left[ (2n)^{\frac{1}{2}} \cos(t \ln(2n) - \frac{1}{4}\pi) - (2n-1)^{\frac{1}{2}} \cos(t \ln(2n-1) - \frac{1}{4}\pi) + C \right]_1^{\infty} = 0 \quad (3)$$

457 With exact DA homogeneity, Eq. (3) is  $F(n)$  DSPL at  $\sigma = \frac{1}{2}$  that will incorporate all  
 458 NTZ [as Pseudo-zeroes to Zeroes conversion]. There is total absence of (non-existent)  
 459 virtual NTZ [as virtual Pseudo-zeroes to virtual Zeroes conversion].

$$\frac{1}{2^{\frac{1}{2}}} \left( t^2 + \frac{9}{25} \right)^{\frac{1}{2}} \cdot \left[ (2n)^{\frac{3}{5}} \cos(t \ln(2n) - \frac{1}{4}\pi) - (2n-1)^{\frac{3}{5}} \cos(t \ln(2n-1) - \frac{1}{4}\pi) + C \right]_1^{\infty} = 0 \quad (4)$$

460 With inexact DA homogeneity, Eq. (4) is F(n) DSPL at  $\sigma = \frac{2}{5}$  that will incorporate  
 461 all (non-existent) virtual NTZ [as virtual Pseudo-zeroes to virtual Zeroes conversion].  
 462 There is total absence of NTZ [as Pseudo-zeroes to Zeroes conversion].

$$\sum_{n=1}^{\infty} (2n)^{-\frac{1}{2}} \sin(t \ln(2n)) - \sum_{n=1}^{\infty} (2n-1)^{-\frac{1}{2}} \sin(t \ln(2n-1)) = 0 \quad (5)$$

463 Eq. (5) can also be equivalently written as

$$464 \sum_{n=1}^{\infty} (2n)^{-\frac{1}{2}} \cos(t \ln(2n) - \frac{1}{2}\pi) - \sum_{n=1}^{\infty} (2n-1)^{-\frac{1}{2}} \cos(t \ln(2n-1) - \frac{1}{2}\pi) = 0.$$

465 With exact DA homogeneity, Eq. (5) is f(n) Gram[y=0] points-sim- $\eta$ (s) at  $\sigma = \frac{1}{2}$   
 466 that will incorporate all Gram[y=0] points [as Zeroes]. There is total absence of virtual  
 467 Gram[y=0] points [as virtual Zeroes].

$$-\frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}} \cdot \left[ (2n)^{\frac{1}{2}} (\cos(t \ln(2n) - \frac{1}{4}\pi) - \cos(t \ln(2n-1) - \frac{1}{4}\pi)) + C \right]_1^{\infty} = 0 \quad (6)$$

468 Eq. (6) can also be equivalently written as

$$469 \frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}} \cdot \left[ (2n)^{\frac{1}{2}} (\cos(t \ln(2n) + \frac{3}{4}\pi) - \cos(t \ln(2n-1) + \frac{3}{4}\pi)) + C \right]_1^{\infty} = 0.$$

470 With exact DA homogeneity, Eq. (6) is F(n) Gram[y=0] points-DSPL at  $\sigma = \frac{1}{2}$  that  
 471 will incorporate all Gram[y=0] points [as Pseudo-zeroes to Zeroes conversion]. There  
 472 is total absence of virtual Gram[y=0] points [as virtual Pseudo-zeroes to virtual Zeroes  
 473 conversion].

$$\sum_{n=1}^{\infty} (2n)^{-\frac{1}{2}} \cos(t \ln(2n)) - \sum_{n=1}^{\infty} (2n-1)^{-\frac{1}{2}} \cos(t \ln(2n-1)) = 0 \quad (7)$$

474 With exact DA homogeneity, Eq. (7) is f(n) Gram[x=0] points-sim- $\eta$ (s) at  $\sigma = \frac{1}{2}$   
 475 that will incorporate all Gram[x=0] points [as Zeroes]. There is total absence of virtual  
 476 Gram[x=0] points [as virtual Zeroes].

$$\frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}} \cdot \left[ (2n)^{\frac{1}{2}} (\cos(t \ln(2n) - \frac{3}{4}\pi) - \cos(t \ln(2n-1) - \frac{3}{4}\pi)) + C \right]_1^{\infty} = 0 \quad (8)$$

477 With exact DA homogeneity, Eq. (8) is F(n) Gram[x=0] points-DSPL at  $\sigma = \frac{1}{2}$  that  
 478 will incorporate all Gram[x=0] points [as Pseudo-zeroes to Zeroes conversion]. There  
 479 is total absence of virtual Gram[x=0] points [as virtual Pseudo-zeroes to virtual Zeroes  
 480 conversion].

481 We outline sim- $\eta$ (s) as Eq. (2) and DSPL as Eq. (4) that comply with inexact DA  
 482 homogeneity at  $\sigma = \frac{2}{5}$  non-critical line (depicted by Figure 3) whereby  $\sigma = \frac{2}{5}$  [instead of  
 483  $\sigma = \frac{1}{2}$ ] is substituted into these two equations. Using [selective] trigonometric identity  
 484 for linear combination of sine and cosine function whenever applicable to relevant f(n) =  
 485 0 and F(n) = 0 equations, we outline exact DA homogeneity at  $\sigma = \frac{1}{2}$  critical line (depicted  
 486 by Figure 2) for Gram[x=0,y=0] points (NTZ) as Eq. (1), Gram[y=0] points as Eq. (5)  
 487 and Gram[x=0] points as Eq. (7). However, f(n) = 0 equations for Gram[y=0] points as  
 488 Eq. (5) and Gram[x=0] points as Eq. (7) with exact DA homogeneity at  $\sigma = \frac{1}{2}$  critical  
 489 line are not amendable to treatments using trigonometric identity with implication that  
 490 their corollary situation endowed with inexact DA homogeneity at  $\sigma \neq \frac{1}{2}$  non-critical  
 491 lines (depicted by Figures 3 and 4) will only manifest solitary [unmixed]  $\neq \frac{1}{2}$  fractional  
 492 exponents. We provide [self-explanatory] corresponding f(n) = 0 equations below for  
 493 Gram[y=0] points and Gram[x=0] points corollary situation when  $\sigma = \frac{2}{5}$ .

$$494 \quad \sum_{n=1}^{\infty} (2n)^{-\frac{2}{5}} \sin(t \ln(2n)) - \sum_{n=1}^{\infty} (2n-1)^{-\frac{2}{5}} \sin(t \ln(2n-1)) = 0$$

$$495 \quad \sum_{n=1}^{\infty} (2n)^{-\frac{2}{5}} \cos(t \ln(2n)) - \sum_{n=1}^{\infty} (2n-1)^{-\frac{2}{5}} \cos(t \ln(2n-1)) = 0$$

496 We arbitrarily chose single cosine wave with format  $R \cos(n \pm \alpha)$  to use above where  
 497  $R$  is scaled amplitude and  $\alpha$  is phase shift. For equations regarding NTZ, Gram[ $y=0$ ]  
 498 points and Gram[ $x=0$ ] points; all their approximate Areas of Varying Loops  $\propto$  precise  
 499 Areas of Varying Loops with  $R$  validly treated as a proportionality factor. We analyze  $f(n)$   
 500  $= 0$  and  $F(n) = 0$  equations at  $\sigma = \frac{1}{2}$  critical line for NTZ situation where  $R = 2^{\frac{1}{2}}(2n)^{-\frac{1}{2}}$   
 501 or  $2^{\frac{1}{2}}(2n-1)^{-\frac{1}{2}}$  in  $f(n)$ 's Eq. (1) and  $R = \frac{1}{2^{\frac{1}{2}}(t^2 + \frac{1}{4})^{\frac{1}{2}}}(2n)^{\frac{1}{2}}$  or  $\frac{1}{2^{\frac{1}{2}}(t^2 + \frac{1}{4})^{\frac{1}{2}}}(2n-1)^{\frac{1}{2}}$

502 in  $F(n)$ 's Eq. (3).

503

504 **Remark 3.** Whereas for NTZ  $F(n)$  Eq. (3) that exactly represent precise Areas of  
 505 Varying Loops and  $f(n)$  Eq. (1) [when interpreted as Riemann sum] that exactly represent  
 506 approximate Areas of Varying Loops in a proportionate manner; so must the associated  
 507 scaled amplitude  $R$  from Eq. (3) **which is dependent on parameter  $t$**  and Eq. (1) **which**  
 508 **is independent of parameter  $t$**  represent [in a surrogate manner] corresponding precise  
 509 and approximate Areas of Varying Loops in a proportionate manner.

510

511 We analyze  $f(n) = 0$  equations [relevant to approximate Areas of Varying Loops]  
 512 at  $\sigma = \frac{1}{2}$  critical line for Gram[ $y=0$ ] points as Eq. (5) and Gram[ $x=0$ ] points as Eq. (7)  
 513 whereby we validly designate  $R = (2n)^{-\frac{1}{2}}$  or  $(2n-1)^{-\frac{1}{2}}$  as the assigned scaled amplitude  
 514 and [unwritten]  $\alpha = 0$  as the assigned phase shift.

515 Relevant to precise Areas of Varying Loops at  $\sigma = \frac{1}{2}$  critical line for Gram[ $y=0$ ]  
 516 points  $F(n)$  Eq. (6) with  $R = -\frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}}(2n)^{\frac{1}{2}}$  or  $-\frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}}(2n-1)^{\frac{1}{2}}$  and Gram[ $x=0$ ]  
 517 points  $F(n)$  Eq. (8) with  $R = \frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}}(2n)^{\frac{1}{2}}$  or  $\frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}}(2n-1)^{\frac{1}{2}}$ , we observe the

518 former  $R$  to be the negative of the later  $R$ . However, this observation is context-sensitive  
 519 because when Eq. (6) is written in its equivalent format above, the former  $R$  is identi-  
 520 cal to the later  $R$ . Both  $R$  are now just given by  $\frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}}(2n)^{\frac{1}{2}}$  or  $\frac{1}{2(t^2 + \frac{1}{4})^{\frac{1}{2}}}(2n-1)^{\frac{1}{2}}$ .

521

522 **Remark 4.** Whereas for Gram[ $y=0$ ] points  $F(n)$  Eq. (6) that exactly represent precise  
 523 Areas of Varying Loops and  $f(n)$  Eq. (5) [when interpreted as Riemann sum] that exactly  
 524 represent approximate Areas of Varying Loops in a proportionate manner; so must the  
 525 associated scaled amplitude  $R$  in Eq. (6) **which is dependent on parameter  $t$**  and Eq. (5)  
 526 **which is independent of parameter  $t$**  represent [in a surrogate manner] corresponding  
 527 precise and approximate Areas of Varying Loops in a proportionate manner.

528

529 **Remark 5.** Whereas for Gram[ $x=0$ ] points  $F(n)$  Eq. (8) that exactly represent precise  
 530 Areas of Varying Loops and  $f(n)$  Eq. (7) [when interpreted as Riemann sum] that exactly  
 531 represent approximate Areas of Varying Loops in a proportionate manner; so must the  
 532 associated scaled amplitude  $R$  in Eq. (8) **which is dependent on parameter  $t$**  and Eq. (7)  
 533 **which is independent of parameter  $t$**  represent [in a surrogate manner] corresponding  
 534 precise and approximate Areas of Varying Loops in a proportionate manner.

535

536 Finally, we analyze  $f(n) = 0$  and  $F(n) = 0$  equations at  $\sigma = \frac{1}{2}$  critical line for NTZ  
 537 situation where phase shift  $\alpha = \frac{1}{4}\pi$  in NTZ  $f(n)$  Eq. (1) and  $-\frac{1}{4}\pi$  in NTZ  $F(n)$  Eq. (3);

538 and  $F(n) = 0$  equations at  $\sigma = \frac{1}{2}$  critical line for Gram[ $y=0$ ] points and Gram[ $x=0$ ] points  
 539 situations where phase shift  $\alpha = -\frac{1}{4}\pi$  (or  $\frac{3}{4}\pi$  when written in its equivalent format  
 540 above) in Gram[ $y=0$ ] points  $F(n)$  Eq. (6) and  $-\frac{3}{4}\pi$  in Gram[ $x=0$ ] points  $F(n)$  Eq. (8).  
 541 Always being  $\frac{1}{2}\pi$  out-of-phase with each other, trigonometric functions sine and cosine  
 542 are cofunctions with  $\sin n = \cos(\frac{\pi}{2} - n)$  or  $\cos(n - \frac{\pi}{2})$ ,  $\cos n = \sin(\frac{\pi}{2} - n)$  or  $\sin(n + \frac{\pi}{2})$ ,  
 543  $\frac{d(\sin n)}{dn} = \cos n$ ,  $\frac{d(\cos n)}{dn} = -\sin n$ ,  $\int \sin n \cdot dn = -\cos n + C [= \sin(n - \frac{\pi}{2}) + C]$  and  
 544  $\int \cos n \cdot dn = \sin n + C [= \cos(n - \frac{\pi}{2}) + C]$ . Last two integrals explain relation between  
 545  $f(n)$ 's Zeroes and  $F(n)$ 's Pseudo-zeroes when they involve simple sine and/or cosine  
 546 terms viz,  $f(n)$ 's CP Zeroes =  $F(n)$ 's CP Pseudo-zeroes  $-\frac{1}{2}\pi$  with CP Zeroes and CP  
 547 Pseudo-zeroes being  $\frac{1}{2}\pi$  out-of-phase with each other.

548  
 549 **Lemma 1.** NTZ obtained directly from IP Zeroes and indirectly from IP Pseudo-  
 550 zeroes behave in accordance with complex sine and/or cosine terms present in their  
 551 equations that are  $\frac{1}{2}\pi$  out-of-phase with each other.

552 **Proof.** Involving trigonometric functions as complex sine and/or cosine terms:  
 553  $f(n)$ 's IP NTZ or [non-existent]  $f(n)$ 's IP virtual NTZ (in  $t$  values) =  $F(n)$ 's IP Pseudo-NTZ  
 554 or [non-existent]  $F(n)$ 's IP virtual Pseudo-NTZ (in  $t$  values)  $-\frac{1}{2}\pi$ ;  $f(n)$ 's IP Gram[ $y=0$ ]  
 555 points or  $f(n)$ 's IP virtual Gram[ $y=0$ ] points (in  $t$  values) =  $F(n)$ 's IP Pseudo-Gram[ $y=0$ ]  
 556 points or  $F(n)$ 's IP virtual Pseudo-Gram[ $y=0$ ] points (in  $t$  values)  $-\frac{3}{4}\pi$ ; and  $f(n)$ 's IP  
 557 Gram[ $x=0$ ] points or  $f(n)$ 's IP virtual Gram[ $x=0$ ] points (in  $t$  values) =  $F(n)$ 's IP Pseudo-  
 558 Gram[ $x=0$ ] points or  $F(n)$ 's IP virtual Pseudo-Gram[ $x=0$ ] points (in  $t$  values)  $-\frac{3}{4}\pi$ .

559  $\int f(n)dn = F(n) + C$  where  $F'(n) = f(n)$ .  $f(n)$  and  $F(n)$  are literally [connected]  
 560 **bijective (both injective and surjective or a one-to-one correspondence) functions.**  
 561 Underlying  $f(n)$  as equation and  $F(n)$  as law (equation) that generate their CIS of IP  
 562 Zeroes, IP virtual Zeroes, IP Pseudo-zeroes and IP virtual Pseudo-zeroes are precisely  
 563 related as  $\frac{1}{2}\pi$  (for NTZ case) or  $\frac{3}{4}\pi$  (for Gram[ $y=0$ ] points and Gram[ $x=0$ ] points cases)  
 564 out-of-phase with each other. Peculiar to IP NTZ as Origin intercept points, we crucially  
 565 note only they will uniquely behave in accordance with complex sine and/or cosine  
 566 terms present in their equations that generate corresponding IP Zeroes and IP Pseudo-  
 567 zeroes which are  $\frac{1}{2}\pi$  [but not  $\frac{3}{4}\pi$ ] out-of-phase with each other. *The proof is now complete*  
 568 *for Lemma 1*  $\square$ .

569  
 570 **Lemma 2.** Corresponding paired IP two types of Gram points [as Zeroes] situation,  
 571 paired IP two types of virtual Gram points [as virtual Zeroes] situation, paired IP two  
 572 types of Pseudo-Gram points [as Pseudo-zeroes] situation, and paired IP two types of  
 573 virtual Pseudo-Gram points [as virtual Pseudo-zeroes] situation are always  $\frac{1}{2}\pi$  out-of-  
 574 phase with each other in every one of these situations.

575 **Proof.** The  $x$ -axis and  $y$ -axis are orthogonal to each other with angle between them  
 576 =  $\frac{1}{2}\pi$  radian. Involving trigonometric functions as complex sine and/or cosine terms:  
 577  $f(n)$ 's IP Gram[ $y=0$ ] points or  $f(n)$ 's IP virtual Gram[ $y=0$ ] points (in  $t$  values) =  $f(n)$ 's IP  
 578 Gram[ $x=0$ ] points or  $f(n)$ 's IP virtual Gram[ $x=0$ ] points (in  $t$  values)  $+\frac{1}{2}\pi$ ; and  $F(n)$ 's IP  
 579 Pseudo-Gram[ $y=0$ ] points or  $F(n)$ 's IP virtual Pseudo-Gram[ $y=0$ ] points (in  $t$  values) =

580  $F(n)$ 's IP Pseudo-Gram $[x=0]$  points or  $F(n)$ 's IP virtual Pseudo-Gram $[x=0]$  points (in t  
581 values)  $+$   $\frac{1}{2}\pi$ .

582 These observations imply underlying  $f(n)$  as equation and  $F(n)$  as law (equation)  
583 that generate corresponding paired IP two types of Gram points [as Zeroes] situation,  
584 paired IP two types of virtual Gram points [as virtual Zeroes] situation, paired IP two  
585 types of Pseudo-Gram points [as Pseudo-zeroes] situation, and paired IP two types of  
586 virtual Pseudo-Gram points [as virtual Pseudo-zeroes] situation are always  $\frac{1}{2}\pi$  out-  
587 of-phase with each other in every one of these mentioned situations. *The proof is now*  
588 *complete for Lemma 2*□.

589

### 590 3. The Completely Predictable and Incompletely Predictable entities

591 The word "number" [singular noun] or "numbers" [plural noun] used in reference  
592 to CP even and odd numbers, IP prime and composite numbers, IP NTZ and two other  
593 types of Gram points can interchangeably be replaced with the word "entity" [singular  
594 noun] or "entities" [plural noun]. For  $i =$  all integers  $\geq 0$  or  $i =$  all integers  $\geq 1$ ; the  $i^{\text{th}}$   
595 position of  $i^{\text{th}}$  CP numbers and  $i^{\text{th}}$  IP numbers is simply given by  $i$ . Apart from the very  
596 first Gram $[y=0]$  point and the very first virtual Gram $[y=0]$  point being both 0, we note  
597 all Gram points and virtual Gram points will consist of  $t$ -valued transcendental numbers  
598 whose positions are IP with the infinitely many digits after the decimal point in each  
599 transcendental number again being IP.

600 We outline an innovative method to classify appropriately chosen equation or al-  
601 gorithm in two ways by using relevant locational properties of its output. This output  
602 consist of generated entities either from function-based equations or from algorithms.  
603 Our novel [albeit loose] classification systems named "*Mathematics for Completely Pre-*  
604 *dictable problems*" that is associated with conveniently-coined *simple calculations*, and  
605 "*Mathematics for Incompletely Predictable problems*" that is associated with conveniently-  
606 coined *complex calculations*, are respectively formalized by providing formal definitions  
607 for CP entities obtained from CP equations or algorithms, and IP entities obtained from  
608 IP equations or algorithms.

609 CP simple equation or algorithm generates CP numbers. A generated CP number  
610 is **locationally defined** as a number whose  $i^{\text{th}}$  position is *independently* determined  
611 by simple calculations without needing to know related positions of all preceding  
612 numbers. IP complex equation or algorithm generates IP numbers. A generated IP  
613 number is **locationally defined** as a number whose  $i^{\text{th}}$  position is *dependently* determined  
614 by complex calculations with needing to know related positions of all preceding numbers.  
615 Container is a useful analogical term that metaphorically group CP entities (e.g. even  
616 and odd numbers) and IP entities (e.g. nontrivial zeros, prime and composite numbers)  
617 to be exclusively located in, respectively, Simple Container and Complex Container.

618 Simple properties are inferred from a sentence such as "This simple equation or  
619 algorithm by itself will intrinsically incorporate *overall location [and actual positions]* of all  
620 CP numbers". Examples: simple equations  $E = (2 \times i)$  for  $i =$  all integers  $\geq 0$  [or  $i =$  all  
621 real numbers  $\geq 0$ ] and  $O = (2 \times i) - 1$  for  $i =$  all integers  $\geq 1$  [or  $i =$  all real numbers  $\geq 1$ ]  
622 will respectively and intrinsically incorporate or generate CIS of all [non-negative] CP  
623 even number  $E_i = 0, 2, 4, 6, \dots$  and CIS of all [non-negative] CP odd numbers  $O_i = 1, 3,$   
624  $5, 7, \dots$  whereby even number (**n**) is defined as "Any integer that can be divided exactly  
625 by 2 with last digit always being 0, 2, 4, 6 or 8" and odd number (**n**) is defined as "Any  
626 integer that cannot be divided exactly by 2 with last digit always being 1, 3, 5, 7 or 9".  
627 Congruence  $\mathbf{n} \equiv 0 \pmod{2}$  holds for even **n** and congruence  $\mathbf{n} \equiv 1 \pmod{2}$  holds for odd  
628 **n**. Note the zeroth even number is given by  $E_0 = 0$ .

629 Complex properties, or meta-properties, are inferred from a sentence such as "This  
630 complex equation or algorithm by itself will intrinsically incorporate *overall location [but*  
631 *not actual positions]* of all IP numbers". Examples: complex algorithms  $P_{i+1} = P_i + p\text{Gap}_i$

632 and  $C_{i+1} = C_i + cGap_i$  for  $i = 1, 2, 3, \dots, \infty$  with  $P_1 = 2$  and  $C_1 = 4$  will respectively and  
 633 intrinsically incorporate CIS of all IP prime number 2, 3, 5, 7, ... and CIS of all IP composite  
 634 numbers 4, 6, 8, 9, ... whereby prime numbers are defined as "All Natural numbers apart  
 635 from 1 that are evenly divisible by itself and by 1" and composite numbers are defined  
 636 as "All Natural numbers apart from 1 that are evenly divisible by numbers other than  
 637 itself and 1". E.g. via computed Pseudo-zeroes that can be converted to Zeroes at  $\sigma = \frac{1}{2}$   
 638 critical line, complex equation DSPL will intrinsically incorporate the CIS of all IP NTZ  
 639 [given as t values rounded off to six decimal places]: 14.134725, 21.022040, 25.010858,  
 640 30.424876, 32.935062, 37.586178, ... and complex equation Gram[y=0] points-DSPL will  
 641 intrinsically incorporate the CIS of all IP Gram[y=0] points [given as t values rounded off  
 642 to six decimal places]: 0, 3.436218, 9.666908, 17.845599; 23.170282, 27.670182, ... Choice of  
 643 index  $n$  for Gram[y=0] points is crudely chosen in the past to be -3, -2, -1, 0, 1, 2, 3, ... [ $\equiv i$   
 644 = 1, 2, 3, 4, 5, 6, 7, ...] whereby the first Gram[y=0] point is historically denoted by  $n = 1$   
 645 [ $\equiv i = 5$ ] with t value 17.845599 (on critical line) being larger than first NTZ's t value of  
 646 14.134725 (on critical line).

647 **The Even-Odd Pairing.** For  $i = 1, 2, 3, \dots, \infty$ ; let mutually exclusive  $i^{th}$  Even numbers  
 648 =  $E_i$  and  $i^{th}$  Odd numbers =  $O_i$ , and  $i^{th}$  even number gaps =  $eGap_i$  and  $i^{th}$  odd number  
 649 gaps =  $oGap_i$ . The  $i^{th}$  positions of  $E_i$  and  $O_i$  are CP, and are independent from each other.

$E_i$	2		4		6		8		10		12	.....
$eGap_i$		2		2		2		2		2		2

651 We employ simple equations  $E = (2 \times i)$  and  $O = (2 \times i) - 1$ . E.g., we can precisely, easily  
 652 and independently calculate  $E_5 = (2 \times 5) = 10$  and  $O_5 = (2 \times 5) - 1 = 9$ .

$O_i$	1		3		5		7		9		11	.....
$oGap_i$		2		2		2		2		2		2

655 **The Prime-Composite Pairing.** For  $i = 1, 2, 3, \dots, \infty$ ; let mutually exclusive  $i^{th}$  Prime  
 656 numbers =  $P_i$  and  $i^{th}$  Composite numbers =  $C_i$ , and  $i^{th}$  prime number gaps =  $pGap_i$   
 657 and  $i^{th}$  composite number gaps =  $cGap_i$ . The  $i^{th}$  positions of  $P_i$  and  $C_i$  are IP, and are  
 658 dependent on each other.

$P_i$	2		3		5		7		11		13	.....
$pGap_i$		1		2		2		4		2		4

660 We employ complex algorithms  $P_{i+1} = P_i + pGap_i$  and  $C_{i+1} = C_i + cGap_i$ . E.g., we  
 661 precisely, tediously and dependently calculate  $P_6 = 13$  as 2 is 1<sup>st</sup> prime number, 3 is 2<sup>nd</sup>  
 662 prime number, 4 is 1<sup>st</sup> composite number, 5 is 3<sup>rd</sup> prime number, 6 is 2<sup>nd</sup> composite  
 663 number, 7 is 4<sup>th</sup> prime number, 8 is 3<sup>rd</sup> composite number, 9 is 4<sup>th</sup> composite number, 10  
 664 is 5<sup>th</sup> composite number, 11 is 5<sup>th</sup> prime number, 12 is 6<sup>th</sup> composite number, and our  
 665 desired 13 is 6<sup>th</sup> prime number.

$C_i$	4		6		8		9		10		12	.....
$cGap_i$		2		2		1		1		2		2

668 **The  $\sigma = \frac{1}{2}$  NTZ computed from Eq. (1) -  $\sigma \neq \frac{1}{2}$  (non-existent) virtual NTZ**  
 669 **computed from Eq. (2) Pairing.** For  $i = 1, 2, 3, \dots, \infty$ ; let mutually exclusive  $i^{th}$  NTZ =  
 670  $NTZ_i$  and  $i^{th}$  virtual NTZ =  $vNTZ_i$ , and  $i^{th}$  NTZ gaps =  $NTZ-Gap_i$  and  $i^{th}$  virtual NTZ  
 671 gaps =  $vNTZ-Gap_i$ . Eq. (1) and Eq. (2) are dependently identical except for associated  $\sigma$   
 672 values. They are used to precisely, tediously and dependently calculate all  $NTZ_i$  and  
 673  $vNTZ_i$  with their  $i^{th}$  positions being IP.

674 **4. The exact and inexact Dimensional analysis homogeneity for Equations**

675 For 'base quantities' *length, mass and time*; their fundamental SI 'units of measure-  
 676 ment' meter (m) is defined as distance travelled by light in vacuum for time interval  
 677  $1/299\,792\,458$  s with speed of light  $c = 299,792,458 \text{ ms}^{-1}$ , kilogram (kg) is defined by  
 678 taking fixed numerical value Planck constant  $h$  to be  $6.626\,070\,15 \times 10^{-34}$  Joules·second  
 679 (Js) [whereby Js is equal to  $\text{kgm}^2\text{s}^{-1}$ ] and second (s) is defined in terms of  $\Delta\nu\text{Cs} =$   
 680  $\Delta(^{133}\text{Cs})_{hfs} = 9,192,631,770 \text{ s}^{-1}$ . Derived SI units such as J and  $\text{ms}^{-1}$  respectively rep-  
 681 resent 'base quantities' *energy* and *velocity*. 'Dimension' is commonly used to indicate



682 'units of measurement' in well-defined equations. DA is a traditional analytic tool with  
 683 DA homogeneity and DA non-homogeneity (respectively) denoting valid and invalid  
 684 equation occurring when 'units of measurements' for 'base quantities' are "balanced"  
 685 and "unbalanced" across both sides of equation. E.g. equation  $2\text{ m} + 3\text{ m} = 5\text{ m}$  is valid  
 686 but equation  $2\text{ m} + 3\text{ kg} = 5\text{ 'm}\cdot\text{kg}'$  is invalid (respectively) manifesting DA homogeneity  
 687 and non-homogeneity.

688 We conveniently adopt concepts from DA which are mathematically correct and  
 689 valid. Let  $(2n)$  and  $(2n-1)$  be 'base quantities' in equation DSPL. Fractional exponents  
 690 as 'units of measurement' given by  $(1 - \sigma)$  in equation DSPL when  $\sigma = \frac{1}{2}$  coincide with  
 691 exact DA homogeneity; and  $(1 - \sigma)$  in equation DSPL when  $\sigma \neq \frac{1}{2}$  coincide with inexact  
 692 DA homogeneity. Respectively, exact DA homogeneity at  $\sigma = \frac{1}{2}$  denotes  $\Sigma$ (all fractional  
 693 exponents) as  $2(1 - \sigma)$  equates to [exact] integer 1; and inexact DA homogeneity at  $\sigma \neq \frac{1}{2}$   
 694 denotes  $\Sigma$ (all fractional exponents) as  $2(1 - \sigma)$  equates to [inexact] fractional number  
 695  $\neq 1$  [Range:  $0 < 2(1 - \sigma) < 1$  and  $1 < 2(1 - \sigma) < 2$ ]. Computations based on exact and  
 696 inexact DA homogeneity in equation DSPL explicitly give rise to  $\sigma = \frac{1}{2}$  critical line Gram  
 697 points (given indirectly as Pseudo-zeroes t-values which can be converted to Zeroes  
 698 t-values) and  $\sigma \neq \frac{1}{2}$  non-critical lines virtual Gram points (given indirectly as virtual  
 699 Pseudo-zeroes t-values which can be converted to virtual Zeroes t-values).

700 Performing exact and inexact DA homogeneity on equation  $\text{sim-}\eta(s)$  is equally valid.  
 701 With same 'base quantities', fractional exponents as 'units of measurement' are now  
 702 given by  $(-\sigma)$ . Respectively, exact DA homogeneity at  $\sigma = \frac{1}{2}$  denotes  $\Sigma$ (all fractional  
 703 exponents) as  $2(-\sigma)$  equates to [exact] integer  $-1$ ; and inexact DA homogeneity at  $\sigma \neq \frac{1}{2}$   
 704 denotes  $\Sigma$ (all fractional exponents) as  $2(-\sigma)$  equates to [inexact] fractional number  $\neq -1$   
 705 [Range:  $-2 < 2(-\sigma) < -1$  and  $-1 < 2(-\sigma) < 0$ ]. Computations using equation  $\text{sim-}\eta(s)$   
 706 [when interpreted as Riemann sum] explicitly give rise to  $\sigma = \frac{1}{2}$  critical line Gram points  
 707 (given directly as Zeroes t-values) while representing exact DA homogeneity and  $\sigma \neq \frac{1}{2}$   
 708 non-critical lines virtual Gram points (given directly as virtual Zeroes t-values) while  
 709 representing inexact DA homogeneity.

710 *For calculations involving  $2(1 - \sigma)$  or  $2(-\sigma)$ , we note it is inconsequential whether  $\sigma$*   
 711 *values from the fractional exponents of 'base quantities'  $(2n)$  or  $(2n-1)$  are formatted in simplest*  
 712 *form or not. For example, since  $\frac{1}{2} \equiv \frac{2}{4}$ ; performing the  $\sigma = \frac{1}{2}$  exact DA homogeneity on*  
 713 *exponent  $\frac{1}{2}$  in  $(2n)^{\frac{1}{2}}$  when depicted in simplest form will be equivalent to performing*  
 714 *the [same]  $\sigma = \frac{1}{2}$  exact DA homogeneity on exponent  $\frac{1}{4}$  in  $(2^2n^2)^{\frac{1}{4}}$  when not depicted in*  
 715 *simplest form.*

## 716 5. Gauss Circle Problem and Primitive Circle Problem

717 Equation of a circle centered at Origin with radius  $r$  and precise Area =  $\pi r^2$  is  
 718 given in Cartesian coordinates as  $x^2 + y^2 = r^2$ . The number of integer lattice points  
 719  $N(r)$  on and inside a circle [viz, pairs of integers  $(m,n)$  such that  $m^2 + n^2 \leq r^2$ ] can be  
 720 exactly determined by following two equations whereby  $N(r)$  is considered the most  
 721 accurate surrogate marker of approximate Area for a given circle. Named after German  
 722 mathematician Carl Friedrich Gauss (April 30, 1777 - February 23, 1855), Gauss Circle  
 723 Problem is the problem of determining how many integer lattice points as approximate  
 724 Area for a given circle. For  $i$  and  $r = 0, 1, 2, 3, \dots, \infty$  and through which it can be given by  
 725 several series such as in terms of a sum involving the floor function;  $N(r)$  is expressed  
 726 as equation  $N(r) = 1 + 4 \sum_{i=0}^{\infty} \left( \left\lfloor \frac{r^2}{4i+1} \right\rfloor - \left\lfloor \frac{r^2}{4i+3} \right\rfloor \right)$  whereby this equation is a conse-  
 727 quence of Jacobi's two-square theorem which follows almost immediately from Jacobi  
 728 triple product. A much simpler sum appears if sum of squares function  $r_2(n)$  that is  
 729 defined as number of ways of writing number  $n$  as sum of two squares is used. Then,  
 730 we have alternative equation  $N(r) = \sum_{n=0}^{r^2} r_2(n)$ . The first few  $N(r)$  values for  $r = 0, 1,$   
 731  $2, 3, 4, 5, 6, 7, 8, \dots$  are  $1, 5, 13, 29, 49, 81, 113, 149, \dots$  whereby these are Incompletely

732 Predictable entities complying with relationship: [simple] equation for precise Area  
 733 of circle =  $\pi r^2$  is proportional to above two most accurate and equivalent [complex]  
 734 equations for approximate Area of circle =  $N(r)$ .

735 We expect  $N(r) = \pi r^2 + E(r)$  for some error term  $E(r)$  of relatively small absolute  
 736 value. Gauss managed to prove  $|E(r)| \leq 2\sqrt{2}\pi r$ . Modern proofs on upper bound value  
 737 [in 2000] and lower bound value [in 1915] for  $E(r)$  have since been derived. We recognize  
 738  $r$  does not have to be an integer. After  $N(4) = 49$ , we obtain  $N(\sqrt{17}) = 57, N(\sqrt{18}) =$   
 739  $61, N(\sqrt{20}) = 69, N(5) = 81$ . At these places,  $E(r)$  increases by 8, 4, 8, 12 after which it  
 740 decreases at a rate of  $2\pi r$  until the next time it increases.

741 Finally, the identity  $N(x) - \frac{r_2(x^2)}{2} = \pi x^2 + x \sum_{n=1}^{\infty} \frac{r_2(n)}{\sqrt{n}} J_1(2\pi x \sqrt{n})$  has implicitly  
 742 been observed to be related to number of integer lattice points,  $N(r)$ , where  $J_1$  denotes  
 743 Bessel function of first kind with order 1. It was discovered by English mathematician  
 744 Godfrey H. Hardy (February 7, 1877 - December 1, 1947)[5].

745 Primitive Circle Problem as least accurate surrogate marker of approximate Area for  
 746 a given circle involves calculating the number of coprime integer solutions  $(m,n)$  to the  
 747 inequality  $m^2 + n^2 \leq r^2$ . If the number of such solutions is denoted  $V(r)$  then the values  
 748 of  $V(r)$  for  $r$  taking small integer values are 0, 4, 8, 16, 32, 48, 72, 88, 120, 152, 192,.... Using  
 749 the same ideas as usual Gauss Circle Problem and the fact that probability two integers  
 750 are coprime is  $\frac{6}{\pi^2}$ , it is relatively straightforward to show  $V(r) = \frac{6}{\pi} r^2 + O(r^{1+\epsilon})$ . We  
 751 solve problematic part of Primitive Circle Problem by reducing the exponent in the error  
 752 term. This exponent is presently best known to be  $221/304 + \epsilon$  since we can now validly  
 753 assume Riemann hypothesis to be true in this paper.

754  
 755 **Remark 6.** Let  $A$  denote Area of a given circle with radius  $r$ . The computed precise  
 756  $A$  using  $A = \pi r^2$  method, computed approximate  $A$  using [most accurate] approximate  
 757  $N(r)$  method of Gauss Circle Problem and computed approximate  $A$  using [least accu-  
 758 rate] approximate  $A(r)$  method of Primitive Circle Problem will explicitly confirm  $A \propto$   
 759  $r^2$  for all three methods.

760

## 761 6. Gauss Areas of Varying Loops and Principle of Maximum Density for Integer 762 Number Solutions

763 We translate concepts from Gauss Circle Problem and Primitive Circle Problem in  
 764 section 5 onto Gauss Areas of Varying Loops to fully support all materials below.

765

766 **Proposition 1.** We can validly and fully demonstrate that only when  $\sigma = \frac{1}{2}$  [and  
 767 not when  $\sigma \neq \frac{1}{2}$ ] in  $\text{sim-}\eta(s)$  or DSPL will the maximum number of integer solutions  
 768 (constituted by all integers  $\geq 1$ ) arise that must uniquely comply with Principle of  
 769 Maximum Density for Integer Number Solutions.

770 **Proof.** For  $n$  classically involving all integers  $\geq 1$  in  $\text{sim-}\eta(s)$  as  $\Delta n \rightarrow 1$  or  $n$   
 771 classically involving all real numbers  $\geq 1$  in DSPL as  $\Delta n \rightarrow 0$ ; their base quantities  $(2n)$   
 772 and  $(2n-1)$ , respectively, generate CIS even numbers commencing from 2 and CIS odd  
 773 numbers commencing from 1. These base quantities are subjected to algebraic function  
 774 square roots at  $\sigma = \frac{1}{2}$  critical line [viz, when  $\sigma = \frac{1}{2}$ ] and cube roots at  $\sigma = \frac{1}{3}$  non-critical  
 775 line or twice cube roots at  $\sigma = \frac{2}{3}$  non-critical line [viz, when  $\sigma \neq \frac{1}{2}$ ] thus giving rise to  
 776 corresponding subset of rational roots and subset of irrational roots. We now concentrate  
 777 on combined  $(2n)$ 's and  $(2n-1)$ 's obtained integer lattice points  $[\geq 1]$  to derive solitary  
 778 subset of rational roots for  $n = 1$  to 100 range in  $\text{sim-}\eta(s)$  or DSPL when:

779 (I)  $\sigma = \frac{1}{2}$  involving a **neither even nor odd function** with no symmetry viz,  
 780  $f(-n) \neq f(n)$  and  $f(-n) \neq -f(n)$  by applying  $f(n)$  as fractional exponent  $\frac{1}{2}$  or  
 781 square root on  $n =$  ten perfect squares 1, 4, 9, 16, 25, 36, 49, 64, 81, 100 giving rise to the  
 782 (maximum) ten rational roots as consecutive integers 1, 2, 3, 4, 5, 6, 7, 8, 9, 10.

783 (II)  $\sigma = \frac{1}{3}$  involving a **odd function** with Origin symmetry viz,  $f(-n) = -f(n)$  by  
 784 applying  $f(n)$  as fractional exponent  $\frac{1}{3}$  or cube root on  $n =$  four perfect cubes 1, 8, 27, 64  
 785 giving rise to the (non-maximum) four rational roots as consecutive integer solutions 1,  
 786 2, 3, 4.

787 (III)  $\sigma = \frac{2}{3}$  involving an **even function** with y-axis symmetry viz,  $f(-n) = f(n)$  by  
 788 applying  $f(n)$  as fractional exponent  $\frac{2}{3}$  or squared cube root on  $n =$  four perfect cubes 1,  
 789 8, 27, 64 giving rise to the (non-maximum) four rational roots as non-consecutive integer  
 790 solutions 1, 4, 9, 16.

791 Only at  $\sigma = \frac{1}{2}$  critical line which involves applying  $f(n)$  as fractional exponent  
 792  $\frac{1}{2}$  or square root on  $n =$  all perfect squares 1, 4, 9, 16, 25, 36, 49, 64, 81, 100... will we  
 793 obtain maximum number of rational roots as consecutive integer solutions 1, 2, 3, 4, 5, 6,  
 794 7, 8, 9, 10... (viz, all integers  $\geq 1$ ). This observation uniquely comply with **Principle of**  
 795 **Maximum Density for Integer Number Solutions** at  $\sigma = \frac{1}{2}$  critical line. *The proof is now*  
 796 *complete for Proposition 1*  $\square$ .

797  
 798 Notation: **Term-(2n)** denote **(2n)-complex term with algebraic functions X (2n)-**  
 799 **complex term with transcendental functions**; and **Term-(2n-1)** denote **(2n-1)-complex**  
 800 **term with algebraic functions X (2n-1)-complex term with transcendental functions**.  
 801  $\text{sim-}\eta(s)$  or DSPL is complex function or law with single variable  $n$  and parameters  
 802  $\sigma, t$ . Their derived equations [Eqs. (1) to (8)] have **(2n)- or (2n-1)-complex term with**  
 803 **algebraic functions** consisting of powers, fractional powers, root extraction and scaled  
 804 amplitude  $R$  that are **dependent on parameter  $\sigma$** , and **(2n)- or (2n-1)-complex term with**  
 805 **transcendental functions** consisting of sine, cosine, single cosine wave, single sine wave,  
 806 natural logarithm that are **independent of parameter  $\sigma$** .

807  
 808 **Remark 7.** Corresponding to Areas of Varying Loops = 0 in  $f(n)$   $\text{sim-}\eta(s)$  or  $F(n)$   
 809 DSPL, **Term-(2n)** must precisely cancel **Term-(2n-1)** in order to obtain  $\sigma = \frac{1}{2}$   $f(n)$ 's Zeroes  
 810 and  $F(n)$ 's Pseudo-zeroes or to obtain  $\sigma \neq \frac{1}{2}$   $f(n)$ 's virtual Zeroes and  $F(n)$ 's virtual  
 811 Pseudo-zeroes.

812  
 813 Applicable to  $\text{sim-}\eta(s)$  and DSPL, we note the computed CIS rational roots (subset)  
 814 as integers [rational numbers] + CIS irrational roots (subset) as irrational numbers = CIS  
 815 total roots.

816  
 817 **Remark 8.** Complex function  $F(n) = \text{DSPL}$  [representive of precise Area under the  
 818 Curve] generates the most accurate precise Areas of Varying Loops [when all rational  
 819 and irrational roots from combined base quantities (2n) and (2n-1) are utilized] and the  
 820 least accurate precise Areas of Varying Loops [when only rational roots from combined  
 821 base quantities (2n) and (2n-1) are utilized]; and complex function  $f(n) = \text{sim-}\eta(s)$  when  
 822 interpreted as Riemann sum [representive of approximate Area under the Curve] gener-  
 823 ates the most accurate approximate Areas of Varying Loops [when all rational and  
 824 irrational roots from combined base quantities (2n) and (2n-1) are utilized] and the least  
 825 accurate approximate Areas of Varying Loops [when only rational roots from combined  
 826 base quantities (2n) and (2n-1) are utilized].

827  
 828 Our [metaphoric] varying radius  $r$  in  $\text{sim-}\eta(s)$  or DSPL is defined as  $r = \text{Term-(2n)} -$   
 829  $\text{Term-(2n-1)}$  whereby perpetually recurring  $r = 0$  will correspond to Areas of Varying  
 830 Loops = 0 in order to obtain  $\sigma = \frac{1}{2}$   $f(n)$ 's Zeroes and  $F(n)$ 's Pseudo-zeroes or to obtain  
 831  $\sigma \neq \frac{1}{2}$   $f(n)$ 's virtual Zeroes and  $F(n)$ 's virtual Pseudo-zeroes. In effect, Areas of Varying  
 832 Loops is conceptionally synonymous with varying radius  $r$  whereby varying radius  
 833  $r$  could also be visualized as [metaphoric] varying distance  $d$  between **Term-(2n)** and  
 834 **Term-(2n-1)**.

835

836 **Remark 9.** Whether involving the most accurate method using total roots or the  
 837 least accurate method using rational roots to determine DSPL's precise or  $\text{sim-}\eta(s)$ 's  
 838 approximate Areas of Varying Loops, we can explicitly conclude all the infinitely-many  
 839 obtained Areas of Varying Loops are proportional and equal to varying radius  $r$  with  
 840 these Varying Loops being synthesized in a perpetually dynamic, cyclical and Incom-  
 841 pletely Predictable manner.

843 **7. Shift of Varying Loops in  $\zeta(\sigma + it)$  Polar Graph and Principle of Equidistant for**  
 844 **Multiplicative Inverse with General Equations for simplified Dirichlet eta function**  
 845 **and Dirichlet Sigma-Power Law**

846 We reiterate that  $\text{Gram}[x=0, y=0]$  points,  $\text{Gram}[y=0]$  points and  $\text{Gram}[x=0]$  points  
 847 are three types of IP Gram points [Zeroes] occurring at  $\sigma = \frac{1}{2}$  critical line (Figure 2)  
 848 based on, respectively, Origin intercept points, x-axis intercept points and y-axis intercept  
 849 points. They can be dependently computed from relevant types of  $\text{sim-}\eta(s) = 0$  equations  
 850 whereby  $\text{sim-}\eta(s)$  is obtained by applying Euler formula to  $\eta(s)$ .  $\text{Gram}[x=0, y=0]$  points  
 851 are synonymous with NTZ and  $\text{Gram}[y=0]$  points are synonymous with 'usual' Gram  
 852 points. Virtual  $\text{Gram}[y=0]$  points and virtual  $\text{Gram}[x=0]$  points are two types of IP virtual  
 853 Gram points [virtual Zeroes] occurring at  $\sigma \neq \frac{1}{2}$  non-critical lines based on, respectively,  
 854 x-axis intercept points and y-axis intercept points – see Figure 3 for  $\sigma = \frac{2}{5}$  and Figure 4  
 855 for  $\sigma = \frac{3}{5}$ . They are also dependently computed from these same equations.

856 **Proposition 2.** Both  $f(n)$   $\text{sim-}\eta(s)$  and  $F(n)$  DSPL will manifest Principle of Equidis-  
 857 tant for Multiplicative Inverse.

858 **Proof.** Let  $\delta = \frac{1}{10}$ . This will generate in Figure 3 and Figure 4 the  $\delta$  induced shift of  
 859 [infinitely many] Varying Loops in reference to Origin; viz, the simple relationship of  
 860 [more negative] left-shift given by  $\zeta(\frac{1}{2} - \delta + it)$  [Figure 3] < [neutral] nil-shift given by  
 861  $\zeta(\frac{1}{2} + it)$  [Figure 2] < [more positive] right-shifted given by  $\zeta(\frac{1}{2} + \delta + it)$  [Figure 4] will  
 862 always be consistently true.

863 Given  $\delta = \frac{1}{10}$ , the  $\sigma = \frac{1}{2} - \delta$  non-critical line (represented by Figure 3) and  $\sigma = \frac{1}{2} + \delta$   
 864 non-critical line (represented by Figure 4) are equidistant from  $\sigma = \frac{1}{2}$  critical line  
 865 (represented by Figure 2). The additive inverse operation of  $\sin(\delta) + \sin(-\delta) = 0$  indicating  
 866 symmetry with respect to Origin [or  $\cos(\delta) - \cos(-\delta) = 0$  indicating symmetry with respect  
 867 to y-axis] is not applicable to our complex single sine wave [or single cosine wave] since  
 868 **(2n)- or (2n-1)-complex term with transcendental functions** consisting of sine, cosine,  
 869 single sine wave, single cosine wave, natural logarithm are **independent of parameter  $\sigma$** .  
 870 However, **(2n)- or (2n-1)-complex term with algebraic functions** consisting of powers,  
 871 fractional powers, root extraction [and scaled amplitude R as alluded to by Remarks 3, 4  
 872 and 5 on its (in)dependency on parameter t] are **dependent on parameter  $\sigma$** .

873 Let  $x = (2n)$  or  $\frac{1}{(2n)}$  or  $(2n - 1)$  or  $\frac{1}{(2n - 1)}$ . With multiplicative inverse operation  
 874 of  $x^\delta \cdot x^{-\delta} = 1$  or  $\frac{1}{x^\delta} \cdot \frac{1}{x^{-\delta}} = 1$  that is applicable, this imply intrinsic presence of Multi-  
 875 plicative Inverse in  $\text{sim-}\eta(s)$  or DSPL for all  $\sigma$  values with this function or law rigidly  
 876 obeying relevant trigonometric identity. This phenomenon is **Principle of Equidistant**  
 877 **for Multiplicative Inverse**. Finally, we note by letting  $\delta = 0$ , we will always generate  
 878 Figure 2 representing  $\sigma = \frac{1}{2}$  critical line. *The proof is now complete for Proposition 2*  $\square$ .

879 For complex functions and complex equations in this paper,  $s = \sigma \pm it$  whereby  
 880 we commonly invoke  $s = \sigma + it$  for discussion. For all  $f(n)$  and  $F(n)$  general equations  
 881 depicted below without trigonometric identity application, we note presence of mixed  
 882 sine and cosine terms in these general equations except for  $f(n)$ 's  $\text{Gram}[y=0]$  points- $\text{sim-}\eta(s)$   
 883 and  $f(n)$ 's  $\text{Gram}[x=0]$  points- $\text{sim-}\eta(s)$ .

884 **I. NTZ or Gram  $[x=0, y=0]$  points** as geometrical Origin intercept points are mathe-  
 885 matically defined by  $\sum \text{ReIm}\{\eta(s)\} = \text{Re}\{\eta(s)\} + \text{Im}\{\eta(s)\} = 0$ . General equation for

$f(n)$ 's  $\text{sim-}\eta(s)$  as Zeroes is given by

$$\sum_{n=1}^{\infty} -(2n)^{-\sigma} (\sin(t \ln(2n)) - \cos(t \ln(2n))) - \sum_{n=1}^{\infty} -(2n-1)^{-\sigma} (\sin(t \ln(2n-1)) - \cos(t \ln(2n-1))) = 0 \quad (9)$$

General equation for  $F(n)$ 's DSPL with ability for Pseudo-zeroes to Zeroes conversion is given by

$$\frac{1}{2(t^2 + (\sigma - 1)^2)} \cdot \left[ (2n)^{1-\sigma} ((t + \sigma - 1) \sin(t \ln(2n)) + (t - \sigma + 1) \cdot \cos(t \ln(2n))) - (2n-1)^{1-\sigma} ((t + \sigma - 1) \cdot \sin(t \ln(2n-1)) + (t - \sigma + 1) \cos(t \ln(2n-1))) \right] + C \Big|_1^{\infty} = 0 \quad (10)$$

**II. Gram[y=0] points** as geometrical x-axis intercept points are mathematically defined by  $\sum \text{ReIm}\{\eta(s)\} = \text{Re}\{\eta(s)\} + 0$ , or simply  $\text{Im}\{\eta(s)\} = 0$ . General equation for  $f(n)$ 's Gram[y=0] points- $\text{sim-}\eta(s)$  as Zeroes is given by

$$\sum_{n=1}^{\infty} (2n)^{-\sigma} \sin(t \ln(2n)) - \sum_{n=1}^{\infty} (2n-1)^{-\sigma} \sin(t \ln(2n-1)) = 0 \quad (11)$$

General equation for  $F(n)$ 's Gram[y=0] points-DSPL with ability for Pseudo-zeroes to Zeroes conversion is given by

$$-\frac{1}{2(t^2 + (\sigma - 1)^2)} \cdot \left[ (2n)^{1-\sigma} ((\sigma - 1) \sin(t \ln(2n)) + t \cos(t \ln(2n))) - (2n-1)^{1-\sigma} ((\sigma - 1) \sin(t \ln(2n-1)) + t \cos(t \ln(2n-1))) \right] + C \Big|_1^{\infty} = 0 \quad (12)$$

**III. Gram[x=0] points** as geometrical y-axis intercept points are mathematically defined by  $\sum \text{ReIm}\{\eta(s)\} = 0 + \text{Im}\{\eta(s)\}$ , or simply  $\text{Re}\{\eta(s)\} = 0$ . General equation for  $f(n)$ 's Gram[x=0] points- $\text{sim-}\eta(s)$  as Zeroes is given by

$$\sum_{n=1}^{\infty} (2n)^{-\sigma} \cos(t \ln(2n)) - \sum_{n=1}^{\infty} (2n-1)^{-\sigma} \cos(t \ln(2n-1)) = 0 \quad (13)$$

886 General equation for  $F(n)$ 's Gram[x=0] points-DSPL with ability for Pseudo-zeroes  
887 to Zeroes conversion is given by

$$888 \frac{1}{2(t^2 + (\sigma - 1)^2)} \cdot \left[ (2n)^{1-\sigma} (t \sin(t \ln(2n)) - (\sigma - 1) \cos(t \ln(2n))) - (2n-1)^{1-\sigma} (t \sin(t \ln(2n-1)) - (\sigma - 1) \cos(t \ln(2n-1))) \right] + C \Big|_1^{\infty} = 0 \quad (14)$$

889 **Remark 10.** The Cartesian Coordinates  $(x,y)$  is intimately related to Polar Coordi-  
890 nates  $(r,\theta)$  with  $r = \sqrt{x^2 + y^2}$  and  $\theta = \tan^{-1}(\frac{y}{x})$ . In anti-clockwise direction, it has four  
891 quadrants defined by the + or - of  $(x,y)$ ; viz, Quadrant I as  $(+,+)$ , Quadrant II as  $(-,+)$ ,  
892 Quadrant III as  $(-,-)$ , and Quadrant IV as  $(+,-)$ .

893  
894 NTZ are Origin intercept points or Gram  $[x=0,y=0]$  points. With 'gap' being syn-  
895 onymous with 'interval', NTZ gap is given by initial NTZ t-value minus next NTZ  
896 t-value. Running a Full cycle from  $0\pi$  to  $2\pi$ , size of each IP Varying Loop in Figure  
897 2 is proportional to magnitude of its corresponding IP NTZ varying gap. We note the  
898  $2\pi$  here as observed in Figure 2 [on Gram points at  $\sigma = \frac{1}{2}$ ], Figure 3 [on virtual Gram  
899 points at  $\sigma = \frac{2}{5}$ ] and Figure 4 [on virtual Gram points at  $\sigma = \frac{3}{5}$ ] refers to IP Varying Loops  
900 transversed by parameter t with NTZ (Gram  $[x=0,y=0]$  points) corresponding to t values

901 as Origin intercept on Origin's solitary (0,0) part (point); Gram [y=0] points and virtual  
 902 Gram [y=0] points corresponding to t values as x-axis intercept on x-axis' (+ve)  $0\pi$  part  
 903 and (-ve)  $1\pi$  part; and Gram [x=0] points and virtual Gram [x=0] points corresponding  
 904 to t values as y-axis intercept on y-axis' (+ve)  $\frac{\pi}{2}$  part and (-ve)  $\frac{3\pi}{2}$  part. Virtual NTZ  
 905 entities do not exist; viz, Origin intercept points do not occur in Figure 3 and Figure 4.

906 With  $\eta(s)$  being proxy function for  $\zeta(s)$ , NTZ are defined by  $\eta(s) = 0$  or  $\text{sim-}\eta(s) = 0$ .  
 907 This mathematically-defined NTZ (or Gram[x=0,y=0] points) are precisely equivalent  
 908 to the geometrically-defined Origin intercept points. Then, NTZ given by relevant  
 909 computed IP t values are validly deduced to be infinite in magnitude since the  $\text{sim-}\eta(s)$   
 910 = 0 equation contains [complex] sine and/or cosine functions which are well-defined  
 911 continuous functions having infinitely many computed Origin intercept points located  
 912 on infinitely many Varying Loops generated by  $0 < t < +\infty$  or [its complex conjugate]  
 913  $-\infty < t < 0$  domain with unlimited range.

914 Riemann hypothesis is the original 1859-dated conjecture that all NTZ are located  
 915 on  $\sigma = \frac{1}{2}$  critical line of  $\zeta(s)$ . Mathematically proving all NTZ location on critical line as  
 916 denoted by solitary  $\sigma = \frac{1}{2}$  value equates to geometrically proving all Origin intercept  
 917 points occurrence at solitary  $\sigma = \frac{1}{2}$  value. Both result in rigorous proof for Riemann  
 918 hypothesis. Locations of first 10,000,000,000,000 NTZ on critical line have previously  
 919 been computed to be correct. Hardy[6], and with Littlewood[7], showed infinitely many  
 920 NTZ on  $\sigma = \frac{1}{2}$  critical line by considering moments of certain functions related to  $\zeta(s)$ .

921  
 922 **Remark 11.** The discovery by Hardy and Littlewood showing infinitely many NTZ  
 923 on  $\sigma = \frac{1}{2}$  critical line cannot constitute rigorous proof for Riemann hypothesis because  
 924 they have not exclude theoretical existence of NTZ in the region located away from the  
 925 critical line [whereby this region is denoted by the infinitely many  $\sigma \neq \frac{1}{2}$  non-critical  
 926 lines]. Furthermore, it is literally a mathematical impossibility ("mathematical impasse")  
 927 to be able to computationally check [in a successful manner] locations of all the infinitely  
 928 many NTZ are on the critical line.

929  
 930 The monumental task of solving Riemann hypothesis is completed by deriving  $F(n)$   
 931 DSPL from  $f(n)$   $\text{sim-}\eta(s)$  with its computed Pseudo-zeroes and virtual Pseudo-zeroes  
 932 which can all be converted to corresponding Zeroes and virtual Zeroes since  $F(n)$ 's IP  
 933 Pseudo-zeroes and IP virtual Pseudo-zeroes (t values) =  $f(n)$ 's IP Zeroes and IP virtual  
 934 Zeroes (t values) +  $\frac{\pi}{2}$  [for NTZ situation] whereby both  $f(n)$  and  $F(n)$  have parameters  $\sigma$   
 935 and t. Correctly deducing exact DA homogeneity in DSPL symbolizes rigorous proof  
 936 for Riemann hypothesis which is depicted as Pseudo-zeroes to Zeroes conversion that  
 937 obeys relevant trigonometric identities.

938 Three types of [traditionally] finite-interval Riemann Sums: Left / Right / Midpoint  
 939 Riemann Sum uses left endpoints / right endpoints / midpoints of the subintervals.  
 940 With  $n = 1, 2, 3, \dots, \infty$  and therefore  $\Delta n = 1$ , we note  $f(n)$  can analogically be interpreted  
 941 as approximate Area under the Curve (AUC) [right infinite-interval] Riemann sum

$$942 \sum_{n=1}^{\infty} f(n)\Delta n = \sum_{n=1}^{\infty} f(n) = \sum_{n=1}^2 f(n) + \sum_{n=3}^4 f(n) + \sum_{n=5}^6 f(n) + \dots + \sum_{n=\infty-1}^{\infty} f(n).$$

943 solution to exact AUC improper integral  $\int_{n=1}^{n=\infty} f(n)dn$  can be validly expanded as

$$944 \int_{n=1}^{n=2} f(n)dn + \int_{n=2}^{n=3} f(n)dn + \int_{n=3}^{n=4} f(n)dn + \dots + \int_{n=\infty-1}^{n=\infty} f(n)dn = [F(n) + C]_1^2 + [F(n) +$$

945  $C]_2^3 + [F(n) + C]_3^4 + \dots + [F(n) + C]_{\infty-1}^{\infty}$  which, for all sufficiently large n as  $n \rightarrow \infty$ ,

946 will manifest *divergence by oscillation* (viz. for all sufficiently large n as  $n \rightarrow \infty$ , this  
 947 cumulative total will not diverge in a particular direction to a solitary well-defined

948 limit value since the [complex] sine and/or cosine terms present in  $\text{sim-}\eta(s)$  and DSPL

949 are periodic transcendental-type functions). Evaluation of definite integrals Eq. (3) or

950 Eq. (10), Eq. (6) or Eq. (12) and Eq. (8) or Eq. (14) using limit as  $n \rightarrow +\infty$  for  $0 < t < +\infty$

951 enable countless computations resulting in t values for (respectively) CIS NTZ, CIS

952 Gram[y=0] points and CIS Gram[x=0] points [all as Pseudo-zeroes to Zeroes conversion].  
 953 Larger n values used for computations will correspond to increasing accuracy of these  
 954 entities.

955 **Remark 12.** Whereas exact AUC from F(n) given by DSPL =  $\int_{n=1}^{n=\infty} \text{sim} - \eta(s) dn$   
 956 and approximate AUC from f(n) given by  $\text{sim} - \eta(s) = \sum_{n=1}^{\infty} \text{sim} - \eta(s)$  [when interpreted as  
 957 Riemann sum] are proportional; the Zeroes when indirectly derived from DSPL [as  
 958 Pseudo-zeroes converted to Zeroes] and the Zeroes when directly derived from  $\text{sim} - \eta(s)$   
 959 must agree with each other at  $\sigma = \frac{1}{2}$  critical line.

960 **8. Riemann zeta function, Dirichlet eta function, simplified Dirichlet eta function and**  
 961 **Dirichlet Sigma-Power Law**

962  $\zeta(s)$  is a function of complex variable s (=  $\sigma \pm it$ ) that analytically continues sum of  
 963 infinite series  $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \dots$ . The common convention is to write  
 964 s as  $\sigma + it$  with  $i = \sqrt{-1}$ , and with  $\sigma$  and t real. Valid for  $\sigma > 0$ , we write  $\zeta(s)$  as  
 965  $Re\{\zeta(s)\} + iIm\{\zeta(s)\}$  and note that  $\zeta(\sigma + it)$  when  $0 < t < +\infty$  is the complex conjugate  
 966 of  $\zeta(\sigma - it)$  when  $-\infty < t < 0$ .

967 Also known as alternating zeta function,  $\eta(s)$  must act as proxy for  $\zeta(s)$  in crit-  
 968 ical strip (viz.  $0 < \sigma < 1$ ) containing critical line (viz.  $\sigma = \frac{1}{2}$ ) because  $\zeta(s)$  only  
 969 converges when  $\sigma > 1$ . This implies  $\zeta(s)$  is undefined to left of this  $\sigma > 1$  region  
 970 [in the critical strip] which then requires  $\eta(s)$  representation instead. They are re-  
 971 lated to each other as  $\zeta(s) = \gamma \cdot \eta(s)$  with proportionality factor  $\gamma = \frac{1}{(1 - 2^{1-s})}$  and

972 
$$\eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s} = \frac{1}{1^s} - \frac{1}{2^s} + \frac{1}{3^s} - \dots$$
  
 973

$$\begin{aligned} \zeta(s) &= \sum_{n=1}^{\infty} \frac{1}{n^s} && (15) \\ &= \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \dots \\ &= \prod_{p \text{ prime}} \frac{1}{(1 - p^{-s})} \\ &= \frac{1}{(1 - 2^{-s})} \cdot \frac{1}{(1 - 3^{-s})} \cdot \frac{1}{(1 - 5^{-s})} \cdot \frac{1}{(1 - 7^{-s})} \cdot \frac{1}{(1 - 11^{-s})} \dots \frac{1}{(1 - p^{-s})} \dots \end{aligned}$$

974  
 975 Eq. (15) is defined for only  $1 < \sigma < \infty$  region where  $\zeta(s)$  is absolutely convergent with  
 976 no zeros located here. In Eq. (15), equivalent Euler product formula with product over  
 977 prime numbers [instead of summation over natural numbers] also represents  $\zeta(s) \implies$   
 978 all prime and, by default, composite numbers are (intrinsically) encoded in  $\zeta(s)$ . Brief  
 979 diversion: On April 17, 2013, Zhang[8] announced a ground-breaking proof stating  
 980 there are infinitely many pairs of prime numbers that differ by 70 million or less. This  
 981 result implies the existence of an infinitely repeatable prime 2-tuple, thus establishing a  
 982 theorem akin to the twin prime conjecture.

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \cdot \Gamma(1 - s) \cdot \zeta(1 - s) \tag{16}$$

983  
 984 With  $\sigma = \frac{1}{2}$  as symmetry line of reflection, Eq. (16) is Riemann's functional equation  
 985 valid for  $-\infty < \sigma < \infty$ . It can be used to find all trivial zeros on horizontal line at  $it =$   
 986 0 occurring when  $\sigma = -2, -4, -6, -8, -10, \dots, \infty$  whereby  $\zeta(s) = 0$  because factor  $\sin\left(\frac{\pi s}{2}\right)$   
 987 vanishes.  $\Gamma$  is gamma function, an extension of factorial function [a product function

988 denoted by ! notation whereby  $n! = n(n-1)(n-2)\dots(n-(n-1))$  with its argument  
 989 shifted down by 1, to real and complex numbers. That is, if  $n$  is a positive integer,  
 990  $\Gamma(n) = (n-1)!$

$$\begin{aligned}\zeta(s) &= \frac{1}{(1-2^{1-s})} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s} \\ &= \frac{1}{(1-2^{1-s})} \left( \frac{1}{1^s} - \frac{1}{2^s} + \frac{1}{3^s} - \dots \right)\end{aligned}\quad (17)$$

991 Eq. (17) is defined for all  $\sigma > 0$  values except for simple pole at  $\sigma = 1$ . As alluded  
 992 to above,  $\zeta(s)$  without  $\frac{1}{(1-2^{1-s})}$  viz.  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s}$  is  $\eta(s)$ . It is a holomorphic function  
 993 of  $s$  defined by analytic continuation and is mathematically defined at  $\sigma = 1$  whereby  
 994 analogous trivial zeros with presence for  $\eta(s)$  [but not for  $\zeta(s)$ ] on vertical straight line  $\sigma$   
 995  $= 1$  are found at  $s = 1 \pm i \frac{2\pi k}{\ln(2)}$  where  $k = 1, 2, 3, 4, \dots, \infty$ .

996 Euler formula can be stated as  $e^{in} = \cos n + i \cdot \sin n$ . Euler identity (where  $n = \pi$ )  
 997 is  $e^{i\pi} = \cos \pi + i \cdot \sin \pi = -1 + 0$  [or stated as  $e^{i\pi} + 1 = 0$ ]. The  $n^s$  of  $\zeta(s)$  is expanded  
 998 to  $n^s = n^{(\sigma+it)} = n^\sigma e^{t \ln(n) \cdot i}$  since  $n^t = e^{t \ln(n)}$ . Apply Euler formula to  $n^s$  result in  
 999  $n^s = n^\sigma (\cos(t \ln(n)) + i \cdot \sin(t \ln(n)))$ . This is written in trigonometric form [designated  
 1000 by short-hand notation  $n^s(Euler)$ ] whereby  $n^\sigma$  is modulus and  $t \ln(n)$  is polar angle  
 1001 (argument).

1002 We apply  $n^s(Euler)$  to Eq. (17) to obtain  $f(n)$  general  $\text{sim-}\eta(s)$  for determining  $\sigma = \frac{1}{2}$   
 1003 NTZ versus (non-existent)  $\sigma \neq \frac{1}{2}$  virtual NTZ[4], section 4, p. 24 - 28. At  $\sigma = \frac{1}{2}$ , this  
 1004 is given as Eq. (9) and with the trigonometric identity application as Eq. (1). Integrate  
 1005  $f(n)$  general  $\text{sim-}\eta(s)$  to obtain  $F(n)$  general DSPL for determining  $\sigma = \frac{1}{2}$  Pseudo-zeros  
 1006 versus (non-existent)  $\sigma \neq \frac{1}{2}$  virtual Pseudo-zeros. Pseudo-zeros and (non-existent)  
 1007 virtual Pseudo-zeros can be converted to Zeroes (NTZ) and (non-existent) virtual Zeroes  
 1008 (virtual NTZ). At  $\sigma = \frac{1}{2}$ , this is given as Eq. (10) and with the trigonometric identity  
 1009 application as Eq. (3).

1010 We provide  $f(n)$  general Gram[ $y=0$ ] points- $\text{sim-}\eta(s)$  for determining  $\sigma = \frac{1}{2}$  Gram[ $y=0$ ]  
 1011 points versus  $\sigma \neq \frac{1}{2}$  virtual Gram[ $y=0$ ] points[4], section 5, p. 28 - 30. At  $\sigma = \frac{1}{2}$ , this  
 1012 is given as Eq. (11) but we are unable to apply trigonometric identity. Integrate  $f(n)$   
 1013 general Gram[ $y=0$ ] points- $\text{sim-}\eta(s)$  to obtain  $F(n)$  general Gram[ $y=0$ ] points-DSPL for  
 1014 determining  $\sigma = \frac{1}{2}$  Pseudo-zeros versus  $\sigma \neq \frac{1}{2}$  virtual Pseudo-zeros. Pseudo-zeros  
 1015 and virtual Pseudo-zeros can be converted to Zeroes (Gram[ $y=0$ ] points) and virtual  
 1016 Zeroes (virtual Gram[ $y=0$ ] points). At  $\sigma = \frac{1}{2}$ , this is given as Eq. (12) and with the  
 1017 trigonometric identity application as Eq. (6).

1018 We provide  $f(n)$  general Gram[ $x=0$ ] points- $\text{sim-}\eta(s)$  for determining  $\sigma = \frac{1}{2}$  Gram[ $x=0$ ]  
 1019 points versus  $\sigma \neq \frac{1}{2}$  virtual Gram[ $x=0$ ] points[4], section 5, p. 28 - 30. At  $\sigma = \frac{1}{2}$ , this  
 1020 is given as Eq. (13) but we are unable to apply trigonometric identity. Integrate  $f(n)$   
 1021 general Gram[ $x=0$ ] points- $\text{sim-}\eta(s)$  to obtain  $F(n)$  general Gram[ $x=0$ ] points-DSPL for  
 1022 determining  $\sigma = \frac{1}{2}$  Pseudo-zeros versus  $\sigma \neq \frac{1}{2}$  virtual Pseudo-zeros. Pseudo-zeros  
 1023 and virtual Pseudo-zeros can be converted to Zeroes (Gram[ $x=0$ ] points) and virtual  
 1024 Zeroes (virtual Gram[ $x=0$ ] points). At  $\sigma = \frac{1}{2}$ , this is given as Eq. (14) and with the  
 1025 trigonometric identity application as Eq. (8).

## 1026 9. Conclusions

1027 Previously regarded as **primary spin-offs**[4], correct and complete mathematical  
 1028 arguments for solving the 1859 Riemann hypothesis, and explaining the closely related  
 1029 Gram[ $y=0$ ] points and Gram[ $x=0$ ] points, can inherently be classified as belonging to  
 1030 Mathematics for Incompletely Predictable problems.



1031 "With this one solution [for Riemann hypothesis], we have proven five hundred theorems  
 1032 or more at once". Previously regarded as **secondary spin-offs**[4] arising out of solving  
 1033 Riemann hypothesis, this profound statement apply to many important theorems in  
 1034 Number theory (mostly on prime numbers) that rely on properties of Riemann zeta  
 1035 functions such as where trivial zeros and nontrivial zeros are / are not located.

1036 Derived innovative *Fic-Fac Ratio* was previously regarded as **tertiary spin-offs**[4]  
 1037 serving as medical or epidemiological tool to assist understanding of SARS-CoV-2 caus-  
 1038 ing COVID-19 and 2020 Coronavirus Pandemic. Unprecedented negative global health  
 1039 and economic impacts have arised from this event. Fic-Fac Ratio connects seemingly  
 1040 unrelated subject of Medicine with frontier Mathematics from Number theory.

1041 There are concrete analogies between the Completely Predictable entities even  
 1042 and odd numbers [that are all located on unique 'linear' lines] versus the Incompletely  
 1043 Predictable entities prime and odd numbers, NTZ, Gram[y=0] points and Gram[x=0]  
 1044 points [that are all located on unique 'non-linear' lines]. We can either conceptionally  
 1045 or mathematically derive valid intrinsic properties such as the actual gaps/intervals  
 1046 between any two adjacent entities, and the various slope/gradient (involving the calcu-  
 1047 lus of differentiation) and Area-under-the-Curve (involving the calculus of integration)  
 1048 of these lines which will all be given by continuous functions that are always defined  
 1049 for any arbitrarily chosen intervals [a,b] except the following. The corresponding lines  
 1050 computed from complex algorithms that generate all prime and composite numbers  
 1051 are only defined at two end-points a,b but not for interval [a,b] as these algorithms are  
 1052 simply not well-defined functions. We immediately recognize these complex algorithms  
 1053 [which are not functions] are not amendable to differentiation or integration. Then as  
 1054 succinctly outlined below, the previously published quantitative and qualitative rigorous  
 1055 proofs[4] for Polignac's and Twin prime conjectures cannot be stated using functions.

1056 **Quantitative proof:** We validly exclude first and only even prime number (**P**) '2', and  
 1057 show from following mathematical arguments that Polignac's and Twin prime conjec-  
 1058 tures are true with appearance of  $\aleph_0$  cardinality 'uniformity' conforming to Dimensional  
 1059 analysis homogeneity. Let (i) cardinality  $T = \aleph_0$  for Set **all odd P** derived from even  
 1060 number (**E**) prime gaps 2, 4, 6,...,  $\infty$ , (ii) cardinality  $T_2 = \aleph_0$  for Subset **odd P** derived  
 1061 from **E** prime gap 2, cardinality  $T_4 = \aleph_0$  for Subset **odd P** derived from **E** prime gap 4,  
 1062 cardinality  $T_6 = \aleph_0$  for Subset **odd P** derived from **E** prime gap 6, etc. Paradoxically, (as  
 1063 sets)  $T = T_2 + T_4 + T_6 + \dots + T_\infty$  equation is valid despite (their cardinality)  $T = T_2 = T_4 =$   
 1064  $T_6 = \dots = T_\infty$ ; and **E** prime gaps are 'infinite in magnitude' can justifiably be perceived  
 1065 instead as 'arbitrarily large in magnitude' since cumulative sum total of **E** prime gaps  
 1066 is relatively much slower to attain the 'infinite in magnitude' status when compared to  
 1067 cumulative sum total of **P** which rapidly attain this status.

1068 **Qualitative proof:** Plus-Minus Gap 2 Composite Number Alternating Law has built-in  
 1069 intrinsic mechanism to automatically generate all prime gaps  $\geq 4$  in a mathematically  
 1070 consistent *ad infinitum* manner. Plus Gap 2 Composite Number Continuous Law has  
 1071 built-in intrinsic mechanism to automatically generate prime gap = 2 appearances in a  
 1072 mathematically consistent *ad infinitum* manner. These two deduced Laws "**that must**  
 1073 **crucially involve both prime and composite numbers being dependently and algorithm-**  
 1074 **ically tabulated together with subsequent analysis on their [consequently combined]**  
 1075 **corresponding gaps"** will qualitatively confirm Polignac's and Twin prime conjectures to  
 1076 be true.

1077 In this paper, we have intrinsically treated and analyzed in a *de novo* fashion  
 1078 simple and complex single-variable function  $f(n)$  or  $F(n)$  and their simple and complex  
 1079 single-variable equation  $f(n) = 0$  or  $F(n) = 0$  as Completely Predictable or Incompletely  
 1080 Predictable mathematical objects. Being mutually exclusive and Incompletely Predictable  
 1081 entities with  $\sigma = \frac{1}{2}$  critical line depicted in Figure 2, and  $\sigma \neq \frac{1}{2}$  non-critical lines as  
 1082 exemplified by  $\sigma = \frac{2}{5}$  depicted in Figure 3 and  $\sigma = \frac{3}{5}$  depicted in Figure 4; **the  $\sigma = \frac{1}{2}$**   
 1083 **NTZ computed from Eq. (1) -  $\sigma \neq \frac{1}{2}$  (non-existent) virtual NTZ computed from Eq. (2)**  
 1084 **Pairing** outlined at the end of section 3 mathematically serve to validly distinguish and

1085 separate the unique complete set of nontrivial zeros from the infinitely many non-unique  
 1086 complete sets of (non-existent) virtual nontrivial zeros. The critical line of Riemann zeta  
 1087 function is denoted by  $\sigma = \frac{1}{2}$  whereby all nontrivial zeros are proposed to be located in the  
 1088 1859 Riemann hypothesis. Our Dirichlet Sigma-Power Law symbolizes the end-product  
 1089 proof on Riemann hypothesis.

1090 We reiterate the following important criteria: The three types (three separate "con-  
 1091 tainers") of Gram points at  $\sigma = \frac{1}{2}$  and two types (two separate "containers") of virtual  
 1092 Gram points at  $\sigma \neq \frac{1}{2}$  are labelled together as *Zeroes*. After performing integration on  
 1093 relevant  $f(n)$  resulting in  $F(n)$ , we obtain corresponding three types (three separate  
 1094 "containers") of Pseudo-Gram points at  $\sigma = \frac{1}{2}$  and two types (two separate "containers")  
 1095 of virtual Pseudo-Gram points at  $\sigma \neq \frac{1}{2}$  which are labelled together as *Pseudo-zeroes*.

1096 With groundings in *Mathematics for Incompletely Predictable problems*, we advocate  
 1097 that we have now provided a comparatively elementary and rigorous proof on Riemann  
 1098 hypothesis while explaining existence of mutually exclusive three types of [Incompletely  
 1099 Predictable] Gram points and two types of [Incompletely Predictable] virtual Gram  
 1100 points. These achievements are completed with appropriate analysis on complex (meta-)  
 1101 properties present in Dirichlet Sigma-Power Law, Gram[y=0] points-Dirichlet Sigma-  
 1102 Power Law and Gram[x=0] points-Dirichlet Sigma-Power Law that give rise to relevant  
 1103 Pseudo-Gram points; and in virtual Gram[y=0] points-Dirichlet Sigma-Power Law and  
 1104 virtual Gram[x=0] points-Dirichlet Sigma-Power Law that give rise to relevant virtual  
 1105 Pseudo-Gram points. Exact Dimensional analysis homogeneity [occurring only once at  $\sigma$   
 1106  $= \frac{1}{2}$  critical line] in these Laws is endowed with ability to convert their computed Pseudo-  
 1107 zeroes to Zeroes resulting in nontrivial zeros (Origin intercept points or Gram[x=0,y=0]  
 1108 points) as one type of Gram points plus Gram[y=0] points and Gram[x=0] points as two  
 1109 remaining types of Gram points. Inexact Dimensional analysis homogeneity [occurring  
 1110 infinitely often at  $\sigma \neq \frac{1}{2}$  non-critical lines] in these Laws is endowed with ability to  
 1111 convert their computed virtual Pseudo-zeroes to virtual Zeroes resulting in virtual  
 1112 Gram[y=0] points and virtual Gram[x=0] points as two types of virtual Gram points.

#### 1113 *Acknowledgements*

1114  
 1115

#### *References*

1. Koblitz, N. (1984). p-adic interpolation of the Riemann zeta-function. In: p-adic Numbers, p-adic Analysis, and Zeta-Functions. Graduate Texts in Mathematics, vol 58. Springer, New York, NY. [https://doi.org/10.1007/978-1-4612-1112-9\\_2](https://doi.org/10.1007/978-1-4612-1112-9_2)
2. Scholze, P. (2012). Perfectoid Spaces. *Publ. math. IHES* 116, pp. 245–313. <https://doi.org/10.1007/s10240-012-0042-x>
3. Fargues, L. & Scholze, P. (2021). Geometrization of the local Langlands correspondence. *arXiv:2102.13459* pp. 1–350. <https://arxiv.org/abs/2102.13459> [Accessed 4 October 2021]
4. Ting, J.Y.C. (2020). Mathematical Modelling of COVID-19 and Solving Riemann Hypothesis, Polignac's and Twin Prime Conjectures Using Novel Fic-Fac Ratio With Manifestations of Chaos-Fractal Phenomena. *J. Math. Res.*, 12(6) pp. 1-49. <https://doi.org/10.5539/jmr.v12n6p1>
5. Landau, E. Vorlesungen uber Zahlentheorie. *New York: Chelsea*, (2) pp. 183-308.
6. Hardy, G. H. (1914). Sur les Zeros de la Fonction  $\zeta(s)$  de Riemann. *C. R. Acad. Sci. Paris*, 158, pp. 1012-1014. JFM 45.0716.04 Reprinted in (Borwein et al., 2008)
7. Hardy, G. H. & Littlewood, J. E. (1921). The zeros of Riemann's zeta-function on the critical line. *Math. Z.*, 10 (3-4), pp. 283-317. <http://dx.doi:10.1007/BF01211614>

8. Zhang, Y. (2014). Bounded gaps between primes. *Ann. of Math.*, 179, pp. 1121-1174. <http://dx.doi.org/10.4007/annals.2014.179>.
9. Noe, T. (2004). A100967. The On-line Encyclopedia of Integer Sequences. <https://oeis.org/A100967>
10. Ting, J. (2013). A228186. The On-Line Encyclopedia of Integer Sequences. <https://oeis.org/A228186>
11. Weisstein, E. W. (2006). A114856. The On-line Encyclopedia of Integer Sequences. <https://oeis.org/A114856>
12. Greathouse IV, C. R. (2012). A216700. The On-line Encyclopedia of Integer Sequences. <https://oeis.org/A216700>
13. Rosser, J. B.; Yohe, J. M.; Schoenfeld, L. (1969). Rigorous computation and the zeros of the Riemann zeta-function. (With discussion). Information Processing 68 (Proc. IFIP Congress, Edinburgh, 1968), Vol. 1: Mathematics, Software, Amsterdam: North-Holland, pp. 70–76, MR 0258245
14. Trudgian, T. (2011). On the success and failure of Gram's Law and the Rosser Rule. *Acta Arithmetica*, 148 (3), pp. 225-256. <http://dx.doi.org/10.4064/aa148-3-2>
15. Trudgian, T. (2014), An improved upper bound for the argument of the Riemann zeta function on the critical line II. *J. Number Theory*, 134, pp. 280-292. <http://dx.doi.org/10.1016/j.jnt.2013.07.017>

### Appendix A. Gram's Law and Rosser Rule for Gram points

Named after Danish mathematician Jørgen Pedersen Gram (June 27, 1850 – April 29, 1916), ['traditional'/'usual'] Gram points or (mathematical) Gram[y=0] points or (geometrical) x-axis intercept points are other conjugate pairs values in Riemann zeta function  $\zeta(s)$  on  $\sigma = \frac{1}{2}$  critical line. Then  $s = \frac{1}{2} + it$  gives rise to  $\zeta(\frac{1}{2} + it)$  on critical line; and Gram points when defined in terms of  $\zeta(s)$  is given by  $\sum ReIm\{\zeta(s)\} = Re\{\zeta(s)\} + 0$ , or simply  $Im\{\zeta(s)\} = 0$ . Alternatively defined using expression denoting  $\zeta(s)$  on critical line  $\zeta(\frac{1}{2} + it) = Z(t)e^{-i\theta(t)}$  whereby Hardy's function, Z, is real for real t, and  $\theta$  is Riemann–Siegel theta function given in terms of gamma function as  $\theta(t) = \arg\left(\Gamma\left(\frac{1}{4} + \frac{it}{2}\right)\right) - \frac{\log \pi}{2}t$  for real values of t; we note that  $\zeta(s)$  is real when  $\sin(\theta(t)) = 0$ . This implies that  $\theta(t)$  is an integer multiple of  $\pi$  which allows for location of Gram points to be calculated easily by inverting the formula for  $\theta$ . Gram points are historically [crudely] numbered as  $g_n$  for  $n = 0, 1, 2, 3, \dots$ , whereby  $g_n$  is the unique solution of  $\theta(t) = n\pi$ . Here,  $n = 0$  is the [first]  $g_0$  value of 17.8455995405... which is larger than the smallest [first] positive nontrivial zeros (NTZ) value of 14.13472515.... Thus,  $n = -3$  correspond to  $g_{-3} = 0$ ,  $n = -2$  correspond to  $g_{-2} = 3.4362182261\dots$ , and  $n = -1$  correspond to  $g_{-1} = 9.6669080561\dots$

*Paired [infinite-length] integer sequences with prestigious connections:*

A100967+0, which is A100967[9], is precisely defined as "Least k such that  $\text{binomial}(2k+1, k-n-1) \geq \text{binomial}(2k, k)$  viz.  $(2k+1)!k! \geq (2k)!(k-n-1)!(k+n+2)!$ ". The terms commencing from Position 0, 1, 2, 3,... of A100967+0 are listed below: 3, 9, 18, 29, 44, 61, 81, 104, 130, 159, 191, 225, 263, 303, 347, 393, 442, 494, 549, 606, 667, 730, 797, 866, 938, 1013, 1091, 1172, 1255, 1342, 1431, 1524, 1619, 1717, 1818, 1922, 2029, 2138, 2251, 2366, 2485, 2606, 2730, 2857, 2987, 3119, 3255, 3394, 3535,....

A100967+1 is precisely defined as "Add 1 to each and every terms from A100967+0". The terms commencing from Position 0, 1, 2, 3,... of A100967+1 are listed below: 4, 10, 19, 30, 45, 62, 82, 105, 131, 160, 192, 226, 264, 304, 348, 394, 443, 495, 550, 607, 668, 731, 798, 867, 939, 1014, 1092, 1173, 1256, 1343, 1432, 1525, 1620, 1718, 1819, 1923, 2030, 2139, 2252, 2367, 2486, 2607, 2731, 2858, 2988, 3120, 3256, 3395, 3536,....

A228186[10] is precisely defined as "Smallest natural number  $k > n$  such that  $(k+n+1)!(k-n-2)! < 2k!(k-1)!$ " or alternatively defined as "Greatest natural number  $k > n$  such that calculated peak values for ratio  $R = \frac{\text{CombinationsWithRepetition}}{\text{CombinationsWithoutRepetition}}$

$= \frac{(k+n-1)!(n-k)!}{n!(n-1)!}$  belong to maximal rational numbers  $< 2$ ". The terms commencing from Position 0, 1, 2, 3,... of A228186 are listed below: 4, 9, 18, 29, 44, 61, 81, 104, 130, 159, 191, 226, 263, 304, 347, 393, 442, 494, 549, 607, 667, 731, 797, 866, 938, 1013, 1091, 1172, 1256, 1342, 1432, 1524, 1619, 1717, 1818, 1922, 2029, 2139, 2251, 2367, 2485, 2606, 2730, 2857, 2987, 3120, 3255, 3394, 3535,....

Unexpected connection [and unrelated to NTZ and Gram points]: A228186 can be considered an innovative [infinite-length] "Hybrid integer sequence" identical to "non-Hybrid integer sequence" A100967+0 except for the interspersed [finite] 21 'exceptional' terms located at Position 0, 11, 13, 19, 21, 28, 30, 37, 39, 45, 50, 51, 52, 55, 57, 62, 66, 70, 73, 77, and 81 with their corresponding 21 values exactly specified by [infinite-length] "non-Hybrid integer sequence" A100967+1.

A114856-"bad"-Gram-points, which is A114856[11], is precisely defined as "Indices n of Gram points  $g_n$  for which  $(-1)^n Z(g_n) < 0$  with Z(t) being Riemann-Siegel Z-function and full given range of values  $n = 0, 1, 2, 3, \dots$ ". The terms of A114856-"bad"-Gram-points are listed below: 126, 134, 195, 211, 232, 254, 288, 367, 377, 379, 397, 400, 461, 507, 518, 529,

567, 578, 595, 618, 626, 637, 654, 668, 692, 694, 703, 715, 728, 766, 777, 793, 795, 807, 819, 848, 857, 869, 887, 964, 992, 995, 1016, 1028, 1034, 1043, 1046, 1071, 1086,....

A114856-"good"-Gram-points, given by "total"-Gram points minus A114856-"bad"-Gram-points, is precisely defined as "Indices  $n$  of Gram points  $g_n$  for which  $(-1)^n Z(g_n) > 0$  with  $Z(t)$  being Riemann-Siegel Z-function and full given range of values  $n = 0, 1, 2, 3, \dots$ ". The derived terms of A114856-"good"-Gram-points: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50,....

A216700[12] is precisely defined as "Violations of Rosser Rule: numbers  $n$  such that the Gram block  $[g_n, g_{n+k}]$  contains fewer than  $k$  points  $t$  such that  $Z(t) = 0$  with  $Z(t)$  being Riemann-Siegel Z-function and full given range of values  $n = 0, 1, 2, 3, \dots$ ". The terms of A216700 are listed below: 13999525, 30783329, 30930927, 37592215, 40870156, 43628107, 46082042, 46875667, 49624541, 50799238, 55221454, 56948780, 60515663, 61331766, 69784844, 75052114, 79545241, 79652248, 83088043, 83689523, 85348958, 86513820, 87947597,....

Expected connection: All NTZ (as conjectured by Riemann hypothesis) and Gram points (by definition) are located on the same critical line of Riemann zeta function. Counting NTZ can be validly reduced to counting all Gram points where Gram's Law is satisfied and adding count of NTZ inside each Gram block. With this process, we need not locate NTZ but just have to accurately compute  $Z(t)$  to show that it changes sign.

Gram's Law is the observation that there is [usually] exactly one NTZ (Gram $[x=0,y=0]$  points or Origin intercept points) between any two "good" Gram points. Examples of closely related statements equivalent to Gram's law are:  $(-1)^n Z(g_n)$  is [usually] positive or  $Z(t)$  [usually] has opposite sign at consecutive Gram points. Thus, a  $t$ -valued Gram point is called a "good" Gram point if  $\zeta(s)$  is positive at  $\frac{1}{2} + it$  with  $(-1)^n Z(g_n) > 0$  and a "bad" Gram point if  $\zeta(s)$  is negative at  $\frac{1}{2} + it$  with  $(-1)^n Z(g_n) < 0$ . The indices of "bad" Gram points where  $Z$  has the 'wrong' sign are given by A114856 in OEIS. A Gram block  $[g_n, g_{n+k}]$  is a half-open interval bounded by two "good" Gram points  $g_n$  and  $g_{n+k}$  such that all Gram points  $g_{n+1}, \dots, g_{n+k-1}$  between them are "bad" Gram points. A refinement of Gram's Law is known as Rosser Rule[13] which stated that Gram blocks [usually] have the expected number of NTZ in them (identical to number of Gram intervals), even though some of the individual Gram intervals in the block may not have exactly one NTZ in them. Example, the interval bounded by  $g_{125}$  and  $g_{127}$  is a Gram block containing a unique "bad" Gram point  $g_{126}$  and the expected number 2 of NTZ although neither of its two Gram intervals contains a unique NTZ.

Gram's Law and Rosser Rule both imply that in some sense NTZ do not stray too far from their expected positions, and that they hold most of the time but are violated infinitely often (in an Incompletely Predictable manner)[14],[15]. Professor Timothy Trudgian in 2011 explicitly showed that both Gram's Law and Rosser Rule fail in a positive proportion of cases. In particular, it is expected that in about 73% [ $\approx \frac{3}{4}$ ] one NTZ is enclosed by two successive Gram points [and thus Gram's Law fails for about 27% [ $\approx \frac{1}{4}$ ] of all Gram intervals to contain exactly one NTZ], but in about 14% no NTZ and in about 13% two NTZ are in such a Gram interval on the long run.

## Appendix B. Miscellaneous Materials

**Cardinality:** With increasing size, arbitrary Set  $X$  can be CFS, CIS or UIS. Cardinality of Set  $X$ ,  $|X|$ , measures *number of elements* in Set  $X$ . E.g. Set **negative Gram $[y=0]$  point** has CFS of negative Gram $[y=0]$  point with **|negative Gram $[y=0]$  point|** = 1, Set **even Prime number** has CFS of even **Prime number** with **|even Prime number|** = 1, Set **Natural numbers** has CIS of **Natural numbers** with **|Natural numbers|** =  $\aleph_0$ , and Set **Real numbers** has UIS of **Real numbers** with **|Real numbers|** =  $\mathfrak{c}$  (cardinality of the continuum). Let  $\mathbb{C}$  = UIS complex numbers,  $\mathbb{R}$  = UIS real numbers,  $\mathbb{Q}$  = CIS rational numbers that include fractional numbers and rational roots,  $\mathbb{R}-\mathbb{Q}$  = UIS total irrational numbers,  $\mathbb{A}$  = CIS algebraic numbers,  $\mathbb{R}-\mathbb{A}$  = UIS transcendental irrational numbers,  $\mathbb{Z}$  = CIS integers which are literally fractional numbers with denominator 1,  $\mathbb{W}$  = CIS whole numbers,  $\mathbb{N}$  = CIS natural numbers,  $\mathbb{E}$  = CIS even numbers,  $\mathbb{O}$  = CIS odd numbers,  $\mathbb{P}$  = CIS prime numbers, and  $\mathbb{C}$  = CIS composite numbers. CIS  $\mathbb{N}$  = Set  $\mathbb{E}$  [whereby we did not include the zeroth even number  $E_0 = 0$ ] + Set  $\mathbb{O}$ ; CIS  $\mathbb{N}$  = CIS  $\mathbb{P}$  + CIS  $\mathbb{C}$  + CFS Number 1; and CIS  $\mathbb{N} \subset$  CIS  $\mathbb{W} \subset$  CIS  $\mathbb{Z} \subset$  CIS  $\mathbb{Q} \subset$  UIS  $\mathbb{R} \subset$  UIS  $\mathbb{C}$ . CIS  $\mathbb{A}$  as  $\mathbb{C}$  (including  $\mathbb{R}$ ) = CIS  $\mathbb{Q}$  that include fractional numbers and rational roots + CIS irrational roots whereby both rational and irrational roots are derived from non-zero polynomials.

The following refined definitions are useful: UIS total irrational numbers = CIS irrational roots (numbers) + UIS transcendental irrational numbers whereby transcendental irrational numbers  $\gg$  [algebraic] irrational numbers. Whereas CIS rational roots (numbers), CIS irrational roots (numbers) and UIS transcendental numbers are treated separately as mutually exclusive numbers; so must the existing algebraic functions that generate CIS rational roots (numbers) and CIS irrational roots (numbers), and the existing transcendental functions that generate UIS transcendental numbers be treated separately as mutually exclusive functions.

**Certain types of infinite series:** An algebraic function [such as rational functions, square root, cube root function, etc] satisfies a polynomial equation. A transcendental function [such as exponential function, natural logarithm, trigonometric

functions, hyperbolic functions, gamma, elliptic, zeta functions, etc] is an analytic function that does not satisfy a polynomial equation. Thus a transcendental function "transcends" algebra since it cannot be expressed in terms of a finite sequence of algebraic operations consisting of addition, subtraction, multiplication, division, powers, and fractional powers or root extraction. All integers, rational numbers, rational or irrational roots of real and complex numbers are algebraic numbers e.g. a root of polynomial  $x^2 - x - 1 =$  golden ratio  $\varphi = \frac{1 + \sqrt{5}}{2} = 1.618033\dots$ , square root of 2 viz,  $\sqrt[2]{2}$  or  $\sqrt{2} = 2^{\frac{1}{2}} = 1.414213\dots$ , or cube root of 2 viz,  $\sqrt[3]{2} = 2^{\frac{1}{3}} \approx 1.259921$ . Real and complex numbers that are not algebraic numbers e.g.  $\pi$  and  $e$  are transcendental numbers. However, we note sine and cosine as transcendental functions generally give rise to mutually exclusive sets of transcendental numbers except at discrete points such as  $\sin \frac{\pi}{6} = \sin 30^\circ = \cos \frac{2\pi}{6} = \cos 60^\circ = \frac{\sqrt{1}}{2} = \frac{1}{2}$  [viz, transcendental functions generating an algebraic number as rational root (number) at certain discrete points].

Following [side-note] treatise of interest involve infinite series. A property of irrational number  $\sqrt{2}$  is  $\frac{1}{\sqrt{2}-1} = \sqrt{2} + 1$  since  $(\sqrt{2} + 1)(\sqrt{2} - 1) = 2 - 1 = 1$ . This is related to the property of silver ratios.  $\sqrt{2}$  can also be expressed in terms of copies of imaginary unit  $i$  using only square root and arithmetic operations, if the square root symbol is interpreted suitably for complex numbers  $i$  and  $-i$ :  $\frac{\sqrt{i} + i\sqrt{i}}{i}$  and  $\frac{\sqrt{-i} - i\sqrt{-i}}{-i}$ . Multiplicative inverse (reciprocal) of  $(2)^{\frac{1}{2}}$  or  $\sqrt{2}$  is  $(2)^{-\frac{1}{2}}$  or  $\sqrt{\frac{1}{2}}$  which is a unique [irrational number] constant since  $\sqrt{\frac{1}{2}} = \frac{1}{\sqrt{2}} = \frac{1}{2}\sqrt{2} = \cos \frac{\pi}{4} = \sin \frac{\pi}{4}$ . Transcendental numbers such as  $\frac{\pi}{4}$  (given by Leibniz series  $\frac{1}{1} - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \dots \approx 0.78539816$ ); and  $\frac{\pi^2}{6}$  (given by  $\zeta(2) = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots \approx 1.6449340668482$ ), respectively, encode complete set of alternating odd and, by default, alternating even numbers; and natural numbers. Also known as alternating zeta function, Dirichlet eta function  $\eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s}$  when expanded, will intrinsically encode complete set of alternating natural numbers e.g.  $\eta(1) = \ln(2)$  (given by  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} = \sum_{n=2}^{\infty} \frac{1}{2^n} [\zeta(n) - 1] + \frac{1}{2} = \frac{1}{1} - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots \approx 0.69314718056$ ). Equivalent Euler product formula for  $\zeta(s)$  with product over prime numbers [instead of summation over natural numbers] will intrinsically encode complete set of prime and, by default, composite numbers. As an extra point, complete set of alternating prime and, by default, alternating composite numbers is encoded in converging alternating series  $\sum_{k=1}^{\infty} \frac{(-1)^k}{p_k} \approx -0.2696063519$  (transcendental number) when fully expanded whereby  $p_k$  is  $k^{\text{th}}$  prime number.

**Zeroes and Pseudo-zeroes:** There are three types of stationary points in a given [simple] periodic  $f(n)$  involving sine and/or cosine functions that act as  $x$ -axis intercept points via three types of  $f(n)$ 's Zeroes with corresponding three types of  $F(n)$ 's Pseudo-zeroes: maximum points e.g. with  $f(n)$  or  $F(n) = \sin n - 1$ ; minimum points e.g. with  $f(n)$  or  $F(n) = \sin n + 1$ ; and points of inflection e.g. with  $f(n)$  or  $F(n) = \sin n$  [which also has Origin intercept point as a Zero or Pseudo-zero]. A fourth type of  $f(n)$ 's Zeroes and  $F(n)$ 's Pseudo-zeroes consist of non-stationary points occurring e.g. with  $f(n)$  or  $F(n) = \sin n + 0.5$ . One can analogically assimilate these concepts to aesthetically explain the more "exotic" characteristics manifested by [complex] periodic  $f(n)$  or  $F(n)$  involving sine and/or cosine functions that are present in  $f(n)$  sim- $\eta(s)$  or  $F(n)$  DSPL at (solitary)  $\sigma = \frac{1}{2}$  critical line and (infinitely many)  $\sigma \neq \frac{1}{2}$  non-critical lines.

With  $(j - i) = (l - k) = 2\pi$  [viz, one Full cycle], let a given Zero be located in  $f(n)$ 's interval  $[i, j]$  viz,  $i < \text{Zero} < j$ ; and its corresponding Pseudo-zero be located in  $F(n)$ 's Pseudo-interval  $[k, l]$  viz,  $k < \text{Pseudo-zero} < l$ . For this Zero and Pseudo-zero characterized by either point of inflection or non-stationary point; both will comply with preserving positivity [going from (-ve) below  $x$ -axis to (+ve) above  $x$ -axis] as explained using the Zero case [with the Pseudo-zero case following similar lines of explanations]. This can be stated as follow for interval  $[i, j]$ : If  $j > i$ , then computed  $f(j) >$  computed  $f(i)$ . In particular, the condition "If  $i \geq 0$ , then computed  $f(i) \geq 0$ " must not be present for these two particular types of Zero to validly exist in interval  $[i, j]$ . With reversal of inequality signs, converse situation for  $j < \text{Zero} < i$  and corresponding  $l < \text{Pseudo-zero} < k$  is equally true in preserving negativity [going from (+ve) above  $x$ -axis to (-ve) below  $x$ -axis]. These are useful properties on Zeroes and Pseudo-zeroes.

**Preservation or conservation of Net Area Value and Total Area Value with definitions[4], p. 10 - 13:**  $\int f(n)dn = F(n) + C$  with  $F'(n) = f(n)$ . Consider a nominated function  $f(n)$  for interval  $[a, b]$ . We define Net Area Value (NAV) calculated using its antiderivative  $F(n)$  as the net difference between positive area value(s) [above horizontal  $x$ -axis] and

negative area value(s) [below horizontal x-axis] in interval [a,b]; viz, NAV = all +ve value(s) + all -ve value(s). Again calculated using  $F(n)$ , we define Total Area Value (TAV) as the total sum of (absolute value) positive area value(s) [above horizontal x-axis] and (absolute value) negative area value(s) [below horizontal x-axis] in interval [a,b]; viz, TAV = all |+ve value(s)| + all |-ve value(s)|. Calculated NAV and TAV are precise using antiderivative  $F(n)$  obtained from integration of  $f(n)$  but are only approximate when using Riemann sum on  $f(n)$ . For  $f(n)$ 's interval [a,b] whereby a = initial Zero and b = next Zero, and  $F(n)$ 's Pseudo interval [c,d] whereby c = initial Pseudo-zero and d = next Pseudo-zero; then compliance with preservation or conservation of NAV and TAV will simultaneously occur in both  $f(n)$ 's Zeroes and  $F(n)$ 's Pseudo-zeroes given by their sine and/or cosine functions only when Zero gap = (b - a) = Pseudo-zero gap = (d - c) =  $2\pi$  [viz, involving one Full cycle]. For our purpose, NAV = 0 condition is validly preserved or conserved for  $f(n)$   $\sin\text{-}\eta(s)$ 's IP Zeroes and  $F(n)$  DSPL's IP Pseudo-zeroes at parameter  $\sigma = \frac{1}{2}$ . Ditto for  $f(n)$   $\sin\text{-}\eta(s)$ 's IP virtual Zeroes and  $F(n)$  DSPL's IP virtual Pseudo-zeroes at parameter  $\sigma \neq \frac{1}{2}$ ; viz, NAV = 0 condition is validly preserved or conserved for  $f(n)$   $\sin\text{-}\eta(s)$ 's IP virtual Zeroes and  $F(n)$  DSPL's IP virtual Pseudo-zeroes.

For single-term trigonometric function  $f(n) = \sin(n)$ , it is an odd function with Origin symmetry since  $-f(n) = f(-n)$  for all n. The  $f(n) = \sin(n)$  has an infinite number of CP x-axis intercept points (Zeroes) and a solitary unique Origin intercept point (Zero) since it belong to a class of odd functions that is defined at  $n = 0$  and must pass through the Origin. Otherwise, the other class of odd functions such as  $f(n) = \sin(\frac{1}{n})$  with infinite number of CP x-axis intercept points (Zeroes) but without Origin intercept point [since  $\sin(\frac{1}{n})$  is undefined at  $n = 0$ ] can remain symmetrical about the Origin without actually passing through it. For single term trigonometric function  $f(n) = \cos(n)$  with symmetry about the y-axis, it is an even function since  $f(n) = f(-n)$  for all n. It has an infinite number of CP x-axis intercept points (Zeroes). Being undefined at  $n = 0$ , it will never have Origin intercept point.

For dual terms trigonometric functions  $f(n) = \cos(n) - \sin(n)$  and  $f(n) = \cos(n) + \sin(n)$ , they are neither even nor odd without any symmetry. They both have an infinite number of CP x-axis intercept points (Zeroes). Being undefined at  $n = 0$ , they will never have Origin intercept point. Special properties for Addition and Multiplication: The sum or difference of two even functions is even. The sum or difference of two odd functions is odd. The sum or difference of an even and odd function is neither even nor odd unless one function is zero; viz, there is (exactly) one function that is both even and odd, and it is the zero function  $f(n) = 0$ . The product of two even functions is an even function. The product of two odd functions is an even function. The product of an even function and an odd function is an odd function.

**Trigonometric identity for the linear combination of sine and cosine functions:** Here, we again use simple single-variable function  $f(n)$  or  $F(n)$ . The trigonometric identity for linear combination of sine and cosine  $a\cos(n) + b\sin(n)$  can be freely, arbitrarily and interchangeably written as either [simple] single cosine wave  $R\cos(n - \alpha)$  or [simple] single sine wave  $R\sin(n + \alpha)$  whereby R is the scaled amplitude and  $\alpha$  is the phase shift.  $R = \sqrt{a^2 + b^2} = (a^2 + b^2)^{\frac{1}{2}}$ . Since  $\sin(\alpha) = \frac{b}{\sqrt{a^2 + b^2}}$  and  $\cos(\alpha) = \frac{a}{\sqrt{a^2 + b^2}}$ , then  $\alpha = \tan^{-1}\frac{b}{a}$ . Below, we assign  $\sqrt{2}$  to equivalently denote  $2^{\frac{1}{2}}$ .

$$\text{With } a = 1, b = -1, R = \sqrt{2}; \cos(n) - \sin(n) = \sqrt{2} \cos\left(n + \frac{1}{4}\pi\right) = \sqrt{2} \sin\left(n + \frac{3}{4}\pi\right).$$

$$\text{With } a = -1, b = 1, R = \sqrt{2}; -\cos(n) + \sin(n) = \sqrt{2} \sin\left(n - \frac{1}{4}\pi\right) = \sqrt{2} \cos\left(n - \frac{3}{4}\pi\right).$$

$$\text{With } a = 1, b = 1, R = \sqrt{2}; \cos(n) + \sin(n) = \sqrt{2} \cos\left(n - \frac{1}{4}\pi\right) = \sqrt{2} \sin\left(n + \frac{1}{4}\pi\right).$$

$$\text{With } a = -1, b = -1, R = \sqrt{2}; -\cos(n) - \sin(n) = \sqrt{2} \cos\left(n + \frac{3}{4}\pi\right) = \sqrt{2} \sin\left(n - \frac{3}{4}\pi\right).$$

$\int f(n)dn = F(n) + C$  with  $F'(n) = f(n)$ . With  $|a| = 1$  and  $|b| = 1$ , consider single-term [simple] trigonometric functions:  $f(n) = a\cos(n)$  which belongs to an even function and  $f(n) = b\sin(n)$  which belongs to an odd function. Whereas all linear combination of [simple]  $\cos(n)$  and [simple]  $\sin(n)$  as sum or difference such as  $f(n) = \cos(n) + \sin(n)$  and  $f(n) = \cos(n) - \sin(n)$  belong to neither even nor odd functions, then their corresponding  $F(n)$  being linear combination of [simple]  $\cos(n)$  and [simple]  $\sin(n)$  as sum or difference must also belong to neither even nor odd functions. With both  $f(n)$  and corresponding  $F(n)$  considered as [simple] functions and relevant trigonometric identities being applied, they can intrinsically and arbitrarily be expressed as either [simple] single cosine wave or [simple] single sine wave containing a phase shift  $\frac{1}{4}\pi$  or  $\frac{3}{4}\pi$  and a scaled amplitude  $\sqrt{2} [= 2^{\frac{1}{2}}$  which is base 2 endowed with exponent  $\frac{1}{2}$ ]. Respectively,  $F(n)$  and  $f(n)$  have an infinite number of x-axis intercept points called Pseudo-zeroes and Zeroes but nil Origin intercept points.