

Funcoids and Reloids*

a generalization of proximities and uniformities

BY VICTOR PORTON

August 10, 2011

Abstract

It is a part of my Algebraic General Topology research.

In this article, I introduce the concepts of *funcoids*, which generalize proximity spaces and *reloids*, which generalize uniform spaces. The concept of funcoid is generalized concept of proximity, the concept of reloid is cleared from superfluous details (generalized) concept of uniformity. Also funcoids generalize pretopologies and preclosures. Also funcoids and reloids are generalizations of binary relations whose domains and ranges are filters (instead of sets).

Also funcoids and reloids can be considered as a generalization of (oriented) graphs, this provides us with a common generalization of analysis and discrete mathematics.

The concept of continuity is defined by an algebraic formula (instead of old messy epsilon-delta notation) for arbitrary morphisms (including funcoids and reloids) of a partially ordered category. In one formula are generalized continuity, proximity continuity, and uniform continuity.

Keywords: algebraic general topology, quasi-uniform spaces, generalizations of proximity spaces, generalizations of nearness spaces, generalizations of uniform spaces

A.M.S. subject classification: 54J05, 54A05, 54D99, 54E05, 54E15, 54E17, 54E99

Table of contents

1 Common	2
1.1 Draft status	2
1.2 Earlier works	2
1.3 Used concepts, notation and statements	3
1.3.1 Filters	3
2 Partially ordered dagger categories	4
2.1 Partially ordered categories	4
2.2 Dagger categories	4
2.2.1 Some special classes of morphisms	5
3 Funcoids	6
3.1 Informal introduction into funcoids	6
3.2 Basic definitions	7
3.2.1 Composition of funcoids	8
3.3 Funcoid as continuation	8
3.4 Lattices of funcoids	11
3.5 More on composition of funcoids	12
3.6 Domain and range of a funcoid	13
3.7 Categories of funcoids	14
3.8 Specifying funcoids by functions or relations on atomic filter objects	14
3.9 Direct product of filter objects	16
3.10 Atomic funcoids	18
3.11 Complete funcoids	20
3.12 Completion of funcoids	23
3.13 Monovalued and injective funcoids	25

*. This document has been written using the GNU \TeX _{MACS} text editor (see www.texmacs.org).

3.14	T_0 -, T_1 - and T_2 -separable functors	26
3.15	Filter objects closed regarding a functor	26
4	Reloids	27
4.1	Composition of reloids	27
4.2	Direct product of filter objects	28
4.3	Restricting reloid to a filter object. Domain and image	29
4.4	Categories of reloids	31
4.5	Monovalued and injective reloids	31
4.6	Complete reloids and completion of reloids	32
5	Relationships between functors and reloids	35
5.1	Functor induced by a reloid	35
5.2	Reloids induced by functor	38
5.3	Galois connections of functors and reloids	39
6	Continuous morphisms	40
6.1	Traditional definitions of continuity	40
6.1.1	Pre-topology	40
6.1.2	Proximity spaces	41
6.1.3	Uniform spaces	41
6.2	Our three definitions of continuity	41
6.3	Continuousness of a restricted morphism	42
7	Connectedness regarding functors and reloids	43
7.1	Some lemmas	43
7.2	Endomorphism series	44
7.3	Connectedness regarding binary relations	44
7.4	Connectedness regarding functors and reloids	45
7.5	Algebraic properties of S and S^*	47
8	Postface	48
8.1	Misc	48
Appendix A Some counter-examples		48
9	Second product. Oblique product	51
Bibliography		51

1 Common

1.1 Draft status

This article is a draft, an almost ready preprint.

This text refers to a preprint edition of [14]. Theorem number clashes may appear due editing both of these manuscripts.

1.2 Earlier works

Some mathematicians researched generalizations of proximities and uniformities before me but they have failed to reach the right degree of generalization which is presented in this work allowing to represent properties of spaces with algebraic (or categorical) formulas.

Proximity structures were introduced by Smirnov in [4].

Some references to predecessors:

- In [5], [6], [11], [2], [18] are studied generalized uniformities and proximities.
- Proximities and uniformities are also studied in [9], [10], [17], [19], [20].
- [7] and [8] contains recent progress in quasi-uniform spaces. [8] has a very long list of related literature.

Some works ([16]) about proximity spaces consider relationships of proximities and compact topological spaces. In this work is not done the attempt to define or research their generalization, compactness of funcoids or reloids. It seems potentially productive to attempt to borrow the definitions and procedures from the above mentioned works. I hope to do this study in a separate article.

[3] studies mappings between proximity structures. (In this work no attempt to research mappings between funcoids is done.) [12] researches relationships of quasi-uniform spaces and topological spaces. [1] studies how proximity structures can be treated as uniform structures and compactification regarding proximity and uniform spaces.

1.3 Used concepts, notation and statements

The set of functions from a set A to a set B is denoted as B^A .

I will often skip parentheses and write fx instead of $f(x)$ to denote the result of a function f acting on the argument x .

I will call *small* sets members of some Grothendieck universe. (Let us assume the axiom of existence of a Grothendieck universe.)

Let f is a small binary relation.

I will denote $\langle f \rangle X = \{f\alpha \mid \alpha \in X\}$ and $X[f]Y \Leftrightarrow \exists x \in X, y \in Y: xfy$ for small sets X, Y .

By just $\langle f \rangle$ and $[f]$ I will denote the corresponding function and relation on small sets.

$\lambda x \in D: f(x) = \{(x; f(x)) \mid x \in D\}$ for every formula $f(x)$ depended on a variable x and set D .

I will denote the least and the greatest element of a poset \mathfrak{A} as $0^{\mathfrak{A}}$ and $1^{\mathfrak{A}}$ respectively.

For elements a and b of a lattice with a minimal element I will denote $a \asymp b$ when $a \cap b$ is the minimal element of the lattice and $a \not\asymp b$ otherwise. See [14] for a more general notion.

1.3.1 Filters

In this work the word *filter* will refer to a filter on a set (in contrast to [14] where filters are considered on arbitrary posets). Note that I do not require filters to be proper.

I will call the set of filters on a set A (*base set*) ordered reverse to set-theoretic inclusion of filters *the set of filter objects* on A and denote it $\mathfrak{F}(A)$ or just \mathfrak{F} when the base set is implied and call its element *filter objects* (f.o. for short). I will denote $\uparrow \mathcal{F}$ the filter corresponding to a filter object \mathcal{F} . So we have $\mathcal{A} \subseteq \mathcal{B} \Leftrightarrow \uparrow \mathcal{A} \supseteq \uparrow \mathcal{B}$ for every filter objects \mathcal{A} and \mathcal{B} on the same set.

In this particular manuscript, we will *not* equate principal filter objects with corresponding sets as it is done in [14]. Instead we will have $\text{Base}(\mathcal{A})$ equal to the unique base of a f.o. \mathcal{A} . I will denote $\uparrow^A X$ (or just $\uparrow X$ when A is implied) the principal filter object on A corresponding to the set X .

Filters are studied in the work [14].

Every set $\mathfrak{F}(A)$ is a complete lattice and we will apply lattice operations to subsets of such sets without explicitly mentioning $\mathfrak{F}(A)$.

Prior reading of [14] is needed to fully understand this work.

Filter objects corresponding to ultrafilters are atoms of the lattice $\mathfrak{F}(A)$ and will be called *atomic filter objects* (on A).

Also we will need to introduce the concept of *generalized filter base*.

Definition 1. *Generalized filter base* is a set $S \in \mathcal{P}\mathfrak{F} \setminus \{0^{\mathfrak{F}}\}$ such that

$$\forall \mathcal{A}, \mathcal{B} \in S \exists \mathcal{C} \in S: \mathcal{C} \subseteq \mathcal{A} \cap \mathcal{B}.$$

Proposition 2. Let S is a generalized filter base. If $\mathcal{A}_1, \dots, \mathcal{A}_n \in S$ ($n \in \mathbb{N}$), then

$$\exists \mathcal{C} \in S: \mathcal{C} \subseteq \mathcal{A}_1 \cap \dots \cap \mathcal{A}_n.$$

Proof. Can be easily proved by induction. □

Theorem 3. If S is a generalized filter base, then $\text{up} \bigcap S = \bigcup \langle \text{up} \rangle S$.

Proof. Obviously $\text{up} \bigcap S \supseteq \bigcup \langle \text{up} \rangle S$. Reversely, let $K \in \text{up} \bigcap S$; then $K = A_1 \cap \dots \cap A_n$ where $A_i \in \text{up} \mathcal{A}_i$ where $\mathcal{A}_i \in S$, $i = 1, \dots, n$, $n \in \mathbb{N}$; so exists $\mathcal{C} \in S$ such that $\mathcal{C} \subseteq A_1 \cap \dots \cap A_n \subseteq \uparrow(A_1 \cap \dots \cap A_n) = \uparrow K$, $K \in \text{up} \mathcal{C}$, $K \in \bigcup \langle \text{up} \rangle S$. \square

Corollary 4. If S is a generalized filter base, then $\bigcap S = 0^{\mathfrak{F}} \Leftrightarrow 0^{\mathfrak{S}} \in S$.

Proof. $\bigcap S = 0^{\mathfrak{F}} \Leftrightarrow \emptyset \in \text{up} \bigcap S \Leftrightarrow \emptyset \in \bigcup \langle \text{up} \rangle S \Leftrightarrow \exists \mathcal{X} \in S: \emptyset \in \text{up} \mathcal{X} \Leftrightarrow 0^{\mathfrak{S}} \in S$. \square

Obvious 5. If S is a filter base on a set A then $\langle \uparrow^A \rangle S$ is a generalized filter base.

Definition 6. I will call *shifted filtrator* a triple $(\mathfrak{A}; \mathfrak{B}; \uparrow)$ where \mathfrak{A} and \mathfrak{B} are posets and \uparrow is an order embedding from \mathfrak{B} to \mathfrak{A} .

Some concepts and notation can be defined for shifted filtrators through similar concepts for filtrators: $\langle \uparrow \rangle \text{up} a = \text{up}^{(\mathfrak{A}; \langle \uparrow \rangle \mathfrak{B})} a$; $\langle \uparrow \rangle \text{Cor} a = \text{Cor}^{(\mathfrak{A}; \langle \uparrow \rangle \mathfrak{B})} a$, etc.

For a set \mathfrak{A} and the set of f.o. \mathfrak{F} on this set we will consider the shifted filtrator $(\mathfrak{F}; \mathfrak{A}; \uparrow)$.

2 Partially ordered dagger categories

2.1 Partially ordered categories

Definition 7. I will call a *partially ordered (pre)category* a (pre)category together with partial order \subseteq on each of its Hom-sets with the additional requirement that

$$f_1 \subseteq f_2 \wedge g_1 \subseteq g_2 \Rightarrow g_1 \circ f_1 \subseteq g_2 \circ f_2$$

for every morphisms f_1, g_1, f_2, g_2 such that $\text{Src } f_1 = \text{Src } f_2 \wedge \text{Dst } f_1 = \text{Dst } f_2 = \text{Src } g_1 = \text{Src } g_2 \wedge \text{Dst } g_1 = \text{Dst } g_2$.

2.2 Dagger categories

Definition 8. I will call a *dagger precategory* a precategory together with an involutive contravariant identity-on-objects prefunctor $x \mapsto x^\dagger$.

In other words, a *dagger precategory* is a precategory equipped with a function $x \mapsto x^\dagger$ on its set of morphisms which reverses the source and the destination and is subject to the following identities for every morphisms f and g :

1. $f^{\dagger\dagger} = f$;
2. $(g \circ f)^\dagger = f^\dagger \circ g^\dagger$.

Definition 9. I will call a *dagger category* a category together with an involutive contravariant identity-on-objects functor $x \mapsto x^\dagger$.

In other words, a *dagger category* is a category equipped with a function $x \mapsto x^\dagger$ on its set of morphisms which reverses the source and the destination and is subject to the following identities for every morphisms f and g and object A :

1. $f^{\dagger\dagger} = f$;
2. $(g \circ f)^\dagger = f^\dagger \circ g^\dagger$;
3. $(1_A)^\dagger = 1_A$.

Theorem 10. If a category is a dagger precategory then it is a dagger category.

Proof. We need to prove only that $(1_A)^\dagger = 1_A$. Really

$$(1_A)^\dagger = (1_A)^\dagger \circ 1_A = (1_A)^\dagger \circ (1_A)^{\dagger\dagger} = ((1_A)^\dagger \circ 1_A)^\dagger = (1_A)^{\dagger\dagger} = 1_A. \quad \square$$

For a partially ordered dagger (pre)category I will additionally require (for every morphisms f and g)

$$f^\dagger \subseteq g^\dagger \Leftrightarrow f \subseteq g.$$

An example of dagger category is the category **Rel** whose objects are sets and whose morphisms are binary relations between these sets with usual composition of binary relations and with $f^\dagger = f^{-1}$.

Definition 11. A morphism f of a dagger category is called *unitary* when it is an isomorphism and $f^\dagger = f^{-1}$.

Definition 12. *Symmetric* (endo)morphism of a dagger precategory is such a morphism f that $f = f^\dagger$.

Definition 13. *Transitive* (endo)morphism of a precategory is such a morphism f that $f = f \circ f$.

Theorem 14. The following conditions are equivalent for a morphism f of a dagger precategory:

1. f is symmetric and transitive.
2. $f = f^\dagger \circ f$.

Proof.

(1) \Rightarrow (2). If f is symmetric and transitive then $f^\dagger \circ f = f \circ f = f$.

(2) \Rightarrow (1). $f^\dagger = (f^\dagger \circ f)^\dagger = f^\dagger \circ f^{\dagger\dagger} = f^\dagger \circ f = f$, so f is symmetric. $f = f^\dagger \circ f = f \circ f$, so f is transitive. \square

2.2.1 Some special classes of morphisms

Definition 15. For a partially ordered dagger category I will call *monovalued* morphism such a morphism f that $f \circ f^\dagger \subseteq 1_{\text{Dst } f}$.

Definition 16. For a partially ordered dagger category I will call *entirely defined* morphism such a morphism f that $f^\dagger \circ f \supseteq 1_{\text{Src } f}$.

Definition 17. For a partially ordered dagger category I will call *injective* morphism such a morphism f that $f^\dagger \circ f \subseteq 1_{\text{Src } f}$.

Definition 18. For a partially ordered dagger category I will call *surjective* morphism such a morphism f that $f \circ f^\dagger \supseteq 1_{\text{Dst } f}$.

Remark 19. It's easy to show that this is a generalization of monovalued, entirely defined, injective, and surjective binary relations as morphisms of the category **Rel**.

Obvious 20. “Injective morphism” is a dual of “monovalued morphism” and “surjective morphism” is a dual of “entirely defined morphism”.

Definition 21. For a given partially ordered dagger category C the *category of monovalued (entirely defined, injective, surjective) morphisms* of C is the category with the same set of objects as of C and the set of morphisms being the set of monovalued (entirely defined, injective, surjective) morphisms of C with the composition of morphisms the same as in C .

We need to prove that these are really categories, that is that composition of monovalued (entirely defined) morphisms is monovalued (entirely defined) and that identity morphisms are monovalued and entirely defined.

Proof. We will prove only for monovalued morphisms and entirely defined morphisms, as injective and surjective morphisms are their duals.

Monovalued. Let f and g are monovalued morphisms, $\text{Dst } f = \text{Src } g$. $(g \circ f) \circ (g \circ f)^\dagger = g \circ f \circ f^\dagger \circ g^\dagger \subseteq g \circ 1_{\text{Dst } f} \circ g^\dagger = g \circ 1_{\text{Src } g} \circ g^\dagger = g \circ g^\dagger \subseteq 1_{\text{Dst } g} = 1_{\text{Dst } (g \circ f)}$. So $g \circ f$ is monovalued.

That identity morphisms are monovalued follows from the following: $1_A \circ (1_A)^\dagger = 1_A \circ 1_A = 1_A = 1_{\text{Dst } 1_A} \subseteq 1_{\text{Dst } 1_A}$.

Entirely defined. Let f and g are entirely defined morphisms, $\text{Dst } f = \text{Src } g$. $(g \circ f)^\dagger \circ (g \circ f) = f^\dagger \circ g^\dagger \circ g \circ f \supseteq f^\dagger \circ 1_{\text{Src } g} \circ f = f^\dagger \circ 1_{\text{Dst } f} \circ f = f^\dagger \circ f \supseteq 1_{\text{Src } f} = 1_{\text{Src}(g \circ f)}$. So $g \circ f$ is entirely defined.

That identity morphisms are entirely defined follows from the following: $(1_A)^\dagger \circ 1_A = 1_A \circ 1_A = 1_A = 1_{\text{Src } 1_A} \supseteq 1_{\text{Src } 1_A}$. \square

Definition 22. I will call a *bijjective* morphism a morphism which is entirely defined, monovalued, injective, and surjective.

Obvious 23. Bijjective morphisms form a full subcategory.

Proposition 24. If a morphism is bijjective then it is an isomorphism.

Proof. Let f is bijjective. Then $f \circ f^\dagger \subseteq 1_{\text{Dst } f}$, $f^\dagger \circ f \supseteq 1_{\text{Src } f}$, $f^\dagger \circ f \subseteq 1_{\text{Src } f}$, $f \circ f^\dagger \supseteq 1_{\text{Dst } f}$. Thus $f \circ f^\dagger = 1_{\text{Dst } f}$ and $f^\dagger \circ f = 1_{\text{Src } f}$ that is f^\dagger is an inverse of f . \square

3 Funcoids

3.1 Informal introduction into funcoids

Funcoids are a generalization of proximity spaces and a generalization of pretopological spaces. Also funcoids are a generalization of binary relations.

That funcoids are a common generalization of “spaces” (proximity spaces, (pre)topological spaces) and binary relations (including monovalued functions) makes them smart for describing properties of functions in regard of spaces. For example the statement “ f is a continuous function from a space μ to a space ν ” can be described in terms of funcoids as the formula $f \circ \mu \subseteq \nu \circ f$ (see below for details).

Most naturally funcoids appear as a generalization of proximity spaces.

Let δ be a proximity that is certain binary relation so that $A \delta B$ is defined for every sets A and B . We will extend it from sets to filter objects by the formula:

$$\mathcal{A} \delta' \mathcal{B} \Leftrightarrow \forall A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B}: A \delta B.$$

Then (as it will be proved below) there exist two functions $\alpha, \beta \in \mathfrak{F}^\delta$ such that

$$\mathcal{A} \delta' \mathcal{B} \Leftrightarrow \mathcal{B} \cap^\delta \alpha \mathcal{A} \neq 0^\delta \Leftrightarrow \mathcal{A} \cap^\delta \beta \mathcal{B} \neq 0^\delta.$$

The pair $(\alpha; \beta)$ is called *funcoid* when $\mathcal{B} \cap^\delta \alpha \mathcal{A} \neq 0^\delta \Leftrightarrow \mathcal{A} \cap^\delta \beta \mathcal{B} \neq 0^\delta$. So funcoids are a generalization of proximity spaces.

Funcoids consist of two components the first α and the second β . The first component of a funcoid f is denoted as $\langle f \rangle$ and the second component is denoted as $\langle f^{-1} \rangle$. (The similarity of this notation with the notation for the image of a set under a function is not a coincidence, we will see that in the case of discrete funcoids (see below) these coincide.)

One of the most important properties of a funcoid is that it is uniquely determined by just one of its components. That is a funcoid f is uniquely determined by the function $\langle f \rangle$. Moreover a funcoid f is uniquely determined by $\langle f \rangle|_{\mathcal{P} \cup \text{dom } \langle f \rangle}$ that is by values of function $\langle f \rangle$ on sets.

Next we will consider some examples of funcoids determined by specified values of the first component on sets.

Funcoids as a generalization of pretopological spaces: Let α be a pretopological space that is a map $\alpha \in \mathfrak{F}^\delta$ for some set \mathcal{U} . Then we define $\alpha' X \stackrel{\text{def}}{=} \bigcup^\delta \{\alpha x \mid x \in X\}$ for every set $X \in \mathcal{P} \mathcal{U}$. We will prove that there exists a unique funcoid f such that $\alpha' = \langle f \rangle|_{\mathcal{P} \mathcal{U}}$. So funcoids are a generalization of pretopological spaces. Funcoids are also a generalization of preclosure operators: For every preclosure operator p on a set \mathcal{U} exists a unique funcoid f such that $\langle f \rangle|_{\mathcal{P} \mathcal{U}} = \uparrow \circ p$.

For every binary relation p on a set \mathcal{U} it exists unique funcooid f such that $\forall X \in \mathcal{P}\mathcal{U}: \langle f \rangle \uparrow X = \uparrow \langle p \rangle X$ (where $\langle p \rangle$ is defined in the introduction), recall that a funcooid is uniquely determined by the values of its first component on sets. I will call such funcooids *discrete*. So funcooids are a generalization of binary relations.

Composition of binary relations (i.e. of discrete funcooids) complies with the formulas:

$$\langle g \circ f \rangle = \langle g \rangle \circ \langle f \rangle \quad \text{and} \quad \langle (g \circ f)^{-1} \rangle = \langle f^{-1} \rangle \circ \langle g^{-1} \rangle.$$

By the same formulas we can define composition of every two funcooids. Funcooids with this composition form a category (*the category of funcooids*).

Also funcooids can be reversed (like reversal of X and Y in a binary relation) by the formula $\langle \alpha; \beta \rangle^{-1} = \langle \beta; \alpha \rangle$. In particular case if μ is a proximity we have $\mu^{-1} = \mu$ because proximities are symmetric.

Funcooids behave similarly to (multivalued) functions but acting on filter objects instead of acting on sets. Below these will be defined domain and image of a funcooid (the domain and the image of a funcooid are filter objects).

3.2 Basic definitions

Definition 25. Let's call a *funcooid* from a set A to a set B a quadruple $(A; B; \alpha; \beta)$ where $\alpha \in \mathfrak{F}(B)^{\mathfrak{F}(A)}$, $\beta \in \mathfrak{F}(A)^{\mathfrak{F}(B)}$ such that

$$\forall \mathcal{X} \in \mathfrak{F}(A), \mathcal{Y} \in \mathfrak{F}(B): (\mathcal{Y} \not\prec \alpha \mathcal{X} \Leftrightarrow \mathcal{X} \not\prec \beta \mathcal{Y}).$$

Futher we will assume that all funcooids in consideration are small without mentioning it explicitly.

Definition 26. *Source* and *destination* of every funcooid $(A; B; \alpha; \beta)$ are defined as

$$\text{Src}(A; B; \alpha; \beta) = A \quad \text{and} \quad \text{Dst}(A; B; \alpha; \beta) = B.$$

I will denote $\text{FCD}(A; B)$ the set of funcooids from A to B .

I will denote FCD the set of all funcooids (for small sets).

Definition 27. $\langle (A; B; \alpha; \beta) \rangle \stackrel{\text{def}}{=} \alpha$ for a funcooid $(A; B; \alpha; \beta)$.

Definition 28. $(A; B; \alpha; \beta)^{-1} = (B; A; \beta; \alpha)$ for a funcooid $(A; B; \alpha; \beta)$.

Proposition 29. If f is a funcooid then f^{-1} is also a funcooid.

Proof. It follows from symmetry in the definition of funcooid. □

Obvious 30. $(f^{-1})^{-1} = f$ for a funcooid f .

Definition 31. The relation $[f] \in \mathcal{P}(\mathfrak{F}(\text{Src } f) \times \mathfrak{F}(\text{Dst } f))$ is defined (for every funcooid f and $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{Y} \in \mathfrak{F}(\text{Dst } f)$) by the formula $\mathcal{X}[f]\mathcal{Y} \stackrel{\text{def}}{=} \mathcal{Y} \not\prec \langle f \rangle \mathcal{X}$.

Obvious 32. $\mathcal{X}[f]\mathcal{Y} = \mathcal{Y} \not\prec \langle f \rangle \mathcal{X} \Leftrightarrow \mathcal{X} \not\prec \langle f^{-1} \rangle \mathcal{Y}$ for every funcooid f and $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{Y} \in \mathfrak{F}(\text{Dst } f)$.

Obvious 33. $[f^{-1}] = [f]^{-1}$ for a funcooid f .

Theorem 34. Let A, B are small sets.

1. For given value of $\langle f \rangle$ exists no more than one funcooid $f \in \text{FCD}(A; B)$.
2. For given value of $[f]$ exists no more than one funcooid $f \in \text{FCD}(A; B)$.

Proof. Let $f, g \in \text{FCD}(A; B)$.

Obviously $\langle f \rangle = \langle g \rangle \Rightarrow [f] = [g]$ and $\langle f^{-1} \rangle = \langle g^{-1} \rangle \Rightarrow [f] = [g]$. So enough to prove that $[f] = [g] \Rightarrow \langle f \rangle = \langle g \rangle$.

Provided that $[f] = [g]$ we have $\mathcal{Y} \not\star \langle f \rangle \mathcal{X} \Leftrightarrow \mathcal{X}[f]\mathcal{Y} \Leftrightarrow \mathcal{X}[g]\mathcal{Y} \Leftrightarrow \mathcal{Y} \not\star \langle g \rangle \mathcal{X}$ and consequently $\langle f \rangle \mathcal{X} = \langle g \rangle \mathcal{X}$ for every $\mathcal{X} \in \mathfrak{F}(A)$ and $\mathcal{Y} \in \mathfrak{F}(B)$ because a set of filter objects is separable [14], thus $\langle f \rangle = \langle g \rangle$. \square

Proposition 35. $\langle f \rangle 0^{\mathfrak{F}(\text{Src } f)} = 0^{\mathfrak{F}(\text{Dst } f)}$ for every functor f .

Proof. $\mathcal{Y} \not\star \langle f \rangle 0^{\mathfrak{F}(\text{Src } f)} \Leftrightarrow 0^{\mathfrak{F}(\text{Src } f)} \not\star \langle f^{-1} \rangle \mathcal{Y} \Leftrightarrow 0 \Leftrightarrow \mathcal{Y} \not\star 0^{\mathfrak{F}(\text{Dst } f)}$. Thus $\langle f \rangle 0^{\mathfrak{F}(\text{Src } f)} = 0^{\mathfrak{F}(\text{Dst } f)}$ by separability of filter objects. \square

Proposition 36. $\langle f \rangle (\mathcal{I} \cup \mathcal{J}) = \langle f \rangle \mathcal{I} \cup \langle f \rangle \mathcal{J}$ for every functor f and $\mathcal{I}, \mathcal{J} \in \mathfrak{F}(\text{Src } f)$.

Proof.

$$\begin{aligned} \star \langle f \rangle (\mathcal{I} \cup \mathcal{J}) &= \\ \{\mathcal{Y} \in \mathfrak{F} \mid \mathcal{Y} \not\star \langle f \rangle (\mathcal{I} \cup \mathcal{J})\} &= \\ \{\mathcal{Y} \in \mathfrak{F} \mid \mathcal{I} \cup \mathcal{J} \not\star \langle f^{-1} \rangle \mathcal{Y}\} &= \text{(by corollary 10 in [14])} \\ \{\mathcal{Y} \in \mathfrak{F} \mid \mathcal{I} \not\star \langle f^{-1} \rangle \mathcal{Y} \vee \mathcal{J} \not\star \langle f^{-1} \rangle \mathcal{Y}\} &= \\ \{\mathcal{Y} \in \mathfrak{F} \mid \mathcal{Y} \not\star \langle f \rangle \mathcal{I} \vee \mathcal{Y} \not\star \langle f \rangle \mathcal{J}\} &= \\ \{\mathcal{Y} \in \mathfrak{F} \mid \mathcal{Y} \not\star \langle f \rangle \mathcal{I} \cup \langle f \rangle \mathcal{J}\} &= \\ \star \langle \langle f \rangle \mathcal{I} \cup \langle f \rangle \mathcal{J} \rangle. & \end{aligned}$$

Thus $\langle f \rangle (\mathcal{I} \cup \mathcal{J}) = \langle f \rangle \mathcal{I} \cup \langle f \rangle \mathcal{J}$ because $\mathfrak{F}(\text{Dst } f)$ is separable. \square

3.2.1 Composition of functors

Definition 37. Functors f and g are *composable* when $\text{Dst } f = \text{Src } g$.

Definition 38. *Composition* of composable functors is defined by the formula

$$(B; C; \alpha_2; \beta_2) \circ (A; B; \alpha_1; \beta_1) = (A; C; \alpha_2 \circ \alpha_1; \beta_1 \circ \beta_2).$$

Proposition 39. If f, g are composable functors then $g \circ f$ is a functor.

Proof. Let $f = (A; B; \alpha_1; \beta_1)$, $g = (B; C; \alpha_2; \beta_2)$. For every $\mathcal{X} \in \mathfrak{F}(A)$, $\mathcal{Y} \in \mathfrak{F}(C)$ we have

$$\mathcal{Y} \not\star (\alpha_2 \circ \alpha_1) \mathcal{X} \Leftrightarrow \mathcal{Y} \not\star \alpha_2 \alpha_1 \mathcal{X} \Leftrightarrow \alpha_1 \mathcal{X} \not\star \beta_2 \mathcal{Y} \Leftrightarrow \mathcal{X} \not\star \beta_1 \beta_2 \mathcal{Y} \Leftrightarrow \mathcal{X} \not\star (\beta_1 \circ \beta_2) \mathcal{Y}.$$

So $(A; C; \alpha_2 \circ \alpha_1; \beta_1 \circ \beta_2)$ is a functor. \square

Obvious 40. $\langle g \circ f \rangle = \langle g \rangle \circ \langle f \rangle$ for every composable functors f and g .

Proposition 41. $(h \circ g) \circ f = h \circ (g \circ f)$ for every composable functors f, g, h .

Proof.

$$\langle \langle h \circ g \rangle \circ f \rangle = \langle h \circ g \rangle \circ \langle f \rangle = \langle \langle h \rangle \circ \langle g \rangle \rangle \circ \langle f \rangle = \langle h \rangle \circ \langle \langle g \rangle \circ \langle f \rangle \rangle = \langle h \rangle \circ \langle g \circ f \rangle = \langle h \circ (g \circ f) \rangle. \quad \square$$

Theorem 42. $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ for every composable functors f and g .

Proof. $\langle \langle g \circ f \rangle^{-1} \rangle = \langle f^{-1} \rangle \circ \langle g^{-1} \rangle = \langle f^{-1} \circ g^{-1} \rangle. \quad \square$

3.3 Functor as continuation

Let f is a functor.

Definition 43. $\langle f \rangle^*$ is the function $\mathcal{P}(\text{Src } f) \rightarrow \mathfrak{F}(\text{Dst } f)$ defined by the formula

$$\langle f \rangle^* X = \langle f \rangle \uparrow^{\text{Src } f} X.$$

Definition 44. $[f]^*$ is the relation between $\mathcal{P}(\text{Src } f)$ and $\mathcal{P}(\text{Dst } f)$ defined by the formula

$$X[f]^* Y = \uparrow^{\text{Src } f} X[f] \uparrow^{\text{Dst } f} Y.$$

Obvious 45.

1. $\langle f \rangle^* = \langle f \rangle \circ \uparrow^{\text{Src } f}$;
2. $[f]^* = (\uparrow^{\text{Dst } f})^{-1} \circ [f] \circ \uparrow^{\text{Src } f}$.

Theorem 46. For every functor f and $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$ and $\mathcal{Y} \in \mathfrak{F}(\text{Dst } f)$

1. $\langle f \rangle \mathcal{X} = \bigcap \langle \langle f \rangle^* \rangle \text{up } \mathcal{X}$;
2. $\mathcal{X}[f] \mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: X[f]^* Y$.

Proof. 2. $\mathcal{X}[f] \mathcal{Y} \Leftrightarrow \mathcal{Y} \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \forall Y \in \text{up } \mathcal{Y}: \uparrow^{\text{Dst } f} Y \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \forall Y \in \text{up } \mathcal{Y}: \mathcal{X}[f] \uparrow^{\text{Dst } f} Y$.

Analogously $\mathcal{X}[f] \mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}: \uparrow^{\text{Src } f} X[f] \mathcal{Y}$. Combining these two equivalences we get

$$\mathcal{X}[f] \mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: \uparrow^{\text{Src } f} X[f] \uparrow^{\text{Dst } f} Y \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: X[f]^* Y.$$

1. $\mathcal{Y} \cap \langle f \rangle \mathcal{X} \neq \emptyset \Leftrightarrow \mathcal{X}[f] \mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}: \uparrow^{\text{Src } f} X[f] \mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}: \mathcal{Y} \cap \langle f \rangle^* X \neq 0^{\mathfrak{F}(\text{Dst } f)}$.

Let's denote $W = \{\mathcal{Y} \cap \langle f \rangle^* X \mid X \in \text{up } \mathcal{X}\}$. We will prove that W is a generalized filter base. To prove this enough to show that $V = \{\langle f \rangle^* X \mid X \in \text{up } \mathcal{X}\}$ is a generalized filter base.

Let $\mathcal{P}, \mathcal{Q} \in V$. Then $\mathcal{P} = \langle f \rangle^* A$, $\mathcal{Q} = \langle f \rangle^* B$ where $A, B \in \text{up } \mathcal{X}$; $A \cap B \in \text{up } \mathcal{X}$ and $\mathcal{R} \subseteq \mathcal{P} \cap \mathcal{Q}$ for $\mathcal{R} = \langle f \rangle^*(A \cap B) \in V$. So V is a generalized filter base and thus W is a generalized filter base.

$0^{\mathfrak{F}(\text{Dst } f)} \notin W \Leftrightarrow \bigcap W \neq 0^{\mathfrak{F}(\text{Dst } f)}$ by the corollary 4 of the theorem 3. That is

$$\forall X \in \text{up } \mathcal{X}: \mathcal{Y} \cap \langle f \rangle^* X \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \mathcal{Y} \cap \bigcap \langle \langle f \rangle^* \rangle \text{up } \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)}.$$

Comparing with the above, $\mathcal{Y} \cap \langle f \rangle \mathcal{X} \neq \emptyset \Leftrightarrow \mathcal{Y} \cap \bigcap \langle \langle f \rangle^* \rangle \text{up } \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)}$. So $\langle f \rangle \mathcal{X} = \bigcap \langle \langle f \rangle^* \rangle \text{up } \mathcal{X}$ because the lattice of filter objects is separable. \square

Proposition 47. For every $f \in \text{FCD}(A; B)$ we have (for every $I, J \in \mathcal{P}A$)

$$\langle f \rangle^* \emptyset = 0^{\mathfrak{F}(B)}, \quad \langle f \rangle^*(I \cup J) = \langle f \rangle^* I \cup \langle f \rangle^* J$$

and

$$\begin{aligned} \neg(\emptyset[f]^* I), \quad I \cup J[f]^* K &\Leftrightarrow I[f]^* K \vee J[f]^* K \quad (\text{for every } I, J \in \mathcal{P}A, K \in \mathcal{P}B), \\ \neg(I[f]^* \emptyset), \quad K[f]^* I \cup J &\Leftrightarrow K[f]^* I \vee K[f]^* J \quad (\text{for every } I, J \in \mathcal{P}B, K \in \mathcal{P}A). \end{aligned}$$

Proof. $\langle f \rangle^* \emptyset = \langle f \rangle \uparrow^A \emptyset = \langle f \rangle 0^{\mathfrak{F}(A)} = 0^{\mathfrak{F}(B)}$; $\langle f \rangle^*(I \cup J) = \langle f \rangle \uparrow^A (I \cup J) = \langle f \rangle (\uparrow^A I \cup \uparrow^A J) = \langle f \rangle \uparrow^A I \cup \langle f \rangle \uparrow^A J = \langle f \rangle^* I \cup \langle f \rangle^* J$.

$I[f]^* \emptyset \Leftrightarrow 0^{\mathfrak{F}(B)} \not\neq \langle f \rangle \uparrow^A I \Leftrightarrow 0$; $I \cup J[f]^* K \Leftrightarrow \uparrow^A (I \cup J)[f] \uparrow^B K \Leftrightarrow \uparrow^B K \neq \langle f \rangle^*(I \cup J) \Leftrightarrow \uparrow^B K \neq \langle f \rangle^* I \cup \langle f \rangle^* J \Leftrightarrow \uparrow^B K \neq \langle f \rangle^* I \vee \uparrow^B K \neq \langle f \rangle^* J \Leftrightarrow I[f]^* K \vee J[f]^* K$.

The rest follows from symmetry. \square

Theorem 48. Fix two small sets A and B . Let $L_F = \lambda f \in \text{FCD}(A; B): \langle f \rangle^*$ and $L_R = \lambda f \in \text{FCD}(A; B): [f]^*$.

1. L_F is a bijection from the set $\text{FCD}(A; B)$ to the set of functions $\alpha \in \mathfrak{F}(B)^{\mathcal{P}A}$ that obey the conditions (for every $I, J \in \mathcal{P}A$)

$$\alpha \emptyset = 0^{\mathfrak{F}(B)}, \quad \alpha(I \cup J) = \alpha I \cup \alpha J. \quad (1)$$

For such α it holds (for every $\mathcal{X} \in \mathfrak{F}(A)$)

$$\langle L_F^{-1} \alpha \rangle \mathcal{X} = \bigcap \langle \alpha \rangle \text{up } \mathcal{X} \quad (2)$$

2. L_R is a bijection from the set $\text{FCD}(A; B)$ to the set of binary relations $\delta \in \mathcal{P}(\mathcal{P}A \times \mathcal{P}B)$ that obey the conditions

$$\begin{aligned} \neg(\emptyset \delta I), \quad I \cup J \delta K &\Leftrightarrow I \delta K \vee J \delta K \quad (\text{for every } I, J \in \mathcal{P}A, K \in \mathcal{P}B), \\ \neg(I \delta \emptyset), \quad K \delta I \cup J &\Leftrightarrow K \delta I \vee K \delta J \quad (\text{for every } I, J \in \mathcal{P}B, K \in \mathcal{P}A). \end{aligned} \quad (3)$$

For such δ it holds (for every $\mathcal{X} \in \mathfrak{F}(A)$, $\mathcal{Y} \in \mathfrak{F}(B)$)

$$\mathcal{X}[L_R^{-1}\delta]\mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: X \delta Y. \quad (4)$$

Proof. Injectivity of L_F and L_R , formulas (2) (for $\alpha \in \text{im } L_F$) and (4) (for $\delta \in \text{im } L_R$), formulas (1) and (3) follow from two previous theorems. The only thing remained to prove is that for every α and δ that obey the above conditions exists a corresponding funcoïd f .

2. Let define $\alpha \in \mathfrak{F}(B)^{\mathcal{P}A}$ by the formula $\partial(\alpha X) = \{Y \in \mathcal{P}B \mid X \delta Y\}$ for every $X \in \mathcal{P}A$. (It is obvious that $\{Y \in \mathcal{P}B \mid X \delta Y\}$ is a free star.) Analogously it can be defined $\beta \in \mathfrak{F}(A)^{\mathcal{P}B}$ by the formula $\partial(\beta X) = \{X \in \mathcal{P}A \mid X \delta Y\}$. Let's continue α and β to $\alpha' \in \mathfrak{F}(B)^{\mathfrak{F}(A)}$ and $\beta' \in \mathfrak{F}(A)^{\mathfrak{F}(B)}$ by the formulas

$$\alpha' \mathcal{X} = \bigcap \langle \alpha \rangle \text{up } \mathcal{X} \quad \text{and} \quad \beta' \mathcal{X} = \bigcap \langle \beta \rangle \text{up } \mathcal{X}$$

and δ to $\delta' \in \mathcal{P}(\mathfrak{F}(A) \times \mathfrak{F}(B))$ by the formula

$$\mathcal{X} \delta' \mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: X \delta Y.$$

$\mathcal{Y} \cap \alpha' \mathcal{X} \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \mathcal{Y} \cap \bigcap \langle \alpha \rangle \text{up } \mathcal{X} \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \bigcap \langle \mathcal{Y} \cap \rangle \langle \alpha \rangle \text{up } \mathcal{X} \neq 0^{\mathfrak{F}(B)}$. Let's prove that

$$W = \langle \mathcal{Y} \cap \rangle \langle \alpha \rangle \text{up } \mathcal{X}$$

is a generalized filter base: To prove it is enough to show that $\langle \alpha \rangle \text{up } \mathcal{X}$ is a generalized filter base. If $\mathcal{A}, \mathcal{B} \in \langle \alpha \rangle \text{up } \mathcal{X}$ then exist $X_1, X_2 \in \text{up } \mathcal{X}$ such that $\mathcal{A} = \alpha X_1$ and $\mathcal{B} = \alpha X_2$.

Then $\alpha(X_1 \cap X_2) \in \langle \alpha \rangle \text{up } \mathcal{X}$. So $\langle \alpha \rangle \text{up } \mathcal{X}$ is a generalized filter base and thus W is a generalized filter base.

Accordingly the corollary 4 of the theorem 3, $\bigcap \langle \mathcal{Y} \cap \rangle \langle \alpha \rangle \text{up } \mathcal{X} \neq 0^{\mathfrak{F}(B)}$ is equivalent to

$$\forall X \in \text{up } \mathcal{X}: \mathcal{Y} \cap \alpha X \neq 0^{\mathfrak{F}(B)},$$

what is equivalent to $\forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: \uparrow^{BY} \cap \alpha X \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: Y \in \partial(\alpha X) \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: X \delta Y$. Combining the equivalencies we get $\mathcal{Y} \cap \alpha' \mathcal{X} \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \mathcal{X} \delta' \mathcal{Y}$. Analogously $\mathcal{X} \cap \beta' \mathcal{Y} \neq 0^{\mathfrak{F}(A)} \Leftrightarrow X \delta' Y$. So $\mathcal{Y} \cap \alpha' \mathcal{X} \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \mathcal{X} \cap \beta' \mathcal{Y} \neq 0^{\mathfrak{F}(A)}$, that is $(A; B; \alpha'; \beta')$ is a funcoïd. From the formula $\mathcal{Y} \cap \alpha' \mathcal{X} \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \mathcal{X} \delta' \mathcal{Y}$ it follows that

$$\mathcal{X}[(A; B; \alpha'; \beta')]^* \mathcal{Y} \Leftrightarrow \uparrow^{BY} \cap \alpha' \uparrow^A X \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \uparrow^B X \delta' \uparrow^A Y \Leftrightarrow X \delta Y.$$

1. Let define the relation $\delta \in \mathcal{P}(\mathcal{P}A \times \mathcal{P}B)$ by the formula $X \delta Y \Leftrightarrow \uparrow^{BY} \cap \alpha X \neq 0^{\mathfrak{F}(B)}$.

That $\neg(\emptyset \delta I)$ and $\neg(I \delta \emptyset)$ is obvious. We have $I \cup J \delta K \Leftrightarrow \uparrow^B K \cap \alpha(I \cup J) \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \uparrow^B K \cap (\alpha I \cup \alpha J) \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \uparrow^B K \cap \alpha I \neq 0^{\mathfrak{F}(B)} \vee \uparrow^B K \cap \alpha J \neq 0^{\mathfrak{F}(B)} \Leftrightarrow I \delta K \vee J \delta K$ and

$K \delta I \cup J \Leftrightarrow \uparrow^B(I \cup J) \cap \alpha K \neq 0^{\mathfrak{F}(B)} \Leftrightarrow (\uparrow^B I \cup \uparrow^B J) \cap \alpha K \neq 0^{\mathfrak{F}(B)} \Leftrightarrow (\uparrow^B I \cap \alpha K) \cup (\uparrow^B J \cap \alpha K) \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \uparrow^B I \cap \alpha K \neq 0^{\mathfrak{F}(B)} \vee \uparrow^B J \cap \alpha K \neq 0^{\mathfrak{F}(B)} \Leftrightarrow K \delta I \vee K \delta J$.

That is the formulas (3) are true.

Accordingly the above there exist a funcoïd f such that

$$\mathcal{X}[f]\mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: X \delta Y.$$

$\forall X \in \mathcal{P}A, Y \in \mathcal{P}B: (\uparrow^{BY} \cap \langle f \rangle \uparrow^A X \neq \emptyset \Leftrightarrow \uparrow^A X[f] \uparrow^{BY} \Leftrightarrow X \delta Y \Leftrightarrow \uparrow^{BY} \cap \alpha X \neq 0^{\mathfrak{F}(B)})$, consequently $\forall X \in \mathcal{P}A: \alpha X = \langle f \rangle \uparrow^A X = \langle f \rangle^* X$. \square

Note that by the last theorem to every proximity δ corresponds a unique funcoïd. So funcoïds are a generalization of (quasi-)proximity structures.

Reverse funcoïds can be considered as a generalization of conjugate quasi-proximity.

Definition 49. Any small (multivalued) function $F: A \rightarrow B$ corresponds to a funcoïd $\uparrow^{\text{FCD}(A; B)} f \in \text{FCD}(A; B)$, where by definition $\langle \uparrow^{\text{FCD}(A; B)} f \rangle \mathcal{X} = \bigcap \langle \uparrow^B \rangle \langle \langle F \rangle \rangle \text{up } \mathcal{X}$ for every $\mathcal{X} \in \mathfrak{F}(A)$.

Using the last theorem it is easy to show that this definition is monovalued and does not contradict to former stuff. (Take $\alpha = \uparrow^B \circ \langle F \rangle$.)

Definition 50. Funcoïds corresponding to a binary relation (= multivalued function) are called *discrete funcoïds*.

We may equate discrete functors with corresponding binary relations by the method of appendix B in [14]. This is useful for describing relationships of functors and binary relations, such as for the formulas of continuous functions and continuous functors (see below).

Theorem 51. If S is a generalized filter base on $\text{Src } f$ then $\langle f \rangle \cap S = \bigcap \langle \langle f \rangle \rangle S$ for every functor f .

Proof. $\langle f \rangle \cap S \subseteq \langle f \rangle X$ for every $X \in S$ and thus $\langle f \rangle \cap S \subseteq \bigcap \langle \langle f \rangle \rangle S$.

By properties of generalized filter bases:

$$\langle f \rangle \cap S = \bigcap \langle \langle f \rangle \rangle \text{up } \bigcap S = \bigcap \langle \langle f \rangle \rangle \{X \mid \exists \mathcal{P} \in S: X \in \text{up } \mathcal{P}\} = \bigcap \{\langle f \rangle^* X \mid \exists \mathcal{P} \in S: X \in \text{up } \mathcal{P}\} \supseteq \bigcap \{\langle f \rangle \mathcal{P} \mid \mathcal{P} \in S\} = \bigcap \langle \langle f \rangle \rangle S. \quad \square$$

3.4 Lattices of functors

Definition 52. $f \subseteq g \stackrel{\text{def}}{=} [f] \subseteq [g]$ for $f, g \in \text{FCD}$.

Thus every $\text{FCD}(A; B)$ is a poset. (Taken into account that $[f] \neq [g]$ if $f \neq g$.)

Definition 53. I will call a *shifted filtrator of functors* the shifted filtrator

$$(\text{FCD}(A; B); \mathcal{P}(\mathcal{P}A \times \mathcal{P}B); \uparrow^{\text{FCD}(A; B)})$$

for some small sets A, B .

$$\text{up } f \stackrel{\text{def}}{=} \text{up}^{(\text{FCD}(A; B); \mathcal{P}(\mathcal{P}A \times \mathcal{P}B); \uparrow^{\text{FCD}(A; B)})} f \text{ for every functor } f \in \text{FCD}(A; B).$$

Lemma 54. $\langle f \rangle^* X = \bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid F \in \text{up } f\}$ for every functor f and set $X \in \mathcal{P}(\text{Src } f)$.

Proof. Obviously $\langle f \rangle^* X \subseteq \bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid F \in \text{up } f\}$.

Let $B \in \text{up } \langle f \rangle^* X$. Let $F_B = X \times B \cup ((\text{Src } f) \setminus X) \times (\text{Dst } f)$.

$$\langle F_B \rangle X = B.$$

We have $\emptyset \neq P \subseteq X \Rightarrow \langle F_B \rangle P = B \supseteq \langle f \rangle^* P$ and $P \not\subseteq X \Rightarrow \langle F_B \rangle P = \text{Dst } f \supseteq \langle f \rangle^* P$. Thus $\langle F_B \rangle P \supseteq \langle f \rangle^* P$ for every set $P \in \mathcal{P}(\text{Src } f)$ and so $\uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \supseteq f$ that is $F_B \in \text{up } f$.

Thus $\forall B \in \text{up } \langle f \rangle^* X: B \in \text{up } \bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid F \in \text{up } f\}$ because $B \in \text{up } \uparrow^{\text{Dst } f} \langle F_B \rangle X$.

$$\text{So } \bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid F \in \text{up } f\} \subseteq \langle f \rangle^* X. \quad \square$$

Theorem 55. $\langle f \rangle \mathcal{X} = \bigcap \{\uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \mathcal{X} \mid F \in \text{up } f\}$ for every functor f and $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$.

Proof. $\bigcap \{\uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \mathcal{X} \mid F \in \text{up } f\} = \bigcap \{\bigcap \langle \uparrow^{\text{Dst } f} \rangle \langle \langle F \rangle \rangle \text{up } \mathcal{X} \mid F \in \text{up } f\} = \bigcap \{\bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid X \in \text{up } \mathcal{X}\} \mid F \in \text{up } f\} = \bigcap \{\bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid F \in \text{up } f\} \mid X \in \text{up } \mathcal{X}\} = \bigcap \{\uparrow^{\text{Dst } f} \langle f \rangle^* X \mid X \in \text{up } \mathcal{X}\} = \langle f \rangle \mathcal{X}$ (the lemma used). \square

Conjecture 56. Every filtrator of functors is:

1. with separable core;
2. with co-separable core.

Theorem 57. $\text{FCD}(A; B)$ is a complete lattice (for every small sets A and B). For every $R \in \mathcal{P}\text{FCD}(A; B)$ and $X \in \mathcal{P}A, Y \in \mathcal{P}B$

1. $X[\bigcup R]^* Y \Leftrightarrow \exists f \in R: X[f]^* Y$;
2. $\langle \bigcup R \rangle^* X = \bigcup \{\langle f \rangle^* X \mid f \in R\}$.

Proof. Accordingly [13] to prove that it is a complete lattice enough to prove existence of all joins.

2. $\alpha X \stackrel{\text{def}}{=} \bigcup \{\langle f \rangle^* X \mid f \in R\}$. We have $\alpha \emptyset = \emptyset$;

$$\begin{aligned} \alpha(I \cup J) &= \bigcup \{\langle f \rangle^*(I \cup J) \mid f \in R\} \\ &= \bigcup \{\langle f \rangle^* I \cup \langle f \rangle^* J \mid f \in R\} \\ &= \bigcup \{\langle f \rangle^* I \mid f \in R\} \cup \bigcup \{\langle f \rangle^* J \mid f \in R\} \\ &= \alpha I \cup \alpha J. \end{aligned}$$

So $\langle h \rangle \circ \uparrow^A = \alpha$ for some funcoid h . Obviously

$$\forall f \in R: h \supseteq f. \quad (5)$$

And h is the least funcoid for which holds the condition (5). So $h = \bigcup R$.

1. $X[\bigcup R]^*Y \Leftrightarrow \uparrow^{\text{Dst}} fY \cap \langle \bigcup R \rangle^*X \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \uparrow^{\text{Dst}} fY \cap \bigcup \{ \langle f \rangle^*X \mid f \in R \} \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \exists f \in R: \uparrow^{\text{Dst}} fY \cap \langle f \rangle^*X \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \exists f \in R: X[f]^*Y$ (used the theorem 40 in [14]). \square

In the next theorem, compared to the previous one, the class of infinite unions is replaced with lesser class of finite unions and simultaneously class of sets is changed to more wide class of filter objects.

Theorem 58. For every $f, g \in \text{FCD}(A; B)$ and $\mathcal{X} \in \mathfrak{F}(A)$ (for every small sets A, B)

1. $\langle f \cup g \rangle \mathcal{X} = \langle f \rangle \mathcal{X} \cup \langle g \rangle \mathcal{X}$;
2. $[f \cup g] = [f] \cup [g]$.

Proof.

1. Let $\alpha \mathcal{X} \stackrel{\text{def}}{=} \langle f \rangle \mathcal{X} \cup \langle g \rangle \mathcal{X}$; $\beta \mathcal{Y} \stackrel{\text{def}}{=} \langle f^{-1} \rangle \mathcal{Y} \cup \langle g^{-1} \rangle \mathcal{Y}$ for every $\mathcal{X} \in \mathfrak{F}(A)$, $\mathcal{Y} \in \mathfrak{F}(B)$. Then

$$\begin{aligned} \mathcal{Y} \cap \alpha \mathcal{X} \neq \emptyset &\Leftrightarrow \mathcal{Y} \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(B)} \vee \mathcal{Y} \cap \langle g \rangle \mathcal{X} \neq 0^{\mathfrak{F}(B)} \\ &\Leftrightarrow \mathcal{X} \cap \langle f^{-1} \rangle \mathcal{Y} \neq 0^{\mathfrak{F}(A)} \vee \mathcal{X} \cap \langle g^{-1} \rangle \mathcal{Y} \neq 0^{\mathfrak{F}(A)} \\ &\Leftrightarrow \mathcal{X} \cap \beta \mathcal{Y} \neq 0^{\mathfrak{F}(A)}. \end{aligned}$$

So $h = (A; B; \alpha; \beta)$ is a funcoid. Obviously $h \supseteq f$ and $h \supseteq g$. If $p \supseteq f$ and $p \supseteq g$ for some funcoid p then $\langle p \rangle \mathcal{X} \supseteq \langle f \rangle \mathcal{X} \cup \langle g \rangle \mathcal{X} = \langle h \rangle \mathcal{X}$ that is $p \supseteq h$. So $f \cup g = h$.

2. $\mathcal{X}[f \cup g] \mathcal{Y} \Leftrightarrow \mathcal{Y} \cap \langle f \cup g \rangle \mathcal{X} \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \mathcal{Y} \cap (\langle f \rangle \mathcal{X} \cup \langle g \rangle \mathcal{X}) \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \mathcal{Y} \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(B)} \vee \mathcal{Y} \cap \langle g \rangle \mathcal{X} \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \mathcal{X}[f] \mathcal{Y} \vee \mathcal{X}[g] \mathcal{Y}$ for every $\mathcal{X} \in \mathfrak{F}(A)$, $\mathcal{Y} \in \mathfrak{F}(B)$. \square

3.5 More on composition of funcoids

Proposition 59. $[g \circ f] = [g] \circ \langle f \rangle = \langle g^{-1} \rangle^{-1} \circ [f]$ for every composable funcoids f and g .

Proof. $\mathcal{X}[g \circ f] \mathcal{Y} \Leftrightarrow \mathcal{Y} \cap \langle g \circ f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } g)} \Leftrightarrow \mathcal{Y} \cap \langle g \rangle \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } g)} \Leftrightarrow \langle f \rangle \mathcal{X}[g] \mathcal{Y} \Leftrightarrow \mathcal{X}([g] \circ \langle f \rangle) \mathcal{Y}$ for every $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{Y} \in \mathfrak{F}(\text{Dst } g)$. $[g \circ f] = [(f^{-1} \circ g^{-1})^{-1}] = [f^{-1} \circ g^{-1}]^{-1} = ([f^{-1}] \circ \langle g^{-1} \rangle)^{-1} = \langle g^{-1} \rangle^{-1} \circ [f]$. \square

The following theorem is a variant for funcoids of the statement (which defines compositions of relations) that $x(g \circ f)z \Leftrightarrow \exists y(xfy \wedge ygz)$ for every x and z and every binary relations f and g .

Theorem 60. For every small sets A, B, C and $f \in \text{FCD}(A; B)$, $g \in \text{FCD}(B; C)$ and $\mathcal{X} \in \mathfrak{F}(A)$, $\mathcal{Z} \in \mathfrak{F}(C)$.

$$\mathcal{X}[g \circ f] \mathcal{Z} \Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: (\mathcal{X}[f]y \wedge y[g] \mathcal{Z}).$$

Proof.

$$\begin{aligned} \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: (\mathcal{X}[f]y \wedge y[g] \mathcal{Z}) &\Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: (\mathcal{Z} \cap \langle g \rangle y \neq 0^{\mathfrak{F}(C)} \wedge y \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(B)}) \\ &\Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: (\mathcal{Z} \cap \langle g \rangle y \neq 0^{\mathfrak{F}(C)} \wedge y \subseteq \langle f \rangle \mathcal{X}) \\ &\Rightarrow \mathcal{Z} \cap \langle g \rangle \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(C)} \\ &\Leftrightarrow \mathcal{X}[g \circ f] \mathcal{Z}. \end{aligned}$$

Reversely, if $\mathcal{X}[g \circ f] \mathcal{Z}$ then $\langle f \rangle \mathcal{X}[g] \mathcal{Z}$, consequently exists $y \in \text{atoms } \langle f \rangle \mathcal{X}$ such that $y[g] \mathcal{Z}$; we have $\mathcal{X}[f]y$. \square

Theorem 61. For every small sets A, B, C

1. $f \circ (g \cup h) = f \circ g \cup f \circ h$ for $g, h \in \text{FCD}(A; B)$ and $f \in \text{FCD}(B; C)$;

2. $(g \cup h) \circ f = g \circ f \cup h \circ f$ for $g, h \in \text{FCD}(B; C)$ and $f \in \text{FCD}(A; B)$.

Proof. I will prove only the first equality because the other is analogous.

For every $\mathcal{X}, \mathcal{Z} \in \mathfrak{F}$

$$\begin{aligned}
\mathcal{X}[f \circ (g \cup h)]\mathcal{Z} &\Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: (\mathcal{X}[g \cup h]y \wedge y[f]\mathcal{Z}) \\
&\Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: ((\mathcal{X}[g]y \vee \mathcal{X}[h]y) \wedge y[f]\mathcal{Z}) \\
&\Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: (\mathcal{X}[g]y \wedge y[f]\mathcal{Z} \vee \mathcal{X}[h]y \wedge y[f]\mathcal{Z}) \\
&\Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: (\mathcal{X}[g]y \wedge y[f]\mathcal{Z}) \vee \exists y \in \text{atoms } 1^{\mathfrak{F}(B)}: (\mathcal{X}[h]y \wedge y[f]\mathcal{Z}) \\
&\Leftrightarrow \mathcal{X}[f \circ g]\mathcal{Z} \vee \mathcal{X}[f \circ h]\mathcal{Z} \\
&\Leftrightarrow \mathcal{X}[f \circ g \cup f \circ h]\mathcal{Z}.
\end{aligned}$$

□

3.6 Domain and range of a funcooid

Definition 62. Let A is a small set. The *identity funcooid* $I^{\text{FCD}(A)} = ((=)|_{\mathfrak{F}(A)}; (=)|_{\mathfrak{F}(A)})$.

Obvious 63. The identity funcooid is a funcooid.

Definition 64. Let A is a small set, $\mathcal{A} \in \mathfrak{F}(A)$. The *restricted identity funcooid*

$$I_{\mathcal{A}}^{\text{FCD}} = (\mathcal{A} \cap; \mathcal{A} \cap).$$

Proposition 65. The restricted identity funcooid is a funcooid.

Proof. We need to prove that $(\mathcal{A} \cap \mathcal{X}) \cap \mathcal{Y} \neq \emptyset \Leftrightarrow (\mathcal{A} \cap \mathcal{Y}) \cap \mathcal{X} \neq \emptyset$ what is obvious. □

Obvious 66.

1. $(I^{\text{FCD}(A)})^{-1} = I^{\text{FCD}(A)}$;
2. $(I_{\mathcal{A}}^{\text{FCD}})^{-1} = I_{\mathcal{A}}^{\text{FCD}}$.

Obvious 67. For every $\mathcal{X}, \mathcal{Y} \in \mathfrak{F}(A)$

1. $\mathcal{X}[I^{\text{FCD}(A)}]\mathcal{Y} \Leftrightarrow \mathcal{X} \cap \mathcal{Y} \neq \emptyset$.
2. $\mathcal{X}[I_{\mathcal{A}}^{\text{FCD}}]\mathcal{Y} \Leftrightarrow \mathcal{A} \cap \mathcal{X} \cap \mathcal{Y} \neq \emptyset$.

Definition 68. I will define *restricting* of a funcooid f to a filter object $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$ by the formula

$$f|_{\mathcal{A}} \stackrel{\text{def}}{=} f \circ I_{\mathcal{A}}^{\text{FCD}}.$$

Definition 69. *Image* of a funcooid f will be defined by the formula $\text{im } f = \langle f \rangle 1^{\mathfrak{F}(\text{Src } f)}$.

Domain of a funcooid f is defined by the formula $\text{dom } f = \text{im } f^{-1}$.

Proposition 70. $\langle f \rangle \mathcal{X} = \langle f \rangle (\mathcal{X} \cap \text{dom } f)$ for every $f \in \text{FCD}$, $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$.

Proof. For every $\mathcal{Y} \in \mathfrak{F}(\text{Dst } f)$ we have $\mathcal{Y} \cap \langle f \rangle (\mathcal{X} \cap \text{dom } f) \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \mathcal{X} \cap \text{dom } f \cap \langle f^{-1} \rangle \mathcal{Y} \neq 0^{\mathfrak{F}(\text{Src } f)} \Leftrightarrow \mathcal{X} \cap \text{im } f^{-1} \cap \langle f^{-1} \rangle \mathcal{Y} \neq 0^{\mathfrak{F}(\text{Src } f)} \Leftrightarrow \mathcal{X} \cap \langle f^{-1} \rangle \mathcal{Y} \neq 0^{\mathfrak{F}(\text{Src } f)} \Leftrightarrow \mathcal{Y} \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)}$. Thus $\langle f \rangle \mathcal{X} = \langle f \rangle (\mathcal{X} \cap \text{dom } f)$ because the lattice of filter objects is separable. □

Proposition 71. $\mathcal{X} \cap \text{dom } f \neq 0^{\mathfrak{F}(\text{Src } f)} \Leftrightarrow \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)}$ for every $f \in \text{FCD}$, $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$.

Proof. $\mathcal{X} \cap \text{dom } f \neq 0^{\mathfrak{F}(\text{Src } f)} \Leftrightarrow \mathcal{X} \cap \langle f^{-1} \rangle 1^{\mathfrak{F}(\text{Dst } f)} \neq 0^{\mathfrak{F}(\text{Src } f)} \Leftrightarrow 1^{\mathfrak{F}(\text{Dst } f)} \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)}$. □

Corollary 72. $\text{dom } f = \bigcup \{a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)} \mid \langle f \rangle a \neq 0^{\mathfrak{F}(\text{Dst } f)}\}$.

Proof. This follows from the fact that $\mathfrak{F}(\text{Src } f)$ is an atomistic lattice. \square

Proposition 73. $\text{dom } f|_{\mathcal{A}} = \mathcal{A} \cap \text{dom } f$ for every funcoid f and $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$.

Proof. $\text{dom } f|_{\mathcal{A}} = \text{im}(I_{\mathcal{A}}^{\text{FCD}} \circ f^{-1}) = \langle I_{\mathcal{A}}^{\text{FCD}} \rangle \langle f^{-1} \rangle 1^{(\text{Dst } f)} = \mathcal{A} \cap \langle f^{-1} \rangle 1^{(\text{Dst } f)} = \mathcal{A} \cap \text{dom } f$. \square

Theorem 74. $\text{im } f = \bigcap \langle \uparrow^{\text{Dst } f} \rangle \langle \text{im} \rangle \text{up } f$ and $\text{dom } f = \bigcap \langle \uparrow^{\text{Src } f} \rangle \langle \text{dom} \rangle \text{up } f$ for every funcoid f .

Proof. $\text{im } f = \langle f \rangle 1^{\mathfrak{F}(\text{Src } f)} = \bigcap \{ \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \rangle 1^{\mathfrak{F}(\text{Src } f)} \mid F \in \text{up } f \} = \bigcap \{ \uparrow^{\text{Dst } f} \text{im } F \mid F \in \text{up } f \} = \bigcap \langle \uparrow^{\text{Dst } f} \rangle \langle \text{im} \rangle \text{up } f$ (used the theorem 55).

The second formula follows from symmetry. \square

Proposition 75. For every composable funcoids f, g :

1. If $\text{im } f \supseteq \text{dom } g$ then $\text{im}(g \circ f) = \text{im } g$.
2. If $\text{im } f \subseteq \text{dom } g$ then $\text{dom}(g \circ f) = \text{dom } f$.

Proof.

1. $\text{im}(g \circ f) = \langle g \circ f \rangle 1^{\mathfrak{F}(\text{Src } f)} = \langle g \rangle \langle f \rangle 1^{\mathfrak{F}(\text{Src } f)} = \langle g \rangle \text{im } f = \langle g \rangle (\text{im } f \cap \text{dom } g) = \langle g \rangle \text{dom } g = \langle g \rangle 1^{\mathfrak{F}(\text{Src } g)} = \text{im } g$.
2. $\text{dom}(g \circ f) = \text{im}(f^{-1} \circ g^{-1})$ what by proved above is equal to $\text{im } f^{-1}$ that is $\text{dom } f$. \square

3.7 Categories of funcoids

I will define two categories, the *category of funcoids* and the *category of funcoid triples*.

The *category of funcoids* is defined as follows:

- Objects are small sets.
- The set of morphisms from a set A to a set B is $\text{FCD}(A; B)$.
- The composition is the composition of funcoids.
- Identity morphism for a set is the identity funcoid for that set.

To show it is really a category is trivial.

The *category of funcoid triples* is defined as follows:

- Objects are filter objects on small sets.
- The morphisms from a f.o. \mathcal{A} to a f.o. \mathcal{B} are triples $(f; \mathcal{A}; \mathcal{B})$ where $f \in \text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))$ and $\text{dom } f \subseteq \mathcal{A} \wedge \text{im } f \subseteq \mathcal{B}$.
- The composition is defined by the formula $(g; \mathcal{B}; \mathcal{C}) \circ (f; \mathcal{A}; \mathcal{B}) = (g \circ f; \mathcal{A}; \mathcal{C})$.
- Identity morphism for an f.o. \mathcal{A} is $I_{\mathcal{A}}^{\text{FCD}}$.

To prove that it is really a category is trivial.

3.8 Specifying funcoids by functions or relations on atomic filter objects

Theorem 76. For every funcoid f and $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{Y} \in \mathfrak{F}(\text{Dst } f)$

1. $\langle f \rangle \mathcal{X} = \bigcup \langle \langle f \rangle \rangle \text{atoms } \mathcal{X}$;
2. $\mathcal{X}[f]\mathcal{Y} \Leftrightarrow \exists x \in \text{atoms } \mathcal{X}, y \in \text{atoms } \mathcal{Y}: x[f]y$.

Proof. 1.

$$\begin{aligned} \mathcal{Y} \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)} &\Leftrightarrow \mathcal{X} \cap \langle f^{-1} \rangle \mathcal{Y} \neq 0^{\mathfrak{F}(\text{Src } f)} \\ &\Leftrightarrow \exists x \in \text{atoms } \mathcal{X}: x \cap \langle f^{-1} \rangle \mathcal{Y} \neq 0^{\mathfrak{F}(\text{Src } f)} \\ &\Leftrightarrow \exists x \in \text{atoms } \mathcal{X}: \mathcal{Y} \cap \langle f \rangle x \neq 0^{\mathfrak{F}(\text{Dst } f)}. \end{aligned}$$

$\partial\langle f\rangle\mathcal{X} = \bigcup \langle \partial \rangle \langle \langle f \rangle \rangle \text{atoms } \mathcal{X} = \partial \bigcup \langle \langle f \rangle \rangle \text{atoms } \mathcal{X}$.

2. If $\mathcal{X}[f]\mathcal{Y}$, then $\mathcal{Y} \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{S}(\text{Dst } f)}$, consequently exists $y \in \text{atoms } \mathcal{Y}$ such that $y \cap \langle f \rangle \mathcal{X} \neq 0^{\mathfrak{S}(\text{Dst } f)}$, $\mathcal{X}[f]y$. Repeating this second time we get that there exist $x \in \text{atoms } \mathcal{X}$ such that $x[f]y$. From this follows

$$\exists x \in \text{atoms } \mathcal{X}, y \in \text{atoms } \mathcal{Y}: x[f]y.$$

The reverse is obvious. □

Theorem 77. Let A and B be small sets.

1. A function $\alpha \in \mathfrak{F}(B)^{\text{atoms } 1^{\mathfrak{S}(A)}}$ such that (for every $a \in \text{atoms } 1^{\mathfrak{S}(A)}$)

$$\alpha a \subseteq \bigcap \langle \bigcup \circ \langle \alpha \rangle \circ \text{atoms} \circ \uparrow^A \rangle \text{up } a \quad (6)$$

can be continued to the function $\langle f \rangle$ for a unique $f \in \text{FCD}(A; B)$;

$$\langle f \rangle \mathcal{X} = \bigcup \langle \alpha \rangle \text{atoms } \mathcal{X} \quad (7)$$

for every $\mathcal{X} \in \mathfrak{F}(A)$.

2. A relation $\delta \in \mathcal{P}(\text{atoms } A \times \text{atoms } B)$ such that (for every $a \in \text{atoms } A, b \in \text{atoms } B$)

$$\forall X \in \text{up } a, Y \in \text{up } b \exists x \in \text{atoms } \uparrow^A X, y \in \text{atoms } \uparrow^B Y: x \delta y \Rightarrow a \delta b \quad (8)$$

can be continued to the relation $[f]$ for a unique $f \in \text{FCD}(A; B)$;

$$\mathcal{X}[f]\mathcal{Y} \Leftrightarrow \exists x \in \text{atoms } \mathcal{X}, y \in \text{atoms } \mathcal{Y}: x \delta y \quad (9)$$

for every $\mathcal{X} \in \mathfrak{F}(A), \mathcal{Y} \in \mathfrak{F}(B)$.

Proof. Existence of no more than one such funcoids and formulas (7) and (9) follow from the previous theorem.

1. Consider the function $\alpha' \in \mathfrak{F}(B)^{\mathcal{P}A}$ defined by the formula (for every $X \in \mathcal{P}A$)

$$\alpha' X = \bigcup \langle \alpha \rangle \text{atoms } \uparrow^A X.$$

Obviously $\alpha' \emptyset = 0^{\mathfrak{S}(B)}$. For every $I, J \in \mathcal{P}A$

$$\begin{aligned} \alpha'(I \cup J) &= \bigcup \langle \alpha' \rangle \text{atoms } \uparrow^A (I \cup J) \\ &= \bigcup \langle \alpha' \rangle (\text{atoms } \uparrow^A I \cup \text{atoms } \uparrow^A J) \\ &= \bigcup (\langle \alpha' \rangle \text{atoms } \uparrow^A I \cup \langle \alpha' \rangle \text{atoms } \uparrow^A J) \\ &= \bigcup \langle \alpha' \rangle \text{atoms } \uparrow^A I \cup \bigcup \langle \alpha' \rangle \text{atoms } \uparrow^A J. \\ &= \alpha' I \cup \alpha' J. \end{aligned}$$

Let continue α' till a funcoid f (by the theorem 48): $\langle f \rangle \mathcal{X} = \bigcap \langle \alpha' \rangle \text{up } \mathcal{X}$.

Let's prove the reverse of (6):

$$\begin{aligned} \bigcap \langle \bigcup \circ \langle \alpha \rangle \circ \text{atoms} \circ \uparrow^A \rangle \text{up } a &= \bigcap \langle \bigcup \circ \langle \alpha \rangle \rangle \langle \text{atoms} \rangle \langle \uparrow^A \rangle \text{up } a \\ &\subseteq \bigcap \langle \bigcup \circ \langle \alpha \rangle \rangle \{ \{a\} \} \\ &= \bigcap \{ \langle \bigcup \circ \langle \alpha \rangle \rangle \{a\} \} \\ &= \bigcap \{ \bigcup \langle \alpha \rangle \{a\} \} \\ &= \bigcap \{ \bigcup \{ \alpha a \} \} = \bigcap \{ \alpha a \} = \alpha a. \end{aligned}$$

Finally,

$$\alpha a = \bigcap \langle \bigcup \circ \langle \alpha \rangle \circ \text{atoms} \circ \uparrow^A \rangle \text{up } a = \bigcap \langle \alpha' \rangle \text{up } a = \langle f \rangle a,$$

so $\langle f \rangle$ is a continuation of α .

2. Consider the relation $\delta' \in \mathcal{P}(\mathcal{P}A \times \mathcal{P}B)$ defined by the formula (for every $X \in \mathcal{P}A, Y \in \mathcal{P}B$)

$$X \delta' Y \Leftrightarrow \exists x \in \text{atoms } \uparrow^A X, y \in \text{atoms } \uparrow^B Y: x \delta y.$$

Obviously $\neg(X \delta' \emptyset)$ and $\neg(\emptyset \delta' Y)$.

For suitable I and J we have:

$$\begin{aligned} (I \cup J) \delta' Y &\Leftrightarrow \exists x \in \text{atoms} \uparrow^A(I \cup J), y \in \text{atoms} \uparrow^B Y: x \delta y \\ &\Leftrightarrow \exists x \in \text{atoms} \uparrow^A I \cup \text{atoms} \uparrow^A J, y \in \text{atoms} \uparrow^B Y: x \delta y \\ &\Leftrightarrow \exists x \in \text{atoms} \uparrow^A I, y \in \text{atoms} \uparrow^B Y: x \delta y \vee \exists x \in \text{atoms} \uparrow^A J, y \in \text{atoms} \uparrow^B Y: x \delta y \\ &\Leftrightarrow I \delta' Y \vee J \delta' Y; \end{aligned}$$

analogously $X \delta' (I \cup J) \Leftrightarrow X \delta' I \vee X \delta' J$ for suitable I and J . Let's continue δ' till a funcoïd f (by the theorem 48):

$$\mathcal{X}[f]\mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: X \delta' Y$$

The reverse of (8) implication is trivial, so

$$\forall X \in \text{up } a, Y \in \text{up } b \exists x \in \text{atoms} \uparrow^A X, y \in \text{atoms} \uparrow^B Y: x \delta y \Leftrightarrow a \delta b.$$

$$\forall X \in \text{up } a, Y \in \text{up } b \exists x \in \text{atoms} \uparrow^A X, y \in \text{atoms} \uparrow^B Y: x \delta y \Leftrightarrow \forall X \in \text{up } a, Y \in \text{up } b: X \delta' Y \Leftrightarrow a[f]b.$$

So $a \delta b \Leftrightarrow a[f]b$, that is $[f]$ is a continuation of δ . \square

One of uses of the previous theorem is the proof of the following theorem:

Theorem 78. If A, B are small sets, $R \in \mathcal{P}\text{FCD}(A; B)$, $x \in \text{atoms } 1^{\mathfrak{F}(A)}$, $y \in \text{atoms } 1^{\mathfrak{F}(B)}$, then

1. $\langle \bigcap R \rangle x = \bigcap \{ \langle f \rangle x \mid f \in R \}$;
2. $x[\bigcap R]y \Leftrightarrow \forall f \in R: x[f]y$.

Proof. 2. Let denote $x \delta y \Leftrightarrow \forall f \in R: x[f]y$.

$$\begin{aligned} \forall X \in \text{up } a, Y \in \text{up } b \exists x \in \text{atoms} \uparrow^A X, y \in \text{atoms} \uparrow^B Y: x \delta y &\Leftrightarrow \\ \forall f \in R, X \in \text{up } a, Y \in \text{up } b \exists x \in \text{atoms} \uparrow^A X, y \in \text{atoms} \uparrow^B Y: x[f]y &\Rightarrow \\ \forall f \in R, X \in \text{up } a, Y \in \text{up } b: X[f]^* Y &\Rightarrow \\ \forall f \in R: a[f]b &\Leftrightarrow \\ a \delta b. & \end{aligned}$$

So, by the theorem 77, δ can be continued till $[p]$ for some funcoïd $p \in \text{FCD}(A; B)$.

For every funcoïd $q \in \text{FCD}(A; B)$ such that $\forall f \in R: q \subseteq f$ we have $x[q]y \Rightarrow \forall f \in R: x[f]y \Leftrightarrow x \delta y \Leftrightarrow x[p]y$, so $q \subseteq p$. Consequently $p = \bigcap R$.

From this $x[\bigcap R]y \Leftrightarrow \forall f \in R: x[f]y$.

1. From the former $y \in \text{atoms} \langle \bigcap R \rangle x \Leftrightarrow y \cap \langle \bigcap R \rangle x \neq 0^{\mathfrak{F}(B)} \Leftrightarrow \forall f \in R: y \cap \langle f \rangle x \neq \emptyset \Leftrightarrow y \in \bigcap \langle \text{atoms} \rangle \{ \langle f \rangle x \mid f \in R \} \Leftrightarrow y \in \text{atoms} \bigcap \{ \langle f \rangle x \mid f \in R \}$ for every $y \in \text{atoms } 1^{\mathfrak{F}(A)}$. From this follows $\langle \bigcap R \rangle x = \bigcap \{ \langle f \rangle x \mid f \in R \}$. \square

3.9 Direct product of filter objects

A generalization of direct (Cartesian) product of two sets is funcoïdal product of two filter objects:

Definition 79. *Funcoïdal product* of filter objects \mathcal{A} and \mathcal{B} is such a funcoïd $\mathcal{A} \times^{\text{FCD}} \mathcal{B} \in \text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))$ that for every $\mathcal{X} \in \mathfrak{F}(\text{Base}(\mathcal{A}))$, $\mathcal{Y} \in \mathfrak{F}(\text{Base}(\mathcal{B}))$

$$\mathcal{X}[\mathcal{A} \times^{\text{FCD}} \mathcal{B}]\mathcal{Y} \Leftrightarrow \mathcal{X} \cap \mathcal{A} \neq \emptyset \wedge \mathcal{Y} \cap \mathcal{B} \neq \emptyset.$$

Proposition 80. $\mathcal{A} \times^{\text{FCD}} \mathcal{B}$ is really a funcoïd and

$$\langle \mathcal{A} \times^{\text{FCD}} \mathcal{B} \rangle \mathcal{X} = \begin{cases} \mathcal{B} & \text{if } \mathcal{X} \not\asymp \mathcal{A}; \\ \emptyset & \text{if } \mathcal{X} \asymp \mathcal{A}. \end{cases}$$

Proof. Obvious. \square

Obvious 81. $\uparrow^{\text{FCD}(U;V)}(A \times B) = \uparrow^U A \times^{\text{FCD}} \uparrow^V B$ for sets $A \subseteq U$ and $B \subseteq V$ (for some small sets U and V).

Proposition 82. $f \subseteq \mathcal{A} \times^{\text{FCD}} \mathcal{B} \Leftrightarrow \text{dom } f \subseteq \mathcal{A} \wedge \text{im } f \subseteq \mathcal{B}$ for every $f \in \text{FCD}(A; B)$ and $\mathcal{A} \in \mathfrak{F}(A)$, $\mathcal{B} \in \mathfrak{F}(B)$.

Proof. If $f \subseteq \mathcal{A} \times^{\text{FCD}} \mathcal{B}$ then $\text{dom } f \subseteq \text{dom}(\mathcal{A} \times^{\text{FCD}} \mathcal{B}) \subseteq \mathcal{A}$, $\text{im } f \subseteq \text{im}(\mathcal{A} \times^{\text{FCD}} \mathcal{B}) \subseteq \mathcal{B}$. If $\text{dom } f \subseteq \mathcal{A} \wedge \text{im } f \subseteq \mathcal{B}$ then

$$\forall \mathcal{X} \in \mathfrak{F}(A), \mathcal{Y} \in \mathfrak{F}(B): (\mathcal{X}[f]\mathcal{Y} \Rightarrow \mathcal{X} \cap \mathcal{A} \neq 0^{\mathfrak{F}(A)} \wedge \mathcal{Y} \cap \mathcal{B} \neq 0^{\mathfrak{F}(B)});$$

consequently $f \subseteq \mathcal{A} \times^{\text{FCD}} \mathcal{B}$. □

The following theorem gives a formula for calculating an important particular case of intersection on the lattice of functors:

Theorem 83. $f \cap (\mathcal{A} \times^{\text{FCD}} \mathcal{B}) = I_{\mathcal{B}}^{\text{FCD}} \circ f \circ I_{\mathcal{A}}^{\text{FCD}}$ for every functor f and $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{B} \in \mathfrak{F}(\text{Dst } f)$.

Proof. $h \stackrel{\text{def}}{=} I_{\mathcal{B}}^{\text{FCD}} \circ f \circ I_{\mathcal{A}}^{\text{FCD}}$. For every $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$

$$\langle h \rangle \mathcal{X} = \langle I_{\mathcal{B}}^{\text{FCD}} \rangle \langle f \rangle \langle I_{\mathcal{A}}^{\text{FCD}} \rangle \mathcal{X} = \mathcal{B} \cap \langle f \rangle (\mathcal{A} \cap \mathcal{X}).$$

From this, as easy to show, $h \subseteq f$ and $h \subseteq \mathcal{A} \times^{\text{FCD}} \mathcal{B}$. If $g \subseteq f \wedge g \subseteq \mathcal{A} \times^{\text{FCD}} \mathcal{B}$ for a $g \in \text{FCD}(\text{Src } f; \text{Dst } f)$ then $\text{dom } g \subseteq \mathcal{A}$, $\text{im } g \subseteq \mathcal{B}$,

$$\langle g \rangle \mathcal{X} = \mathcal{B} \cap \langle g \rangle (\mathcal{A} \cap \mathcal{X}) \subseteq \mathcal{B} \cap \langle f \rangle (\mathcal{A} \cap \mathcal{X}) = \langle I_{\mathcal{B}}^{\text{FCD}} \rangle \langle f \rangle \langle I_{\mathcal{A}}^{\text{FCD}} \rangle \mathcal{X} = \langle h \rangle \mathcal{X},$$

$g \subseteq h$. So $h = f \cap (\mathcal{A} \times^{\text{FCD}} \mathcal{B})$. □

Corollary 84. $f|_{\mathcal{A}} = f \cap (\mathcal{A} \times^{\text{FCD}} 1^{\mathfrak{F}(\text{Dst } f)})$ for every $f \in \text{FCD}$ and $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$.

Proof. $f \cap (\mathcal{A} \times^{\text{FCD}} 1^{\mathfrak{F}(\text{Dst } f)}) = I_{1^{\mathfrak{F}(\text{Dst } f)}}^{\text{FCD}} \circ f \circ I_{\mathcal{A}}^{\text{FCD}} = f \circ I_{\mathcal{A}}^{\text{FCD}} = f|_{\mathcal{A}}$. □

Corollary 85. $f \not\subseteq (\mathcal{A} \times^{\text{FCD}} \mathcal{B}) \Leftrightarrow \mathcal{A}[f]\mathcal{B}$ for every $f \in \text{FCD}$, $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{B} \in \mathfrak{F}(\text{Dst } f)$.

Proof. $f \not\subseteq (\mathcal{A} \times^{\text{FCD}} \mathcal{B}) \Leftrightarrow \langle f \cap (\mathcal{A} \times^{\text{FCD}} \mathcal{B}) \rangle (\text{Src } f) \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \langle I_{\mathcal{B}}^{\text{FCD}} \circ f \circ I_{\mathcal{A}} \rangle (\text{Src } f) \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \langle I_{\mathcal{B}}^{\text{FCD}} \rangle \langle f \rangle \langle I_{\mathcal{A}}^{\text{FCD}} \rangle 1^{\mathfrak{F}(\text{Src } f)} \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \mathcal{B} \cap \langle f \rangle (\mathcal{A} \cap 1^{\mathfrak{F}(\text{Src } f)}) \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \mathcal{B} \cap \langle f \rangle \mathcal{A} \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \mathcal{A}[f]\mathcal{B}$. □

Corollary 86. Every filter of functors is star-separable.

Proof. The set of direct products of principal filter objects is a separation subset of the lattice of functors. □

Theorem 87. Let A, B are small sets. If $S \in \mathcal{P}(\mathfrak{F}(A) \times \mathfrak{F}(B))$ then

$$\bigcap \{ \mathcal{A} \times^{\text{FCD}} \mathcal{B} \mid (\mathcal{A}; \mathcal{B}) \in S \} = \bigcap \text{dom } S \times^{\text{FCD}} \bigcap \text{im } S.$$

Proof. If $x \in \text{atoms } 1^{\mathfrak{F}(A)}$ then by the theorem 78

$$\langle \bigcap \{ \mathcal{A} \times^{\text{FCD}} \mathcal{B} \mid (\mathcal{A}; \mathcal{B}) \in S \} \rangle x = \bigcap \{ \langle \mathcal{A} \times^{\text{FCD}} \mathcal{B} \rangle x \mid (\mathcal{A}; \mathcal{B}) \in S \}.$$

If $x \not\subseteq \bigcap \text{dom } S$ then

$$\begin{aligned} \forall (\mathcal{A}; \mathcal{B}) \in S: (x \cap \mathcal{A} \neq 0^{\mathfrak{F}(A)} \wedge \langle \mathcal{A} \times^{\text{FCD}} \mathcal{B} \rangle x = \mathcal{B}); \\ \{ \langle \mathcal{A} \times^{\text{FCD}} \mathcal{B} \rangle x \mid (\mathcal{A}; \mathcal{B}) \in S \} = \text{im } S; \end{aligned}$$

if $x \subseteq \bigcap \text{dom } S$ then

$$\begin{aligned} \exists (\mathcal{A}; \mathcal{B}) \in S: (x \cap \mathcal{A} = 0^{\mathfrak{F}(A)} \wedge \langle \mathcal{A} \times^{\text{FCD}} \mathcal{B} \rangle x = 0^{\mathfrak{F}(B)}); \\ \{ \langle \mathcal{A} \times^{\text{FCD}} \mathcal{B} \rangle x \mid (\mathcal{A}; \mathcal{B}) \in S \} \ni 0^{\mathfrak{F}(B)}. \end{aligned}$$

So

$$\langle \bigcap \{ \mathcal{A} \times^{\text{FCD}} \mathcal{B} \mid (\mathcal{A}; \mathcal{B}) \in S \} \rangle x = \begin{cases} \bigcap \text{im } S & \text{if } x \not\prec \bigcap \text{dom } S; \\ 0^{\mathfrak{F}(B)} & \text{if } x \succ \bigcap \text{dom } S. \end{cases}$$

From this follows the statement of the theorem. \square

Corollary 88. For every $\mathcal{A}_0, \mathcal{A}_1 \in \mathfrak{F}(A)$, $\mathcal{B}_0, \mathcal{B}_1 \in \mathfrak{F}(B)$ (for every small sets A, B)

$$(\mathcal{A}_0 \times^{\text{FCD}} \mathcal{B}_0) \cap (\mathcal{A}_1 \times^{\text{FCD}} \mathcal{B}_1) = (\mathcal{A}_0 \cap \mathcal{A}_1) \times^{\text{FCD}} (\mathcal{B}_0 \cap \mathcal{B}_1).$$

Proof. $(\mathcal{A}_0 \times^{\text{FCD}} \mathcal{B}_0) \cap (\mathcal{A}_1 \times^{\text{FCD}} \mathcal{B}_1) = \bigcap \{ \mathcal{A}_0 \times^{\text{FCD}} \mathcal{B}_0, \mathcal{A}_1 \times^{\text{FCD}} \mathcal{B}_1 \}$ what is by the last theorem equal to $(\mathcal{A}_0 \cap \mathcal{A}_1) \times^{\text{FCD}} (\mathcal{B}_0 \cap \mathcal{B}_1)$. \square

Theorem 89. If A, B are small sets and $\mathcal{A} \in \mathfrak{F}(A)$ then $\mathcal{A} \times^{\text{FCD}}$ is a complete homomorphism of the lattice $\mathfrak{F}(B)$ to a complete sublattice of the lattice $\text{FCD}(A; B)$, if also $\mathcal{A} \neq 0^{\mathfrak{F}(A)}$ then it is an isomorphism.

Proof. Let $S \in \mathcal{P}\mathfrak{F}(B)$, $X \in \mathcal{P}A$, $x \in \text{atoms } 1^{\mathfrak{F}(A)}$.

$$\begin{aligned} \langle \bigcup \langle \mathcal{A} \times^{\text{FCD}} \rangle S \rangle X &= \bigcup \{ \langle \mathcal{A} \times^{\text{FCD}} \mathcal{B} \rangle X \mid \mathcal{B} \in S \} \\ &= \begin{cases} \bigcup S & \text{if } X \in \star \partial \mathcal{A} \\ 0^{\mathfrak{F}(B)} & \text{if } X \notin \star \partial \mathcal{A} \end{cases} \\ &= \langle \mathcal{A} \times^{\text{FCD}} \bigcup S \rangle X; \\ \langle \bigcap \langle \mathcal{A} \times^{\text{FCD}} \rangle S \rangle x &= \bigcap \{ \langle \mathcal{A} \times^{\text{FCD}} \mathcal{B} \rangle x \mid \mathcal{B} \in S \} \\ &= \begin{cases} \bigcap S & \text{if } x \not\prec \mathcal{A} \\ 0^{\mathfrak{F}(B)} & \text{if } x \succ \mathcal{A} \end{cases} \\ &= \langle \mathcal{A} \times^{\text{FCD}} \bigcap S \rangle x. \end{aligned}$$

If $\mathcal{A} \neq 0^{\mathfrak{F}(A)}$ then obviously the function $\mathcal{A} \times^{\text{FCD}}$ is injective. \square

The following proposition states that cutting a rectangle of atomic width from a funcoid always produces a rectangular (representable as a direct product of filter objects) funcoid (of atomic width).

Proposition 90. If $f \in \text{FCD}$ and a is an atomic filter object on $\text{Src } f$ then

$$f|_a = a \times^{\text{FCD}} \langle f \rangle a.$$

Proof. Let $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$.

$$\mathcal{X} \not\prec a \Rightarrow \langle f|_a \rangle \mathcal{X} = \langle f \rangle a, \quad \mathcal{X} \succ a \Rightarrow \langle f|_a \rangle \mathcal{X} = 0^{\mathfrak{F}(\text{Dst } f)}. \quad \square$$

3.10 Atomic funcoids

Theorem 91. An $f \in \text{FCD}(A; B)$ is an atom of the lattice $\text{FCD}(A; B)$ iff it is direct product of two atomic filter objects.

Proof.

\Rightarrow . Let $f \in \text{FCD}(A; B)$ is an atom of the lattice $\text{FCD}(A; B)$. Let's get elements $a \in \text{atoms dom } f$ and $b \in \text{atoms } \langle f \rangle a$. Then for every $\mathcal{X} \in \mathfrak{F}(A)$

$$\mathcal{X} \succ a \Rightarrow \langle a \times^{\text{FCD}} b \rangle \mathcal{X} = 0^{\mathfrak{F}(B)} \subseteq \langle f \rangle \mathcal{X}, \quad \mathcal{X} \not\prec a \Rightarrow \langle a \times^{\text{FCD}} b \rangle \mathcal{X} = b \subseteq \langle f \rangle \mathcal{X}.$$

So $a \times^{\text{FCD}} b \subseteq f$; because f is atomic we have $f = a \times^{\text{FCD}} b$.

\Leftarrow . Let $a \in \text{atoms } 1^{\mathfrak{F}(A)}$, $b \in \text{atoms } 1^{\mathfrak{F}(B)}$, $f \in \text{FCD}(A; B)$. If $b \asymp \langle f \rangle a$ then $\neg(a[f]b)$, $f \asymp a \times^{\text{FCD}} b$; if $b \subseteq \langle f \rangle a$ then $\forall \mathcal{X} \in \mathfrak{F}(A): (\mathcal{X} \not\asymp a \Rightarrow \langle f \rangle \mathcal{X} \supseteq b)$, $f \supseteq a \times^{\text{FCD}} b$. Consequently $f \asymp a \times^{\text{FCD}} b \vee f \supseteq a \times^{\text{FCD}} b$; that is $a \times^{\text{FCD}} b$ is an atomic filter object. \square

Theorem 92. The lattice $\text{FCD}(A; B)$ is atomic (for every small sets A, B).

Proof. Let f is a non-empty funcoid. Then $\text{dom } f \neq \emptyset$, thus by the theorem 46 in [14] exists $a \in \text{atoms dom } f$. So $\langle f \rangle a \neq 0^{\mathfrak{F}(B)}$ thus exists $b \in \text{atoms } \langle f \rangle a$. Finally the atomic funcoid $a \times^{\text{FCD}} b \subseteq f$. \square

Theorem 93. The lattice $\text{FCD}(A; B)$ is separable (for every small sets A, B).

Proof. Let $f, g \in \text{FCD}(A; B)$, $f \subset g$. Then exists $a \in \text{atoms } 1^{\mathfrak{F}(A)}$ such that $\langle f \rangle a \subset \langle g \rangle a$. So because the lattice $\mathfrak{F}(B)$ is atomically separable then exists $b \in \text{atoms } 1^{\mathfrak{F}(B)}$ such that $\langle f \rangle a \cap b = 0^{\mathfrak{F}(B)}$ and $b \subseteq \langle g \rangle a$. For every $x \in \text{atoms } 1^{\mathfrak{F}(A)}$

$$\begin{aligned} \langle f \rangle a \cap \langle a \times^{\text{FCD}} b \rangle a &= \langle f \rangle a \cap b = 0^{\mathfrak{F}(B)}, \\ x \neq a \Rightarrow \langle f \rangle x \cap \langle a \times^{\text{FCD}} b \rangle x &= \langle f \rangle x \cap 0^{\mathfrak{F}(B)} = 0^{\mathfrak{F}(B)}. \end{aligned}$$

Thus $\langle f \rangle x \cap \langle a \times^{\text{FCD}} b \rangle x = 0^{\mathfrak{F}(B)}$ and consequently $f \asymp a \times^{\text{FCD}} b$.

$$\begin{aligned} \langle a \times^{\text{FCD}} b \rangle a &= b \subseteq \langle g \rangle a, \\ x \neq a \Rightarrow \langle a \times^{\text{FCD}} b \rangle x &= 0^{\mathfrak{F}(B)} \subseteq \langle g \rangle a. \end{aligned}$$

Thus $\langle a \times^{\text{FCD}} b \rangle x \subseteq \langle g \rangle x$ and consequently $a \times^{\text{FCD}} b \subseteq g$.

So the lattice $\text{FCD}(A; B)$ is separable by the theorem 19 in [14]. \square

Corollary 94. The lattice $\text{FCD}(A; B)$ is:

1. separable;
2. atomically separable;
3. conforming to Wallman's disjunction property.

Proof. By the theorem 22 in [14]. \square

Remark 95. For more ways to characterize (atomic) separability of the lattice of funcoids see [14], subsections "Separation subsets and full stars" and "Atomically separable lattices".

Corollary 96. The lattice $\text{FCD}(A; B)$ is an atomistic lattice.

Proof. Let $f \in \text{FCD}(A; B)$. Suppose contrary to the statement to be proved that $\bigcup \text{atoms } f \subset f$. Then it exists $a \in \text{atoms } f$ such that $a \cap \bigcup \text{atoms } f = \emptyset$ what is impossible. \square

Proposition 97. $\text{atoms}(f \cup g) = \text{atoms } f \cup \text{atoms } g$ for every funcoids $f, g \in \text{FCD}(A; B)$ (for every small sets A and B).

Proof. $a \times^{\text{FCD}} b \not\asymp f \cup g \Leftrightarrow a[f \cup g]b \Leftrightarrow a[f]b \vee a[g]b \Leftrightarrow a \times^{\text{FCD}} b \not\asymp f \vee a \times^{\text{FCD}} b \not\asymp g$ for every atomic filter objects a and b . \square

Theorem 98. For every $f, g, h \in \text{FCD}(A; B)$, $R \in \mathcal{P}\text{FCD}(A; B)$ (for every small sets A and B)

1. $f \cap (g \cup h) = (f \cap g) \cup (f \cap h)$;
2. $f \cup \bigcap R = \bigcap \langle f \cup \rangle R$.

Proof. We will take in account that the lattice of funcoids is an atomistic lattice.

1. $\text{atoms}(f \cap (g \cup h)) = \text{atoms } f \cap \text{atoms}(g \cup h) = \text{atoms } f \cap (\text{atoms } g \cup \text{atoms } h) = (\text{atoms } f \cap \text{atoms } g) \cup (\text{atoms } f \cap \text{atoms } h) = \text{atoms}(f \cap g) \cup \text{atoms}(f \cap h) = \text{atoms}((f \cap g) \cup (f \cap h))$.

2. $\text{atoms}(f \cup \bigcap R) = \text{atoms } f \cup \text{atoms } \bigcap R = \text{atoms } f \cup \bigcap \langle \text{atoms} \rangle R = \bigcap \langle (\text{atoms } f) \cup \rangle \langle \text{atoms} \rangle R = \bigcap \langle \text{atoms} \rangle \langle f \cup \rangle R = \text{atoms } \bigcap \langle f \cup \rangle R$. (Used the following equality.)

$$\begin{aligned}
& \langle (\text{atoms } f) \cup \rangle \langle \text{atoms} \rangle R = \\
& \{ (\text{atoms } f) \cup A \mid A \in \langle \text{atoms} \rangle R \} = \\
& \{ (\text{atoms } f) \cup A \mid \exists C \in R: A = \text{atoms } C \} = \\
& \{ (\text{atoms } f) \cup (\text{atoms } C) \mid C \in R \} = \\
& \{ \text{atoms}(f \cup C) \mid C \in R \} = \\
& \{ \text{atoms } B \mid \exists C \in R: B = f \cup C \} = \\
& \{ \text{atoms } B \mid B \in \langle f \cup \rangle R \} = \\
& \langle \text{atoms} \rangle \langle f \cup \rangle R.
\end{aligned}$$

□

Note that distributivity of the lattice of funcoids is proved through using atoms of this lattice. I have never seen such method of proving distributivity.

Corollary 99. The lattice $\text{FCD}(A; B)$ is co-brouwerian (for every small sets A and B).

The next proposition is one more (among the theorem 60) generalization for funcoids of composition of relations.

Proposition 100. For every composable funcoids f, g

$$\text{atoms}(g \circ f) = \{ x \times^{\text{FCD}} z \mid x \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, z \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } g)}, \exists y \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)}: (x \times^{\text{FCD}} y \in \text{atoms } f \wedge y \times^{\text{FCD}} z \in \text{atoms } g) \}.$$

Proof. $(x \times^{\text{FCD}} z) \cap (g \circ f) \neq \emptyset \Leftrightarrow x[g \circ f]z \Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)}: (x[f]y \wedge y[g]z) \Leftrightarrow \exists y \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)}: ((x \times^{\text{FCD}} y) \cap f \neq \emptyset \wedge (y \times^{\text{FCD}} z) \cap g \neq \emptyset)$ (it was used the theorem 60). □

3.11 Complete funcoids

Definition 101. I will call *co-complete* such a funcoid f that $\langle f \rangle^* X$ is a principal f.o. for every $X \in \mathcal{P}(\text{Src } f)$.

Remark 102. I will call *generalized closure* such a function $\alpha \in \mathcal{P}B^{\mathcal{P}A}$ (for some small sets A, B) that

1. $\alpha \emptyset = \emptyset$;
2. $\forall I, J \in \mathcal{P}A: \alpha(I \cup J) = \alpha I \cup \alpha J$.

Obvious 103. A funcoid f is co-complete iff $\langle f \rangle^* = \uparrow^{\text{Dst } f} \circ \alpha$ for a generalized closure α .

Remark 104. Thus funcoids can be considered as a generalization of generalized closures. A topological space in Kuratowski sense is the same as reflexive and transitive generalized closure. So topological spaces can be considered as a special case of funcoids.

Definition 105. I will call a *complete funcoid* a funcoid whose reverse is co-complete.

Theorem 106. The following conditions are equivalent for every funcoid f :

1. funcoid f is complete;
2. $\forall S \in \mathcal{P}\mathfrak{F}(\text{Src } f), J \in \mathcal{P}(\text{Dst } f): (\bigcup S[f] \uparrow^{\text{Dst } f} J \Leftrightarrow \exists I \in S: I[f] \uparrow^{\text{Dst } f} J)$;
3. $\forall S \in \mathcal{P}\mathcal{P}(\text{Src } f), J \in \mathcal{P}(\text{Dst } f): (\bigcup S[f]^* J \Leftrightarrow \exists I \in S: I[f]^* J)$;
4. $\forall S \in \mathcal{P}\mathfrak{F}(\text{Src } f): \langle f \rangle \cup S = \bigcup \langle \langle f \rangle \rangle S$;

5. $\forall S \in \mathcal{P}\mathcal{P}(\text{Src } f): \langle f \rangle^* \cup S = \bigcup \langle \langle f \rangle^* \rangle S;$
 6. $\forall A \in \mathcal{P}(\text{Src } f): \langle f \rangle^* A = \bigcup \{ \langle f \rangle^* \{a\} \mid a \in A \}.$

Proof.

(3) \Rightarrow (1). For every $S \in \mathcal{P}\mathcal{P}(\text{Src } f), J \in \mathcal{P}(\text{Dst } f)$

$$\uparrow^{\text{Src } f} \bigcup S \cap \langle f^{-1} \rangle^* J \neq \emptyset \Leftrightarrow \exists I \in S: \uparrow^{\text{Src } f} I \cap \langle f^{-1} \rangle^* J \neq \emptyset, \quad (10)$$

consequently by the theorem 52 in [14] we have that $\langle f^{-1} \rangle^* J$ is a principal f.o.

(1) \Rightarrow (2). For every $S \in \mathcal{P}\mathcal{F}(\text{Src } f), J \in \mathcal{P}(\text{Dst } f)$ we have $\langle f^{-1} \rangle^* J$ a principal f.o., consequently

$$\bigcup S \cap \langle f^{-1} \rangle^* J \neq \emptyset \Leftrightarrow \exists I \in S: I \cap \langle f^{-1} \rangle^* J \neq \emptyset.$$

From this follows (2).

(6) \Rightarrow (5). $\langle f \rangle^* \cup S = \bigcup \{ \langle f \rangle^* \{a\} \mid a \in \bigcup S \} = \bigcup \{ \bigcup \{ \langle f \rangle^* \{a\} \mid a \in A \} \mid A \in S \} = \bigcup \{ \langle f \rangle^* A \mid A \in S \} = \bigcup \langle \langle f \rangle^* \rangle S.$

(2) \Rightarrow (4). $\uparrow^{\text{Dst } f} J \neq \langle f \rangle \cup S \Leftrightarrow \bigcup S [f] \uparrow^{\text{Dst } f} J \Leftrightarrow \exists I \in S: I [f] \uparrow^{\text{Dst } f} J \Leftrightarrow \exists I \in S: \uparrow^{\text{Dst } f} J \neq \langle f \rangle I \Leftrightarrow \uparrow^{\text{Dst } f} J \neq \bigcup \langle \langle f \rangle \rangle S$ (used the theorem 53 in [14]).

(2) \Rightarrow (3), (4) \Rightarrow (5), (5) \Rightarrow (3), (5) \Rightarrow (6). Obvious. \square

The following proposition shows that complete funcoids are a direct generalization of pre-topological spaces.

Proposition 107. To specify a complete funcoid f it is enough to specify $\langle f \rangle^*$ on one-element sets, values of $\langle f \rangle^*$ on one element sets can be specified arbitrarily.

Proof. From the above theorem is clear that knowing $\langle f \rangle^*$ on one-element sets $\langle f \rangle^*$ can be found on every set and then its value can be inferred for every filter objects.

Choosing arbitrarily the values of $\langle f \rangle^*$ on one-element sets we can define a complete funcoid the following way: $\langle f \rangle^* X \stackrel{\text{def}}{=} \bigcup \{ \langle f \rangle^* \{ \alpha \} \mid \alpha \in X \}$ for every $X \in \mathcal{P}(\text{Src } f)$. Obviously it is really a complete funcoid. \square

Theorem 108. A funcoid is discrete iff it is both complete and co-complete.

Proof.

\Rightarrow . Obvious.

\Leftarrow . Let f is both a complete and co-complete funcoid. Consider the relation g defined by that $\uparrow^{\text{Dst } f} \langle g \rangle \{ \alpha \} = \langle f \rangle^* \{ \alpha \}$ (g is correctly defined because f corresponds to a generalized closure). Because f is a complete funcoid f is the funcoid corresponding to g . \square

Theorem 109. If $R \in \mathcal{P}\mathcal{FCD}(A; B)$ is a set of (co-)complete funcoids then $\bigcup R$ is a (co-)complete funcoid (for every small sets A and B).

Proof. It is enough to prove only for co-complete funcoids. Let $R \in \mathcal{P}\mathcal{FCD}(A; B)$ is a set of co-complete funcoids. Then for every $X \in \mathcal{P}(\text{Src } f)$

$$\langle \bigcup R \rangle^* X = \bigcup \{ \langle f \rangle^* X \mid f \in R \}$$

is a principal f.o. (used the theorem 57). \square

Corollary 110. If R is a set of binary relations between small sets A and B then $\bigcup \langle \uparrow^{\text{FCD}(A; B)} \rangle R = \uparrow^{\text{FCD}(A; B)} \bigcup R.$

Proof. From two last theorems. \square

Theorem 111. Filtrators of functors are filtered.

Proof. It's enough to prove that every functor is representable as (infinite) meet (on the lattice $\text{FCD}(A; B)$) of some set of discrete functors.

Let $f \in \text{FCD}(A; B)$, $X \in \mathcal{P}A$, $Y \in \text{up}\langle f \rangle A$, $g(X; Y) \stackrel{\text{def}}{=} \uparrow^A X \times^{\text{FCD}} \uparrow^B Y \cup \uparrow^A \bar{X} \times^{\text{FCD}} 1^{\mathfrak{F}(B)}$. For every $K \in \mathcal{P}A$

$$\langle g(X; Y) \rangle^* K = \langle \uparrow^A X \times^{\text{FCD}} \uparrow^B Y \rangle^* K \cup \langle \uparrow^A \bar{X} \times^{\text{FCD}} 1^{\mathfrak{F}(B)} \rangle^* K = \left(\begin{array}{l} 0^{\mathfrak{F}(B)} \text{ if } K = \emptyset \\ Y \text{ if } \emptyset \neq K \subseteq X \\ 1^{\mathfrak{F}(B)} \text{ if } K \not\subseteq X \end{array} \right) \supseteq \langle f \rangle^* K;$$

so $g(X; Y) \supseteq f$. For every $X \in \mathcal{P}A$

$$\bigcap \{ \langle g(X; Y) \rangle^* X \mid Y \in \text{up}\langle f \rangle X \} = \bigcap \{ Y \mid Y \in \text{up}\langle f \rangle^* X \} = \langle f \rangle^* X;$$

consequently

$$\bigcap \{ g(X; Y) \mid X \in \mathcal{P}A, Y \in \text{up}\langle f \rangle^* X \} = f. \quad \square$$

Conjecture 112. If $f \in \text{FCD}(B; C)$ is a complete functor and $R \in \mathcal{P}\text{FCD}(A; B)$ then $f \circ \bigcup R = \bigcup \langle f \circ \rangle R$.

This conjecture can be weakened:

Conjecture 113. If f is a discrete functor from B to C and $R \in \mathcal{P}\text{FCD}(A; B)$ then $f \circ \bigcup R = \bigcup \langle f \circ \rangle R$.

I will denote ComplFCD and CoComplFCD the sets of complete and co-complete functors correspondingly. $\text{ComplFCD}(A; B)$ are complete functors from A to B and likewise with $\text{CoComplFCD}(A; B)$.

Obvious 114. ComplFCD and CoComplFCD are closed regarding composition of functors.

Proposition 115. $\text{ComplFCD}(A; B)$ and $\text{CoComplFCD}(A; B)$ (with induced order) are complete lattices.

Proof. It follows from the theorem 109. □

Theorem 116. Atoms of the lattice $\text{ComplFCD}(A; B)$ are exactly direct products of the form $\uparrow^A \{\alpha\} \times^{\text{FCD}} b$ where $\alpha \in A$ and b is an atomic f.o. on B

Proof. First, it's easy to see that $\{\alpha\} \times^{\text{FCD}} b$ are elements of $\text{ComplFCD}(A; B)$. Also $0^{\text{FCD}(A; B)}$ is an element of $\text{ComplFCD}(A; B)$.

$\uparrow^A \{\alpha\} \times^{\text{FCD}} b$ are atoms of $\text{ComplFCD}(A; B)$ because these are atoms of $\text{FCD}(A; B)$.

It remains to prove that if f is an atom of $\text{ComplFCD}(A; B)$ then $f = \{\alpha\} \times^{\text{FCD}} b$ for some $\alpha \in A$ and an atomic f.o. b on B .

Suppose $f \in \text{FCD}(A; B)$ is a non-empty complete functor. Then exists $\alpha \in A$ such that $\langle f \rangle^* \{\alpha\} \neq 0^{\mathfrak{F}(B)}$. Thus $\uparrow^A \{\alpha\} \times^{\text{FCD}} b \subseteq f$ for some atomic f.o. b on B . If f is an atom then $f = \uparrow^A \{\alpha\} \times^{\text{FCD}} b$. □

Theorem 117.

1. A functor $f \in \text{FCD}(A; B)$ is complete iff there exists a function $G: \text{Src } f \rightarrow \mathfrak{F}(\text{Dst } f)$ such that

$$f = \bigcup \{ \uparrow^A \{\alpha\} \times^{\text{FCD}} G(\alpha) \mid \alpha \in \text{Src } f \}. \quad (11)$$

2. A functor $f \in \text{FCD}(A; B)$ is co-complete iff there exists a function $G: \text{Dst } f \rightarrow \mathfrak{F}(\text{Src } f)$ such that

$$f = \bigcup \{ G(\alpha) \times^{\text{FCD}} \uparrow^A \{\alpha\} \mid \alpha \in \text{Dst } f \}.$$

Proof. We will prove only the first as the second is symmetric.

\Rightarrow . Let f is complete. Then take

$$G(\alpha) = \bigcup \{b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} \mid \exists \alpha \in \text{Src } f: \uparrow^A \{\alpha\} \times^{\text{FCD}} b \subseteq f\}$$

and we have (11) obviously.

\Leftarrow . Let (11) holds. Then $G(\alpha) = \bigcup \text{atoms } G(\alpha)$ and thus

$$f = \bigcup \{\{\alpha\} \times^{\text{FCD}} b \mid \alpha \in \text{Src } f, b \in \text{atoms } G(\alpha)\}$$

and so f is complete. \square

Theorem 118.

1. For a complete funcoid f there exist exactly one function $F \in \mathfrak{F}(\text{Dst } f)^{\text{Src } f}$ such that

$$f = \bigcup \{\uparrow^{\text{Src } f} \{\alpha\} \times^{\text{FCD}} F(\alpha) \mid \alpha \in \text{Src } f\}.$$

2. For a co-complete funcoid f there exist exactly one function $F \in \mathfrak{F}(\text{Src } f)^{\text{Dst } f}$ such that

$$f = \bigcup \{F(\alpha) \times^{\text{FCD}} \uparrow^{\text{Dst } f} \{\alpha\} \mid \alpha \in \text{Dst } f\}.$$

Proof. We will prove only the first as the second is similar. Let

$$f = \bigcup \{\uparrow^{\text{Src } f} \{\alpha\} \times^{\text{FCD}} F(\alpha) \mid \alpha \in \text{Src } f\} = \bigcup \{\uparrow^{\text{Src } f} \{\alpha\} \times^{\text{FCD}} G(\alpha) \mid \alpha \in \text{Src } f\}$$

for some $F, G \in \mathfrak{F}(\text{Dst } f)^{\text{Src } f}$. We need to prove $F = G$. Let $\beta \in \text{Src } f$.

$$\langle f \rangle \{\beta\} = \bigcup \{\langle \uparrow^{\text{Src } f} \{\alpha\} \times^{\text{FCD}} F(\alpha) \rangle \{\beta\} \mid \alpha \in \text{Src } f\} = F(\beta).$$

Similarly $\langle f \rangle \{\beta\} = G(\beta)$. So $F(\beta) = G(\beta)$. \square

3.12 Completion of funcoids

Theorem 119. $\text{Cor } f = \text{Cor}' f$ for an element f of a filtrator of funcoids. (Core part is taken for the shifted filtrator of funcoids.)

Proof. From the theorem 26 in [14] and the corollary 110 and theorem 111. \square

Definition 120. *Completion* of a funcoid $f \in \text{FCD}(A; B)$ is the complete funcoid $\text{Compl } f \in \text{FCD}(A; B)$ defined by the formula $\langle \text{Compl } f \rangle^* \{\alpha\} = \langle f \rangle^* \{\alpha\}$ for $\alpha \in \text{Src } f$.

Definition 121. *Co-completion* of a funcoid f is defined by the formula

$$\text{CoCompl } f = (\text{Compl } f^{-1})^{-1}.$$

Obvious 122. $\text{Compl } f \subseteq f$ and $\text{CoCompl } f \subseteq f$ for every funcoid f .

Proposition 123. The filtrator $(\text{FCD}(A; B); \text{ComplFCD}(A; B))$ is filtered.

Proof. Because the shifted filtrator $(\text{FCD}(A; B); \mathcal{P}(A \times B); \uparrow^{\text{FCD}(A; B)})$ is filtered. \square

Theorem 124. $\text{Compl } f = \text{Cor}^{(\text{FCD}(A; B); \text{ComplFCD}(A; B))} f = \text{Cor}'^{(\text{FCD}(A; B); \text{ComplFCD}(A; B))} f$ for every funcoid $f \in \text{FCD}(A; B)$.

Proof. $\text{Cor}^{(\text{FCD}(A; B); \text{ComplFCD}(A; B))} f = \text{Cor}'^{(\text{FCD}(A; B); \text{ComplFCD}(A; B))} f$ since (the theorem 26 in [14]) the filtrator $(\text{FCD}(A; B); \text{ComplFCD}(A; B))$ is filtered and with join closed core (the theorem 109).

Let $g \in \text{up}^{(\text{FCD}(A; B); \text{ComplFCD}(A; B))} f$. Then $g \in \text{ComplFCD}(A; B)$ and $g \supseteq f$. Thus $g = \text{Compl } g \supseteq \text{Compl } f$.

Thus $\forall g \in \text{up}^{\text{FCD}(A;B); \text{ComplFCD}(A;B)} f: g \supseteq \text{Compl } f$.

Let $\forall g \in \text{up}^{\text{FCD}(A;B); \text{ComplFCD}(A;B)} f: h \subseteq g$ for some $h \in \text{ComplFCD}(A;B)$.

Then $h \subseteq \bigcap^{\text{FCD}(A;B)} \text{up}^{\text{FCD}(A;B); \text{ComplFCD}(A;B)} f = f$ and consequently $h = \text{Compl } h \subseteq \text{Compl } f$.

Thus

$$\text{Compl } f = \bigcap^{\text{ComplFCD}(A;B)} \text{up}^{\text{FCD}(A;B); \text{ComplFCD}(A;B)} f = \text{Cor}^{\text{FCD}(A;B); \text{ComplFCD}(A;B)} f. \quad \square$$

Theorem 125. $\langle \text{CoCompl } f \rangle^* X = \text{Cor } \langle f \rangle^* X$ for every funcooid f and set $X \in \mathcal{P}(\text{Src } f)$.

Proof. $\text{CoCompl } f \subseteq f$ thus $\langle \text{CoCompl } f \rangle^* X \subseteq \langle f \rangle^* X$, but $\langle \text{CoCompl } f \rangle^* X$ is a principal f.o. thus $\langle \text{CoCompl } f \rangle^* X \subseteq \text{Cor } \langle f \rangle^* X$.

Let $\alpha X = \text{Cor } \langle f \rangle^* X$. Then $\alpha \emptyset = 0^{\mathfrak{F}(\text{Dst } f)}$ and

$$\alpha(X \cup Y) = \text{Cor } \langle f \rangle^*(X \cup Y) = \text{Cor}(\langle f \rangle^* X \cup \langle f \rangle^* Y) = \text{Cor } \langle f \rangle^* X \cup \text{Cor } \langle f \rangle^* Y = \alpha X \cup \alpha Y.$$

(used the theorem 64 from [14]). Thus α can be continued till $\langle g \rangle$ for some funcooid g . This funcooid is co-complete.

Evidently g is the greatest co-complete element of $\text{FCD}(\text{Src } f; \text{Dst } f)$ which is lower than f .

Thus $g = \text{CoCompl } f$ and so $\text{Cor } \langle f \rangle^* X = \alpha X = \langle g \rangle^* X = \langle \text{CoCompl } f \rangle^* X$. \square

Theorem 126. $\text{ComplFCD}(A;B)$ is an atomistic lattice.

Proof. Let $f \in \text{ComplFCD}(A;B)$. $\langle f \rangle^* X = \bigcup \{ \langle f \rangle^* \{x\} \mid x \in X \} = \bigcup \{ \langle f|_{\uparrow^{\text{Src } f} \{x\}} \rangle^* \{x\} \mid x \in X \} = \langle f|_{\uparrow^{\text{Src } f} \{x\}} \rangle^* X$, thus $f = f|_{\uparrow^{\text{Src } f} \{x\}}$. It is trivial that every $f|_{\uparrow^{\text{Src } f} \{x\}}$ is a union of atoms of $\text{ComplFCD}(A;B)$. \square

Theorem 127. A funcooid f is complete iff it is a join (on the lattice $\text{FCD}(\text{Src } f; \text{Dst } f)$) of atomic complete funcooids.

Proof. Follows from the theorem 109 and the previous theorem. \square

Corollary 128. $\text{ComplFCD}(A;B)$ is join-closed.

Theorem 129. $\text{Compl}(\bigcup R) = \bigcup \langle \text{Compl} \rangle R$ for every $R \in \mathcal{P}\text{FCD}(A;B)$ (for every small sets A, B).

Proof. $\langle \text{Compl}(\bigcup R) \rangle^* X = \bigcup \{ \langle \bigcup R \rangle^* \{\alpha\} \mid \alpha \in X \} = \bigcup \{ \bigcup \{ \langle f \rangle^* \{\alpha\} \mid f \in R \} \mid \alpha \in X \} = \bigcup \{ \bigcup \{ \langle f \rangle^* \{\alpha\} \mid \alpha \in X \} \mid f \in R \} = \bigcup \{ \langle \text{Compl } f \rangle^* X \mid f \in R \} = \langle \bigcup \langle \text{Compl} \rangle R \rangle^* X$ for every set X . \square

Proposition 130. $\text{Compl } f = \bigcup \{ f|_{\uparrow^{\text{Src } f} \{\alpha\}} \mid \alpha \in \text{Src } f \}$ for every funcooid f .

Proof. Let denote R the right part of the equality to prove.

$\langle R \rangle^* \{\beta\} = \bigcup \{ \langle f|_{\uparrow^{\text{Src } f} \{\alpha\}} \rangle^* \{\beta\} \mid \alpha \in \text{Src } f \} = \langle f \rangle^* \{\beta\}$ for every $\beta \in A$ and R is complete as a join of complete funcooids.

Thus R is the completion of f . \square

Corollary 131. Compl is a lower adjoint.

Conjecture 132. Compl is not an upper adjoint (in general).

Conjecture 133. $\text{Compl } f = f \setminus^* (\Omega \times^{\text{FCD}} \mathcal{U})$ for every funcooid f .

This conjecture may be proved by considerations similar to these in the section ‘‘Fréchet filter’’ in [14].

Lemma 134. Co-completion of a complete funcooid is complete.

Proof. Let f is a complete funcooid.

$\langle \text{CoCompl } f \rangle^* X = \text{Cor } \langle f \rangle^* X = \text{Cor } \bigcup \{ \langle f \rangle^* \{x\} \mid x \in X \} = \bigcup \{ \text{Cor } \langle f \rangle^* \{x\} \mid x \in X \} = \bigcup \{ \langle \text{CoCompl } f \rangle^* \{x\} \mid x \in X \}$ for every set X . Thus $\text{CoCompl } f$ is complete. \square

Theorem 135. $\text{Compl CoCompl } f = \text{CoCompl Compl } f = \text{Cor } f$ for every funcoid f .

Proof. $\text{Compl CoCompl } f$ is co-complete since (used the lemma) $\text{CoCompl } f$ is co-complete. Thus $\text{Compl CoCompl } f$ is a discrete funcoid. $\text{CoCompl } f$ is the the greatest co-complete funcoid under f and $\text{Compl CoCompl } f$ is the greatest complete funcoid under $\text{CoCompl } f$. So $\text{Compl CoCompl } f$ is greater than any discrete funcoid under $\text{CoCompl } f$ which is greater than any discrete funcoid under f . Thus $\text{Compl CoCompl } f$ it is the greatest discrete funcoid under f . Thus $\text{Compl CoCompl } f = \text{Cor } f$. Similarly $\text{CoCompl Compl } f = \text{Cor } f$. \square

Question 136. Is $\text{ComplFCD}(A; B)$ a co-brouwerian lattice for every small sets A, B ?

3.13 Monovalued and injective funcoids

Following the idea of definition of monovalued morphism let's call *monovalued* such a funcoid f that $f \circ f^{-1} \subseteq I_{\text{im } f}^{\text{FCD}}$.

Similarly, I will call a funcoid *injective* when $f^{-1} \circ f \subseteq I_{\text{dom } f}^{\text{FCD}}$.

Obvious 137. A funcoid f is

- monovalued iff $f \circ f^{-1} \subseteq I^{\text{FCD}(\text{Dst } f)}$;
- injective iff $f^{-1} \circ f \subseteq I^{\text{FCD}(\text{Src } f)}$.

In other words, a funcoid is monovalued (injective) when it is a monovalued (injective) morphism of the category of funcoids.

Monovaluedness is dual of injectivity.

Obvious 138.

1. A morphism $(f; \mathcal{A}; \mathcal{B})$ of the category of funcoid triples is monovalued iff the funcoid f is monovalued.
2. A morphism $(f; \mathcal{A}; \mathcal{B})$ of the category of funcoid triples is injective iff the funcoid f is injective.

Theorem 139. The following statements are equivalent for a funcoid f :

1. f is monovalued.
2. $\forall a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}: \langle f \rangle a \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} \cup \{0^{\mathfrak{F}(\text{Dst } f)}\}$.
3. $\forall \mathcal{I}, \mathcal{J} \in \mathfrak{F}(\text{Dst } f): \langle f^{-1} \rangle (\mathcal{I} \cap \mathcal{J}) = \langle f^{-1} \rangle \mathcal{I} \cap \langle f^{-1} \rangle \mathcal{J}$.
4. $\forall I, J \in \mathcal{P}(\text{Dst } f): \langle f^{-1} \rangle^* (I \cap J) = \langle f^{-1} \rangle^* I \cap \langle f^{-1} \rangle^* J$.

Proof.

(2) \Rightarrow (3). Let $a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}$, $\langle f \rangle a = b$. Then because $b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} \cup \{0^{\mathfrak{F}(\text{Dst } f)}\}$

$$\begin{aligned} (\mathcal{I} \cap \mathcal{J}) \cap b \neq \emptyset &\Leftrightarrow \mathcal{I} \cap b \neq \emptyset \wedge \mathcal{J} \cap b \neq \emptyset; \\ a[f](\mathcal{I} \cap \mathcal{J}) &\Leftrightarrow a[f]\mathcal{I} \wedge a[f]\mathcal{J}; \\ (\mathcal{I} \cap \mathcal{J})[f^{-1}]a &\Leftrightarrow \mathcal{I}[f^{-1}]a \wedge \mathcal{J}[f^{-1}]a; \\ a \cap \langle f^{-1} \rangle (\mathcal{I} \cap \mathcal{J}) \neq \emptyset &\Leftrightarrow a \cap \langle f^{-1} \rangle \mathcal{I} \neq \emptyset \wedge a \cap \langle f^{-1} \rangle \mathcal{J} \neq \emptyset; \\ \langle f^{-1} \rangle (\mathcal{I} \cap \mathcal{J}) &= \langle f^{-1} \rangle \mathcal{I} \cap \langle f^{-1} \rangle \mathcal{J}. \end{aligned}$$

(3) \Rightarrow (1). $\langle f^{-1} \rangle a \cap \langle f^{-1} \rangle b = \langle f^{-1} \rangle (a \cap b) = \langle f^{-1} \rangle 0^{\mathfrak{F}(\text{Dst } f)} = 0^{\mathfrak{F}(\text{Src } f)}$ for every two distinct atomic filter objects a and b on $\text{Dst } f$. This is equivalent to $\neg(\langle f^{-1} \rangle a[f]b)$; $b \asymp \langle f \rangle \langle f^{-1} \rangle a$; $b \asymp \langle f \circ f^{-1} \rangle a$; $\neg(a[f \circ f^{-1}]b)$. So $a[f \circ f^{-1}]b \Rightarrow a = b$ for every atomic filter objects a and b . This is possible only when $f \circ f^{-1} \subseteq I^{\text{FCD}(\text{Dst } f)}$.

(4) \Rightarrow (3). $\langle f^{-1} \rangle(\mathcal{I} \cap \mathcal{J}) = \bigcap \langle \langle f^{-1} \rangle^* \rangle \text{up}(\mathcal{I} \cap \mathcal{J}) = \bigcap \langle \langle f^{-1} \rangle^* \rangle \{I \cap J \mid I \in \text{up } \mathcal{I}, J \in \text{up } \mathcal{J}\} = \bigcap \{\langle f^{-1} \rangle^*(I \cap J) \mid I \in \text{up } \mathcal{I}, J \in \text{up } \mathcal{J}\} = \bigcap \{\langle f^{-1} \rangle^* I \cap \langle f^{-1} \rangle^* J \mid I \in \text{up } \mathcal{I}, J \in \text{up } \mathcal{J}\} = \bigcap \{\langle f^{-1} \rangle^* I \mid I \in \text{up } \mathcal{I}\} \cap \bigcap \{\langle f^{-1} \rangle^* J \mid J \in \text{up } \mathcal{J}\} = \langle f^{-1} \rangle \mathcal{I} \cap \langle f^{-1} \rangle \mathcal{J}.$

(3) \Rightarrow (4). Obvious.

\neg (2) \Rightarrow \neg (1). Suppose $\langle f \rangle a \notin \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} \cup \{0^{\mathfrak{F}(\text{Dst } f)}\}$ for some $a \in \text{atoms } \mathcal{A}$. Then there exist two atomic filter objects p and q on $\text{Dst } f$ such that $p \neq q$ and $\langle f \rangle a \supseteq p \wedge \langle f \rangle a \supseteq q$. Consequently $p \not\prec \langle f \rangle a$; $a \not\prec \langle f^{-1} \rangle p$; $a \subseteq \langle f^{-1} \rangle p$; $\langle f \circ f^{-1} \rangle p = \langle f \rangle \langle f^{-1} \rangle p \supseteq \langle f \rangle a \supseteq q$; $\langle f \circ f^{-1} \rangle p \not\subseteq p$ and $\langle f \circ f^{-1} \rangle p \neq 0^{\mathfrak{F}(\text{Dst } f)}$. So it cannot be $f \circ f^{-1} \subseteq I^{\text{FCD}(\text{Dst } f)}$. \square

Corollary 140. A binary relation corresponds to a monovalued funcoid iff it is a function.

Proof. Because $\forall I, J \in \mathcal{P}(\text{im } f): \langle f^{-1} \rangle^*(I \cap J) = \langle f^{-1} \rangle^* I \cap \langle f^{-1} \rangle^* J$ is true for a funcoid f corresponding to a binary relation if and only if it is a function. \square

Remark 141. This corollary can be reformulated as follows: For binary relations (discrete funcoids) the classic concept of monovaluedness and monovaluedness in the above defined sense of monovaluedness of a funcoid are the same.

3.14 T_0 -, T_1 - and T_2 -separable funcoids

For funcoids it can be generalized T_0 -, T_1 - and T_2 - separability. Worthwhile note that T_0 and T_2 separability is defined through T_1 separability.

Definition 142. Let call T_1 -separable such funcoid f that for every $\alpha \in \text{Src } f$, $\beta \in \text{Dst } f$ is true

$$\alpha \neq \beta \Rightarrow \neg(\{\alpha\}[f]^*\{\beta\}).$$

Definition 143. Let call T_0 -separable such funcoid $f \in \text{FCD}(A; A)$ that $f \cap f^{-1}$ is T_1 -separable.

Definition 144. Let call T_2 -separable such funcoid f that the funcoid $f^{-1} \circ f$ is T_1 -separable.

For symmetric transitive funcoids T_1 - and T_2 -separability are the same (see theorem 14).

Obvious 145. A funcoid f is T_2 -separable iff $\alpha \neq \beta \Rightarrow \langle f \rangle^* \{\alpha\} \succ \langle f \rangle^* \{\beta\}$ for every $\alpha, \beta \in \text{Src } f$.

3.15 Filter objects closed regarding a funcoid

Definition 146. Let's call *closed* regarding a funcoid $f \in \text{FCD}(A; A)$ such filter object $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$ that $\langle f \rangle \mathcal{A} \subseteq \mathcal{A}$.

This is a generalization of closedness of a set regarding an unary operation.

Proposition 147. If \mathcal{I} and \mathcal{J} are closed (regarding some funcoid f), S is a set of closed filter objects on $\text{Src } f$, then

1. $\mathcal{I} \cup \mathcal{J}$ is a closed filter object;
2. $\bigcap S$ is a closed filter object.

Proof. Let denote the given funcoid as f . $\langle f \rangle(\mathcal{I} \cup \mathcal{J}) = \langle f \rangle \mathcal{I} \cup \langle f \rangle \mathcal{J} \subseteq \mathcal{I} \cup \mathcal{J}$, $\langle f \rangle \bigcap S \subseteq \bigcap \langle \langle f \rangle \rangle S \subseteq \bigcap S$. Consequently the filter objects $\mathcal{I} \cup \mathcal{J}$ and $\bigcap S$ are closed. \square

Proposition 148. If S is a set of filter objects closed regarding a complete funcoid, then the filter object $\bigcup S$ is also closed regarding our funcoid.

Proof. $\langle f \rangle \bigcup S = \bigcup \langle \langle f \rangle \rangle S \subseteq \bigcup S$ where f is the given funcoid. \square

4 Reloids

Definition 149. I will call a *reloid* from a small set A to a small set B a triple $(A; B; F)$ where $F \in \mathfrak{F}(A \times B)$.

Definition 150. *Source* and *destination* of every reloid $(A; B; F)$ are defined as

$$\text{Src}(A; B; F) = A \quad \text{and} \quad \text{Dst}(A; B; F) = B.$$

I will denote $\text{RLD}(A; B)$ the set of reloids from A to B .

I will denote RLD the set of all reloids (for small sets).

Further we will assume that all reloids in consideration are small.

Reloids are a generalization of uniform spaces. Also reloids are generalization of binary relations (I will call a reloid $(A; B; F)$ *discrete* when F is a principal filter on $A \times B$.)

I will denote $\text{up}(A; B; F) = \text{up } F$ for every reloid $(A; B; F)$.

Definition 151. The *reverse* reloid of a reloid f is defined by the formula

$$(A; B; F)^{-1} = (B; A; \{F^{-1} \mid F \in \text{up } f^{-1}\}).$$

Reverse reloid is a generalization of conjugate quasi-uniformity.

I will denote $\uparrow^{\text{RLD}(A; B)} f = (A; B; \uparrow^{A \times B} f)$ for every small sets A, B and a binary relation $f \subseteq A \times B$.

4.1 Composition of reloids

Definition 152. Reloids f and g are *composable* when $\text{Dst } f = \text{Src } g$.

Definition 153. Composition of (composable) reloids is defined by the formula

$$g \circ f = \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } g)} (G \circ F) \mid F \in \text{up } f, G \in \text{up } g \}.$$

Composition of reloids is a reloid.

Theorem 154. $(h \circ g) \circ f = h \circ (g \circ f)$ for every composable reloids f, g, h .

Proof. For two nonempty collections A and B of sets I will denote

$$A \sim B \Leftrightarrow (\forall K \in A \exists L \in B: L \subseteq K) \wedge (\forall K \in B \exists L \in A: L \subseteq K).$$

It is easy to see that \sim is a transitive relation.

I will denote $B \circ A = \{L \circ K \mid K \in A, L \in B\}$.

Let first prove that for every nonempty collections of relations A, B, C

$$A \sim B \Rightarrow A \circ C \sim B \circ C.$$

Suppose $A \sim B$ and $P \in A \circ C$ that is $K \in A$ and $M \in C$ such that $P = K \circ M$. $\exists K' \in B: K' \subseteq K$ because $A \sim B$. We have $P' = K' \circ M \in B \circ C$. Obviously $P' \subseteq P$. So for every $P \in A \circ C$ exist $P' \in B \circ C$ such that $P' \subseteq P$; the vice versa is analogous. So $A \circ C \sim B \circ C$.

$\text{up}((h \circ g) \circ f) \sim \text{up}(h \circ g) \circ \text{up } f$, $\text{up}(h \circ g) \sim (\text{up } h) \circ (\text{up } g)$. By proven above $\text{up}((h \circ g) \circ f) \sim (\text{up } h) \circ (\text{up } g) \circ (\text{up } f)$.

Analogously $\text{up}(h \circ (g \circ f)) \sim (\text{up } h) \circ (\text{up } g) \circ (\text{up } f)$.

So $\text{up}((h \circ g) \circ f) \sim \text{up}(h \circ (g \circ f))$ what is possible only if $\text{up}((h \circ g) \circ f) = \text{up}(h \circ (g \circ f))$. \square

Theorem 155. For every reloid f :

1. $f \circ f = \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} (F \circ F) \mid F \in \text{up } f \}$ if $\text{Src } f = \text{Dst } f$;
2. $f^{-1} \circ f = \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Src } f)} (F^{-1} \circ F) \mid F \in \text{up } f \}$;
3. $f \circ f^{-1} = \bigcap \{ \uparrow^{\text{RLD}(\text{Dst } f; \text{Dst } f)} (F \circ F^{-1}) \mid F \in \text{up } f \}$.

Proof. I will prove only (1) and (2) because (3) is analogous to (2).

1. It's enough to show that $\forall F, G \in \text{up } f \exists H \in \text{up } f: H \circ H \subseteq G \circ F$. To prove it take $H = F \cap G$.
2. It's enough to show that $\forall F, G \in \text{up } f \exists H \in \text{up } f: H^{-1} \circ H \subseteq G^{-1} \circ F$. To prove it take $H = F \cap G$. Then $H^{-1} \circ H = (F \cap G)^{-1} \circ (F \cap G) \subseteq G^{-1} \circ F$. \square

Theorem 156. For every small sets A, B, C if $g, h \in \text{RLD}(A; B)$ then

1. $f \circ (g \cup h) = f \circ g \cup f \circ h$ for every $f \in \text{RLD}(B; C)$;
2. $(g \cup h) \circ f = g \circ f \cup h \circ f$ for every $f \in \text{RLD}(C; A)$.

Proof. We'll prove only the first as the second is dual.

By the infinite distributivity law for filters we have

$$\begin{aligned} f \circ g \cup f \circ h &= \bigcap \{ \uparrow^{\text{RLD}(A; C)}(F \circ G) \mid F \in \text{up } f, G \in \text{up } g \} \cup \bigcap \{ \uparrow^{\text{RLD}(A; C)}(F \circ H) \mid F \in \text{up } f, \\ &\quad H \in \text{up } h \} \\ &= \bigcap \{ \uparrow^{\text{RLD}(A; C)}(F_1 \circ G) \cup \uparrow^{\text{RLD}(A; C)}(F_2 \circ H) \mid F_1, F_2 \in \text{up } f, G \in \text{up } g, H \in \text{up } h \} \\ &= \bigcap \{ \uparrow^{\text{RLD}(A; C)}((F_1 \circ G) \cup (F_2 \circ H)) \mid F_1, F_2 \in \text{up } f, G \in \text{up } g, H \in \text{up } h \}. \end{aligned}$$

Obviously

$$\begin{aligned} &\bigcap \{ \uparrow^{\text{RLD}(A; C)}((F_1 \circ G) \cup (F_2 \circ H)) \mid F_1, F_2 \in \text{up } f, G \in \text{up } g, H \in \text{up } h \} \supseteq \\ &\bigcap \{ \uparrow^{\text{RLD}(A; C)}(((F_1 \cap F_2) \circ G) \cup ((F_1 \cap F_2) \circ H)) \mid F_1, F_2 \in \text{up } f, G \in \text{up } g, H \in \text{up } h \} = \\ &\quad \bigcap \{ \uparrow^{\text{RLD}(A; C)}((F \circ G) \cup (F \circ H)) \mid F \in \text{up } f, G \in \text{up } g, H \in \text{up } h \} = \\ &\quad \bigcap \{ \uparrow^{\text{RLD}(A; C)}(F \circ (G \cup H)) \mid F \in \text{up } f, G \in \text{up } g, H \in \text{up } h \}. \end{aligned}$$

Because $G \in \text{up } g \wedge H \in \text{up } h \Rightarrow G \cup H \in \text{up}(g \cup h)$ we have

$$\begin{aligned} &\bigcap \{ \uparrow^{\text{RLD}(A; C)}(F \circ (G \cup H)) \mid F \in \text{up } f, G \in \text{up } g, H \in \text{up } h \} \supseteq \\ &\quad \bigcap \{ \uparrow^{\text{RLD}(A; C)}(F \circ K) \mid F \in \text{up } f, K \in \text{up}(g \cup h) \} = \\ &\quad f \circ (g \cup h). \end{aligned}$$

Thus we proved $f \circ g \cup f \circ h \supseteq f \circ (g \cup h)$. But obviously $f \circ (g \cup h) \supseteq f \circ g$ and $f \circ (g \cup h) \supseteq f \circ h$ and so $f \circ (g \cup h) \supseteq f \circ g \cup f \circ h$. Thus $f \circ (g \cup h) = f \circ g \cup f \circ h$. \square

Conjecture 157. If f and g are reloids, then

$$g \circ f = \bigcup \{ G \circ F \mid F \in \text{atoms } f, G \in \text{atoms } g \}.$$

4.2 Direct product of filter objects

Definition 158. *Reloidal product* of filter objects \mathcal{A} and \mathcal{B} is defined by the formula

$$\mathcal{A} \times^{\text{RLD}} \mathcal{B} \stackrel{\text{def}}{=} \bigcap \{ \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))}(A \times B) \mid A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B} \}.$$

Obvious 159. $\uparrow^U \mathcal{A} \times^{\text{RLD}} \uparrow^V \mathcal{B} = \uparrow^{\text{RLD}(U; V)}(A \times B)$ for every small sets $A \subseteq U$ and $B \subseteq V$.

Theorem 160. $\mathcal{A} \times^{\text{RLD}} \mathcal{B} = \bigcup \{ a \times^{\text{RLD}} b \mid a \in \text{atoms } \mathcal{A}, b \in \text{atoms } \mathcal{B} \}$ for every filter objects \mathcal{A}, \mathcal{B} .

Proof. Obviously

$$\mathcal{A} \times^{\text{RLD}} \mathcal{B} \supseteq \bigcup \{ a \times^{\text{RLD}} b \mid a \in \text{atoms } \mathcal{A}, b \in \text{atoms } \mathcal{B} \}$$

Reversely, let

$$K \in \text{up} \bigcup \{ a \times^{\text{RLD}} b \mid a \in \text{atoms } \mathcal{A}, b \in \text{atoms } \mathcal{B} \}.$$

Then $K \in \text{up}(a \times^{\text{RLD}} b)$ for every $a \in \text{atoms } \mathcal{A}, b \in \text{atoms } \mathcal{B}$; $K \supseteq X_a \times Y_b$ for some $X_a \in \text{up } a, Y_b \in \text{up } b$; $K \supseteq \bigcup \{ X_a \times Y_b \mid a \in \text{atoms } \mathcal{A}, b \in \text{atoms } \mathcal{B} \} = \bigcup \{ X_a \mid a \in \text{atoms } \mathcal{A} \} \times \bigcup \{ Y_b \mid b \in \text{atoms } \mathcal{B} \} \supseteq A \times B$ where $A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B}$; $K \in \text{up}(A \times^{\text{RLD}} B)$. \square

Theorem 161. If $\mathcal{A}_0, \mathcal{A}_1 \in \mathfrak{F}(A)$, $\mathcal{B}_0, \mathcal{B}_1 \in \mathfrak{F}(B)$ for some small sets A, B then

$$(\mathcal{A}_0 \times^{\text{RLD}} \mathcal{B}_0) \cap (\mathcal{A}_1 \times^{\text{RLD}} \mathcal{B}_1) = (\mathcal{A}_0 \cap \mathcal{A}_1) \times^{\text{RLD}} (\mathcal{B}_0 \cap \mathcal{B}_1).$$

Proof.

$$\begin{aligned} (\mathcal{A}_0 \times^{\text{RLD}} \mathcal{B}_0) \cap (\mathcal{A}_1 \times^{\text{RLD}} \mathcal{B}_1) &= \bigcap \{ \uparrow^{\text{RLD}(A;B)}(P \cap Q) \mid P \in \text{up}(\mathcal{A}_0 \times^{\text{RLD}} \mathcal{B}_0), Q \in \\ &\quad \text{up}(\mathcal{A}_1 \times^{\text{RLD}} \mathcal{B}_1) \} \\ &= \bigcap \{ \uparrow^{\text{RLD}(A;B)}((A_0 \times B_0) \cap (A_1 \times B_1)) \mid A_0 \in \text{up } \mathcal{A}_0, B_0 \in \text{up } \mathcal{B}_0, \\ &\quad A_1 \in \text{up } \mathcal{A}_1, B_1 \in \text{up } \mathcal{B}_1 \} \\ &= \bigcap \{ \uparrow^{\text{RLD}(A;B)}((A_0 \cap A_1) \times (B_0 \cap B_1)) \mid A_0 \in \text{up } \mathcal{A}_0, B_0 \in \text{up } \mathcal{B}_0, \\ &\quad A_1 \in \text{up } \mathcal{A}_1, B_1 \in \text{up } \mathcal{B}_1 \} \\ &= \bigcap \{ \uparrow^{\text{RLD}(A;B)}(K \times L) \mid K \in \text{up}(\mathcal{A}_0 \cap \mathcal{A}_1), L \in \text{up}(\mathcal{B}_0 \cap \mathcal{B}_1) \} \\ &= (\mathcal{A}_0 \cap \mathcal{A}_1) \times^{\text{RLD}} (\mathcal{B}_0 \cap \mathcal{B}_1). \end{aligned}$$

□

Theorem 162. If $S \in \mathcal{P}(\mathfrak{F}(A) \times \mathfrak{F}(B))$ for some small sets A, B then

$$\bigcap \{ \mathcal{A} \times^{\text{RLD}} \mathcal{B} \mid (\mathcal{A}; \mathcal{B}) \in S \} = \bigcap \text{dom } S \times^{\text{RLD}} \bigcap \text{im } S.$$

Proof. Let $\mathcal{P} = \bigcap \text{dom } S$, $\mathcal{Q} = \bigcap \text{im } S$; $l = \bigcap \{ \mathcal{A} \times^{\text{RLD}} \mathcal{B} \mid (\mathcal{A}; \mathcal{B}) \in S \}$.

$\mathcal{P} \times^{\text{RLD}} \mathcal{Q} \subseteq l$ is obvious.

Let $F \in \text{up}(\mathcal{P} \times^{\text{RLD}} \mathcal{Q})$. Then exist $P \in \text{up } \mathcal{P}$ and $Q \in \text{up } \mathcal{Q}$ such that $F \supseteq P \times Q$.

$P = P_1 \cap \dots \cap P_n$ where $P_i \in \langle \text{up} \rangle \text{dom } S$ and $Q = Q_1 \cap \dots \cap Q_m$ where $Q_i \in \langle \text{up} \rangle \text{im } S$.

$P \times Q = \bigcap_{i,j} (P_i \times Q_j)$.

$P_i \times Q_j \in \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{B})$ for some $(\mathcal{A}; \mathcal{B}) \in S$. $P \times Q = \bigcap_{i,j} (P_i \times Q_j) \in \text{up } l$. So $F \in \text{up } l$. □

Conjecture 163. If $\mathcal{A} \in \mathfrak{F}$ then $\mathcal{A} \times^{\text{RLD}}$ is a complete homomorphism of every lattice $\mathfrak{F}(B)$ to a complete sublattice of the lattice $\text{RLD}(\text{Base}(A); \text{Base}(B))$, if also $\mathcal{A} \neq \emptyset$ then it is an isomorphism.

Definition 164. I will call a reloid *convex* iff it is a union of direct products.

Example 165. Non-convex reloids exist.

Proof. Let a is a non-trivial atomic f.o. Then $\uparrow^{\text{RLD}(\text{Base}(a); \text{Base}(a))}(=)|_a$ is non-convex. This follows from the fact that only direct products which are below $(=)$ are direct products of atomic f.o. and $\uparrow^{\text{RLD}(\text{Base}(a); \text{Base}(a))}(=)|_a$ is not their join. □

4.3 Restricting reloid to a filter object. Domain and image

Definition 166. *Identity reloid* for a small set A is defined by the formula $I^{\text{RLD}(A)} = \uparrow^{\text{RLD}(A;A)} I_A$.

Definition 167. I call *restricting* a reloid f to a filter object \mathcal{A} as $f|_{\mathcal{A}} = f \cap (\mathcal{A} \times^{\text{RLD}} 1_{\mathfrak{F}(\text{Dst } f)})$.

Definition 168. *Domain* and *image* of a reloid f are defined as follows:

$$\text{dom } f = \bigcap \langle \uparrow^{\text{Dst } f} \rangle \langle \text{dom} \rangle \text{up } f; \quad \text{im } f = \bigcap \langle \uparrow^{\text{Src } f} \rangle \langle \text{im} \rangle \text{up } f.$$

Proposition 169. $f \subseteq \mathcal{A} \times^{\text{RLD}} \mathcal{B} \Leftrightarrow \text{dom } f \subseteq \mathcal{A} \wedge \text{im } f \subseteq \mathcal{B}$ for every reloid f and filter objects $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{B} \in \mathfrak{F}(\text{Dst } f)$.

Proof.

\Rightarrow . Follows from $\text{dom}(\mathcal{A} \times^{\text{RLD}} \mathcal{B}) \subseteq \mathcal{A} \wedge \text{im}(\mathcal{A} \times^{\text{RLD}} \mathcal{B}) \subseteq \mathcal{B}$.

\Leftarrow . $\text{dom } f \subseteq \mathcal{A} \Leftrightarrow \forall A \in \text{up } \mathcal{A} \exists F \in \text{up } f: \text{dom } F \subseteq A$. Analogously

$$\text{im } f \subseteq \mathcal{B} \Leftrightarrow \forall B \in \text{up } \mathcal{B} \exists G \in \text{up } f: \text{im } G \subseteq B.$$

Let $\text{dom } f \subseteq \mathcal{A} \wedge \text{im } f \subseteq \mathcal{B}$, $A \in \text{up } \mathcal{A}$, $B \in \text{up } \mathcal{B}$. Then exist $F \in \text{up } f$, $G \in \text{up } f$ such that $\text{dom } F \subseteq A \wedge \text{im } G \subseteq B$. Consequently $F \cap G \in \text{up } f$, $\text{dom}(F \cap G) \subseteq A$, $\text{im}(F \cap G) \subseteq B$ that is $F \cap G \subseteq A \times B$. So exists $H \in \text{up } f$ such that $H \subseteq A \times B$ for every $A \in \text{up } \mathcal{A}$, $B \in \text{up } \mathcal{B}$. So $f \subseteq \mathcal{A} \times^{\text{RLD}} \mathcal{B}$. \square

Definition 170. I call *restricted identity reloid* for a filter object \mathcal{A} the reloid

$$I_{\mathcal{A}}^{\text{RLD}} \stackrel{\text{def}}{=} (I^{\text{RLD}(\mathcal{A})})|_{\mathcal{A}}.$$

Theorem 171. $I_{\mathcal{A}}^{\text{RLD}} = \bigcap \{\uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} I_A \mid A \in \text{up } \mathcal{A}\}$ where I_A is the identity relation on a set A .

Proof. Let $K \in \text{up } \bigcap \{\uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} I_A \mid A \in \text{up } \mathcal{A}\}$, then exists $A \in \text{up } \mathcal{A}$ such that $K \supseteq I_A$. Then

$$\begin{aligned} I_{\mathcal{A}}^{\text{RLD}} &\subseteq \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} (I_{\text{Base}(\mathcal{A})}) \cap (\mathcal{A} \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Base}(\mathcal{A}))}) \subseteq \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} (I_{\text{Base}(\mathcal{A})}) \cap \\ &(\uparrow^{\text{Base}(\mathcal{A})} A \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Base}(\mathcal{A}))}) = \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} (I_{\text{Base}(\mathcal{A})}) \cap \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} (A \times \\ &\text{Base}(\mathcal{A})) = \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} (I_{\text{Base}(\mathcal{A})} \cap (A \times \text{Base}(\mathcal{A}))) = \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} I_A \subseteq \\ &\uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} K, \end{aligned}$$

Thus $K \in \text{up } I_{\mathcal{A}}^{\text{RLD}}$.

Reversely let $K \in \text{up } I_{\mathcal{A}}^{\text{RLD}} = \text{up}(I^{\text{RLD}(\mathcal{A})} \cap (\mathcal{A} \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Base}(\mathcal{A}))}))$, then exists $A \in \text{up } \mathcal{A}$ such that $K \in \text{up}(I^{\text{RLD}(\mathcal{A})} \cap (A \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Base}(\mathcal{A}))})) = \text{up } \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} I_A \subseteq \text{up } \bigcap \{\uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} I_A \mid A \in \text{up } \mathcal{A}\}$. \square

Proposition 172. $(I_{\mathcal{A}}^{\text{RLD}})^{-1} = I_{\mathcal{A}}^{\text{RLD}}$.

Proof. Follows from the previous theorem. \square

Theorem 173. $f|_{\mathcal{A}} = f \circ I_{\mathcal{A}}^{\text{RLD}}$ for every reloid f and $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$.

Proof. We need to prove that $f \cap (\mathcal{A} \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Dst } f)}) = f \circ \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Src } f)} I_A \mid A \in \text{up } \mathcal{A}\}$. We have $f \circ \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Src } f)} I_A \mid A \in \text{up } \mathcal{A}\} = \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} (F \circ I_A) \mid F \in \text{up } f, A \in \text{up } \mathcal{A}\} = \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} (F|_A) \mid F \in \text{up } f, A \in \text{up } \mathcal{A}\} = \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} (F \cap (A \times \text{Dst } f)) \mid F \in \text{up } f, A \in \text{up } \mathcal{A}\} = \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} F \mid F \in \text{up } f\} \cap \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} (A \times \text{Dst } f) \mid A \in \text{up } \mathcal{A}\} = f \cap (\mathcal{A} \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Dst } f)})$. \square

Theorem 174. $(g \circ f)|_{\mathcal{A}} = g \circ (f|_{\mathcal{A}})$ for every composable reلودs f and g and $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$.

Proof. $(g \circ f)|_{\mathcal{A}} = (g \circ f) \circ I_{\mathcal{A}}^{\text{RLD}} = g \circ (f \circ I_{\mathcal{A}}^{\text{RLD}}) = g \circ (f|_{\mathcal{A}})$. \square

Theorem 175. $f \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{B}) = I_{\mathcal{B}}^{\text{RLD}} \circ f \circ I_{\mathcal{A}}^{\text{RLD}}$ for every reloid f and $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{B} \in \mathfrak{F}(\text{Dst } f)$.

Proof. $f \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{B}) = f \cap (\mathcal{A} \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Dst } f)}) \cap (1^{\mathfrak{F}(\text{Src } f)} \times^{\text{RLD}} \mathcal{B}) = f|_{\mathcal{A}} \cap (1^{\mathfrak{F}(\text{Src } f)} \times^{\text{RLD}} \mathcal{B}) = (f \circ I_{\mathcal{A}}^{\text{RLD}}) \cap (1^{\mathfrak{F}(\text{Src } f)} \times^{\text{RLD}} \mathcal{B}) = ((f \circ I_{\mathcal{A}}^{\text{RLD}})^{-1} \cap (1^{\mathfrak{F}(\text{Src } f)} \times^{\text{RLD}} \mathcal{B})^{-1})^{-1} = ((I_{\mathcal{A}}^{\text{RLD}} \circ f^{-1}) \cap (\mathcal{B} \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Src } f)}))^{-1} = (I_{\mathcal{A}}^{\text{RLD}} \circ f^{-1} \circ I_{\mathcal{B}}^{\text{RLD}})^{-1} = I_{\mathcal{B}}^{\text{RLD}} \circ f \circ I_{\mathcal{A}}^{\text{RLD}}$. \square

Theorem 176. $f|_{\uparrow^{\text{Src } f} \{\alpha\}} = \uparrow^{\text{Src } f} \{\alpha\} \times^{\text{RLD}} \text{im}(f|_{\uparrow^{\text{Src } f} \{\alpha\}})$ for every reloid f and $\alpha \in \text{Src } f$.

Proof. First,

$$\begin{aligned} \text{im}(f|_{\uparrow^{\text{Src } f} \{\alpha\}}) &= \\ &\bigcap \langle \uparrow^{\text{Dst } f} \rangle \langle \text{im} \rangle \text{up}(f|_{\uparrow^{\text{Src } f} \{\alpha\}}) = \\ &\bigcap \langle \uparrow^{\text{Dst } f} \rangle \langle \text{im} \rangle \text{up}(f \cap (\uparrow^{\text{Src } f} \{\alpha\} \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Dst } f)})) = \\ &\bigcap \langle \uparrow^{\text{Dst } f} \rangle \text{im}(F \cap (\{\alpha\} \times \text{Dst } f)) \mid F \in \text{up } f = \\ &\bigcap \langle \uparrow^{\text{Dst } f} \rangle \text{im}(F|_{\{\alpha\}}) \mid F \in \text{up } f. \end{aligned}$$

Taking this into account we have:

$$\begin{aligned}
& \uparrow^{\text{Src } f} \{\alpha\} \times^{\text{RLD}} \text{im}(f|_{\uparrow^{\text{Src } f} \{\alpha\}}) = \\
& \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(\{\alpha\} \times K) \mid K \in \text{up im}(f|_{\uparrow^{\text{Src } f} \{\alpha\}}) \} = \\
& \quad \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(\{\alpha\} \times \text{im}(F|_{\{\alpha\}})) \mid F \in \text{up } f \} = \\
& \quad \quad \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(F|_{\{\alpha\}}) \mid F \in \text{up } f \} = \\
& \quad \quad \quad \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(F \cap (\{\alpha\} \times \text{Dst } f)) \mid F \in \text{up } f \} = \\
& \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} F \mid F \in \text{up } f \} \cap \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(\{\alpha\} \times \text{Dst } f) = \\
& \quad f \cap \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(\{\alpha\} \times \text{Dst } f) = \\
& \quad \quad f|_{\uparrow^{\text{Src } f} \{\alpha\}}.
\end{aligned}$$

□

4.4 Categories of reloids

I will define two categories, the *category of reloids* and the *category of reloid triples*.

The *category of reloids* is defined as follows:

- Objects are small sets.
- The set of morphisms from a set A to a set B is $\text{RLD}(A; B)$.
- The composition is the composition of reloids.
- Identity morphism for a set is the identity reloid for that set.

To show it is really a category is trivial.

The *category of reloid triples* is defined as follows:

- Objects are filter objects on small sets.
- The morphisms from a f.o. \mathcal{A} to a f.o. \mathcal{B} are triples $(f; \mathcal{A}; \mathcal{B})$ where $f \in \text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))$ and $\text{dom } f \subseteq \mathcal{A} \wedge \text{im } f \subseteq \mathcal{B}$.
- The composition is defined by the formula $(g; \mathcal{B}; \mathcal{C}) \circ (f; \mathcal{A}; \mathcal{B}) = (g \circ f; \mathcal{A}; \mathcal{C})$.
- Identity morphism for an f.o. \mathcal{A} is $I_{\mathcal{A}}^{\text{RLD}}$.

To prove that it is really a category is trivial.

4.5 Monovalued and injective reloids

Following the idea of definition of monovalued morphism let's call *monovalued* such a reloid f that $f \circ f^{-1} \subseteq I_{\text{im } f}^{\text{RLD}}$.

Similarly, I will call a reloid *injective* when $f^{-1} \circ f \subseteq I_{\text{dom } f}^{\text{RLD}}$.

Obvious 177. A reloid f is

- monovalued iff $f \circ f^{-1} \subseteq I^{\text{RLD}}(\text{Dst } f)$;
- injective iff $f^{-1} \circ f \subseteq I^{\text{RLD}}(\text{Src } f)$.

In other words, a funcooid is monovalued (injective) when it is a monovalued (injective) morphism of the category of funcooids.

Monovaluedness is dual of injectivity.

Obvious 178.

1. A morphism $(f; \mathcal{A}; \mathcal{B})$ of the category of reloid triples is monovalued iff the reloid f is monovalued.
2. A morphism $(f; \mathcal{A}; \mathcal{B})$ of the category of reloid triples is injective iff the reloid f is injective.

Theorem 179.

1. A reloid f is a monovalued iff it exists a function (monovalued binary relation) $F \in \text{up } f$.
2. A reloid f is a injective iff it exists an injective binary relation $F \in \text{up } f$.
3. A reloid f is a both monovalued and injective iff exists an injection (a monovalued and injective binary relation = injective function) $F \in \text{up } f$.

Proof. The reverse implications are obvious. Let's prove the direct implications:

1. Let f is a monovalued reloid. Then $f \circ f^{-1} \subseteq I^{\text{RLD}(\text{Dst } f)}$. So exists

$$h \in \text{up}(f \circ f^{-1}) = \text{up} \bigcap \{ \uparrow^{\text{RLD}(\text{Dst } f; \text{Dst } f)}(F \circ F^{-1}) \mid F \in \text{up } f \}$$

such that $h \subseteq I^{\text{RLD}(\text{Dst } f)}$. It's simple to show that $\{F \circ F^{-1} \mid F \in \text{up } f\}$ is a filter base. Consequently it exists $F \in \text{up } f$ such that $F \circ F^{-1} \subseteq I_{\text{Dst } f}$ that is F is a function.

2. Similar.

3. Let f is a both monovalued and injective reloid. Then by proved above there exist $F, G \in \text{up } f$ such that F is monovalued and G is injective. Thus $F \cap G \in \text{up } f$ is both monovalued and injective. \square

Conjecture 180. A reloid f is monovalued iff

$$\forall g \in \text{RLD}(\text{Src } f; \text{Dst } f): (g \subseteq f \Rightarrow \exists \mathcal{A} \in \mathfrak{F}(\text{Src } f): g = f|_{\mathcal{A}}).$$

4.6 Complete reloids and completion of reloids

Definition 181. A *complete* reloid is a reloid representable as join of direct products $\uparrow^A \{\alpha\} \times^{\text{RLD}} b$ where $\alpha \in A$ and b is an atomic f.o. on B for some small sets A and B .

Definition 182. A *co-complete* reloid is a reloid representable as join of direct products $a \times^{\text{RLD}} \uparrow^B \{\beta\}$ where $\beta \in B$ and a is an atomic f.o. on A for some small sets A and B .

I will denote the sets of complete and co-complete reloids correspondingly as CompRLD and CoCompRLD .

Obvious 183. Complete and co-complete are dual.

Theorem 184.

1. A reloid f is complete iff there exists a function $G: \text{Src } f \rightarrow \mathfrak{F}(\text{Dst } f)$ such that

$$f = \bigcup \{ \uparrow^{\text{Src } f} \{\alpha\} \times^{\text{RLD}} G(\alpha) \mid \alpha \in \text{Src } f \}. \quad (12)$$

2. A reloid f is co-complete iff there exists a function $G: \text{Dst } f \rightarrow \mathfrak{F}(\text{Src } f)$ such that

$$f = \bigcup \{ G(\alpha) \times^{\text{RLD}} \uparrow^{\text{Dst } f} \{\alpha\} \mid \alpha \in \text{Dst } f \}.$$

Proof. We will prove only the first as the second is symmetric.

\Rightarrow . Let f is complete. Then take

$$G(\alpha) = \bigcup \{ b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} \mid \exists \alpha \in \text{Src } f: \uparrow^{\text{Src } f} \{\alpha\} \times^{\text{RLD}} b \subseteq f \}$$

and we have (12) obviously.

\Leftarrow . Let (12) holds. Then $G(\alpha) = \bigcup \text{atoms } G(\alpha)$ and thus

$$f = \bigcup \{ \uparrow^{\text{Src } f} \{\alpha\} \times^{\text{RLD}} b \mid \alpha \in \text{Src } f, b \in \text{atoms } G(\alpha) \}$$

and so f is complete. \square

Obvious 185. Complete and co-complete reloids are convex.

Obvious 186. Discrete reloids are complete and co-complete.

Obvious 187. Join (on the lattice of reloids) of complete reloids is complete.

Corollary 188. ComplRLD (with the induced order) is a complete lattice.

Theorem 189. A reloid which is both complete and co-complete is discrete.

Proof. Let f is a complete and co-complete reloid. We have

$$f = \bigcup \{ \uparrow^{\text{Src } f} \{ \alpha \} \times^{\text{RLD}} G(\alpha) \mid \alpha \in \text{Src } f \} \quad \text{and} \quad f = \bigcup \{ H(\beta) \times^{\text{RLD}} \uparrow^{\text{Dst } f} \{ \beta \} \mid \beta \in \text{Dst } f \}$$

for some functions $G: \text{Src } f \rightarrow \mathfrak{F}(\text{Dst } f)$, $H: \text{Dst } f \rightarrow \mathfrak{F}(\text{Src } f)$. For every $\alpha \in \text{Src } f$ we have

$$\begin{aligned} G(\alpha) &= \\ \text{im } f \upharpoonright_{\uparrow^{\text{Src } f} \{ \alpha \}} &= \\ \text{im} (f \cap (\uparrow^{\text{Src } f} \{ \alpha \} \times^{\text{RLD}} 1_{\mathfrak{F}(\text{Dst } f)})) &= (*) \\ \text{im} \bigcup \{ (H(\beta) \times^{\text{RLD}} \uparrow^{\text{Dst } f} \{ \beta \}) \cap (\uparrow^{\text{Src } f} \{ \alpha \} \times^{\text{RLD}} 1_{\mathfrak{F}(\text{Dst } f)}) \mid \beta \in \text{Dst } f \} &= \\ \text{im} \bigcup \{ (H(\beta) \cap \uparrow^{\text{Src } f} \{ \alpha \}) \times^{\text{RLD}} \uparrow^{\text{Dst } f} \{ \beta \} \mid \beta \in \text{Dst } f \} &= \\ \text{im} \bigcup \left\{ \left(\begin{array}{ll} \uparrow^{\text{Src } f} \{ \alpha \} \times \uparrow^{\text{Dst } f} \{ \beta \} & \text{if } H(\beta) \not\prec \uparrow^{\text{Src } f} \{ \alpha \} \\ 0_{\text{RLD}(\text{Src } f; \text{Dst } f)} & \text{if } H(\beta) \prec \uparrow^{\text{Src } f} \{ \alpha \} \end{array} \right) \mid \beta \in \text{Dst } f \right\} &= \\ \text{im} \bigcup \{ \uparrow^{\text{Src } f} \{ \alpha \} \times \uparrow^{\text{Dst } f} \{ \beta \} \mid \beta \in \text{Dst } f, H(\beta) \not\prec \uparrow^{\text{Src } f} \{ \alpha \} \} &= \\ \text{im} \bigcup \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} \{ (\alpha; \beta) \} \mid \beta \in \text{Dst } f, H(\beta) \not\prec \uparrow^{\text{Src } f} \{ \alpha \} \} &= \\ \bigcup \{ \uparrow^{\text{Dst } f} \{ \beta \} \mid \beta \in \text{Dst } f, H(\beta) \not\prec \uparrow^{\text{Src } f} \{ \alpha \} \}. & \end{aligned}$$

* the theorem 40 from [14] is used.

Thus $G(\alpha)$ is a principal f.o. that is $G(\alpha) = \uparrow^{\text{Dst } f} g(\alpha)$ for some $g: \text{Dst } f \rightarrow \text{Src } f$; $\uparrow^{\text{Src } f} \{ \alpha \} \times^{\text{RLD}} G(\alpha) = \uparrow^{\text{Src } f} (\{ \alpha \} \times g(\alpha))$; f is discrete as a join of discrete reloids. \square

Conjecture 190. Composition of complete reloids is complete.

Theorem 191.

1. For a complete reloid f there exist exactly one function $F \in \mathfrak{F}(\text{Dst } f)^{\text{Src } f}$ such that

$$f = \bigcup \{ \uparrow^{\text{Src } f} \{ \alpha \} \times^{\text{RLD}} F(\alpha) \mid \alpha \in \text{Src } f \}.$$

2. For a co-complete reloid f there exist exactly one function $F \in \mathfrak{F}(\text{Src } f)^{\text{Dst } f}$ such that

$$f = \bigcup \{ F(\alpha) \times^{\text{RLD}} \uparrow^{\text{Dst } f} \{ \alpha \} \mid \alpha \in \text{Dst } f \}.$$

Proof. We will prove only the first as the second is similar. Let

$$f = \bigcup \{ \uparrow^{\text{Src } f} \{ \alpha \} \times^{\text{RLD}} F(\alpha) \mid \alpha \in \text{Src } f \} = \bigcup \{ \uparrow^{\text{Src } f} \{ \alpha \} \times^{\text{RLD}} G(\alpha) \mid \alpha \in \text{Src } f \}$$

for some $F, G \in \mathfrak{F}(\text{Dst } f)^{\text{Src } f}$. We need to prove $F = G$. Let $\beta \in \text{Src } f$.

$$\begin{aligned} f \cap (\uparrow^{\text{Src } f} \{ \beta \} \times^{\text{RLD}} 1_{\mathfrak{F}(\text{Dst } f)}) &= (\text{theorem 40 in [14]}) \\ \bigcup^{\text{RLD}} \{ (\uparrow^{\text{Src } f} \{ \alpha \} \times^{\text{RLD}} F(\alpha)) \cap (\uparrow^{\text{Src } f} \{ \beta \} \times^{\text{RLD}} 1_{\mathfrak{F}(\text{Dst } f)}) \mid \alpha \in \text{Src } f \} &= \\ \uparrow^{\text{Src } f} \{ \beta \} \times^{\text{RLD}} F(\beta). & \end{aligned}$$

Similarly $f \cap (\uparrow^{\text{Src } f} \{ \beta \} \times^{\text{RLD}} 1_{\mathfrak{F}(\text{Dst } f)}) = \uparrow^{\text{Src } f} \{ \beta \} \times^{\text{RLD}} G(\beta)$. Thus $\uparrow^{\text{Src } f} \{ \beta \} \times^{\text{RLD}} F(\beta) = \uparrow^{\text{Src } f} \{ \beta \} \times^{\text{RLD}} G(\beta)$ and so $F(\beta) = G(\beta)$. \square

Definition 192. *Completion* and *co-completion* of a reloid $f \in \text{RLD}(A; B)$ are defined by the formulas:

$$\text{Compl } f = \text{Cor}^{(\text{RLD}(A; B); \text{ComplRLD}(A; B))} f \quad \text{and} \quad \text{CoCompl } f = \text{Cor}^{(\text{RLD}(A; B); \text{CoComplRLD}(A; B))} f.$$

Theorem 193. Atoms of the lattice $\text{ComplRLD}(A; B)$ are exactly direct products of the form $\uparrow^A\{\alpha\} \times^{\text{RLD}} b$ where $\alpha \in A$ and b is an atomic f.o. on B .

Proof. First, it's easy to see that $\uparrow^A\{\alpha\} \times^{\text{FCD}} b$ are elements of $\text{ComplRLD}(A; B)$. Also $0^{\text{RLD}(A; B)}$ is an element of ComplRLD .

$\uparrow^A\{\alpha\} \times^{\text{RLD}} b$ are atoms of ComplFCD because these are atoms of RLD .

It remains to prove that if f is an atom of $\text{ComplRLD}(A; B)$ then $f = \uparrow^A\{\alpha\} \times^{\text{RLD}} b$ for some $\alpha \in A$ and an atomic f.o. b on B .

Suppose f is a non-empty complete reloid. Then $\uparrow^A\{\alpha\} \times^{\text{RLD}} b \subseteq f$ for some $\alpha \in A$ and atomic f.o. b on B . If f is an atom then $f = \uparrow^A\{\alpha\} \times^{\text{FCD}} b$. \square

Obvious 194. ComplRLD is an atomistic lattice.

Proposition 195. $\text{Compl } f = \bigcup \{f|_{\uparrow^{\text{Src } f}\{\alpha\}} \mid \alpha \in \text{Src } f\}$ for every reloid f .

Proof. Let's denote R the right part of the equality to be proven.

That R is a complete reloid follows from the equality

$$f|_{\uparrow^{\text{Src } f}\{\alpha\}} = \uparrow^{\text{Src } f}\{\alpha\} \times^{\text{RLD}} \text{im}(f|_{\uparrow^{\text{Src } f}\{\alpha\}}).$$

The only thing left to prove is that $g \subseteq R$ for every complete reloid g such that $g \subseteq f$.

Really let g is a complete reloid such that $g \subseteq f$. Then

$$g = \bigcup \{\uparrow^{\text{Src } f}\{\alpha\} \times^{\text{RLD}} G(\alpha) \mid \alpha \in \text{Src } f\}$$

for some function $G: \text{Src } f \rightarrow \mathfrak{F}(\text{Dst } f)$.

We have $\uparrow^{\text{Src } f}\{\alpha\} \times^{\text{RLD}} G(\alpha) = g|_{\uparrow^{\text{Src } f}\{\alpha\}} \subseteq f|_{\uparrow^{\text{Src } f}\{\alpha\}}$. Thus $g \subseteq R$. \square

Conjecture 196. $\text{Compl } f \cap \text{Compl } g = \text{Compl}(f \cap g)$ for every reloids f and g .

Theorem 197. $\text{Compl}(\bigcup R) = \bigcup \langle \text{Compl} \rangle R$ for every set $R \in \text{RLD}(A; B)$ for every small sets A, B .

Proof.

$$\begin{aligned} \text{Compl}(\bigcup R) &= \\ \bigcup \{(\bigcup R)|_{\uparrow^A\{\alpha\}} \mid \alpha \in A\} &= \text{(theorem 40 in [14])} \\ \bigcup \{\bigcup \{f|_{\uparrow^A\{\alpha\}} \mid \alpha \in A\} \mid f \in R\} &= \\ \bigcup \langle \text{Compl} \rangle R. & \end{aligned}$$

\square

Lemma 198. Completion of a co-complete reloid is discrete.

Proof. Let f is a co-complete reloid. Then there is a function $F: \text{Dst } f \rightarrow \mathfrak{F}(\text{Src } f)$ such that

$$f = \bigcup \{F(\alpha) \times^{\text{RLD}} \uparrow^{\text{Dst } f}\{\alpha\} \mid \alpha \in \text{Dst } f\}.$$

So

$$\begin{aligned} \text{Compl } f &= \\ \bigcup \{(\bigcup^{\text{RLD}} \{F(\alpha) \times^{\text{RLD}} \uparrow^{\text{Dst } f}\{\alpha\} \mid \alpha \in \text{Dst } f\})|_{\uparrow^{\text{Src } f}\{\beta\}} \mid \beta \in \text{Src } f\} &= \\ \bigcup \{(\bigcup \{F(\alpha) \times^{\text{RLD}} \uparrow^{\text{Dst } f}\{\alpha\} \mid \alpha \in \text{Dst } f\}) \cap (\uparrow^{\text{Src } f}\{\beta\} \times 1^{\mathfrak{F}(\text{Dst } f)}) \mid \beta \in \text{Src } f\} &= (*) \\ \bigcup \{ \bigcup \{ (F(\alpha) \times^{\text{RLD}} \uparrow^{\text{Dst } f}\{\alpha\}) \cap (\uparrow^{\text{Src } f}\{\beta\} \times 1^{\mathfrak{F}(\text{Dst } f)}) \mid \alpha \in \text{Dst } f\} \mid \beta \in \text{Src } f\} &= \\ \bigcup \{ \bigcup \{ \uparrow^{\text{Src } f}\{\beta\} \times^{\text{RLD}} \uparrow^{\text{Dst } f}\{\alpha\} \mid \alpha \in \text{Dst } f\} \mid \beta \in \text{Src } f, \{\beta\} \subseteq F(\alpha)\} &\in \\ \langle \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} \rangle \mathcal{P}(\text{Src } f \times \text{Dst } f). & \end{aligned}$$

* theorem 40 in [14]. \square

Theorem 199. $\text{Compl } \text{CoCompl } f = \text{CoCompl } \text{Compl } f = \text{Cor } f$ for every reloid f .

Proof. We will prove only $\text{Compl CoCompl } f = \text{Cor } f$. The rest follows from symmetry.

From the lemma $\text{Compl CoCompl } f$ is discrete. It is obvious $\text{Compl CoCompl } f \subseteq f$. So to finish the proof we need to show only that for every discrete reloid $F \subseteq f$ we have $F \subseteq \text{Compl CoCompl } f$.

Really, obviously $F \subseteq \text{CoCompl } f$ and thus $F = \text{Compl } F \subseteq \text{Compl CoCompl } f$. \square

Question 200. Is $\text{ComplRLD}(A; B)$ a distributive lattice? Is $\text{ComplRLD}(A; B)$ a co-brouwerian lattice?

Conjecture 201. Let A, B, C are small sets. If $f \in \text{RLD}(B; C)$ is a complete reloid and $R \in \mathcal{P}\text{RLD}(A; B)$ then

$$f \circ \bigcup R = \bigcup \langle f \circ \rangle R.$$

This conjecture can be weakened:

Conjecture 202. Let A, B, C are small sets. If $f \in \text{RLD}(B; C)$ is a discrete reloid and $R \in \mathcal{P}\text{RLD}(A; B)$ then

$$f \circ \bigcup R = \bigcup \langle f \circ \rangle R.$$

Conjecture 203. $\text{Compl } f = f \setminus *_{\text{RLD}(\text{Src } f; \text{Dst } f)} (\Omega^{\text{Src } f} \times^{\text{RLD}} 1^{\mathfrak{F}(\text{Dst } f)})$ for every reloid f .

5 Relationships between funcoids and reloids

5.1 Funcoid induced by a reloid

Every reloid f induces a funcoid $(\text{FCD})f \in \text{FCD}(\text{Src } f; \text{Dst } f)$ by the following formulas (for every $\mathcal{X} \in \mathfrak{F}(\text{Src } f)$, $\mathcal{Y} \in \mathfrak{F}(\text{Dst } f)$):

$$\begin{aligned} \mathcal{X}[(\text{FCD})f]\mathcal{Y} &\Leftrightarrow \forall F \in \text{up } f: \mathcal{X}[\uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F]\mathcal{Y} \\ \langle (\text{FCD})f \rangle \mathcal{X} &= \bigcap \{ \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \rangle \mathcal{X} \mid F \in \text{up } f \}. \end{aligned}$$

We should prove that $(\text{FCD})f$ is really a funcoid.

Proof. We need to prove that

$$\mathcal{X}[(\text{FCD})f]\mathcal{Y} \Leftrightarrow \mathcal{Y} \cap \langle (\text{FCD})f \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \mathcal{X} \cap \langle (\text{FCD})f^{-1} \rangle \mathcal{Y} \neq 0^{\mathfrak{F}(\text{Dst } f)}.$$

The above formula is equivalent to:

$$\begin{aligned} \forall F \in \text{up } f: \mathcal{X}[\uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F]\mathcal{Y} &\Leftrightarrow \\ \mathcal{Y} \cap \bigcap \{ \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \rangle \mathcal{X} \mid F \in \text{up } f \} &\neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow \\ \mathcal{X} \cap \bigcap \{ \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F^{-1} \rangle \mathcal{Y} \mid F \in \text{up } f \} &\neq 0^{\mathfrak{F}(\text{Src } f)} \end{aligned}$$

We have $\mathcal{Y} \cap \bigcap \{ \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \rangle \mathcal{X} \mid F \in \text{up } f \} = \bigcap \{ \mathcal{Y} \cap \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \rangle \mathcal{X} \mid F \in \text{up } f \}$.

Let's denote $W = \{ \mathcal{Y} \cap \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \rangle \mathcal{X} \mid F \in \text{up } f \}$.

$\forall F \in \text{up } f: \mathcal{X}[\uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F]\mathcal{Y} \Leftrightarrow \forall F \in \text{up } f: \mathcal{Y} \cap \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \rangle \mathcal{X} \neq 0^{\mathfrak{F}(\text{Dst } f)} \Leftrightarrow 0^{\mathfrak{F}(\text{Dst } f)} \notin W$.

We need to prove that $0^{\mathfrak{F}(\text{Dst } f)} \notin W \Leftrightarrow \bigcap W \neq 0^{\mathfrak{F}(\text{Dst } f)}$. (The rest follows from symmetry.)

This follows from the fact that W is a generalized filter base.

Let's prove that W is a generalized filter base. For this enough to prove that $V = \{ \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} F \rangle \mathcal{X} \mid F \in \text{up } f \}$ is a generalized filter base. Let $\mathcal{A}, \mathcal{B} \in V$ that is $\mathcal{A} = \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} P \rangle \mathcal{X}$, $\mathcal{B} = \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} Q \rangle \mathcal{X}$ where $P, Q \in \text{up } f$. Then for $\mathcal{C} = \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} (P \cap Q) \rangle \mathcal{X}$ is true both $\mathcal{C} \in V$ and $\mathcal{C} \subseteq \mathcal{A}, \mathcal{B}$. So V is a generalized filter base and thus W is a generalized filter base. \square

Proposition 204. $(\text{FCD})\uparrow^{\text{RLD}(A;B)}f = \uparrow^{\text{FCD}(A;B)}f$ for every small sets A, B and binary relation $f \subseteq A \times B$.

Proof. $\mathcal{X}[(\text{FCD})\uparrow^{\text{RLD}(A;B)}f]\mathcal{Y} \Leftrightarrow \forall F \in \text{up } \uparrow^{\text{RLD}(A;B)}f: \mathcal{X}[\uparrow^{\text{FCD}(A;B)}F]\mathcal{Y} \Leftrightarrow \mathcal{X}[\uparrow^{\text{FCD}(A;B)}f]\mathcal{Y}$ (for every $\mathcal{X}, \mathcal{Y} \in \mathfrak{F}$). \square

Theorem 205. $\mathcal{X}[(\text{FCD})f]\mathcal{Y} \Leftrightarrow (\mathcal{X} \times^{\text{RLD}}\mathcal{Y}) \not\prec f$ for every $f \in \text{RLD}$ and $\mathcal{X} \in \mathfrak{F}(\text{Src } f), \mathcal{Y} \in \mathfrak{F}(\text{Dst } f)$.

Proof.

$$\begin{aligned} (\mathcal{X} \times^{\text{RLD}}\mathcal{Y}) \not\prec f &\Leftrightarrow \forall F \in \text{up } f, P \in \text{up } (\mathcal{X} \times^{\text{RLD}}\mathcal{Y}): P \not\prec F \\ &\Leftrightarrow \forall F \in \text{up } f, X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: (X \times Y) \not\prec F \\ &\Leftrightarrow \forall F \in \text{up } f, X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y}: \uparrow^{\text{Src } f}X[\uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)}F]\uparrow^{\text{Dst } f}Y \\ &\Leftrightarrow \forall F \in \text{up } f: \mathcal{X}[\uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)}F]\mathcal{Y} \\ &\Leftrightarrow \mathcal{X}[(\text{FCD})f]\mathcal{Y}. \end{aligned}$$

\square

Theorem 206. $(\text{FCD})f = \bigcap \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} \rangle \text{up } f$ for every reloid f .

Proof. Let a is an atomic filter object.

$$\begin{aligned} \langle (\text{FCD})f \rangle a &= \bigcap \{ \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)}F \rangle a \mid F \in \text{up } f \} \text{ by the definition of (FCD).} \\ \langle \bigcap \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} \rangle \text{up } f \rangle a &= \bigcap \{ \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)}F \rangle a \mid F \in \text{up } f \} \text{ by the theorem 78.} \\ \text{So } \langle (\text{FCD})f \rangle a &= \langle \bigcap \langle \uparrow^{\text{FCD}(\text{Src } f; \text{Dst } f)} \rangle \text{up } f \rangle a \text{ for every atomic filter object } a. \end{aligned}$$

\square

Lemma 207. For every two filter bases S and T of binary relations on $U \times V$ for some small sets U, V and every set $A \subseteq U$

$$\bigcap \uparrow^{\text{RLD}(U;V)}S = \bigcap \uparrow^{\text{RLD}(U;V)}T \Rightarrow \bigcap \{ \uparrow^V \langle F \rangle A \mid F \in S \} = \bigcap \{ \uparrow^V \langle G \rangle A \mid G \in T \}$$

Proof. Let $\bigcap \uparrow^{\text{RLD}(U;V)}S = \bigcap \uparrow^{\text{RLD}(U;V)}T$.

First let prove that $\{ \langle F \rangle A \mid F \in S \}$ is a filter base. Let $X, Y \in \{ \langle F \rangle A \mid F \in S \}$. Then $X = \langle F_X \rangle A$ and $Y = \langle F_Y \rangle A$ for some $F_X, F_Y \in S$. Because S is a filter base, we have $S \ni F_Z \subseteq F_X \cap F_Y$. So $\langle F_Z \rangle A \subseteq X \cap Y$ and $\langle F_Z \rangle A \in \{ \langle F \rangle A \mid F \in S \}$. So $\{ \langle F \rangle A \mid F \in S \}$ is a filter base.

Suppose $X \in \text{up } \bigcap \{ \uparrow^V \langle F \rangle A \mid F \in S \}$. Then exists $X' \in \{ \langle F \rangle A \mid F \in S \}$ where $X \supseteq X'$ because $\{ \langle F \rangle A \mid F \in S \}$ is a filter base. That is $X' = \langle F \rangle A$ for some $F \in S$. There exists $G \in T$ such that $G \subseteq F$ because T is a filter base. Let $Y' = \langle G \rangle A$. We have $Y' \subseteq X' \subseteq X$; $Y' \in \{ \langle G \rangle A \mid G \in T \}$; $Y' \in \text{up } \bigcap \{ \uparrow^V \langle G \rangle A \mid G \in T \}$; $X \in \text{up } \bigcap \{ \uparrow^V \langle G \rangle A \mid G \in T \}$. The reverse is symmetric. \square

Lemma 208. $\{ G \circ F \mid F \in \text{up } f, G \in \text{up } g \}$ is a filter base for every reloids f and g .

Proof. Let denote $D = \{ G \circ F \mid F \in \text{up } f, G \in \text{up } g \}$. Let $A \in D \wedge B \in D$. Then $A = G_A \circ F_A \wedge B = G_B \circ F_B$ for some $F_A, F_B \in \text{up } f$ and $G_A, G_B \in \text{up } g$. So $A \cap B \supseteq (G_A \cap G_B) \circ (F_A \cap F_B) \in D$ because $F_A \cap F_B \in \text{up } f$ and $G_A \cap G_B \in \text{up } g$. \square

Theorem 209. $(\text{FCD})(g \circ f) = ((\text{FCD})g) \circ ((\text{FCD})f)$ for every composable reloids f and g .

Proof.

$$\begin{aligned} \langle (\text{FCD})(g \circ f) \rangle X &= \bigcap \{ \uparrow^{\text{Dst } g} \langle H \rangle X \mid H \in \text{up } (g \circ f) \} \\ &= \bigcap \{ \uparrow^{\text{Dst } g} \langle H \rangle X \mid H \in \text{up } \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } g)}(G \circ F) \mid F \in \text{up } f, G \in \text{up } g \} \}. \end{aligned}$$

Obviously

$$\bigcap \langle \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } g)} \rangle \text{up } \bigcap \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } g)}(G \circ F) \mid F \in \text{up } f, G \in \text{up } g \} =$$

from this by the lemma 207 (taking in account that $\{G \circ F \mid F \in \text{up } f, G \in \text{up } g\}$ and $\text{up } \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Dst } g)}(G \circ F) \mid F \in \text{up } f, G \in \text{up } g\}$ are filter bases)

$$\bigcap \{\uparrow^{\text{Dst } g} \langle H \rangle X \mid H \in \text{up } \bigcap \{\uparrow^{\text{RLD}(\text{Src } f; \text{Dst } g)}(G \circ F) \mid F \in \text{up } f, G \in \text{up } g\}\} = \bigcap \{\uparrow^{\text{Dst } g} \langle G \circ F \rangle X \mid F \in \text{up } f, G \in \text{up } g\}.$$

On the other side

$$\begin{aligned} \langle ((\text{FCD})g) \circ ((\text{FCD})f) \rangle^* X &= \langle (\text{FCD})g \rangle \langle (\text{FCD})f \rangle^* X \\ &= \langle (\text{FCD})g \rangle \bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid F \in \text{up } f\} \\ &= \bigcap \{\langle \uparrow^{\text{FCD}(\text{Src } g; \text{Dst } g)} G \rangle \bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid F \in \text{up } f\} \mid G \in \text{up } g\}. \end{aligned}$$

Let's prove that $\{\langle F \rangle X \mid F \in \text{up } f\}$ is a filter base. If $A, B \in \{\langle F \rangle X \mid F \in \text{up } f\}$ then $A = \langle F_1 \rangle X$ and $B = \langle F_2 \rangle X$ where $F_1, F_2 \in \text{up } f$. $A \cap B \supseteq \langle F_1 \cap F_2 \rangle X \in \{\langle F \rangle X \mid F \in \text{up } f\}$. So $\{\langle F \rangle X \mid F \in \text{up } f\}$ is really a filter base.

By the theorem 51 we have

$$\langle \uparrow^{\text{FCD}(\text{Src } g; \text{Dst } g)} G \rangle \bigcap \{\uparrow^{\text{Dst } f} \langle F \rangle X \mid F \in \text{up } f\} = \bigcap \{\uparrow^{\text{Dst } g} \langle G \rangle \langle F \rangle X \mid F \in \text{up } f\}.$$

So continuing the above equalities,

$$\begin{aligned} \langle ((\text{FCD})g) \circ ((\text{FCD})f) \rangle^* X &= \bigcap \left\{ \bigcap \{\uparrow^{\text{Dst } g} \langle G \rangle \langle F \rangle X \mid F \in \text{up } f\} \mid G \in \text{up } g \right\} \\ &= \bigcap \{\uparrow^{\text{Dst } g} \langle G \rangle \langle F \rangle X \mid F \in \text{up } f, G \in \text{up } g\} \\ &= \bigcap \{\uparrow^{\text{Dst } g} \langle G \circ F \rangle X \mid F \in \text{up } f, G \in \text{up } g\}. \end{aligned}$$

Combining these equalities we get $\langle (\text{FCD})(g \circ f) \rangle^* X = \langle ((\text{FCD})g) \circ ((\text{FCD})f) \rangle^* X$ for every set X . \square

Corollary 210.

1. $(\text{FCD})f$ is a monovalued funcoid if f is a monovalued reloid.
2. $(\text{FCD})f$ is an injective funcoid if f is an injective reloid.

Proof. We will prove only the first as the second is dual. Let f is a monovalued reloid. Then $f \circ f^{-1} \subseteq I^{\text{RLD}(\text{Dst } f)}$; $(\text{FCD})(f \circ f^{-1}) \subseteq I^{\text{FCD}(\text{Dst } f)}$; $(\text{FCD})f \circ ((\text{FCD})f)^{-1} \subseteq I^{\text{FCD}(\text{Dst } f)}$ that is $(\text{FCD})f$ is a monovalued funcoid. \square

Proposition 211. $(\text{FCD})I_{\mathcal{A}}^{\text{RLD}} = I_{\mathcal{A}}^{\text{FCD}}$ for every f.o. \mathcal{A} .

Proof. Recall that $I_{\mathcal{A}}^{\text{RLD}} = \bigcap \{\uparrow^{\text{Base}(\mathcal{A})} I_A \mid A \in \text{up } \mathcal{A}\}$. For every $\mathcal{X}, \mathcal{Y} \in \mathfrak{F}(\text{Base}(\mathcal{A}))$ we have:

$\mathcal{X}[(\text{FCD})I_{\mathcal{A}}^{\text{RLD}}] \mathcal{Y} \Leftrightarrow \mathcal{X} \times^{\text{RLD}} \mathcal{Y} \not\neq I_{\mathcal{A}}^{\text{RLD}} \Leftrightarrow \forall A \in \text{up } \mathcal{A}: \mathcal{X} \times^{\text{RLD}} \mathcal{Y} \not\neq \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} I_A \Leftrightarrow \forall A \in \text{up } \mathcal{A}: \mathcal{X}[\uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{A}))} I_A] \mathcal{Y} \Leftrightarrow \forall A \in \text{up } \mathcal{A}: \mathcal{X} \cap \mathcal{Y} \not\neq \uparrow^{\text{Base}(\mathcal{A})} A \Leftrightarrow \mathcal{X} \cap \mathcal{Y} \not\neq \mathcal{A} \Leftrightarrow \mathcal{X}[(\text{FCD})I_{\mathcal{A}}^{\text{FCD}}] \mathcal{Y}$ (used properties of generalized filter bases). \square

Proposition 212. $(\text{FCD})(\mathcal{A} \times^{\text{RLD}} \mathcal{B}) = \mathcal{A} \times^{\text{FCD}} \mathcal{B}$ for every f.o. \mathcal{A} and \mathcal{B} .

Proof. $\mathcal{X}[(\text{FCD})(\mathcal{A} \times^{\text{RLD}} \mathcal{B})] \mathcal{Y} \Leftrightarrow \forall F \in \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{B}): \mathcal{X}[\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} F] \mathcal{Y}$ (for every $\mathcal{X}, \mathcal{Y} \in \mathfrak{F}$).

Evidently

$\forall F \in \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{B}): \mathcal{X}[\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} F] \mathcal{Y} \Rightarrow \forall A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B}: \mathcal{X}[\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} (A \times B)] \mathcal{Y}$.

Let $\forall A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B}: \mathcal{X}[\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} (A \times B)] \mathcal{Y}$. Then if $F \in \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{B})$ then there are $A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B}$ such that $F \supseteq A \times B$. So $\mathcal{X}[\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} F] \mathcal{Y}$.

We proved $\forall F \in \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{B}): \mathcal{X}[\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} F] \mathcal{Y} \Leftrightarrow \forall A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B}: \mathcal{X}[\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} (A \times B)] \mathcal{Y}$.

Further $\forall A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B}: \mathcal{X}[\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} (A \times B)] \mathcal{Y} \Leftrightarrow \forall A \in \text{up } \mathcal{A}, B \in \text{up } \mathcal{B}: \mathcal{X} \not\neq \uparrow^{\text{Base}(\mathcal{A})} A \wedge \mathcal{Y} \not\neq \uparrow^{\text{Base}(\mathcal{B})} B \Leftrightarrow \mathcal{X} \not\neq \mathcal{A} \wedge \mathcal{Y} \not\neq \mathcal{B} \Leftrightarrow \mathcal{X}[\mathcal{A} \times^{\text{FCD}} \mathcal{B}] \mathcal{Y}$.

Thus $\mathcal{X}[(\text{FCD})(\mathcal{A} \times^{\text{RLD}} \mathcal{B})] \mathcal{Y} \Leftrightarrow \mathcal{X}[\mathcal{A} \times^{\text{FCD}} \mathcal{B}] \mathcal{Y}$. \square

Proposition 213. $\text{dom}(\text{FCD})f = \text{dom} f$ and $\text{im}(\text{FCD})f = \text{im} f$ for every reloid f .

Proof. $\text{im}(\text{FCD})f = \langle (\text{FCD})f \rangle 1^{\mathfrak{F}(\text{Src } f)} = \bigcap \{ \uparrow^{\text{Dst } f} \langle F \rangle (\text{Src } f) \mid F \in \text{up } f \} = \bigcap \{ \uparrow^{\text{Dst } f} \text{im } F \mid F \in \text{up } f \} = \bigcap \langle \uparrow^{\text{Dst } f} \rangle \langle \text{im} \rangle \text{up } f = \text{im } f$.
 $\text{dom}(\text{FCD})f = \text{dom} f$ is similar. \square

Proposition 214. $(\text{FCD})(f \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{B})) = (\text{FCD})f \cap (\mathcal{A} \times^{\text{FCD}} \mathcal{B})$ for every reloid f and $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$ and $\mathcal{B} \in \mathfrak{F}(\text{Dst } f)$.

Proof. $(\text{FCD})(f \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{B})) = (\text{FCD})(I_{\mathcal{B}}^{\text{RLD}} \circ f \circ I_{\mathcal{A}}^{\text{RLD}}) = (\text{FCD})I_{\mathcal{B}}^{\text{RLD}} \circ (\text{FCD})f \circ (\text{FCD})I_{\mathcal{A}}^{\text{RLD}} = I_{\mathcal{B}}^{\text{FCD}} \circ (\text{FCD})f \circ I_{\mathcal{A}}^{\text{FCD}} = (\text{FCD})f \cap (\mathcal{A} \times^{\text{FCD}} \mathcal{B})$. \square

Corollary 215. $(\text{FCD})(f|_{\mathcal{A}}) = ((\text{FCD})f)|_{\mathcal{A}}$ for every reloid f and f.o. \mathcal{A} .

Proposition 216. $\langle (\text{FCD})f \rangle \mathcal{X} = \text{im}(f|_{\mathcal{X}})$ for every reloid f and f.o. \mathcal{X} .

Proof. $\text{im}(f|_{\mathcal{X}}) = \text{im}(\text{FCD})(f|_{\mathcal{X}}) = \text{im}((\text{FCD})f|_{\mathcal{X}}) = \langle (\text{FCD})f \rangle \mathcal{X}$. \square

5.2 Reloids induced by funcoid

Every funcoid $f \in \text{FCD}(A; B)$ induces a reloid from A to B in two ways, intersection of *outward* relations and union of *inward* direct products of filter objects:

$$\begin{aligned} (\text{RLD})_{\text{out}} f &\stackrel{\text{def}}{=} \bigcap \langle \uparrow^{\text{RLD}(A;B)} \rangle \text{up } f; \\ (\text{RLD})_{\text{in}} f &\stackrel{\text{def}}{=} \bigcup \{ \mathcal{A} \times^{\text{RLD}} \mathcal{B} \mid \mathcal{A} \in \mathfrak{F}(A), \mathcal{B} \in \mathfrak{F}(B), \mathcal{A} \times^{\text{FCD}} \mathcal{B} \subseteq f \} \end{aligned}$$

Theorem 217. $(\text{RLD})_{\text{in}} f = \bigcup^{\text{RLD}} \{ a \times^{\text{RLD}} b \mid a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)}, a \times^{\text{FCD}} b \subseteq f \}$.

Proof. Follows from the theorem 160. \square

Remark 218. It seems that $(\text{RLD})_{\text{in}}$ has smoother properties and is more important than $(\text{RLD})_{\text{out}}$. (However see also the exercise below for $(\text{RLD})_{\text{in}}$ not preserving identities.)

Proposition 219. $(\text{RLD})_{\text{out}} \uparrow^{\text{FCD}(A;B)} f = \uparrow^{\text{RLD}(A;B)} f$ for every small sets A, B and binary relation $f \subseteq A \times B$.

Proof. $(\text{RLD})_{\text{out}} \uparrow^{\text{FCD}(A;B)} f = \bigcap \langle \uparrow^{\text{RLD}(A;B)} \rangle \text{up } \uparrow^{\text{FCD}(A;B)} f = \uparrow^{\text{RLD}(A;B)} \text{min up } \uparrow^{\text{FCD}(A;B)} f = \uparrow^{\text{RLD}(A;B)} f$. \square

Lemma 220. $F \in \text{up}(\text{RLD})_{\text{in}} f \Leftrightarrow \forall a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} : (a[f]b \Rightarrow \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} F \supseteq a \times^{\text{RLD}} b)$ for a funcoid f .

Proof.

$$\begin{aligned} F \in \text{up}(\text{RLD})_{\text{in}} f &\Leftrightarrow F \in \text{up} \bigcup \{ a \times^{\text{RLD}} b \mid a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)}, a \times^{\text{FCD}} b \subseteq f \} \\ &\Leftrightarrow \forall a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} : (a \times^{\text{FCD}} b \subseteq f \Rightarrow F \in \text{up}(a \times^{\text{RLD}} b)) \\ &\Leftrightarrow \forall a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} : ((a \times^{\text{FCD}} b) \not\subseteq f \Rightarrow \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} F \not\supseteq a \times^{\text{RLD}} b) \\ &\Leftrightarrow \forall a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)} : (a[f]b \Rightarrow \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} F \supseteq a \times^{\text{RLD}} b). \end{aligned} \quad \square$$

Surprisingly a funcoid is greater inward than outward:

Theorem 221. $(\text{RLD})_{\text{out}} f \subseteq (\text{RLD})_{\text{in}} f$ for every funcoid f .

Proof. We need to prove

$$\bigcap \langle \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} \rangle_{\text{up}} f \subseteq \bigcup \{ \mathcal{A} \times^{\text{RLD}} \mathcal{B} \mid \mathcal{A}, \mathcal{B} \in \mathfrak{F}, \mathcal{A} \times^{\text{FCD}} \mathcal{B} \subseteq f \}.$$

Let

$$K \in \text{up} \bigcup \{ \mathcal{A} \times^{\text{RLD}} \mathcal{B} \mid \mathcal{A}, \mathcal{B} \in \mathfrak{F}, \mathcal{A} \times^{\text{FCD}} \mathcal{B} \subseteq f \}.$$

Then

$$\begin{aligned} K &= \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} \bigcup \{ X_{\mathcal{A}} \times Y_{\mathcal{B}} \mid \mathcal{A}, \mathcal{B} \in \mathfrak{F}, \mathcal{A} \times^{\text{FCD}} \mathcal{B} \subseteq f \} \\ &= \bigcup \{ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(X_{\mathcal{A}} \times Y_{\mathcal{B}}) \mid \mathcal{A}, \mathcal{B} \in \mathfrak{F}, \mathcal{A} \times^{\text{FCD}} \mathcal{B} \subseteq f \} \\ &\supseteq f \end{aligned}$$

where $X_{\mathcal{A}} \in \text{up } \mathcal{A}$, $Y_{\mathcal{B}} \in \text{up } \mathcal{B}$. So $K \in \text{up } f$; $K \in \text{up} \bigcap \langle \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)} \rangle_{\text{up}} f$. \square

Theorem 222. $(\text{FCD})(\text{RLD})_{\text{in}} f = f$ for every funcoid f .

Proof. For every sets $X \in \mathcal{P}(\text{Src } f)$ and $Y \in \mathcal{P}(\text{Dst } f)$

$$\begin{aligned} X[(\text{FCD})(\text{RLD})_{\text{in}} f]^* Y &\Leftrightarrow \\ (\uparrow^{\text{Src } f} X \times^{\text{RLD}} \uparrow^{\text{Dst } f} Y) \not\subseteq (\text{RLD})_{\text{in}} f &\Leftrightarrow \\ \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(X \times Y) \not\subseteq \bigcup \{ a \times^{\text{RLD}} b \mid a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)}, \\ a \times^{\text{FCD}} b \subseteq f \} &\Leftrightarrow (*) \\ \exists a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)}: (a \times^{\text{FCD}} b \subseteq f \wedge \uparrow^{\text{RLD}(\text{Src } f; \text{Dst } f)}(X \times Y) \not\subseteq (a \times^{\text{RLD}} b)) &\Leftrightarrow \\ \exists a \in \text{atoms } 1^{\mathfrak{F}(\text{Src } f)}, b \in \text{atoms } 1^{\mathfrak{F}(\text{Dst } f)}: (a[f]b \wedge a \subseteq \uparrow^{\text{Src } f} X \wedge b \subseteq \uparrow^{\text{Dst } f} Y) &\Leftrightarrow \\ X[f]^* Y. & \end{aligned}$$

* theorem 53 in [14].

Thus $(\text{FCD})(\text{RLD})_{\text{in}} f = f$. \square

Remark 223. The above theorem allows to represent funcoids as reloids.

Obvious 224. $(\text{RLD})_{\text{in}}(\mathcal{A} \times^{\text{FCD}} \mathcal{B}) = \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ for every f.o. \mathcal{A}, \mathcal{B} .

Conjecture 225. $(\text{RLD})_{\text{out}} I_{\mathcal{A}}^{\text{FCD}} = I_{\mathcal{A}}^{\text{RLD}}$ for every f.o. \mathcal{A} .

Exercise 1. Prove that generally $(\text{RLD})_{\text{in}} I_{\mathcal{A}}^{\text{FCD}} \neq I_{\mathcal{A}}^{\text{RLD}}$.

Conjecture 226. $\text{dom}(\text{RLD})_{\text{in}} f = \text{dom } f$ and $\text{im}(\text{RLD})_{\text{in}} f = \text{im } f$ for every funcoid f .

Proposition 227. $\text{dom}(f|_{\mathcal{A}}) = \mathcal{A} \cap \text{dom } f$ for every reloid f and f.o. $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$.

Proof. $\text{dom}(f|_{\mathcal{A}}) = \text{dom}(\text{FCD})f|_{\mathcal{A}} = \text{dom}((\text{FCD})f)|_{\mathcal{A}} = \mathcal{A} \cap \text{dom}(\text{FCD})f = \mathcal{A} \cap \text{dom } f$. \square

Theorem 228. For every reloids f, g :

1. If $\text{im } f \supseteq \text{dom } g$ then $\text{im}(g \circ f) = \text{im } g$.
2. If $\text{im } f \subseteq \text{dom } g$ then $\text{dom}(g \circ f) = \text{dom } f$.

Proof.

1. $\text{im}(g \circ f) = \text{im}(\text{FCD})(g \circ f) = \text{im}((\text{FCD})g \circ (\text{FCD})f) = \text{im}(\text{FCD})g = \text{im } g$.
2. Similar. \square

Corollary 229. $(\text{RLD})_{\text{in}}(f|_{\mathcal{A}}) = ((\text{RLD})_{\text{in}} f)|_{\mathcal{A}}$ for every funcoid f and f.o. \mathcal{A} .

5.3 Galois connections of funcoids and reloids

Theorem 230. $(\text{FCD}): \text{RLD}(A; B) \rightarrow \text{FCD}(A; B)$ is the lower adjoint of $(\text{RLD})_{\text{in}}: \text{FCD}(A; B) \rightarrow \text{RLD}(A; B)$ for every small sets A, B .

Proof. Because (FCD) and $(\text{RLD})_{\text{in}}$ are trivially monotone, it's enough to prove (for every $f \in \text{RLD}(A; B)$, $g \in \text{FCD}(A; B)$)

$$f \subseteq (\text{RLD})_{\text{in}}(\text{FCD})f \text{ and } (\text{FCD})(\text{RLD})_{\text{in}}g \subseteq g.$$

The second formula follows from the fact that $(\text{FCD})(\text{RLD})_{\text{in}}g = g$.

$$\begin{aligned} & (\text{RLD})_{\text{in}}(\text{FCD})f = \\ \bigcup & \{a \times^{\text{RLD}} b \mid a \in \text{atoms } 1^{\mathfrak{F}(A)}, b \in \text{atoms } 1^{\mathfrak{F}(B)}, a \times^{\text{FCD}} b \subseteq (\text{FCD})f\} = \\ & \bigcup \{a \times^{\text{RLD}} b \mid a \in \text{atoms } 1^{\mathfrak{F}(A)}, b \in \text{atoms } 1^{\mathfrak{F}(B)}, a[(\text{FCD})f]b\} = \\ & \bigcup \{a \times^{\text{RLD}} b \mid a \in \text{atoms } 1^{\mathfrak{F}(A)}, b \in \text{atoms } 1^{\mathfrak{F}(B)}, (a \times^{\text{RLD}} b) \not\subseteq f\} \supseteq \\ & \bigcup \{p \in \text{atoms}(a \times^{\text{RLD}} b) \mid a \in \text{atoms } 1^{\mathfrak{F}(A)}, b \in \text{atoms } 1^{\mathfrak{F}(B)}, p \not\subseteq f\} = \\ & \bigcup \{p \in \text{atoms } 1^{\mathfrak{F}(A \times B)} \mid p \not\subseteq f\} = \\ & \bigcup \{p \mid p \in \text{atoms } f\} = f. \end{aligned}$$

□

Corollary 231.

1. $(\text{FCD}) \cup S = \bigcup \langle (\text{FCD}) \rangle S$ if S is a set of reloids.
2. $(\text{RLD})_{\text{in}} \cap S = \bigcap \langle (\text{RLD})_{\text{in}} \rangle S$ if S is a set of functors.

Proposition 232. $(\text{RLD})_{\text{in}}(f \cap (\mathcal{A} \times^{\text{FCD}} \mathcal{B})) = ((\text{RLD})_{\text{in}}f) \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{B})$ for every functor f and f.o. $\mathcal{A} \in \mathfrak{F}(\text{Src } f)$ and $\mathcal{B} \in \mathfrak{F}(\text{Dst } f)$.

Proof. $(\text{RLD})_{\text{in}}(f \cap (\mathcal{A} \times^{\text{FCD}} \mathcal{B})) = ((\text{RLD})_{\text{in}}f) \cap (\text{RLD})_{\text{in}}(\mathcal{A} \times^{\text{FCD}} \mathcal{B}) = ((\text{RLD})_{\text{in}}f) \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{B})$. □

Conjecture 233. $(\text{RLD})_{\text{in}}$ is not a lower adjoint (in general).

Conjecture 234. $(\text{RLD})_{\text{out}}$ is neither a lower adjoint nor an upper adjoint (in general).

See also the corollary 296 below.

6 Continuous morphisms

This section uses the apparatus from the section “Partially ordered dagger categories”.

6.1 Traditional definitions of continuity

In this section we will show that having a functor or reloid $\uparrow f$ corresponding to a function f we can express continuity of it by the formula $\uparrow f \circ \mu \subseteq \nu \circ \uparrow f$ (or similar formulas) where μ and ν are some spaces.

6.1.1 Pre-topology

Let μ and ν be functors representing some pre-topologies. By definition a function f is continuous map from μ to ν in point a iff

$$\forall \epsilon \in \text{up} \langle \nu \rangle f a \exists \delta \in \text{up} \langle \mu \rangle^* \{a\}: \langle f \rangle \delta \subseteq \epsilon.$$

Equivalently transforming this formula we get:

$$\begin{aligned} & \forall \epsilon \in \text{up} \langle \nu \rangle f a: \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle \langle \mu \rangle \uparrow^{\text{Src } \mu} \{a\} \subseteq \epsilon; \\ & \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle \langle \mu \rangle \uparrow^{\text{Src } \mu} \{a\} \subseteq \langle \nu \rangle f a; \\ & \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle \langle \mu \rangle \uparrow^{\text{Src } \mu} \{a\} \subseteq \langle \nu \rangle \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle \uparrow^{\text{Src } \mu} \{a\}; \\ & \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \circ \mu \rangle \uparrow^{\text{Src } \mu} \{a\} \subseteq \langle \nu \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle \uparrow^{\text{Src } \mu} \{a\}. \end{aligned}$$

So f is a continuous map from μ to ν in every point of its domain iff

$$\uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \circ \mu \subseteq \nu \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f.$$

6.1.2 Proximity spaces

Let μ and ν are proximity (nearness) spaces (which I consider a special case of funcoids). By definition a function f is a proximity-continuous map (also called equicontinuous) from μ to ν iff

$$\forall X \in \mathcal{P}(\text{Src } \mu), Y \in \mathcal{P}(\text{Dst } \nu): (X[\mu]^* Y \Rightarrow fX[\nu]^* fY).$$

Equivalently transforming this formula we get:

$$\begin{aligned} \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle Y \cap \langle \nu \rangle \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle X \neq 0^{\mathfrak{F}(\text{Dst } \nu)}); \\ \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle Y \cap \langle \nu \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle X \neq 0^{\mathfrak{F}(\text{Dst } \nu)}); \\ \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow X[\nu \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f]^* \langle f \rangle Y); \\ \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow \langle f \rangle Y [(\nu \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f)^{-1}]^* X); \\ \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow \langle f \rangle Y [(\uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f)^{-1} \circ \nu^{-1}]^* X); \\ \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow \uparrow^{\mathfrak{F}(\text{Src } \mu)} X \cap \langle (\uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f)^{-1} \circ \nu^{-1} \rangle \langle \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle Y \neq 0^{\mathfrak{F}(\text{Src } \mu)}); \\ \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow \uparrow^{\mathfrak{F}(\text{Src } \mu)} X \cap \langle (\uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f)^{-1} \circ \nu^{-1} \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f \rangle Y \neq 0^{\mathfrak{F}(\text{Src } \mu)}); \\ \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow Y [(\uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f)^{-1} \circ \nu^{-1} \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f]^* X); \\ \forall X, Y \in \mathcal{P}\mathcal{U}: (X[\mu]^* Y \Rightarrow X [(\uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f)^{-1} \circ \nu \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f]^* Y); \\ \mu \subseteq (\uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f)^{-1} \circ \nu \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f. \end{aligned}$$

So a function f is proximity-continuous iff $\mu \subseteq (\uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f)^{-1} \circ \nu \circ \uparrow^{\text{FCD}(\text{Src } \mu; \text{Dst } \nu)} f$.

6.1.3 Uniform spaces

Uniform spaces are a special case of reloids.

Let μ and ν are uniform spaces. By definition a function f is a uniformly continuous map from μ to ν iff

$$\forall \epsilon \in \text{up } \nu \exists \delta \in \text{up } \mu \forall (x; y) \in \delta: (fx; fy) \in \epsilon.$$

Equivalently transforming this formula we get:

$$\begin{aligned} \forall \epsilon \in \text{up } \nu \exists \delta \in \text{up } \mu \forall (x; y) \in \delta: \{(fx; fy)\} \subseteq \epsilon \\ \forall \epsilon \in \text{up } \nu \exists \delta \in \text{up } \mu \forall (x; y) \in \delta: f \circ \{(x; y)\} \circ f^{-1} \subseteq \epsilon \\ \forall \epsilon \in \text{up } \nu \exists \delta \in \text{up } \mu: f \circ \delta \circ f^{-1} \subseteq \epsilon \\ \forall \epsilon \in \text{up } \nu: \uparrow^{\text{RLD}(\text{Dst } \mu; \text{Dst } \nu)} f \circ \mu \circ (\uparrow^{\text{RLD}(\text{Dst } \mu; \text{Dst } \nu)} f)^{-1} \subseteq \epsilon \\ \uparrow^{\text{RLD}(\text{Dst } \mu; \text{Dst } \nu)} f \circ \mu \circ (\uparrow^{\text{RLD}(\text{Dst } \mu; \text{Dst } \nu)} f)^{-1} \subseteq \nu. \end{aligned}$$

So a function f is uniformly continuous iff $\uparrow^{\text{RLD}(\text{Dst } \mu; \text{Dst } \nu)} f \circ \mu \circ (\uparrow^{\text{RLD}(\text{Dst } \mu; \text{Dst } \nu)} f)^{-1} \subseteq \nu$.

6.2 Our three definitions of continuity

I have expressed different kinds of continuity with simple algebraic formulas hiding the complexity of traditional epsilon-delta notation behind a smart algebra. Let's summarize these three algebraic formulas:

Let μ and ν are endomorphisms of some partially ordered precategory. Continuous functions can be defined as these morphisms f of this precategory which conform to the following formula:

$$f \in C(\mu; \nu) \Leftrightarrow f \in \text{Mor}(\text{Ob } \mu; \text{Ob } \nu) \wedge f \circ \mu \subseteq \nu \circ f.$$

If the precategory is a partially ordered dagger precategory then continuity also can be defined in two other ways:

$$\begin{aligned} f \in C'(\mu; \nu) &\Leftrightarrow f \in \text{Mor}(\text{Ob } \mu; \text{Ob } \nu) \wedge \mu \subseteq f^\dagger \circ \nu \circ f; \\ f \in C''(\mu; \nu) &\Leftrightarrow f \in \text{Mor}(\text{Ob } \mu; \text{Ob } \nu) \wedge f \circ \mu \circ f^\dagger \subseteq \nu. \end{aligned}$$

Remark 235. In the examples (above) about functors and relicts the “dagger functor” is the inverse of a functor or relict, that is $f^\dagger = f^{-1}$.

Proposition 236. Every of these three definitions of continuity forms a sub-precategory (subcategory if the original precategory is a category).

Proof.

C. Let $f \in C(\mu; \nu)$, $g \in C(\nu; \pi)$. Then $f \circ \mu \subseteq \nu \circ f$, $g \circ \nu \subseteq \pi \circ g$; $g \circ f \circ \mu \subseteq g \circ \nu \circ f \subseteq \pi \circ g \circ f$.
So $g \circ f \in C(\mu; \pi)$. $1_{\text{Ob } \mu} \in C(\mu; \mu)$ is obvious.

C'. Let $f \in C'(\mu; \nu)$, $g \in C'(\nu; \pi)$. Then $\mu \subseteq f^\dagger \circ \nu \circ f$, $\nu \subseteq g^\dagger \circ \pi \circ g$;

$$\mu \subseteq f^\dagger \circ g^\dagger \circ \pi \circ g \circ f; \quad \mu \subseteq (g \circ f)^\dagger \circ \pi \circ (g \circ f).$$

So $g \circ f \in C'(\mu; \pi)$. $1_{\text{Ob } \mu} \in C'(\mu; \mu)$ is obvious.

C''. Let $f \in C''(\mu; \nu)$, $g \in C''(\nu; \pi)$. Then $f \circ \mu \circ f^\dagger \subseteq \nu$, $g \circ \nu \circ g^\dagger \subseteq \pi$;

$$g \circ f \circ \mu \circ f^\dagger \circ g^\dagger \subseteq \pi; \quad (g \circ f) \circ \mu \circ (g \circ f)^\dagger \subseteq \pi.$$

So $g \circ f \in C''(\mu; \pi)$. $1_{\text{Ob } \mu} \in C''(\mu; \mu)$ is obvious. \square

Proposition 237. For a monovalued morphism f of a partially ordered dagger category and its endomorphisms μ and ν

$$f \in C'(\mu; \nu) \Rightarrow f \in C(\mu; \nu) \Rightarrow f \in C''(\mu; \nu).$$

Proof. Let $f \in C'(\mu; \nu)$. Then $\mu \subseteq f^\dagger \circ \nu \circ f$; $f \circ \mu \subseteq f \circ f^\dagger \circ \nu \circ f \subseteq 1_{\text{Dst } f} \circ \nu \circ f = \nu \circ f$; $f \in C(\mu; \nu)$.
Let $f \in C(\mu; \nu)$. Then $f \circ \mu \subseteq \nu \circ f$; $f \circ \mu \circ f^\dagger \subseteq \nu \circ f \circ f^\dagger \subseteq \nu \circ 1_{\text{Dst } f} = \nu$; $f \in C''(\mu; \nu)$. \square

Proposition 238. For an entirely defined morphism f of a partially ordered dagger category and its endomorphisms μ and ν

$$f \in C''(\mu; \nu) \Rightarrow f \in C(\mu; \nu) \Rightarrow f \in C'(\mu; \nu).$$

Proof. Let $f \in C''(\mu; \nu)$. Then $f \circ \mu \circ f^\dagger \subseteq \nu$; $f \circ \mu \circ f^\dagger \circ f \subseteq \nu \circ f$; $f \circ \mu \circ 1_{\text{Src } f} \subseteq \nu \circ f$; $f \circ \mu \subseteq \nu \circ f$;
 $f \in C(\mu; \nu)$.

Let $f \in C(\mu; \nu)$. Then $f \circ \mu \subseteq \nu \circ f$; $f^\dagger \circ f \circ \mu \subseteq f^\dagger \circ \nu \circ f$; $1_{\text{Src } f} \circ \mu \subseteq f^\dagger \circ \nu \circ f$; $\mu \subseteq f^\dagger \circ \nu \circ f$;
 $f \in C'(\mu; \nu)$. \square

For entirely defined monovalued morphisms our three definitions of continuity coincide:

Theorem 239. If f is a monovalued and entirely defined morphism then

$$f \in C'(\mu; \nu) \Leftrightarrow f \in C(\mu; \nu) \Leftrightarrow f \in C''(\mu; \nu).$$

Proof. From two previous propositions. \square

The classical general topology theorem that uniformly continuous function from a uniform space to an other uniform space is near-continuous regarding the proximities generated by the uniformities, generalized for relicts and functors takes the following form:

Theorem 240. If an entirely defined morphism of the category of relicts $f \in C''(\mu; \nu)$ for some endomorphisms μ and ν of the category of relicts, then $(\text{FCD})f \in C'((\text{FCD})\mu; (\text{FCD})\nu)$.

Exercise 2. I leave a simple exercise for the reader to prove the last theorem.

6.3 Continuousness of a restricted morphism

Consider some partially ordered semigroup. (For example it can be the semigroup of functors or semigroup of relicts regarding the composition.) Consider also some lattice (*lattice of objects*). (For example take the lattice of set theoretic filters.)

We will map every object A to *identity element* I_A of the semigroup (for example identity funcoid or identity reloid). For identity elements we will require

1. $I_A \circ I_B = I_{A \cap B}$;
2. $f \circ I_A \subseteq f$; $I_A \circ f \subseteq f$.

In the case when our semigroup is “dagger” (that is a dagger precategory) we will require also $(I_A)^\dagger = I_A$.

We can define *restricting* an element f of our semigroup to an object A by the formula $f|_A = f \circ I_A$.

We can define *rectangular restricting* an element μ of our semigroup to objects A and B as $I_B \circ \mu \circ I_A$. Optionally we can define direct product $A \times B$ of two objects by the formula (true for funcoids and for reloids):

$$\mu \cap (A \times B) = I_B \circ \mu \circ I_A.$$

Square restricting of an element μ to an object A is a special case of rectangular restricting and is defined by the formula $I_A \circ \mu \circ I_A$ (or by the formula $\mu \cap (A \times A)$).

Theorem 241. For every elements f, μ, ν of our semigroup and an object A

1. $f \in C(\mu; \nu) \Rightarrow f|_A \in C(I_A \circ \mu \circ I_A; \nu)$;
2. $f \in C'(\mu; \nu) \Rightarrow f|_A \in C'(I_A \circ \mu \circ I_A; \nu)$;
3. $f \in C''(\mu; \nu) \Rightarrow f|_A \in C''(I_A \circ \mu \circ I_A; \nu)$.

(Two last items are true for the case when our semigroup is dagger.)

Proof.

1. $f|_A \in C(I_A \circ \mu \circ I_A; \nu) \Leftrightarrow f|_A \circ I_A \circ \mu \circ I_A \subseteq \nu \circ f|_A \Leftrightarrow f \circ I_A \circ I_A \circ \mu \circ I_A \subseteq \nu \circ f|_A \Leftrightarrow f \circ I_A \circ \mu \circ I_A \subseteq \nu \circ f \circ I_A \Leftrightarrow f \circ I_A \circ \mu \subseteq \nu \circ f \Leftrightarrow f \circ \mu \subseteq \nu \circ f \Leftrightarrow f \in C(\mu; \nu)$.
2. $f|_A \in C'(I_A \circ \mu \circ I_A; \nu) \Leftrightarrow I_A \circ \mu \circ I_A \subseteq (f|_A)^\dagger \circ \nu \circ f|_A \Leftrightarrow I_A \circ \mu \circ I_A \subseteq (f \circ I_A)^\dagger \circ \nu \circ f \circ I_A \Leftrightarrow I_A \circ \mu \circ I_A \subseteq I_A \circ f^\dagger \circ \nu \circ f \circ I_A \Leftrightarrow \mu \subseteq f^\dagger \circ \nu \circ f \Leftrightarrow f \in C'(\mu; \nu)$.
3. $f|_A \in C''(I_A \circ \mu \circ I_A; \nu) \Leftrightarrow f|_A \circ I_A \circ \mu \circ I_A \circ (f|_A)^\dagger \subseteq \nu \Leftrightarrow f \circ I_A \circ I_A \circ \mu \circ I_A \circ I_A \circ f^\dagger \subseteq \nu \Leftrightarrow f \circ I_A \circ \mu \circ I_A \circ f^\dagger \subseteq \nu \Leftrightarrow f \circ \mu \circ f^\dagger \subseteq \nu \Leftrightarrow f \in C''(\mu; \nu)$. \square

7 Connectedness regarding funcoids and reloids

Definition 242. I will call *endo-reloids* and *endo-funcoids* reloids and funcoids with the same source and destination.

7.1 Some lemmas

Lemma 243. If $\neg(A[f]^*B) \wedge A \cup B \in \text{up}(\text{dom } f \cup \text{im } f)$ then f is closed on $\uparrow^U A$ for a funcoid $f \in \text{FCD}(U; U)$ and sets $A, B \in \mathcal{P}U$ (for every small set U).

Proof. $\neg(A[f]^*B) \Leftrightarrow \uparrow^U B \cap \langle f \rangle \uparrow^U A = 0^{\mathfrak{F}(U)} \Leftrightarrow (\text{dom } f \cup \text{im } f) \cap \uparrow^U B \cap \langle f \rangle^* A = 0^{\mathfrak{F}(U)} \Rightarrow ((\text{dom } f \cup \text{im } f) \setminus \uparrow^U A) \cap \langle f \rangle^* A = 0^{\mathfrak{F}(U)} \Leftrightarrow \langle f \rangle^* A \subseteq \uparrow^U A$. \square

Corollary 244. If $\neg(A[f]^*B) \wedge A \cup B \in \text{up}(\text{dom } f \cup \text{im } f)$ then f is closed on $\uparrow^U(A \setminus B)$ for a funcoid f and sets $A, B \in \mathcal{P}U$ (for every small set U).

Proof. Let $\neg(A[f]^*B) \wedge A \cup B \in \text{up}(\text{dom } f \cup \text{im } f)$. Then $\neg((A \setminus B)[f]^*B) \wedge \uparrow^U((A \setminus B) \cup B) \in \text{up}(\text{dom } f \cup \text{im } f)$. \square

Lemma 245. If $\neg(A[f]^*B) \wedge A \cup B \in \text{up}(\text{dom } f \cup \text{im } f)$ then $\neg(A[f^n]^*B)$ for every whole positive n .

Proof. Let $\neg(A[f]^*B) \wedge A \cup B \in \text{up}(\text{dom } f \cup \text{im } f)$. From the above proposition $\langle f \rangle^*A \subseteq \uparrow^U A$. $\uparrow^U B \cap \langle f \rangle \uparrow^U A = 0^{\mathfrak{B}(U)}$, consequently $\langle f \rangle^*A \subseteq \uparrow^U(A \setminus B)$. Because (by the above corollary) f is closed on $\uparrow^U(A \setminus B)$, then $\langle f \rangle \langle f \rangle \uparrow^U A \subseteq \uparrow^U(A \setminus B)$, $\langle f \rangle \langle f \rangle \langle f \rangle \uparrow^U A \subseteq \uparrow^U(A \setminus B)$, etc. So $\langle f^n \rangle \uparrow^U A \subseteq \uparrow^U(A \setminus B)$, $\uparrow^U B \asymp \langle f^n \rangle \uparrow^U A$, $\neg(A[f^n]^*B)$. \square

7.2 Endomorphism series

Definition 246. $S_1(\mu) \stackrel{\text{def}}{=} \mu \cup \mu^2 \cup \mu^3 \cup \dots$ for an endomorphism μ of a precategory with countable union of morphisms.

Definition 247. $S(\mu) \stackrel{\text{def}}{=} \mu^0 \cup S_1(\mu) = \mu^0 \cup \mu \cup \mu^2 \cup \mu^3 \cup \dots$ where $\mu^0 \stackrel{\text{def}}{=} I_{\text{Ob } \mu}$ (identity morphism for the object $\text{Ob } \mu$) where $\text{Ob } \mu$ is the object of endomorphism μ for an endomorphism μ of a category with countable union of morphisms.

I call S_1 and S *endomorphism series*.

We will consider the collection of all binary relations (on a set \mathcal{U}), as well as the collection of all functors and the collection of all relicts on a fixed set, as categories with single object \mathcal{U} and the identity morphisms $I_{\mathcal{U}}$, $I^{\text{FCD}(\Omega)}$, $I^{\text{RLD}(\Omega)}$.

Proposition 248. The relation $S(\mu)$ is transitive for the category of binary relations.

Proof.

$$\begin{aligned} S(\mu) \circ S(\mu) &= \mu^0 \circ S(\mu) \cup \mu \circ S(\mu) \cup \mu^2 \circ S(\mu) \cup \dots \\ &= (\mu^0 \cup \mu^1 \cup \mu^2 \cup \dots) \cup (\mu^1 \cup \mu^2 \cup \mu^3 \cup \dots) \cup (\mu^2 \cup \mu^3 \cup \mu^4 \cup \dots) \\ &= \mu^0 \cup \mu^1 \cup \mu^2 \cup \dots \\ &= S(\mu). \end{aligned}$$

\square

7.3 Connectedness regarding binary relations

Before going to research connectedness for functors and relicts we will excursion into the basic special case of connectedness regarding binary relations on a set \mathcal{U} .

Definition 249. A set A is called (*strongly*) *connected* regarding a binary relation μ when

$$\forall X \in \mathcal{P}(\text{dom } \mu) \setminus \{\emptyset\}, Y \in \mathcal{P}(\text{im } \mu) \setminus \{\emptyset\}: (X \cup Y = A \Rightarrow X[\mu]Y).$$

Let \mathcal{U} is a set.

Definition 250. *Path* between two elements $a, b \in \mathcal{U}$ in a set $A \subseteq \mathcal{U}$ through binary relation μ is the finite sequence $x_0 \dots x_n$ where $x_0 = a$, $x_n = b$ for $n \in \mathbb{N}$ and $x_i(\mu \cap A \times A)x_{i+1}$ for every $i = 0, \dots, n-1$. n is called *path length*.

Proposition 251. There exists path between every element $a \in \mathcal{U}$ and that element itself.

Proof. It is the path consisting of one vertex (of length 0). \square

Proposition 252. There is a path from element a to element b in a set A through a binary relation μ iff $a(S(\mu \cap A \times A))b$ (that is $(a, b) \in S(\mu \cap A \times A)$).

Proof.

\Rightarrow . If exists a path from a to b , then $\{b\} \subseteq \langle (\mu \cap A \times A)^n \rangle \{a\}$ where n is the path length. Consequently $\{b\} \subseteq \langle S(\mu \cap A \times A) \rangle \{a\}$; $a(S(\mu \cap A \times A))b$.

\Leftarrow . If $a(S(\mu \cap A \times A))b$ then exists $n \in \mathbb{N}$ such that $a(\mu \cap A \times A)^n b$. By definition of composition of binary relations this means that there exist finite sequence $x_0 \dots x_n$ where $x_0 = a$, $x_n = b$ for $n \in \mathbb{N}$ and $x_i(\mu \cap A \times A)x_{i+1}$ for every $i = 0, \dots, n-1$. That is there is path from a to b . \square

Theorem 253. The following statements are equivalent for a relation μ and a set A :

1. For every $a, b \in A$ there is a path between a and b in A through μ .
2. $S(\mu \cap A \times A) \supseteq A \times A$.
3. $S(\mu \cap A \times A) = A \times A$.
4. A is connected regarding μ .

Proof.

(1) \Rightarrow (2). Let for every $a, b \in A$ there is a path between a and b in A through μ . Then $a(S(\mu \cap A \times A))b$ for every $a, b \in A$. It is possible only when $S(\mu \cap A \times A) \supseteq A \times A$.

(3) \Rightarrow (1). For every two vertices a and b we have $a(S(\mu \cap A \times A))b$. So (by the previous theorem) for every two vertices a and b exist path from a to b .

(3) \Rightarrow (4). Suppose that $\neg(X[\mu \cap A \times A]Y)$ for some $X, Y \in \mathcal{P}U \setminus \{\emptyset\}$ such that $X \cup Y = A$. Then by a lemma $\neg(X[(\mu \cap A \times A)^n]Y)$ for every $n \in \mathbb{N}$. Consequently $\neg(X[S(\mu \cap A \times A)]Y)$. So $S(\mu \cap A \times A) \neq A \times A$.

(4) \Rightarrow (3). If $\langle S(\mu \cap A \times A) \rangle \{v\} = A$ for every vertex v then $S(\mu \cap A \times A) = A \times A$. Consider the remaining case when $V \stackrel{\text{def}}{=} \langle S(\mu \cap A \times A) \rangle \{v\} \subset A$ for some vertex v . Let $W = A \setminus V$. If $\text{card } A = 1$ then $S(\mu \cap A \times A) \supseteq (=) = A \times A$; otherwise $W \neq \emptyset$. Then $V \cup W = A$ and so $V[\mu]W$ what is equivalent to $V[\mu \cap A \times A]W$ that is $\langle \mu \cap A \times A \rangle V \cap W \neq \emptyset$. This is impossible because $\langle \mu \cap A \times A \rangle V = \langle \mu \cap A \times A \rangle \langle S(\mu \cap A \times A) \rangle V = \langle S_1(\mu \cap A \times A) \rangle V \subseteq \langle S(\mu \cap A \times A) \rangle V = V$.

(2) \Rightarrow (3). Because $S(\mu \cap A \times A) \subseteq A \times A$. \square

Corollary 254. A set A is connected regarding a binary relation μ iff it is connected regarding $\mu \cap A \times A$.

Definition 255. A *connected component* of a set A regarding a binary relation F is a maximal connected subset of A .

Theorem 256. The set A is partitioned into connected components (regarding every binary relation F).

Proof. Consider the binary relation $a \sim b \Leftrightarrow a(S(F))b \wedge b(S(F))a$. \sim is a symmetric, reflexive, and transitive relation. So all points of A are partitioned into a collection of sets Q . Obviously each component is (strongly) connected. If a set $R \subseteq A$ is greater than one of that connected components A then it contains a point $b \in B$ where B is some other connected component. Consequently R is disconnected. \square

Proposition 257. A set is connected (regarding a binary relation) iff it has one connected component.

Proof. Direct implication is obvious. Reverse is proved by contradiction. \square

7.4 Connectedness regarding funcoids and reloids

Definition 258. $S_1^*(\mu) = \bigcap \{ \uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} S_1(M) \mid M \in \text{up } \mu \}$ for an endo-reloid μ .

Definition 259. *Connectivity reloid* $S^*(\mu)$ for an endo-reloid μ is defined as follows:

$$S^*(\mu) = \bigcap \{ \uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} S(M) \mid M \in \text{up } \mu \}.$$

Remark 260. Do not mess the word *connectivity* with the word *connectedness* which means being connected.¹

Proposition 261. $S^*(\mu) = I^{\text{RLD}(\text{Ob } \mu)} \cup S_1^*(\mu)$ for every endo-reloid μ .

Proof. Follows from the theorem about distributivity of \cup regarding \cap (see [14]). \square

Proposition 262. $S^*(\mu) = S(\mu)$ if μ is a discrete reloid.

Proof. $S^*(\mu) = \bigcap \{S(\mu)\} = S(\mu)$. \square

Definition 263. A filter object $\mathcal{A} \in \mathfrak{F}(\text{Ob } \mu)$ is called *connected* regarding an endo-reloid μ when $S^*(\mu \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{A})) \supseteq \mathcal{A} \times^{\text{RLD}} \mathcal{A}$.

Obvious 264. A filter object $\mathcal{A} \in \text{Ob } \mu$ is connected regarding a reloid μ iff $S^*(\mu \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{A})) = \mathcal{A} \times^{\text{RLD}} \mathcal{A}$.

Definition 265. A filter object \mathcal{A} is called *connected* regarding an endo-funcoid μ when

$$\forall \mathcal{X}, \mathcal{Y} \in \mathfrak{F}(\text{Ob } \mu) \setminus \{0^{\mathfrak{F}(\text{Ob } \mu)}\}: (\mathcal{X} \cup \mathcal{Y} = \mathcal{A} \Rightarrow \mathcal{X}[\mu]\mathcal{Y}).$$

Proposition 266. Let A be a set. The f.o. $\uparrow^{\text{Ob } \mu} A$ is connected regarding an endo-funcoid μ iff

$$\forall \mathcal{X}, \mathcal{Y} \in \mathcal{P}(\text{Ob } \mu) \setminus \{\emptyset\}: (X \cup Y = A \Rightarrow X[\mu]^*Y).$$

Proof.

\Rightarrow . Obvious.

\Leftarrow . Follows from co-separability of filter objects. \square

Theorem 267. The following are equivalent for every set A and binary relation μ :

1. A is connected regarding binary relation μ .
2. $\uparrow^{\text{Ob } \mu} A$ is connected regarding $\uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} \mu$.
3. $\uparrow^{\text{Ob } \mu} A$ is connected regarding $\uparrow^{\text{FCD}(\text{Ob } \mu; \text{Ob } \mu)} \mu$.

Proof.

(1) \Leftrightarrow (2). $S^*(\uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} \mu \cap (\uparrow^{\text{Ob } \mu} A \times^{\text{RLD}} \uparrow^{\text{Ob } \mu} A)) = S^*(\uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} (\mu \cap A \times A)) = \uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} S(\mu \cap A \times A)$. So $S^*(\uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} \mu \cap (\uparrow^{\text{Ob } \mu} A \times^{\text{RLD}} \uparrow^{\text{Ob } \mu} A)) \supseteq \uparrow^{\text{Ob } \mu} A \times^{\text{RLD}} \uparrow^{\text{Ob } \mu} A \Leftrightarrow \uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} S(\mu \cap A \times A) \supseteq \uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} (A \times A) = \uparrow^{\text{Ob } \mu} A \times^{\text{RLD}} \uparrow^{\text{Ob } \mu} A$.

(1) \Leftrightarrow (3). Follows from the previous proposition. \square

Next is conjectured a statement more strong than the above theorem:

Conjecture 268. Let \mathcal{A} is an f.o. and F is a binary relation on $A \times B$ for some sets A, B . \mathcal{A} is connected regarding $\uparrow^{\text{FCD}(A; B)} F$ iff \mathcal{A} is connected regarding $\uparrow^{\text{RLD}(A; B)} F$.

Obvious 269. A filter object \mathcal{A} is connected regarding a reloid μ iff it is connected regarding the reloid $\mu \cap (\mathcal{A} \times^{\text{RLD}} \mathcal{A})$.

Obvious 270. A filter object \mathcal{A} is connected regarding a funcoid μ iff it is connected regarding the funcoid $\mu \cap (\mathcal{A} \times^{\text{FCD}} \mathcal{A})$.

Theorem 271. A filter object \mathcal{A} is connected regarding a reloid f iff $\uparrow^{\text{Ob } f} \mathcal{A}$ is connected regarding every $F \in \langle \uparrow^{\text{RLD}(\text{Ob } f; \text{Ob } f)} \rangle_{\text{up } f}$.

1. In some math literature these two words are used interchangeably.

Proof.

\Rightarrow . Obvious.

\Leftarrow . $\uparrow^{\text{Ob } f} \mathcal{A}$ is connected regarding $\uparrow^{\text{RLD}(\text{Ob } f; \text{Ob } f)} F$ iff $S(F) = F^0 \cup F^1 \cup F^2 \cup \dots \in \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{A})$.
 $S^*(f) = \bigcap \{ \uparrow^{\text{RLD}(\text{Ob } \mu; \text{Ob } \mu)} S(F) \mid F \in \text{up } f \} \supseteq \bigcap \{ \mathcal{A} \times^{\text{RLD}} \mathcal{A} \mid F \in \text{up } f \} = \mathcal{A} \times^{\text{RLD}} \mathcal{A}$. \square

Conjecture 272. A filter object \mathcal{A} is connected regarding a funoid μ iff \mathcal{A} is connected for every $F \in \langle \uparrow^{\text{FCD}(\text{Ob } \mu; \text{Ob } \mu)} \rangle \text{up } \mu$.

The above conjecture is open even for the case when \mathcal{A} is a principal f.o.

Conjecture 273. A filter object \mathcal{A} is connected regarding a reloid f iff it is connected regarding the funoid (FCD) f .

The above conjecture is true in the special case of principal filters:

Proposition 274. A f.o. $\uparrow^{\text{Ob } \mu} A$ (for a set A) is connected regarding an endo-reloid f iff it is connected regarding the endo-funoid (FCD) f .

Proof. $\uparrow^{\text{Ob } f} \mathcal{A}$ is connected regarding a reloid f iff A is connected regarding every $F \in \text{up } f$ that is when (taken in account that connectedness for $\uparrow^{\text{RLD}(\text{Ob } f; \text{Ob } f)} F$ is the same as connectedness of $\uparrow^{\text{FCD}(\text{Ob } f; \text{Ob } f)} F$)

$$\begin{aligned} \forall F \in \text{up } f \forall \mathcal{X}, \mathcal{Y} \in \mathfrak{F}(\text{Ob } f) \setminus \{0^{\mathfrak{F}(\text{Ob } f)}\}: (\mathcal{X} \cup \mathcal{Y} = \uparrow^{\text{Ob } f} \mathcal{A} \Rightarrow \mathcal{X}[\uparrow^{\text{FCD}(\text{Ob } f; \text{Ob } f)} F] \mathcal{Y}) &\Leftrightarrow \\ \forall \mathcal{X}, \mathcal{Y} \in \mathfrak{F}(\text{Ob } f) \setminus \{0^{\mathfrak{F}(\text{Ob } f)}\} \forall F \in \text{up } f: (\mathcal{X} \cup \mathcal{Y} = \uparrow^{\text{Ob } f} \mathcal{A} \Rightarrow \mathcal{X}[\uparrow^{\text{FCD}(\text{Ob } f; \text{Ob } f)} F] \mathcal{Y}) &\Leftrightarrow \\ \forall \mathcal{X}, \mathcal{Y} \in \mathfrak{F}(\text{Ob } f) \setminus \{0^{\mathfrak{F}(\text{Ob } f)}\}: (\mathcal{X} \cup \mathcal{Y} = \uparrow^{\text{Ob } f} \mathcal{A} \Rightarrow \forall F \in \text{up } f: \mathcal{X}[\uparrow^{\text{FCD}(\text{Ob } f; \text{Ob } f)} F] \mathcal{Y}) &\Leftrightarrow \\ \forall \mathcal{X}, \mathcal{Y} \in \mathfrak{F}(\text{Ob } f) \setminus \{0^{\mathfrak{F}(\text{Ob } f)}\}: (\mathcal{X} \cup \mathcal{Y} = \uparrow^{\text{Ob } f} \mathcal{A} \Rightarrow \mathcal{X}[(\text{FCD})f] \mathcal{Y}) &\end{aligned}$$

that is when the set $\uparrow^{\text{Ob } f} \mathcal{A}$ is connected regarding the funoid (FCD) f . \square

7.5 Algebraic properties of S and S^*

Theorem 275. $S^*(S^*(f)) = S^*(f)$ for every endo-reloid f .

Proof. $S^*(S^*(f)) = \bigcap \{ \uparrow^{\text{RLD}(\text{Ob } f; \text{Ob } f)} S(R) \mid R \in \text{up } S^*(f) \} \subseteq \bigcap \{ \uparrow^{\text{RLD}(\text{Ob } f; \text{Ob } f)} S(R) \mid R \in \{S(F) \mid F \in \text{up } f\} \} = \bigcap \{ \uparrow^{\text{RLD}(\text{Ob } f; \text{Ob } f)} S(S(F)) \mid F \in \text{up } f \} = \bigcap \{ \uparrow^{\text{RLD}(\text{Ob } f; \text{Ob } f)} S(F) \mid F \in \text{up } f \} = S^*(f)$.

So $S^*(S^*(f)) \subseteq S^*(f)$. That $S^*(S^*(f)) \supseteq S^*(f)$ is obvious. \square

Corollary 276. $S^*(S(f)) = S(S^*(f)) = S^*(f)$ for any endo-reloid f .

Proof. Obviously $S^*(S(f)) \supseteq S^*(f)$ and $S(S^*(f)) \supseteq S^*(f)$.

But $S^*(S(f)) \subseteq S^*(S^*(f)) = S^*(f)$ and $S(S^*(f)) \subseteq S^*(S^*(f)) = S^*(f)$. \square

Conjecture 277. $S(S(f)) = S(f)$ for

1. every endo-reloid f ;
2. every endo-funoid f .

Conjecture 278. For every endo-reloid f

1. $S(f) \circ S(f) = S(f)$;
2. $S^*(f) \circ S^*(f) = S^*(f)$;
3. $S(f) \circ S^*(f) = S^*(f) \circ S(f) = S^*(f)$.

Conjecture 279. $S(f) \circ S(f) = S(f)$ for every endo-funoid f .

8 Postface

8.1 Misc

See this Web page for my research plans: <http://www.mathematics21.org/agt-plans.html>

I deem that now the most important research topics in Algebraic General Topology are:

- to solve the open problems mentioned in this work;
- define and research compactness of funcoids.
- research are n -ary (where n is an ordinal, or more generally an index set) funcoids and reloids (plain funcoids and reloids are binary by analogy with binary relations).

We should also research relationships between complete funcoids and complete reloids.

All my research of funcoids and reloids is presented at

<http://www.mathematics21.org/algebraic-general-topology.html>

Appendix A Some counter-examples

For further examples we will use the filter object Δ defined by the formula

$$\Delta = \bigcap \{ \uparrow^{\mathfrak{F}(\mathbb{R})}(-\varepsilon; \varepsilon) \mid \varepsilon \in \mathbb{R}, \varepsilon > 0 \}.$$

I also will denote $\Omega(A)$ the Fréchet f.o. on the set A .

Example 280. There exist a funcoid f and a set S of funcoids such that $f \cap \bigcup S \neq \bigcup \langle f \cap \rangle S$.

Proof. Let $f = \Delta \times^{\text{FCD}} \uparrow^{\mathfrak{F}(\mathbb{R})} \{0\}$ and $S = \{ \uparrow^{\text{FCD}(\mathbb{R}; \mathbb{R})}((\varepsilon; +\infty) \times \{0\}) \mid \varepsilon > 0 \}$. Then $f \cap \bigcup S = \uparrow^{\text{FCD}(\mathbb{R}; \mathbb{R})}(\Delta \times \uparrow^{\mathfrak{F}(\mathbb{R})} \{0\}) \cap \uparrow^{\text{FCD}(\mathbb{R}; \mathbb{R})}((0; +\infty) \times \{0\}) = (\Delta \cap \uparrow^{\mathfrak{F}(\mathbb{R})}(0; +\infty)) \times^{\text{FCD}} \uparrow^{\mathfrak{F}(\mathbb{R})} \{0\} \neq 0^{\text{FCD}(\mathbb{R}; \mathbb{R})}$ while $\bigcup \langle f \cap \rangle S = \bigcup \{ 0^{\text{FCD}(\mathbb{R}; \mathbb{R})} \} = 0^{\text{FCD}(\mathbb{R}; \mathbb{R})}$. \square

Conjecture 281. There exist a set R of funcoids and a funcoid f such that $f \circ \bigcup R \neq \bigcup \langle f \circ \rangle R$.

Example 282. There exist a set R of funcoids and f.o. \mathcal{X} and \mathcal{Y} such that

1. $\mathcal{X}[\bigcup R]\mathcal{Y} \wedge \nexists f \in R: \mathcal{X}[f]\mathcal{Y}$;
2. $\langle \bigcup R \rangle \mathcal{X} \supset \bigcup \{ \langle f \rangle \mathcal{X} \mid f \in R \}$.

Proof.

1. Let $\mathcal{X} = \Delta$ and $\mathcal{Y} = 1^{\mathfrak{F}(\mathbb{R})}$. Let $R = \{ \uparrow^{\text{FCD}(\mathbb{R}; \mathbb{R})}((\varepsilon; +\infty) \times \mathbb{R}) \mid \varepsilon \in \mathbb{R}, \varepsilon > 0 \}$. Then $\bigcup R = \uparrow^{\text{FCD}(\mathbb{R}; \mathbb{R})}((0; +\infty) \times \mathbb{R})$. So $\mathcal{X}[\bigcup R]\mathcal{Y}$ and $\forall f \in R: \neg(\mathcal{X}[f]\mathcal{Y})$.
2. With the same \mathcal{X} and R we have $\langle \bigcup R \rangle \mathcal{X} = \mathbb{R}$ and $\langle f \rangle \mathcal{X} = 0^{\mathfrak{F}(\mathbb{R})}$ for every $f \in R$, thus $\bigcup \{ \langle f \rangle \mathcal{X} \mid f \in R \} = 0^{\mathfrak{F}(\mathbb{R})}$. \square

Theorem 283. For a f.o. a we have $a \times^{\text{RLD}} a \subseteq I^{\text{RLD}(\text{Base}(a))}$ only in the case if $a = 0^{\mathfrak{F}(\text{Base}(a))}$ or a is a trivial atomic f.o. (that is corresponds to an one-element set).

Proof. If $a \times^{\text{RLD}} a \subseteq I^{\text{RLD}(\text{Base}(a))}$ then exists $m \in \text{up}(a \times^{\text{RLD}} a)$ such that $m \subseteq I_{\text{Base}(a)}$. Consequently exist $A, B \in \text{up} a$ such that $A \times B \subseteq I_{\text{Base}(a)}$ what is possible only in the case when $A = B = a$ is an one-element set or empty set. \square

Corollary 284. Reloidal product of non-trivial atomic filter objects is non-atomic.

Proof. Obviously $(a \times^{\text{RLD}} a) \cap I^{\text{RLD}(\text{Base}(a))} \neq 0^{\mathfrak{F}(\text{Base}(a))}$ and $(a \times^{\text{RLD}} a) \cap I^{\text{RLD}(\text{Base}(a))} \subseteq a \times^{\text{RLD}} a$. \square

Example 285. $(\text{RLD})_{\text{in}} f \neq (\text{RLD})_{\text{out}} f$ for a funcoid f .

Proof. Let $f = I^{\text{FCD}(\mathbb{N})}$. Then $(\text{RLD})_{\text{in}}f = \bigcup \{a \times^{\text{RLD}} a \mid a \in \text{atoms } 1^{\mathfrak{F}(\mathbb{N})}\}$ and $(\text{RLD})_{\text{out}}f = I^{\text{RLD}(\mathbb{N})}$. But as we shown above $a \times^{\text{RLD}} a \not\subseteq I^{\text{RLD}(\mathbb{N})}$ for non-trivial f.o. a , and so $(\text{RLD})_{\text{in}}f \not\subseteq (\text{RLD})_{\text{out}}f$. \square

Proposition 286. $I^{\text{FCD}(\mathbb{N})} \cap \uparrow^{\text{FCD}(\mathbb{N};\mathbb{N})}((\mathbb{N} \times \mathbb{N}) \setminus I_{\mathbb{N}}) = I_{\Omega(\mathbb{N})}^{\text{FCD}} \neq 0^{\text{FCD}(\mathbb{N};\mathbb{N})}$.

Proof. Note that $\langle I_{\Omega(\mathbb{N})}^{\text{FCD}} \rangle \mathcal{X} = \mathcal{X} \cap \Omega(\mathbb{N})$.

Let $f = I^{\text{FCD}(\mathbb{N})}$, $g = \uparrow^{\text{FCD}(\mathbb{N};\mathbb{N})}((\mathbb{N} \times \mathbb{N}) \setminus I_{\mathbb{N}})$.

Let x is a non-trivial atomic f.o. If $X \in \text{up } x$ then $\text{card } X \geq 2$ (In fact, X is infinite but we don't need this.) and consequently $\langle g \rangle^* X = 1^{\mathfrak{F}(\mathbb{N})}$. Thus $\langle g \rangle x = 1^{\mathfrak{F}(\mathbb{N})}$. Consequently

$$\langle f \cap g \rangle x = \langle f \rangle x \cap \langle g \rangle x = x \cap 1^{\mathfrak{F}(\mathbb{N})} = x.$$

Also $\langle I_{\Omega(\mathbb{N})}^{\text{FCD}} \rangle x = x \cap \Omega(\mathbb{N}) = x$.

Let now x is a trivial f.o. Then $\langle f \rangle x = x$ and $\langle g \rangle x = 1^{\mathfrak{F}(\mathbb{N})} \setminus x$. So

$$\langle f \cap g \rangle x = \langle f \rangle x \cap \langle g \rangle x = x \cap (1^{\mathfrak{F}(\mathbb{N})} \setminus x) = 0^{\mathfrak{F}(\mathbb{N})}.$$

Also $\langle I_{\Omega(\mathbb{N})}^{\text{FCD}} \rangle x = x \cap \Omega(\mathbb{N}) = 0^{\mathfrak{F}(\mathbb{N})}$.

So $\langle f \cap g \rangle x = \langle I_{\Omega(\mathbb{N})}^{\text{FCD}} \rangle x$ for every atomic f.o. x . Thus $f \cap g = I_{\Omega(\mathbb{N})}^{\text{FCD}}$. \square

Example 287. There exist binary relations f and g such that $\uparrow^{\text{FCD}(A;B)} f \cap \uparrow^{\text{FCD}(A;B)} g \neq f \cap g$ for some sets A, B such that $f, g \subseteq A \times B$.

Proof. From the proposition above. \square

Example 288. There exists a discrete funcooid which is not a complemented element of the lattice of funcooids.

Proof. I will prove that quasi-complement (see [14] for the definition of quasi-complement) of the funcooid $I^{\text{FCD}(\mathbb{N})}$ is not its complement. We have:

$$\begin{aligned} (I^{\text{FCD}(\mathbb{N})})^* &= \bigcup \{c \in \text{FCD} \mid c \asymp I^{\text{FCD}(\mathbb{N})}\} \\ &\supseteq \bigcup \{\uparrow^{\mathbb{N}}\{\alpha\} \times^{\text{FCD}} \uparrow^{\mathbb{N}}\{\beta\} \mid \alpha, \beta \in \mathbb{N}, \{\alpha\} \times^{\text{FCD}} \{\beta\} \asymp I^{\text{FCD}(\mathbb{N})}\} \\ &= \bigcup \{\uparrow^{\mathbb{N}}\{\alpha\} \times^{\text{FCD}} \uparrow^{\mathbb{N}}\{\beta\} \mid \alpha, \beta \in \mathbb{N}, \alpha \neq \beta\} \\ &= \uparrow^{\text{FCD}(\mathbb{N};\mathbb{N})} \bigcup \{\{\alpha\} \times \{\beta\} \mid \alpha, \beta \in \mathbb{N}, \alpha \neq \beta\} \\ &= \uparrow^{\text{FCD}(\mathbb{N};\mathbb{N})}(\mathbb{N} \times \mathbb{N} \setminus I_{\mathbb{N}}) \end{aligned}$$

(used the corollary 110). But by proved above

$$(I^{\text{FCD}(\mathbb{N})})^* \cap I^{\text{FCD}(\mathbb{N})} \neq 0^{\mathfrak{F}(\mathbb{N})}. \quad \square$$

Example 289. There exists funcooid h such that $\text{up } h$ is not a filter.

Proof. Consider the funcooid $h = I_{\Omega(\mathbb{N})}^{\text{FCD}}$. We have (from the proposition) that $f \in \text{up } h$ and $g \in \text{up } f$, but $f \cap g = \emptyset \notin \text{up } h$. \square

Example 290. There exists a funcooid $h \neq 0^{\text{FCD}(A;B)}$ such that $(\text{RLD})_{\text{out}}h = 0^{\text{RLD}(A;B)}$.

Proof. Consider $h = I_{\Omega(\mathbb{N})}^{\text{FCD}}$. By proved above $h = f \cap g$ where $f = I^{\text{FCD}(\mathbb{N})}$, $g = \uparrow^{\text{FCD}(\mathbb{N};\mathbb{N})}((\mathbb{N} \times \mathbb{N}) \setminus I_{\mathbb{N}})$.

We have $f, g \in \text{up } h$.

So $(\text{RLD})_{\text{out}}h = \bigcap \langle \uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})} \rangle \text{up } h \subseteq \uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})}(f \cap g) = 0^{\text{RLD}(\mathbb{N};\mathbb{N})}$; and thus $(\text{RLD})_{\text{out}}h = 0^{\text{RLD}(\mathbb{N};\mathbb{N})}$. \square

Example 291. There exists a funcooid h such that $(\text{FCD})(\text{RLD})_{\text{out}}h \neq h$.

Proof. Follows from the previous example. \square

Example 292. $(\text{RLD})_{\text{in}}(\text{FCD})f \neq f$ for some convex reloid f .

Proof. Let $f = I^{\text{RLD}(\mathbb{N})}$. Then $(\text{FCD})f = I^{\text{FCD}(\mathbb{N})}$. Let a is some nontrivial atomic f.o. Then $(\text{RLD})_{\text{in}}(\text{FCD})f \supseteq a \times^{\text{RLD}} a \not\subseteq I^{\text{RLD}(\mathbb{N})}$ and thus $(\text{RLD})_{\text{in}}(\text{FCD})f \not\subseteq f$. \square

Remark 293. Before I found the last counter-example, I thought that $(\text{RLD})_{\text{in}}$ is an isomorphism from the set of of funcoids to the set of convex reloids. As this conjecture failed, we need an other way to characterize the set of reloids isomorphic to funcoids.

Example 294. There exist funcoids f and g such that

$$(\text{RLD})_{\text{out}}(g \circ f) \neq (\text{RLD})_{\text{out}}g \circ (\text{RLD})_{\text{out}}f.$$

Proof. Take $f = I_{\Omega(\mathbb{N})}^{\text{FCD}}$ and $g = 1^{\mathfrak{F}(\mathbb{N})} \times^{\text{FCD}} \uparrow^{\mathbb{N}}\{\alpha\}$ for some $\alpha \in \mathbb{N}$. Then $(\text{RLD})_{\text{out}}f = 0^{\text{RLD}(\mathbb{N};\mathbb{N})}$ and thus $(\text{RLD})_{\text{out}}g \circ (\text{RLD})_{\text{out}}f = 0^{\text{RLD}(\mathbb{N};\mathbb{N})}$.

We have $g \circ f = \Omega(\mathbb{N}) \times^{\text{FCD}} \uparrow^{\mathbb{N}}\{\alpha\}$.

Let's prove $(\text{RLD})_{\text{out}}(\Omega(\mathbb{N}) \times^{\text{FCD}} \uparrow^{\mathbb{N}}\{\alpha\}) = \Omega(\mathbb{N}) \times^{\text{RLD}} \uparrow^{\mathbb{N}}\{\alpha\}$.

Really: $(\text{RLD})_{\text{out}}(\Omega(\mathbb{N}) \times^{\text{FCD}} \uparrow^{\mathbb{N}}\{\alpha\}) = \bigcap \uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})} \text{up}(\Omega(\mathbb{N}) \times^{\text{FCD}} \{\alpha\}) = \bigcap \{\uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})}(K \times \{\alpha\}) \mid K \in \text{up } \Omega(\mathbb{N})\}$.

$F \in \text{up } \bigcap \{\uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})}(K \times \{\alpha\}) \mid K \in \text{up } \Omega(\mathbb{N})\} \Leftrightarrow F \in \text{up}(\bigcap \{\uparrow^{\mathbb{N}}K \mid K \in \text{up } \Omega(\mathbb{N})\} \times^{\text{RLD}} \uparrow^{\mathbb{N}}\{\alpha\})$ for every $F \in \mathcal{P}(\mathbb{N} \times \mathbb{N})$. Thus

$$\bigcap \{\uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})}(K \times \{\alpha\}) \mid K \in \text{up } \Omega(\mathbb{N})\} = \bigcap \{\uparrow^{\mathbb{N}}K \mid K \in \text{up } \Omega(\mathbb{N})\} \times^{\text{RLD}} \uparrow^{\mathbb{N}}\{\alpha\} = \Omega(\mathbb{N}) \times^{\text{RLD}} \uparrow^{\mathbb{N}}\{\alpha\}.$$

So $(\text{RLD})_{\text{out}}(\Omega(\mathbb{N}) \times^{\text{FCD}} \uparrow^{\mathbb{N}}\{\alpha\}) = \Omega(\mathbb{N}) \times^{\text{RLD}} \uparrow^{\mathbb{N}}\{\alpha\}$.

Thus $(\text{RLD})_{\text{out}}(g \circ f) = \Omega(\mathbb{N}) \times^{\text{RLD}} \uparrow^{\mathbb{N}}\{\alpha\} \neq 0^{\text{RLD}(\mathbb{N};\mathbb{N})}$. \square

Example 295. (FCD) does not preserve finite meets.

Proof. $(\text{FCD})(I^{\text{RLD}(\mathbb{N})} \cap (1^{\text{RLD}(\mathbb{N};\mathbb{N})} \setminus I^{\text{RLD}(\mathbb{N})})) = (\text{FCD})0^{\text{RLD}(\mathbb{N};\mathbb{N})} = 0^{\text{FCD}(\mathbb{N};\mathbb{N})}$.

On the other hand

$$(\text{FCD})I^{\text{RLD}(\mathbb{N})} \cap (\text{FCD})(1^{\text{RLD}(\mathbb{N};\mathbb{N})} \setminus I^{\text{RLD}(\mathbb{N})}) = I^{\text{FCD}(\mathbb{N})} \cap (1^{\text{FCD}(\mathbb{N};\mathbb{N})} \setminus I^{\text{FCD}(\mathbb{N})}) = I_{\Omega(\mathbb{N})}^{\text{FCD}} \neq 0^{\text{FCD}(\mathbb{N};\mathbb{N})}$$

(used the proposition 204). \square

Corollary 296. (FCD) is not an upper adjoint (in general).

Considering restricting polynomials (considered as reloids) to atomic filter objects, it is simple to prove that each that restriction is injective if not restricting a constant polynomial. Does this hold in general? No, see the following example:

Example 297. There exists a monovalued reloid with atomic domain which is neither injective nor constant (that is not a restriction of a constant function).

Proof. (based on [15]) Consider the function $F \in \mathbb{N}^{\mathbb{N} \times \mathbb{N}}$ defined by the formula $(x; y) \mapsto x$.

Let ω_x is a non-principal atomic filter object on the vertical line $\{x\} \times \mathbb{N}$ for every $x \in \mathbb{N}$.

Let T is the collection of such sets Y that $Y \cap (\{x\} \times \mathbb{N}) \in \text{up } \omega_x$ for all but finitely many vertical lines. Obviously T is a filter.

Let $\omega \in \text{atoms up}^{-1} T$.

For every $x \in \mathbb{N}$ we have some $Y \in T$ for which $(\{x\} \times \mathbb{N}) \cap Y = \emptyset$ and thus $(\{x\} \times \mathbb{N}) \cap \text{up } \omega = \emptyset$.

Let $g = (\uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})} F)|_{\omega}$. If g is constant, then there exist a constant function $G \in \text{up } g$ and $F \cap G$ is also constant. Obviously $\text{dom } \uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})}(F \cap G) \supseteq \omega$. The function $F \cap G$ cannot be constant because otherwise $\omega \subseteq \text{dom } \uparrow^{\text{RLD}(\mathbb{N};\mathbb{N})}(F \cap G) \subseteq \uparrow^{\mathbb{N}}\{x\} \times \mathbb{N}$ for some $x \in \mathbb{N}$ what is impossible by proved above. So g is not constant.

Suppose there g is injective. Then there exists an injection $G \in \text{up } g$. So $\text{dom } G$ intersects each vertical line by atmost one element that is $\overline{\text{dom } G}$ intersects every vertical line by the whole line or the line without one element. Thus $\overline{\text{dom } G} \in T \subseteq \text{up } \omega$ and consequently $\text{dom } G \notin \text{up } \omega$ what is impossible.

Thus g is neither injective nor constant. \square

9 Second product. Oblique product

Definition 298. $\mathcal{A} \times_F^{\text{RLD}} \mathcal{B} \stackrel{\text{def}}{=} (\text{RLD})_{\text{out}}(\mathcal{A} \times^{\text{FCD}} \mathcal{B})$ for every f.o. \mathcal{A} and \mathcal{B} . I will call it *second direct product* of filter objects \mathcal{A} and \mathcal{B} .

Remark 299. The letter F in the above definition is from the word “funcoid”. It signifies that it seems to be impossible to define $\mathcal{A} \times_F^{\text{RLD}} \mathcal{B}$ directly without referring to the direct product $\mathcal{A} \times^{\text{FCD}} \mathcal{B}$ of funcoids.

Definition 300. *Oblique products* of f.o. \mathcal{A} and \mathcal{B} are defined as

$$\begin{aligned} \mathcal{A} \times \mathcal{B} &= \bigcap \{ \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \mid f \in \mathcal{P}(\text{Base}(\mathcal{A}) \times \text{Base}(\mathcal{B})), \forall B \in \text{up } \mathcal{B}: \\ &\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \supseteq \mathcal{A} \times^{\text{FCD}} \uparrow^{\text{Base}(\mathcal{B})} B \}; \\ \mathcal{A} \times \mathcal{B} &= \bigcap \{ \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \mid f \in \mathcal{P}(\text{Base}(\mathcal{A}) \times \text{Base}(\mathcal{B})), \forall A \in \text{up } \mathcal{A}: \\ &\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \supseteq \uparrow^{\text{Base}(\mathcal{A})} A \times^{\text{FCD}} \mathcal{B} \}. \end{aligned}$$

Proposition 301. $\mathcal{A} \times_F^{\text{RLD}} \mathcal{B} \subseteq \mathcal{A} \times \mathcal{B} \subseteq \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ for every f.o. \mathcal{A}, \mathcal{B} .

Proof. $\mathcal{A} \times \mathcal{B} \subseteq \bigcap \{ \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \mid f \in \mathcal{P}(\text{Base}(\mathcal{A}) \times \text{Base}(\mathcal{B})), \forall A \in \text{up } \mathcal{A}, \forall B \in \text{up } \mathcal{B}: \\ \uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \supseteq \uparrow^{\text{Base}(\mathcal{A})} A \times^{\text{FCD}} \uparrow^{\text{Base}(\mathcal{B})} B \} \subseteq \bigcap \{ \uparrow^{\text{Base}(\mathcal{A})} A \times^{\text{FCD}} \uparrow^{\text{Base}(\mathcal{B})} B \mid A \in \text{up } \mathcal{A}, \\ B \in \text{up } \mathcal{B} \} = \mathcal{A} \times^{\text{RLD}} \mathcal{B}$.

$\mathcal{A} \times \mathcal{B} \supseteq \bigcap \{ \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \mid f \in \mathcal{P}(\text{Base}(\mathcal{A}) \times \text{Base}(\mathcal{B})), \uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \supseteq \\ \mathcal{A} \times^{\text{FCD}} \mathcal{B} \} = \bigcap \{ \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \mid f \in \text{up}(\mathcal{A} \times^{\text{FCD}} \mathcal{B}) \} = (\text{RLD})_{\text{out}}(\mathcal{A} \times^{\text{FCD}} \mathcal{B}) = \mathcal{A} \times_F^{\text{RLD}} \mathcal{B}$. \square

Conjecture 302. $\mathcal{A} \times_F^{\text{RLD}} \mathcal{B} \subset \mathcal{A} \times \mathcal{B}$ for some f.o. \mathcal{A}, \mathcal{B} .

A stronger conjecture:

Conjecture 303. $\mathcal{A} \times_F^{\text{RLD}} \mathcal{B} \subset \mathcal{A} \times \mathcal{B} \subset \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ for some f.o. \mathcal{A}, \mathcal{B} . Particularly, is this formula true for $\mathcal{A} = \mathcal{B} = \Delta \cap \uparrow^{\mathbb{R}}(0; +\infty)$?

The above conjecture is similar to Fermat Last Theorem as having no value by itself but being somehow challenging to prove it.

Example 304. $\mathcal{A} \times \mathcal{B} \subset \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ for some f.o. \mathcal{A}, \mathcal{B} .

Proof. It's enough to prove $\mathcal{A} \times \mathcal{B} \neq \mathcal{A} \times^{\text{RLD}} \mathcal{B}$.

Let $\Delta_+ = \Delta \cap (0; +\infty)$. Let $\mathcal{A} = \mathcal{B} = \Delta_+$.

Let $K = (\geq)_{\mathbb{R} \times \mathbb{R}}$.

Obviously $K \notin \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{B})$.

$\mathcal{A} \times \mathcal{B} \subseteq \uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} K$ and thus $K \in \text{up}(\mathcal{A} \times \mathcal{B})$ because $\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} K \supseteq \\ \Delta_+ \times^{\text{FCD}} \uparrow^{\text{Base}(\mathcal{B})} B = \mathcal{A} \times^{\text{FCD}} \uparrow^{\text{Base}(\mathcal{B})} B$ for $B = (0; +\infty)$.

Thus $\mathcal{A} \times \mathcal{B} \neq \mathcal{A} \times^{\text{RLD}} \mathcal{B}$. \square

Example 305. $\mathcal{A} \times_F^{\text{RLD}} \mathcal{B} \subset \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ for some f.o. \mathcal{A}, \mathcal{B} .

Proof. This follows from the above example. \square

Proposition 306. $(\mathcal{A} \times \mathcal{B}) \cap (\mathcal{A} \times \mathcal{B}) = \mathcal{A} \times_F^{\text{RLD}} \mathcal{B}$ for every f.o. \mathcal{A}, \mathcal{B} .

Proof. $(\mathcal{A} \times \mathcal{B}) \cap (\mathcal{A} \times \mathcal{B}) \subseteq \bigcap \{\uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \mid f \in \mathcal{P}(\text{Base}(\mathcal{A}) \times \text{Base}(\mathcal{B}))\}$,
 $\uparrow^{\text{FCD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \supseteq \mathcal{A} \times^{\text{FCD}} \mathcal{B} = \bigcap \{\uparrow^{\text{RLD}(\text{Base}(\mathcal{A}); \text{Base}(\mathcal{B}))} f \mid f \in \text{up}(\mathcal{A} \times^{\text{FCD}} \mathcal{B})\} =$
 $(\text{RLD})_{\text{out}}(\mathcal{A} \times^{\text{FCD}} \mathcal{B}) = \mathcal{A} \times^{\text{RLD}} \mathcal{B}$.

To finish the proof we need to show $\mathcal{A} \times \mathcal{B} \supseteq \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ and $\mathcal{A} \times \mathcal{B} \supseteq \mathcal{A} \times^{\text{RLD}} \mathcal{B}$. By symmetry it's enough to show $\mathcal{A} \times \mathcal{B} \supseteq \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ what is proved above. \square

Conjecture 307. $(\mathcal{A} \times \mathcal{B}) \cup (\mathcal{A} \times \mathcal{B}) = \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ for every f.o. \mathcal{A}, \mathcal{B} .

Example 308. $(\text{RLD})_{\text{out}}(\text{FCD})f \neq f$ for some convex reloid f .

Proof. Let $f = \mathcal{A} \times^{\text{RLD}} \mathcal{B}$ where \mathcal{A} and \mathcal{B} are from the previous example.

$(\text{FCD})(\mathcal{A} \times^{\text{RLD}} \mathcal{B}) = \mathcal{A} \times^{\text{FCD}} \mathcal{B}$ by the proposition 212.

So $(\text{RLD})_{\text{out}}(\text{FCD})(\mathcal{A} \times^{\text{RLD}} \mathcal{B}) = (\text{RLD})_{\text{out}}(\mathcal{A} \times^{\text{FCD}} \mathcal{B}) = \mathcal{A} \times^{\text{RLD}} \mathcal{B} \neq \mathcal{A} \times^{\text{RLD}} \mathcal{B}$. \square

Bibliography

- [1] E. M. Alfsen and J. E. Fenstad. On the equivalence between proximity structures and totally bounded uniform structures. *Mathematica Scandinavica*, 7:353–360, 1959.
- [2] E. M. Alfsen and Olav Njåstad. Proximity and generalized uniformity. *Fund. Math.*, 52:235–252, 1963.
- [3] C. H. Dowker. Mappings of proximity structures. *General Topology and its Relations to Modern Analysis and Algebra*, Proceedings of the symposium held in Prague in September 1961:139–141, 1961.
- [4] V.A. Efremovich. The geometry of proximity i. *Mat. Sb. (N.S.)*, 31(73):189–200, 1952.
- [5] Miroslav Hušek. Generalized proximity and uniform spaces. i. *Commentationes Mathematicae Universitatis Carolinae*, 5(4):247–266, 1964.
- [6] Miroslav Hušek. Generalized proximity and uniform spaces. ii. *Commentationes Mathematicae Universitatis Carolinae*, 6(1):119–139, 1965.
- [7] Hans-Peter A. Kunzi. Quasi-uniform spaces, 2001.
- [8] Hans-Peter A. Kunzi. Quasi-uniform spaces in the year 2001, 2001.
- [9] Mrówka. On complete proximity spaces. *Doklady Akad. Nauk. SSSR*, N.S. 108:587–590, 1956.
- [10] Mrówka. On the notion of completeness in proximity spaces. *Bull. Acad. Polon. Sci.*, III, 4:201–216, 1956.
- [11] Olav Njåstad. Some properties of proximity and generalized uniformity. *Mathematica Scandinavica*, 12:47–56, 1963.
- [12] William J. Pervin. Quasi-uniformization of topological spaces. *Mathematische Annalen*, 147:316–317, 1962. 10.1007/BF01440953.
- [13] PlanetMath. Criteria for a poset to be a complete lattice. At <http://planetmath.org/encyclopedia/CriteriaForAPosetToBeACompleteLattice.html>.
- [14] Victor Porton. Filters on posets and generalizations. At <http://www.mathematics21.org/binaries/filters.pdf>.
- [15] Mark Sapir (mathoverflow.net/users/6976). An ultrafilter and a partition. MathOverflow. <http://mathoverflow.net/questions/44055> (version: 2010-10-29).
- [16] Yu. M. Smirnov. On proximity spaces. *Mat. Sb. (N.S.)*, 31(73):543–574, 1952.
- [17] Yu. M. Smirnov. On completeness of proximity spaces. *Doklady Acad. Nauk SSSR*, 88:761–764, 1953.
- [18] Yu. M. Smirnov. On completeness of uniform spaces and proximity spaces. *Doklady Acad. Nauk SSSR*, 91:1281–1284, 1953.
- [19] Yu. M. Smirnov. On completeness of proximity spaces i. *Trudy Moscov. Mat. Obsc.*, 3:271–306, 1954.
- [20] Yu. M. Smirnov. On completeness of proximity spaces ii. *Trudy Moscov. Mat. Obsc.*, 4:421–438, 1955.