

The New Prime theorems (45) - (70)

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that the new prime theorems (45)-(70) contain infinitely many prime solutions and no prime solutions.

这是人类有史以来第一次发现素数新天地，用蒋函数证明在什么情况下存在，在什么情况下不存在。简单情况下可用计算机验证。

The New Prime theorem (45)

$$P, jP^{10} + k - j (j = 1, \dots, k - 1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{10} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{10} + k - j (j = 1, \dots, k - 1). \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P - 1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{10} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P - 1 \quad (3)$$

If $\chi(P) \leq P - 2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P - 1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{10} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(10)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 11$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

we prove that for $k = 3, 11$, (1) contain no prime solutions

Example 2. Let $k = 5, 7$ and $k > 11$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 5, 7$ and $k > 11$ (1) contain infinitely many prime solutions

The New Prime theorem (46)

$$P, jP^{12} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang

Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{12} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{12} + k - j (j = 1, \dots, k-1). \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{12} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{12} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(12)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3, 5, 7, 13$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 13$ (1) contain no prime solutions.

Example 2. Let $k = 11$ and $k > 13$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 11$ and $k > 13$, (1) contain infinitely many prime solutions

The New Prime theorem (47)

$$P, jP^{14} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{14} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{14} + k - j (j = 1, \dots, k-1). \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{14} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{14} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(14)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3$. From (2) and(3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$, (1) contain no prime solutions.

Example 2. Let $k > 3$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k > 3$ (1) contain infinitely many prime solutions

The New Prime theorem (48)

$$P, jP^{16} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang

Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{16} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{16} + k - j (j = 1, \dots, k-1). \quad (1)$$

contain infinitely many prime solutions or no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{16} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{16} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(16)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 5, 17$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 17$ (1) contain no prime solutions.

Example 2. Let $k = 7, 11, 13$ and $k > 17$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 7, 11, 13$ and $k > 17$ (1) contain infinitely many prime solutions

The New Prime theorem (49)

$$P, jP^{18} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{18} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{18} + k - j (j = 1, \dots, k-1). \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{18} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{18} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(18)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 7, 19$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 19$, (1) contain no prime solutions.

Example 2. Let $k = 5, 11, 13, 17$ and $k > 19$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 5, 11, 13, 17$ and $k > 19$, (1) contain infinitely many prime solutions

The New Prime theorem (50)

$$P, jP^{20} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{20} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{20} + k - j (j=1, \dots, k-1). \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{20} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{20} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega) \omega^{k-1}}{(20)^{k-1} \phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3, 5, 11$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 11$, (1) contain no prime solutions.

Example 2. Let $k = 7$ and $k > 11$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 7$ and $k > 11$, (1) contain infinitely many prime solutions

The New Prime theorem (51)

$$P, jP^{22} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{22} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{22} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{22} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{22} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(22)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3, 23$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 23$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 23$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 23$, (1) contain infinitely many prime solutions

The New Prime theorem (52)

$$P, jP^{24} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{24} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{24} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P - 1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{24} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{24} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega) \omega^{k-1}}{(24)^{k-1} \phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3, 5, 7, 13$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 13$, (1) contain no prime solutions.

Example 2. Let $k = 11$ and $k \geq 17$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 11$ and $k \geq 17$, (1) contain infinitely many prime solutions

The New Prime theorem (53)

$$P, jP^{26} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{26} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{26} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{26} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{26} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(26)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$, (1) contain no prime solutions.

Example 2. Let $k > 3$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k > 3$, (1) contain infinitely many prime solutions

The New Prime theorem (54)

$$P, jP^{28} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang

Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{28} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{28} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{28} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{28} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(28)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3, 29$. From (2) and(3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 29$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 29$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 29$, (1) contain infinitely many prime solutions

The New Prime theorem (55)

$$P, jP^{30} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{30} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{30} + k - j (j = 1, \dots, k-1). \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{30} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{30} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(30)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 7, 11, 31$. From (2) and(3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 11, 31$, (1) contain no prime solutions.

Example 2. Let $k = 5, 13, 17, 19, 23, 29$ and $k > 31$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 5, 13, 17, 19, 23, 29$ and $k > 31$, (1) contain infinitely many prime solutions

The New Prime theorem (56)

$$P, jP^{32} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{32} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{32} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{32} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{32} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega) \omega^{k-1}}{(32)^{k-1} \phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 5, 17$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 17$, (1) contain no prime solutions.

Example 2. Let $k = 7, 11, 13$ and $k > 17$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 7, 11, 13$ and $k > 17$, (1) contain infinitely many prime solutions

The New Prime theorem (57)

$$P, jP^{34} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{34} + k - j$ contain infinitely many prime solutions and no

prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{34} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{34} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{34} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(34)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3$. From (2) and(3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3$, (1) contain no prime solutions.

Example 2. Let $k > 3$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k > 3$, (1) contain infinitely many prime solutions

The New Prime theorem (58)

$$P, jP^{36} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang

Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{36} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{36} + k - j (j=1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P - 1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{36} + k - j] \equiv 0 \pmod{P}, q=1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{36} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega) \omega^{k-1}}{(36)^{k-1} \phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 7, 13, 19, 37$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 13, 19, 37$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 7, 13, 19, 37$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7, 13, 19, 37$, (1) contain infinitely many prime solutions

The New Prime theorem (59)

$$P, jP^{38} + k - j (j=1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{38} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{38} + k - j (j=1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{38} + k - j] \equiv 0 \pmod{P}, q=1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{38} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega) \omega^{k-1}}{(38)^{k-1} \phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$.

Example 1. Let $k=3$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k=3$, (1) contain no prime solutions.

Example 2. Let $k > 3$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k > 3$, (1) contain infinitely many prime solutions

The New Prime theorem (60)

$$P, jP^{40} + k - j(j = 1, \dots, k - 1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{40} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{40} + k - j(j = 1, \dots, k - 1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P - 1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{40} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P - 1 \quad (3)$$

If $\chi(P) \leq P - 2$ then we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P - 1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{40} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(40)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_p (P - 1)$.

Example 1. Let $k = 3, 5, 11, 41$. From (2) and(3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 11, 41$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 5, 11, 41$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 11, 41$, (1) contain infinitely many prime solutions

The New Prime theorem (61)

$$P, jP^{42} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{42} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{42} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{42} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{42} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(42)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 7, 43$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 43$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 7, 43$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7, 43$, (1) contain infinitely many prime solutions

The New Prime theorem (62)

$$P, jP^{44} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{44} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{44} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{44} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P - 1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{44} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(44)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P - 1)$.

Example 1. Let $k = 3, 5, 23$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 23$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 5, 23$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 23$, (1) contain infinitely many prime solutions

The New Prime theorem (63)

$$P, jP^{46} + k - j (j = 1, \dots, k - 1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{46} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{46} + k - j (j = 1, \dots, k - 1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P - 1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{46} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) \leq P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{46} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(46)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3, 47$. From (2) and(3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 47$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 47$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 47$, (1) contain infinitely many prime solutions

The New Prime theorem (64)

$$P, jP^{48} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{48} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{48} + k - j (j=1, \dots, k-1). \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{48} + k - j] \equiv 0 \pmod{P}, q=1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{48} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(48)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 5, 7, 13, 17$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 13, 17$, (1) contain no prime solutions.

Example 2. Let $k = 11$ and $k > 17$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 11$ and $k > 17$, (1) contain infinitely many prime solutions

The New Prime theorem (65)

$$P, jP^{50} + k - j (j=1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{50} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{50} + k - j (j=1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P - 1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{50} + k - j] \equiv 0 \pmod{P}, q=1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{50} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(50)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 11$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 11$, (1) contain no prime solutions.

Example 2. Let $k = 5, 7$ and $k > 11$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k = 5, 7$ and $k > 11$, (1) contain infinitely many prime solutions

The New Prime theorem (66)

$$P, jP^{52} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{52} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{52} + k - j (j = 1, \dots, k-1). \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)], \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{52} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{52} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(52)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 5, 53$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 53$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 5, 53$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 53$, (1) contain infinitely many prime solutions

The New Prime theorem (67)

$$P, jP^{54} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{54} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{54} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{54} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{54} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(54)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_p (P-1)$.

Example 1. Let $k = 3, 7, 19$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 7, 19$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 7, 19$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 7, 19$, (1) contain infinitely many prime solutions

The New Prime theorem (68)

$$P, jP^{56} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{56} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{56} + k - j (j = 1, \dots, k-1). \quad (1)$$

Contain infinitely many prime solutions or no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_p [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_p P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{56} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{56} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(56)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 5, 29$. From (2) and(3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 29$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 5, 29$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 29$, (1) contain infinitely many prime solutions

The New Prime theorem (69)

$$P, jP^{58} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang

Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{58} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{58} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{58} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{58} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(58)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\varphi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 59$. From (2) and (3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 59$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 59$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 59$, (1) contain infinitely many prime solutions

The New Prime theorem (70)

$$P, jP^{60} + k - j (j = 1, \dots, k-1)$$

Chun-Xuan Jiang
Jiangchunxuan@vip.sohu.com

Abstract

Using Jiang function we prove that $jP^{60} + k - j$ contain infinitely many prime solutions and no prime solutions.

Theorem. Let k be a given odd prime.

$$P, jP^{60} + k - j (j = 1, \dots, k-1), \quad (1)$$

contain infinitely many prime solutions and no prime solutions.

Proof. We have Jiang function [1,2]

$$J_2(\omega) = \prod_P [P-1 - \chi(P)] \quad (2)$$

where $\omega = \prod_P P$, $\chi(P)$ is the number of solutions of congruence

$$\prod_{j=1}^{k-1} [jq^{60} + k - j] \equiv 0 \pmod{P}, q = 1, \dots, P-1 \quad (3)$$

If $\chi(P) \leq P-2$ then from (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (4)$$

We prove that (1) contain infinitely many prime solutions.

If $\chi(P) = P-1$ then from (2) and (3) we have

$$J_2(\omega) = 0 \quad (5)$$

We prove that (1) contain no prime solutions [1,2]

If $J_2(\omega) \neq 0$ then we have asymptotic formula [1,2]

$$\pi_k(N, 2) = \left| \left\{ P \leq N : jP^{60} + k - j = \text{prime} \right\} \right| \sim \frac{J_2(\omega)\omega^{k-1}}{(60)^{k-1}\phi^k(\omega)} \frac{N}{\log^k N} \quad (6)$$

where $\phi(\omega) = \prod_P (P-1)$.

Example 1. Let $k = 3, 5, 7, 11, 13, 31, 61$. From (2) and(3) we have

$$J_2(\omega) = 0 \quad (7)$$

We prove that for $k = 3, 5, 7, 11, 13, 31, 61$, (1) contain no prime solutions.

Example 2. Let $k \neq 3, 5, 7, 11, 13, 31, 61$. From (2) and (3) we have

$$J_2(\omega) \neq 0 \quad (8)$$

We prove that for $k \neq 3, 5, 7, 11, 13, 31, 61$, (1) contain infinitely many prime solutions

Remark. The prime number theory is basically to count the Jiang function $J_{n+1}(\omega)$ and Jiang

prime k -tuple singular series $\sigma(J) = \frac{J_2(\omega)\omega^{k-1}}{\phi^k(\omega)} = \prod_P \left(1 - \frac{1 + \chi(P)}{P} \right) \left(1 - \frac{1}{P} \right)^{-k}$ [1,2], which can count

the number of prime numbers. The prime distribution is not random. But Hardy-Littlewood prime k -tuple

singular series $\sigma(H) = \prod_P \left(1 - \frac{\nu(P)}{P} \right) \left(1 - \frac{1}{P} \right)^{-k}$ is false [3-8], which cannot count the number of prime

numbers[3].

References

- [1] Chun-Xuan Jiang, Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. Inter. Acad. Press, 2002, MR2004c:11001, (<http://www.i-b-r.org/docs/jiang.pdf>) (<http://www.wbabin.net/math/xuan13.pdf>)(<http://vixra.org/numth/>).
- [2] Chun-Xuan Jiang, Jiang's function $J_{n+1}(\omega)$ in prime distribution.(<http://www.wbabin.net/math/xuan2.pdf>.) (<http://wbabin.net/xuan.htm#chun-xuan>.)(<http://vixra.org/numth/>)
- [3] Chun-Xuan Jiang, The Hardy-Littlewood prime k -tuple conjecture is false.(<http://wbabin.net/xuan.htm#chun-xuan>)(<http://vixra.org/numth/>)
- [4] G. H. Hardy and J. E. Littlewood, Some problems of "Partitio Numerorum", III: On the expression of a number as a sum of primes. Acta Math., 44(1923)1-70.
- [5] W. Narkiewicz, The development of prime number theory. From Euclid to Hardy and Littlewood. Springer-Verlag, New York, NY. 2000, 333-353.
- [6] B. Green and T. Tao, Linear equations in primes. To appear, Ann. Math.
- [7] D. Goldston, J. Pintz and C. Y. Yildirim, Primes in tuples I. Ann. Math., 170(2009) 819-862.
- [8] T. Tao. Recent progress in additive prime number theory, preprint. 2009. <http://terrytao.files.wordpress.com/2009/08/prime-number-theory1.pdf>

Szemerédi's theorem does not directly to the primes, because it cannot count the number of primes. Cramér's random model cannot prove any prime problems. The probability of $1/\log N$ of being prime is false. Assuming that the events " P is prime", " $P+2$ is prime" and " $P+4$ is prime" are independent, we conclude that P , $P+2$, $P+4$ are simultaneously prime with probability about $1/\log^3 N$. There are about $N/\log^3 N$ primes less than N . Letting $N \rightarrow \infty$ we obtain the prime conjecture, which is false. The tool of additive prime number theory is basically the Hardy-Littlewood prime tuples conjecture, but cannot prove and count any prime problems[6].

Mathematicians have tried in vain to discover some order in the sequence of prime numbers but we have every reason to believe that there are some mysteries which the human mind will never penetrate.

Leonhard Euler(1707-1783)

It will be another million years, at least, before we understand the primes.

Paul Erdos(1913-1996)

Of course, the primes are a deterministic set of integers, not a random one, so the predictions given by random models are not rigorous (Terence Tao, Structure and randomness in the prime numbers, preprint). Erdos and Turán(1936) contributed to probabilistic number theory, where the primes are treated as if they were random, which generates Szemerédi's theorem (1975) and Green-Tao theorem(2004). But they cannot actually prove and count any simplest prime examples: twin primes and Goldbach's conjecture. They don't know what prime theory means, only conjectures.

