

## GENERALIZATIONS OF DEGARGUES THEOREM\*

Florentin Smarandache  
University of New Mexico  
200 College Road  
Gallup, NM 87301, USA  
E-mail: smarand@unm.edu

Let's consider the points  $A_1, \dots, A_n$  situated on the same plane, and  $B_1, \dots, B_n$  situated on another plane, such that the lines  $A_i B_i$  are concurrent. Let's prove that if the lines  $A_i A_j$  and  $B_i B_j$  are concurrent, then their intersecting points are collinear.

*Solution.* Let  $\alpha$  be the plane that contains the points  $A_1, \dots, A_n$  (in the case in which the points are non-collinear  $\alpha$  is unique), and analogously, let  $\beta = P(B_1, \dots, B_n)$ , and consider  $\alpha \cap \beta = d$ .

Because the lines  $A_i A_j$  and  $B_i B_j$  are concurrent,  $A_i A_j \subset \alpha$ , and  $B_i B_j \subset \beta$ , therefore their intersection belongs to line  $d$ .

### Remark 1.

For  $n = 3$  and  $A_1, A_2, A_3$  non-collinear,  $B_1, B_2, B_3$  non-collinear, and  $A_i \neq B_j$  we obtain Desargues theorem.

### Remark 2.

An extension of this generalization is: If we consider  $A_1, \dots, A_n$  situated in a plane, and  $B_1, \dots, B_m$  situated on another plane, prove that if  $A_i A_j$  and  $B_k B_r$  are concurrent, then their intersection points are concurrent.

### Remark 3.

For  $n = m$ , and  $A_i B_i$  concurrent lines, we obtain the first generalization.

### Remark 4.

If in addition we also have  $n = m = 3$  along with the previous conditions, we obtain the Desargues theorem.

---

\* Gamma, Anul X, nr. 1-2, Oct. 1987.