

# The local fractional Hilbert transform in fractal space

Guang-Sheng Chen

Department of Computer Engineering, Guangxi Modern Vocational Technology College,  
Hechi, Guangxi, 547000, P.R. China

E-mail address: [cgswavelets@126.com](mailto:cgswavelets@126.com)(Chen)

**Abstract:** In this paper, we establish local fractional Hilbert transform in fractal space, consider some properties of local fractional Hilbert Transforms.

**Keywords:** fractal space; local fractional Hilbert transform; local fractional derivative

## 1 Introduction

In the past ten years, Local fractional calculus[1-27] has been widely applied to many fields such as mathematics , image processing and signal processing etc .Some authors have given many definitions of local fractional derivatives and local fractional integrals (also called fractal calculus) [1-16]. Hereby we rewrite the following local fractional derivative which is given by [15,16]

$$f^{(\alpha)}(x_0) = \frac{df(x)}{dx^\alpha} \Big|_{x=x_0} = \lim_{\delta x \rightarrow 0} \frac{\Delta^\alpha(f(x) - f(x_0))}{(x - x_0)^\alpha}, \quad \text{for } 0 < \alpha \leq 1, \quad (1.1)$$

where

$$\Delta^\alpha(f(x) - f(x_0)) \cong \Gamma(1 + \alpha)\Delta(f(x) - f(x_0)),$$

and local fractional integral of  $f(x)$  defined by [15-16]

$${}_a I_b^{(\alpha)} f(t) = \frac{1}{\Gamma(1 + \alpha)} \int_a^b f(t)(dt)^\alpha = \frac{1}{\Gamma(1 + \alpha)} \lim_{\Delta t \rightarrow 0} \sum_{j=0}^{N-1} f(t_j)(\Delta t_j)^\alpha, \quad (1.2)$$

with  $\Delta t_j = t_{j+1} - t_j$  and  $\Delta t = \max\{\Delta t_1, \Delta t_2, \dots, \Delta t_j, \dots\}$ , where for  $j = 1, 2, \dots, N-1$  ,  $[t_j, t_{j+1}]$

is a partition of the interval  $[a, b]$  and  $t_0 = a$  ,  $t_N = b$ .

The aim of this paper is to define local fractional Hilbert transforms based on Local fractional calculus. This paper is organized as follows. In section 2, local fractional Hilbert transforms is defined; Section 3 presents Properties of local fractional Hilbert transforms.

## 2 The local fractional Hilbert transform and some examples

In the section, both local fractional Hilbert transform and its inverse transform are defined.

**Definition 2.1** (The local fractional Hilbert transform). If  $f(x)$  is defined on the real line

$-\infty < x < \infty$  , its local fractional Hilbert transform, denoted by  $f_x^{H,\alpha}(x)$  is defined by

$$H_\alpha\{f(t)\} = \hat{f}_H^\alpha(x) = \frac{1}{\Gamma(1 + \alpha)} \oint_R \frac{f(t)}{(t-x)^\alpha} (dt)^\alpha \quad (2.1)$$

where  $x$  is real and the integral is treated as a Cauchy principal value, that is,

$$\begin{aligned} & \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{f(t)}{(t-x)^\alpha} (dt)^\alpha \\ &= \lim_{\varepsilon \rightarrow 0} \left[ \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{x-\varepsilon} \frac{f(t)}{(t-x)^\alpha} (dt)^\alpha + \frac{1}{\Gamma(1+\alpha)} \int_{x+\varepsilon}^{\infty} \frac{f(t)}{(t-x)^\alpha} (dt)^\alpha \right] \end{aligned} \quad (2.2)$$

To obtain the inverse local fractional Hilbert transform, write again (2.1) as

$$\begin{aligned} \hat{f}_H^\alpha(x) &= \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{f(t)}{(t-x)^\alpha} (dt)^\alpha \\ &= \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} f(t) g(x-t) (dt)^\alpha = f(x) * g(x) \end{aligned} \quad (2.3)$$

where  $g(x) = -\frac{1}{x^\alpha}$ , application of the local fractional Yang-Fourier transform [15,16] with respect to  $x$  gives

$$f_\omega^{F,\alpha}(\omega) = \frac{f_{H,\omega}^{F,\alpha}(\omega)}{g_\omega^{F,\alpha}(\omega)}, \quad g_\omega^{F,\alpha}(\omega) = i^\alpha \pi^\alpha \operatorname{sgn}_\alpha \omega^\alpha \quad (2.4)$$

Taking the inverse Yang-Fourier transform [15,16], we find the solution for  $f(x)$  as

$$f(x) = \frac{1}{(2\pi)^\alpha} \int_{-\infty}^{\infty} i^\alpha \pi^\alpha \operatorname{sgn}_\alpha \omega^\alpha f_{H,\omega}^{F,\alpha}(\omega) E_\alpha(i^\alpha \omega^\alpha x^\alpha) (d\omega)^\alpha$$

which is, by the Convolution Theorem [15,16]

$$f(x) = \frac{1}{(2\pi)^\alpha} \oint_R \frac{\hat{f}_H^\alpha(\xi)}{(x-\xi)^\alpha} (d\xi)^\alpha = -\frac{1}{(2\pi)^\alpha} \oint_R \frac{\hat{f}_H^\alpha(\xi)}{(\xi-x)^\alpha} (d\xi)^\alpha = -H\{\hat{f}_H^\alpha(\xi)\} \quad (2.5)$$

Obviously,  $-H_\alpha^2\{f(t)\} = -H_\alpha[H_\alpha\{f(t)\}] = f(x)$  and hence,  $H_\alpha^{-1} = -H_\alpha$ . Thus, the inverse local fractional Hilbert transform is given by

$$f(t) = H_\alpha^{-1}\{\hat{f}_H^\alpha(x)\} = -H_\alpha\{\hat{f}_H^\alpha(x)\} = -\frac{1}{(2\pi)^\alpha} \oint_R \frac{\hat{f}_H^\alpha(\xi)}{(x-\xi)^\alpha} (d\xi)^\alpha \quad (2.6)$$

**Example 2.1** Find the local fractional Hilbert transform of

$$f(t) = \frac{t^\alpha}{t^{2\alpha} + a^{2\alpha}}, \quad a > 0$$

we get, by definition,

$$\begin{aligned} \hat{f}_H^\alpha(x) &= \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{t^\alpha}{(t^{2\alpha} + a^{2\alpha})(t-x)^\alpha} (dt)^\alpha \\ &= \frac{1}{(a^{2\alpha} + x^{2\alpha})\Gamma(1+\alpha)} \left[ a^{2\alpha} \int_{-\infty}^{\infty} \frac{1}{(t^{2\alpha} + a^{2\alpha})} (dt)^\alpha + x^\alpha \int_{-\infty}^{\infty} \frac{1}{(t-x)^\alpha} (dt)^\alpha - x^\alpha \int_{-\infty}^{\infty} \frac{t^\alpha}{(t^{2\alpha} + a^{2\alpha})} (dt)^\alpha \right] \end{aligned}$$

The second and third integrals as the Cauchy principal value vanish and hence, only the first integral makes a non-zero contribution. Thus, we have

$$\hat{f}_H^\alpha(x) = \frac{a^\alpha}{(a^{2\alpha} + x^{2\alpha})} \arctan_\alpha \frac{t^\alpha}{a^\alpha} \Big|_{-\infty}^{\infty} = \frac{\pi^\alpha a^\alpha}{(a^{2\alpha} + x^{2\alpha})} \quad (2.7)$$

**Example 2.2** Find the local fractional Hilbert transform of

(a)  $f(t) = \cos_\alpha \omega^\alpha t^\alpha$  and (b)  $f(t) = \sin_\alpha \omega^\alpha t^\alpha$

$$\begin{aligned} \hat{f}_H^\alpha(x) &= \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{\cos_\alpha \omega^\alpha (t-x)^\alpha \cos_\alpha \omega^\alpha x^\alpha - \sin_\alpha \omega^\alpha (t-x)^\alpha \sin_\alpha \omega^\alpha x^\alpha}{(t-x)^\alpha} (dt)^\alpha \\ &= \cos_\alpha \omega^\alpha x^\alpha \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{\cos_\alpha \omega^\alpha (t-x)^\alpha}{(t-x)^\alpha} (dt)^\alpha - \sin_\alpha \omega^\alpha x^\alpha \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{\sin_\alpha \omega^\alpha (t-x)^\alpha}{(t-x)^\alpha} (dt)^\alpha \end{aligned}$$

which is, the new variable  $T = t - x$ ,

$$\begin{aligned} \hat{f}_H^\alpha(x) &= \cos_\alpha \omega^\alpha x^\alpha \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{\cos_\alpha \omega^\alpha T^\alpha}{T^\alpha} (dT)^\alpha \\ &\quad - \sin_\alpha \omega^\alpha x^\alpha \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{\sin_\alpha \omega^\alpha T^\alpha}{T^\alpha} (dT)^\alpha \end{aligned} \tag{2.8}$$

Obviously, the first integral vanishes because its integrand is an odd function of  $T$ . On the other hand, the second integral makes a non-zero contribution so that (2.8) gives

$$H_\alpha \{ \cos_\alpha \omega^\alpha t^\alpha \} = -\pi^\alpha \sin_\alpha \omega^\alpha x^\alpha \tag{2.9}$$

Similarly, it can be shown that

$$H_\alpha \{ \sin_\alpha \omega^\alpha t^\alpha \} = \pi^\alpha \cos_\alpha \omega^\alpha x^\alpha \tag{2.10}$$

### 3 Basic Properties of the local fractional Hilbert Transforms

**Theorem 3.1** If  $H_\alpha \{ f(t) \} = \hat{f}_H^\alpha(x)$ , then the following properties hold:

(a)  $H_\alpha \{ f(t+a) \} = \hat{f}_H^\alpha(x+a)$ , (3.1)

(b)  $H_\alpha \{ f(at) \} = \hat{f}_H^\alpha(ax)$ , (3.2)

(c)  $H_\alpha \{ f(-at) \} = -\hat{f}_H^\alpha(-ax)$ , (3.3)

(d)  $H_\alpha \{ f^{(\alpha)}(t) \} = \frac{d^\alpha \hat{f}_H^\alpha(x)}{dx^\alpha}$ , (3.4)

(e)  $H_\alpha \{ tf(t) \} = x^\alpha \hat{f}_H^\alpha(x) + \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} f(t)(dt)^\alpha$ , (3.5)

(f)  $F_\alpha \{ H_\alpha \{ f(t) \} \} = -i^\alpha \operatorname{sgn}_\alpha \omega^\alpha F_\alpha \{ f(x) \}$ , (3.6)

(g)  $\| H_\alpha \{ f(t) \} \|_\alpha = \| f(t) \|_\alpha$  (3.7)

where  $\| f(t) \|_\alpha = \sqrt{\langle f, f \rangle_\alpha}$  denotes the norm in  $L_{2,\alpha}(\mathbb{R})$ ,

(h)  $H_\alpha [f](x) = \hat{f}_H^\alpha(x)$ ,  $H_\alpha [\hat{f}_H^\alpha](x) = -f$  (Reciprocity relations), (3.8)

(i) (Parseval' s formulas).

$$\langle f, H_\alpha g \rangle_\alpha = \langle -H_\alpha f, g \rangle_\alpha \quad \text{and} \quad \langle H_\alpha f, g \rangle_\alpha = \langle f, -H_\alpha g \rangle_\alpha, \quad (3.9)$$

**Proof.** (a) We obtain, by definition ,set  $u = t + a$

$$\begin{aligned} H_\alpha \{f(t+a)\} &= \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{f(t+a)}{(t-x)^\alpha} (dt)^\alpha \\ &= \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{f(u)}{(u-(x+a))^\alpha} (du) = \hat{f}_H^\alpha(x+a) \end{aligned}$$

(b) We find, by definition ,set  $u = at$

$$H_\alpha \{f(at)\} = \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{f(at)}{(t-x)^\alpha} (dt)^\alpha = \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{f(u)}{(u-ax)^\alpha} (du)^\alpha = \hat{f}_H^\alpha(ax)$$

Similarly, result (c) can be proved.

(d) We have, by definition

$$\begin{aligned} H_\alpha \{f^{(\alpha)}(t)\} &= \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{f^{(\alpha)}(t)}{(t-x)^\alpha} (dt)^\alpha \\ &= \frac{f(t)}{(t-x)^\alpha} \Big|_{-\infty}^{\infty} - \frac{1}{\Gamma(1+\alpha)} \oint_R f(t) \left[ \frac{1}{(t-x)^\alpha} \right]^{(\alpha)} (dt)^\alpha \\ &= -\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)} \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{f(t)}{(t-x)^{2\alpha}} (dt)^\alpha = \frac{d^\alpha \hat{f}_H^\alpha(x)}{dx^\alpha} \end{aligned}$$

Proofs of (e) - (i) are similar and consequently,we omit it

**Theorem 3.2** If  $f(t)$  is an even function of  $t$ , then, an alternative form of the local fractional Hilbert transform is

$$\hat{f}_H^\alpha(x) = \frac{x^\alpha}{\Gamma(1+\alpha)} \oint_R \frac{f(t) - f(x)}{t^{2\alpha} - x^{2\alpha}} (dt)^\alpha \quad (3.10)$$

**Proof.** As the Cauchy principal value, we obtain

$$\frac{1}{\Gamma(1+\alpha)} \oint_R \frac{1}{(t-x)^\alpha} (dt)^\alpha = 0$$

consequently,

$$\begin{aligned} \hat{f}_H^\alpha(x) &= \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{f(t) - f(x)}{t^\alpha - x^\alpha} (dt)^\alpha \\ &= \frac{1}{\Gamma(1+\alpha)} \oint_R \frac{t^\alpha (f(t) - f(x))}{t^{2\alpha} - x^{2\alpha}} (dt)^\alpha + \frac{x^\alpha}{\Gamma(1+\alpha)} \oint_R \frac{(f(t) - f(x))}{t^{2\alpha} - x^{2\alpha}} (dt)^\alpha \end{aligned} \quad (3.11)$$

Since  $f(t)$  is an even function, the integrand of the first integral of (3.11) is an odd function; hence, the first integral vanishes, and (3.11) gives (3.10). Since result (2.3) discovers that the local fractional Hilbert transform can be written as a convolution transform, we state the following.

**Theorem 3.3** If  $f$  and  $g \in L_{1,\alpha}(\mathbb{R})$  are such that their local fractional Hilbert transforms are

also in  $L_{1,\alpha}(\mathbb{R})$ , Then

$$H_\alpha(f * g)(x) = (H_\alpha f * g)(x) = (f * H_\alpha g)(x) \quad (3.12)$$

and

$$(f * g)(x) = -(H_\alpha f * H_\alpha g)(x). \quad (3.13)$$

**Proof.** We obtain, by definition,  $\xi = t - y$

$$\begin{aligned} H_\alpha(f * g)(x) &= \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{1}{(t-x)^\alpha} \left[ \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} f(y)g(t-y)(dy)^\alpha \right] (dt)^\alpha \\ &= \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} f(y)(H_\alpha g)(x-y)(dy)^\alpha = (f * H_\alpha g)(x) \end{aligned}$$

Likewise, we can obtain the second result in (3.12).

To prove (3.13), we replace  $g$  by  $H_\alpha g$  in (3.12) and then use  $H_\alpha^2 g = -g$ .

#### 4 discussions

In present paper we give local fractional Hilbert transforms as follows:

$$H_\alpha\{f(t)\} = \hat{f}_H^\alpha(x) = \frac{1}{\Gamma(1+\alpha)} \int_{-\infty}^{\infty} \frac{f(t)}{(t-x)^\alpha} (dt)^\alpha \quad \text{for } 0 < \alpha \leq 1 \quad (4.1)$$

and its inverse transform

$$f(t) = H_\alpha^{-1}\{\hat{f}_H^\alpha(x)\} = -H_\alpha\{\hat{f}_H^\alpha(x)\} = -\frac{1}{(2\pi)^\alpha} \int_{-\infty}^{\infty} \frac{\hat{f}_H^\alpha(\xi)}{(x-\xi)^\alpha} (d\xi)^\alpha \quad \text{for } 0 < \alpha \leq 1 \quad (4.2)$$

The transforming functions are local fractional continuous. That is to say (in other words, it means that), it is fractal function defined on fractal sets. Hilbert transforms in integer space are the special case of fractal dimension  $\alpha = 1$ . It is a tool to deal with differential equation with local fractional derivative.

#### References

- [1] J. Tenreiro Machado, Virginia Kiryakova, Francesco Mainardi, Recent History of Fractional Calculus, Communications in Nonlinear Science and Numerical Simulations, Elsevier, vol. 16, Issue 3, pp. 1140-1153, March 2011.
- [2] J. A. Tenreiro Machado, Manuel F. Silva, Ramiro S. Barbosa, Isabel S. Jesus, Cecília M. Reis, Maria G. Marcos, Alexandra F. Galhano, Some Applications of Fractional Calculus in Engineering, Mathematical Problems in Engineering, Hindawi, vol. 2010, Article ID 639801, 2009. doi:10.1155/2010/639801.
- [3] S.G. Samko, A.A. Kilbas, O.I. Marichev: Fractional Integrals and Derivatives: Theory and Applications. Gordon and Breach, 1993.
- [4] Kolwankar K.M., Gangal A.D., Local Fractional Fokker-Planck Equation. Phys. Rev. Lett., 80, 1998, 214--217.
- [5] Babakhani A., Daftardar-Gejji V., On Calculus of Local Fractional Derivatives. J. Math. Anal. Appl., 270, 2002, 66--79.
- [6] Chen Y., Yan Y., Zhang K., On the local fractional derivative. J. Math. Anal. Appl..

362,2010,17--33.

- [7] Adda F.B., and Cresson J., About Non-differentiable Functions. *J. Math. Anal. Appl.*, 263, 2001, 721--737.
- [8] Carpinteri A., Chiaia B., Cornetti P., Static-kinematic Duality and the Principle of Virtual Work in the Mechanics of Fractal Media. *Comput. Methods Appl. Mech. Engrg.* 191, 2001, 3--19.
- [9] Gao F., Yang X., Kang Z., Local Fractional Newton's Method Derived from Modified Local Fractional Calculus. In: *The 2th Scientific and Engineering Computing Symposium on Computational Sciences and Optimization*, pp.228—232, IEEE Computer Society, 2009.
- [10] Yang X., Gao F., The Fundamentals of Local Fractional Derivative of the One-variable Nondifferentiable Functions, *World SCI-TECH R&D.*, 31, 2009, 920--921.
- [11] Yang X., Li L., Yang R., Problems of Local Fractional Definite Integral of the One-variable Non-differentiable Function. *World SCI-TECH R&D.*, 31, 2009, 722--724.
- [12] X. Yang. Applications of local fractional calculus to engineering in fractal time-space: Local fractional differential equations with local fractional derivative. ArXiv:1106.3010v1 [math-ph].
- [13] Zhong, W., Gao, F., Shen, X.: Applications of Yang-Fourier Transform to Local Fractional Equations with Local Fractional Derivative and Local Fractional Integral. In: *Proc. of 2011 International Conference on Computer Science and Education*, Springer, 2011.
- [14] S. Das. *FunctionaI Fractional Calculus for System Identification and Controls*. Springer, Heidelberg, 2008.
- [15] X. Yang, Local Fractional Integral Transforms, *Progress in Nonlinear Science*, 4(2011): 1-225.
- [16] X. Yang. *Local Fractional Functional Analysis and Its Applications*. Asian Academic Publisher Limited. 2011.
- [17] G-S. Chen, Local fractional Improper imtegral in fractal space, *Advances in Information Technology and Management* Vol. 1, No. 1, March 2012
- [18] G-S. Chen, Mean Value Theorems for Local Fractional Integrals on Fractal Space, *Advances in Mechanical Engineering and its Applications* Vol. 1, No. 1, March 2012.
- [19] G-S. Chen, A generalized Young inequality and some new results on fractal space, *Advances in Computational Mathematics and its Applications* , Vol. 1, No. 1, March 2012.
- [20] G-S. Chen, Local fractional Mellin transform in fractal space, *Advances in Electrical Engineering Systems*, 1(2)(2012) 89-95.
- [21] G-S. Chen, Fourier Cosine and Sine Transform on fractal space, <http://arxiv.org/abs/1110.4756>, 8 pages.
- [22] G-S. Chen, Generalizations of Holder's and some related integral inequalities on fractal space, <http://arxiv.org/abs/1109.5567>, 8 pages.
- [23] G-S. Chen, The finite Yang-Laplace Transform in fractal space, <http://viXra.org/abs/1203.0038> , 4 pages.
- [24] G-S. Chen, The local fractional Stieltjes Transform in fractal space, *Advances in Intelligent Transportation Systems*, 1(1)(2012)29-31.
- [25] Yang, X.J., Local fractional partial differential equations with fractal boundary problems, *Advances in Computational Mathematics and its Applications*, 1(1) (2012) 60-63.
- [27] Yang, X.J. A short introduction to Yang-Laplace Transforms in fractal space, *Advances in Information Technology and Management*, 1(2)(2012) 38-43.