

Can we form Gravitinos by something other than a Higgs Boson in the EW era ?

A. Beckwith¹

1) abeckwith@uh.edu, Chongqing University Department of Physics; Chongqing, PRC, 400044

Abstract

What is the physical nature of gravitinos? As asked before, this question was the template of how to introduce Machian Physics as a way to link gravitinos in the electro weak era and gravitons as of the present. What we wish to do now is to ask how a flaw in the Higgs equation as brought up by Conway shows a branch off from orthodox quantum physics, leading to, with the Machs principle application done earlier a way to embed the beginning of the universe as a semi classical super structure of which QM is sub set of. We argue that this will necessitate a review of the Higgs equation of state for reasons stated in the manuscript. We also finally review a proposal for another form of mass formation mechanism as a replacement for the Higgs mass as introduced by Glinka and Beckwith, 2012, with commentary as to how suitable it may be to get a gravitino mass in fidelity to the Machian proposal introduced by Beckwith previously, to get linkage between electroweak era gravitinos and present day gravitons.

Introduction

We will ask the question here. In an earlier document, the author presented an equivalence between Gravitinos in the electro weak era, and Gravitons today. The motivation of using two types of Machs principle, one for the Gravitinos in the electro weak era, and then the 2nd modern day Mach's principle, as organized by the author are as seen in [1]

$$\frac{GM_{\text{electro-weak}} \Big|_{\text{Super-partner}}}{R_{\text{electro-weak}} c^2} \approx \frac{GM_{\text{today}} \Big|_{\text{Not-Super-Partner}}}{R_0 c^2} \quad (1)$$

are really a statement of information conservation. What we now ask is if the Gravitino can be re stated in terms in fidelity to quantum mechanics, or if some other theoretical construction must be used. The motivation for asking this question will be seen in examining if the Gravitino, as in the mass in the left hand side of Eq. (1) , as it materializes due to Conway's [2] presentation as to defects in the Higgs equation of state, is in fidelity with QM principles. If not, then what would replace it?

In this document, we will review what can be said about the class of equations as given by Conway which are well behaved experimentally, namely the Dirac Equation, and review the current Higgs eqn theory which would underly the creation of Gravitinos, in the electro weak era.

The Conway article [2] has its modus operandi in calling every theory as to field theory not congruent to the Dirac Equation as a failure. The Higgs theory in that sense has a 'leftover' term of $\dot{\phi}^* \cdot \dot{\phi}$, which has to be subtracted from a Hamiltonian H_{Higgs} in order to reduce the formation and evolution of ϕ in Hamiltonian mechanics to be in fidelity to the physics given by a Higgs modified style Hamiltonian , call it $H_{\text{Higgs-changed}}$ so there is a

$$H_{\text{Higgs-changed}} = H_{\text{Higgs}} - \beta \cdot \dot{\phi}^* \cdot \dot{\phi} \quad (2)$$

And

$$i \frac{d\phi_{\text{Higgs-Gravitino}}}{dt} = H_{\text{Higgs-changed}} \phi_{\text{Higgs-Gravitino}} \quad (3)$$

In fidelity with the physics evolution of

$$i \frac{d\phi_{Dirac}}{dt} = H_{Dirac} \phi_{Dirac} \quad (4)$$

Whereas what is observed is, instead [2]

$$i \frac{d\phi_{Higgs-Gravitino}}{dt} \neq H_{Higgs} \phi_{Higgs-Gravitino} \quad (5)$$

To further elucidate this question, we will also ask if there is a way to encapsulate H_{Higgs} in Eq. (2) above in the methodology of constructing QM within a larger, semi classical theory. . As given in the 5th Dice 2010 work shop, as given by Elze, Gambarotta and Vallone [3] there is a speculated ensemble theory involving a “ $\hat{\zeta}$ of

$$i\partial_t \text{"state"} = \hat{\zeta} \text{"state"} \quad (6)$$

The end result is , after a Fourier transform re casting the Eq. (6) in terms of a matrix equation looking like

$$i \cdot \partial_t \rho_{jk} = \sum_{l,m} \zeta_{jk,lm} \rho_{lm} = \sum_{l,m} \left[\zeta_{jk,lm} = H_{jl} \delta_{mk} - \delta_{jl} H_{mk} \right] \rho_{lm} \quad (7)$$

We will discuss Eq. (7) in a generalized incantation in APPENDIX A which will show as that the QM type interactions require a most specialized potential, as either a constant, or a Harmonic potential, with others not suitable, if we wish to extract QM from the results of Eq. (7), and from there to comment upon candidate equations which may be a way to contain H_{Higgs} as far as a generalized theory which may contain QM (Dirac) type behavior . If not, then Eq. (2) does not qualify as far as having H_{Higgs} reduced to a QM sub set and we must then go to the Conway description of the Higgs equation used to define the creation of/ evolution of the Gravitino as faulty physics, needing an abrupt fix to reduce it to the form of Eq. (2) to salvage QM .

Appendix B brings up the relevance of the Dirac Eq to the critique which Conway [2] brings to the discussion of a proper equation for a well behaved experimentally verified equation. We add an example of how early universe Lorentz violation is equivalent to the break up of the fidelity of the Higgs term, and in fact, the Eq. (B12) presented in Appendix B is in its behavior (if it were 10 orders of magnitude larger, i.e. as a Torsion term added in) very similar to the problem outlined in Eq(B6) in the Higgs potential, I .e. note in Eq (B3) with the unwanted $\dot{\phi}^* \dot{\phi}$ term which blocks the Higgs equation of state from having the good behavior postulated by Conway[2] in his Claims 1 , 2 and 3 as given in **Appendix B** below. Note also that the problem as outlined in $\dot{\phi}^* \dot{\phi}$ term shows up in an even more glaring fashion in the incredibly complicated Lagrangian specified for the formation of Gravitinos in the early universe. We will get to that next. It is useful to compare these ideas with what J. Lee published recently [4]

Examining the formation of Gravitinos in the early universe

In [5] the density is given by, if g_* is for early universe degrees of freedom

$$\rho = g_* \left[\frac{\pi^2}{30} \right] \cdot T^4 \quad (8)$$

With a resulting Hubble rate for the radiation era as written as for $H(T)$, radiation era, as

$$H(T) = \sqrt{\frac{g_* \cdot \pi^2}{90}} T^2 \quad (9)$$

The early Gravitino relic density is then given by an expression

$$\Omega_{\tilde{G}}^{TP} h^2 = \sum_{\alpha=1}^3 \left(1 + \frac{M_{\alpha}(T_R)^2}{3 \cdot m_{\tilde{G}}^2} \right) \cdot \omega_{\alpha} \cdot g_{\alpha}(T_R)^2 \cdot \ln \left(\frac{k_{\alpha}}{g_{\alpha}(T_R)} \right) \left(\frac{m_{\tilde{G}}}{100 GeV} \right) \left(\frac{T_R}{10^{10} GeV} \right) \quad (10)$$

This is , in terms of re heating temperature very close to linear in growth due to scaling with a re heating temperature T_R . One obtains an approximately linear growth rate in terms of gravitino density with a most complicated Lagrangian density function which is in the top of section 2.2. of [5] is so complicated that one cannot, even in linear approximations of it get either a classical or a quantum analogy in terms easily identifiable terms of page 5 of this PhD dissertation. We will review in Appendix A the DICE 2010 [3] article treatment of QM in a larger non linear theory [3] , and in Appendix B the Comway [2] treatment in terms of lagrangian density both for the Dirac Eq, and also for the Higgs, and then from there make the case necessary as to if the Gravitino is QM in its construction. Or not.

Getting the template as to keeping information content available for Eq. (10) right and its implications for Eq. (1) and Eq. (4), and Eq. (5). Yielding a new expression of Gravitino mass in the EW regime ?

The Machian hypothesis and actually Eq. (10) are a way to address a serious issue, I.e. how to keep the consistency of physical law intact, in cosmological evolution [1] . Another significant issue is the following. How to reconcile the Comway hypothesis [2] and postulates , as given in Appendix B , and also the DICE 2010 delination of QM as in Appendix A either requiring a zero valued potential, a constant potential, or a potential with quadratic flavor to delineate clear QM behavior [3] . If these potential field requirements are not met, as given by Appendix A [3] , then one has to ask if a Higgs mechanism in fidelity with Appendix B [2] can be constructed for an allegedly optimal experimental modeling of mass formation.

Eq. 10, which has neither a zero valued potential, a linear or a quadratic potential is clearly NOT in sync with the DICE 2010 Appendix A treatments leading to QM , alone.[3]

Eq. 10 does NOT have fidelity with the sort of Comway criteria[2] as given in **Appendix B** as to a potential energy which is most likely to have optimal match up with experimental data as cited by the Dirac equation results as given in the Comway article.

Either Eq. (10) signifies that there is no match up with the sort of evolution equation (for creation of a Gravitino in the electro weak era) as exemplified by the Dirac Equation which Comway likes so much [2] , or we have to go to live with the results as given by Appendix B, that what we are seeing in the Gravitino in the Electro weak era is QM contained in a larger non linear theory, as Elze et al seem to be inferring [3] as brought up in Appendix A .

Another approach. How about a new method for obtaining in the EW era mass without the Higgs ?

What we can look at is the Glinka-Beckwith [6] proposal as to a new mass formation process, which may show a different way to examine potential systems, as opposed to the either-or criteria as given by Appendix A [3] and Appendix B [2] below. To do so, note that the article as given in.

$$|m|^2 = \left(\frac{\hbar \cdot \omega}{c^2} \right)^2 - \left(\frac{\hbar \bar{k}}{c} \right)^2 \quad (11)$$

We can treat the k as a wave ‘vector’, and look at the term $\hbar \cdot \omega$ as an energy term. Dependent upon how we interpret $\hbar \cdot \omega$, i.e. as a per unit interpretation of energy, we could reconcile a treatment of a physically averaged out quantity of the potential energy as given in [5] is contained via the correct effective Lagrangian for light gravitinos $\zeta_{\psi, \text{int}}^{(\alpha)}$, which is Eq.(2.82), page 22 of Pradler’s dissertation [5] for obtaining gravitino interactions with ordinary matter fields.

We can, to first order model the potential to be looked at in the Gravitino-matter field interaction as [5]

$$V_{\psi, \text{int}}^{(\alpha)} = i \frac{\bar{\psi}_{\mu} \cdot [\gamma^{\rho}, \gamma^{\sigma}] \cdot \gamma^{\mu} \cdot \lambda^{(\alpha)a} F_{\rho\sigma}^{(\alpha)a}}{8M_{Pl}} \quad (12)$$

This Eq. (12) is the potential energy term of Eq.(2.82), page 22 of Josef Pradler’s [5] dissertation, and we argue that the physics of the gravitino, as interacting with matter in the electro weak regime, can be to first order, averaged out to be an energy which can be then made equivalent to $\hbar \cdot \omega$ of Eq. (11). We argue then that effectively, in early universe conditions that we are looking at, then [6],

$$|m|^2 \Big|_{EW} \sim \left(\frac{\hbar \cdot \omega}{c^2} \right)^2 \propto \left(\frac{\langle V_{\psi, \text{int}}^{(\alpha)} \rangle}{c^2} \right)^2 + \text{vanishingly - small - terms} \quad (13)$$

$$M = im$$

Then, if we do Eq. (13) in this spirit, we can then go to what Glinka-Beckwith wrote [6] and look at

$$M \Big|_{EW} = \left(\frac{\hbar \cdot \omega = E = \langle V_{\psi, \text{int}}^{(\alpha)} \rangle}{c^2} \right) - \left(\hat{i} \cdot \frac{p_x}{c} + \hat{j} \cdot \frac{p_y}{c} + \hat{k} \cdot \frac{p_z}{c} \right) \sim \left(\frac{\hbar \cdot \omega = E = \langle V_{\psi, \text{int}}^{(\alpha)} \rangle}{c^2} \right) \Big|_{EW} \quad (14)$$

Ultimately, the analysis of terms as specified in a gravitino-EW ‘matter’ regime would specify the exact particulars as to Eq. (12). We will also venture a first order approximate description as to why the mass of the Graviton in the later regime of space time, near the present would be so much smaller than the Gravitino.

Conclusion

Via use of the Glinka-Beckwith approximation for the formation of Mass, we have come up with a criteria where the Gravitino interaction with space-time physics in the electro weak, as outlined above, can be construed as either embedded within a larger theory than QM, as suggested by Elze et al. [3], or a corrected Higgs mass formation [2], or something else, which has to be constructed. As outlined by Beckwith[1] there is room to delineate if such a gravitino, using some of the field theoretic construction as given by [5] will be either classically embedded, or something else. The formalism as to massive graviton distortion of early universe space time, as given in [7], and [8] needs to be developed more fully, and we hope we can experimentally test if t’Hoofts supposition about QM [9] is falsifiable experimentally, and analytically, in this early universe setting, as brought up by the author [1]

BIBLIOGRAPHY

- [1] A.W. Beckwith, "Gravitons, Gravitinos, and Using Machs Principle to Make a Statement of Equivalent Information in a Gravitino-Graviton Correspondence of Critical Information Exchange from Electro-Weak to Present Era", <http://vixra.org/abs/1204.0089>
- [2] Eliahu Comway, "Physical Consequences of Mathematical Principles", Progress in Physics, Vol 4, 91-98, 2009
- [3] H-T Elze, G Gambarotta and F. Vallone, "General Linear Dynamics –quantum, classical or hybrid", part of the IOP conference proceedings of the DICE 2010 meeting, <http://iopscience.iop.org/1742-6596/306/1/012010>
- [4] **Jae-Weon Lee**, "On the Origin of Entropic Gravity and Inertia", <http://arxiv.org/abs/1003.4464>; Foundations of physics, Volume 42, 2012
- [5] J. Pradler, <http://arxiv.org/pdf/0708.2786.pdf> Thesis
- [6] A.Beckwith,L.Glinka, "The Quaternionic Particle Mass", Prespacetime Journal| February 2012 j Vol. 3 j Issue 2 j pp. 126-130
- [7] Kurt Hinterbichler, "Theoretical Aspects of Massive Gravity" <http://arxiv.org/abs/1105.3735.pdf>
- [8] <http://web.mit.edu/redingtn/www/netadv/Xgravitati.html>
- [9] G. t'Hooft, "Quantum Mechanics as a dissipative deterministic system" http://arxiv.org/PS_cache/gr-qc/pdf/9903/9903084v3.pdf
- [10] J. Bjorken, and S. Drell, "Relativistic Quantum Mechanics", McGraw Hill, 1964
- [11] J. Bjorken, "Emergent Photons and Gravitons: The Problem of vacuum structure"; arXIV 1008.0033 v1 [hep-th] , July 30,2010

Appendix A

Elze et al DICE 2010 Summary as to quantum mechanics embedded in a larger non linear classical theory

This discussion serves to bring up a Quantum like version of the Liouville equation and to from there to also make sense of the given equation, as of the main text:[3]

$$i\partial_t \text{"state"} = \hat{\zeta} \text{"state"} \quad (6)\text{-main text}$$

To begin with , look at a generic Hamiltonian as given by

$$H(x, p) = \frac{1}{2} p^2 + V(x) \quad (A1)$$

This Hamiltonian is incorporated in the Liouville equation of motion

$$-\partial_t \rho = \frac{\partial H}{\partial p} \cdot \frac{\partial \rho}{\partial x} - \frac{\partial H}{\partial x} \cdot \frac{\partial \rho}{\partial p} = \left\{ p \cdot \partial_x - \frac{dV}{dx} \cdot \partial_p \right\} \rho \quad (A2)$$

The upshot if a Fourier transform is taken of Eq. (A2) above, and the space like co – ordinates of

$$\begin{aligned} Q &:= x + y / 2 \\ q &:= x - y / 2 \end{aligned} \quad (A3)$$

Eqn(A1) then becomes

$$i\partial_t \rho = \left\{ \hat{H}_Q - \hat{H}_q + \varepsilon(Q, q) \right\} \rho \quad (A4)$$

The term put in, namely $\varepsilon(Q, q)$ which retrieves if we have classical or quantum information, and also, note

$$\hat{H}_\chi = \frac{-1}{2} \partial_\chi^2 + V(\chi) \quad (A5)$$

And

$$\varepsilon(Q, q) := (Q - q) \cdot \left[\frac{d}{dt} V \left(\frac{Q + q}{2} \right) \right] - V(Q) + V(q) = -\varepsilon(q, Q) \quad (A6)$$

Then,

$$\varepsilon(Q, q) := 0 \Leftrightarrow V = \text{const}, V \sim \text{linear}, V \sim \text{harmonic} \quad (A7)$$

If so, then one can write

$$i\partial_t \rho = \{ \hat{H}_Q - \hat{H}_q \} \rho \quad (A8)$$

I.e. then we have that for the potentials represented by Eq. (7), there is an overlap between classical and quantum versions of the Liouville equation as given by the Von Neuman equation as presented by

$$i\partial_t \hat{\rho} = [\hat{H}, \hat{\rho}] \quad (A9)$$

In so many words, we have a QM type situation guaranteed if Eq. (A7) holds, whereas we can solve a more general theoretical construction in which there may be what is known as a super action given by

$$S := \int_{t_0}^t d\tau \cdot \left(\frac{m}{2} \cdot \dot{Q}^2 - V(Q) - \left[\frac{m}{2} \cdot \dot{q}^2 - V(q) \right] - \varepsilon(Q, q) \right) \quad (10)$$

We then will be stuck with working with Eq. (A4)

When the super action is reduced to , with

$$\varepsilon(Q, q) := 0 \Leftrightarrow V = \text{const}, V \sim \text{linear}, V \sim \text{harmonic} \quad (A7)$$

$$S := \int_{t_0}^t d\tau \cdot \left(\frac{m}{2} \cdot \dot{Q}^2 - V(Q) - \left[\frac{m}{2} \cdot \dot{q}^2 - V(q) \right] \right) \quad (A11)$$

We recover Eq.(A9)

In short, the restrictions on the potential energy, as given by Eq. (A7) are essential for the formation of QM, for exact quantum mechanical Hilbert space operators, whereas more general cases with $\varepsilon(Q, q) \neq 0$

APPENDIX B.

Problems with the Higgs eqn., lectured upon in Chongqing University, November 2011.

We summarize the main point of Comway's article [2] in terms of their relationship to the Dirac equation and the question of what is the optimal form of a physics equation most in fidelity to experimental measurements.

The initial points of this borrowing from Comway have already been made in Eq. (2) to Eq.(5) so we will be discussing the action integral interpretation which Comway made, which was his primary way to differentiate between the faulty mathematics as he saw in the Higgs equation and the Dirac equation. We will reproduce his arguments as to that interpretation in this appendix.

$$S = \int \zeta(\psi_1, \psi_2) \cdot d^4x \quad (\text{B1})$$

Here, $\zeta(\psi_1, \psi_2)$ is a Lagrangian density function which is a Lorentz scalar, so then Eq. (B1) is a Lorentz scalar.

The consequences that Eq. (B1) is a Lorentz scalar lead to several claims by Comway to follow upon and to use.

CLAIM 1:

1. A relativistically consistent quantum theory may be derived from Lagrangian density $\zeta(\psi_1, \psi_2)$ which is a Lorentz scalar.
2. An acceptable dimension for a Lagrangian density is of the form $[L^{-4}]$
3. A wave functional $\psi(x^\mu)$ for both $\zeta(\psi_1, \psi_2)$ and S cannot define a composite particle **if** x^μ is for a single four dimensional point in space time

Sub claim to 3 above, and an effective re statement of 3 is : **If** $\psi(x^\mu)$ were for a single (not composite) particle, then

3*. A: $\psi(x^\mu)$ needs space time co – ordinates of its center of energy

3*. B: One needs additional co – ordinates for describing internal structure.

We shall then go to the next specific Comway Claim, namely

CLAIM 2

Use the following procedure to get consistency of a quantum (massive particle) theory with a classical (massive particle) particle theory, namely by using the following field equation, as given by

$$\frac{\partial}{\partial x^\mu} \cdot \frac{\partial \zeta}{\partial \left(\frac{\partial \psi}{\partial x^\mu} \right)} - \frac{\partial \zeta}{\partial x^\mu} = 0 \quad (\text{B2})$$

For energy start off with the equation given by the 2nd order tensor, $T_{\mu\nu}$, with T_{00} the energy density , and $T_{\mu\nu}$ having L^{-4} dimensions , with

$$T_{\mu\nu} = \frac{\partial \zeta}{\partial \left(\frac{\partial \psi}{\partial x^\mu} \right)} \cdot \frac{\partial \zeta}{\partial x^\nu} - \zeta g_{\mu\nu} \quad (\text{B3})$$

Sub set of CLAIM 2. In QM, the Hamiltonian is equal to the total energy, so we can write T_{00} as the Hamiltonian density

$$T_{00} = H_{\text{Hamiltonian-density}} = \dot{\psi} \cdot \frac{\partial \zeta}{\partial \dot{\psi}} - \zeta \quad (\text{B4})$$

Eq. (B4) satisfies the continuity equation as given by

$$j_{,\mu}^\mu = 0 \quad (\text{B5})$$

Then either of the two happen:

- A. Hamiltonian density $H_{\text{Hamiltonian-density}}$ may be used to extract Hamiltonian H so that one can write a Hamiltonian H so that then the following happens: Energy E is an eigenvalue of ψ

$$H\psi = E\psi \quad (\text{B6})$$

And the De Broglie functions hold as given by

$$i \frac{\partial \psi}{\partial t} = E\psi \Rightarrow i \frac{\partial \psi}{\partial t} = H\psi \quad (\text{B7})$$

So then the Hilbert space is formed using all ψ of H (completeness of the Hilbert space, using basis from ψ)

OR

- B. Use expression for density to form inner product for inner product of ψ and construct an orthonormal basis set (often using Gram Schmitz orthogonalization) for orthonormal basis for corresponding Hilbert space.

Then, after B, to then look at a matrix equation given by

$$H_{i,j} = \int H_{\text{Hamiltonian-density}} (\psi_i; \psi_{i,\mu}; \psi_j; \psi_{j,\nu}) \cdot dx^3 \quad (\text{B8})$$

Form a matrix from Eq. B8, and then diagonalize this matrix to get eigenvalues ψ and ENERGY eigenvectors .

CLAIM 3.

Procedures from CLAIM 1 and CLAIM 2, give the same eigenvalues and eigenvectors, SAME information.

CLAIM 4

The following Equations give almost the same information, one QM, and the other CM (Quantum versus Classical)

$$i \frac{\partial \psi}{\partial t} = E\psi \Rightarrow i \frac{\partial \psi}{\partial t} = H\psi \quad (\text{B7})$$

$$\frac{\partial}{\partial x^\mu} \cdot \frac{\partial \zeta}{\partial \left(\frac{\partial \psi}{\partial x^\mu} \right)} - \frac{\partial \zeta}{\partial x^\mu} = 0 \quad (\text{B2})$$

Applications of this formulation. See the Dirac Eq. as given by Bjorken And Drell ,[10] , plus Conway [2]

This example works beautifully. Pion physics, Quark physics and more. There is an excellent match up with experiment.

Next application, Higgs equation, so that

$$E \Big|_{\text{Higgs}} \neq H_{\text{Hamiltonian-density}} \Big|_{\text{Higgs}} = \dot{\phi} \cdot \frac{\partial \zeta}{\partial \dot{\phi}} - \zeta H \sim \{ \} + \dot{\phi}^* \dot{\phi} \quad (\text{B3})$$

Here we see then that

$$i \frac{\partial \psi}{\partial t} = E\psi \text{ DOES NOT } \Rightarrow i \frac{\partial \psi}{\partial t} = H\psi \quad (\text{B4})$$

Specifically, for the Higgs, one has

$$\zeta(\text{Higgs}) \equiv \phi_{,\mu}^* \phi_{,\nu} g_{\mu\nu} + L.O.T. \quad (\text{B5})$$

Eq(B5) will then lead to a Higgs potential energy looking like, in simplest form. Where we only know the ratio of $\mu^\otimes / |\lambda|$.

$$V(\text{Higgs})[\phi^\dagger \phi] = \mu^\otimes (\phi^\dagger \phi) + |\lambda| (\phi^\dagger \phi)^2 \quad (\text{B6})$$

And we get a vacuum state given by

$$\langle \phi \rangle_0 = \begin{pmatrix} 0 \\ \sqrt{-\mu^\otimes / |\lambda|} \end{pmatrix} \quad (\text{B7})$$

For the Higgs nucleation of mass, for a Graviton, we have a huge problem, i.e. many undetermined coefficients.

This is similar to what happens with Bjorken's work with

“Emergent Photons and Gravitons: The Problem of vacuum structure ”;
arXIV 1008.0033 v1 [hep-th] , July 30,2010

Let $H(DE)$ be the Hubble rate of expansion of the cosmos, and set a scale factor as

$$a(t) = \exp(-H(DE) \cdot t) \quad (\text{B8})$$

Here we can re phrase H_{oc} as being the Hubble rate of expansion without torsion added in. Also

$$H(DE) = H_{oc} - \left[(4\pi \cdot \gamma \cdot \rho_A)^2 / M_{Pl}^4 \cdot (1 + \gamma^2) \right] \quad (\text{B9})$$

If we go to the Zeldovich relationship

$$\left[(4\pi \cdot \gamma \cdot \rho_A) / M_{Pl}^0 \cdot (1 + \gamma^2)^{1/2} \right] \sim (\Lambda_{QCD} = 10^{-20} M_{Pl})^3 \quad (\text{B10})$$

Then we get a Lorentz violating ‘‘Lagrangian’’ added on term looking like, if

$$b_\mu = \frac{\eta_\mu 2\pi\rho_A\gamma}{M_{Pl}(1+\gamma^2)} \leq 10^{-33} eV \quad (\text{B11})$$

$$\dot{L} = b_\mu \bar{\psi} \gamma^\mu \gamma_5 \psi \quad (\text{B12})$$

This Eq. (B12) is a ten orders too small Lorentz violation term, in the Potential for a Lagrangian, for space time emergence, but if it were larger, it would be similar in effect to the problem with the Higgs which Comway is outlining. Very close.