

The arithmetic of binary representations of even positive integer $2n$ and its application to the solution of the Goldbach's binary problem

ALEXANDER FEDOROV

Abstract

One of causes why Goldbach's binary problem was unsolved over a long period is that binary representations of even integer $2n$ (BR $2n$) in the view of a sum of two odd primes (VSTOP) are considered separately from other BR $2n$. By purpose of this work is research of connections between different types of BR $2n$. For realization of this purpose by author was developed the "Arithmetic of binary representations of even positive integer $2n$ " (ABR $2n$). In ABR $2n$ are defined four types BR $2n$. As shown in ABR $2n$ all types BR $2n$ are connected with each other by relations which represent distribution of prime and composite positive integers less than $2n$ between them. On the basis of this relations (axioms ABR $2n$) are deduced formulas for computation of the number of BR $2n$ (NBR $2n$) for each types. In ABR $2n$ also is defined and computed Average value of the number of binary sums are formed from odd prime and composite positive integers $< 2n$ (AVNBS). Separately AVNBS for prime and AVNBS for composite positive integers. We also deduced formulas for computation of deviation NBR $2n$ from AVNBS. It was shown that if n go to infinity then NBR $2n$ go to AVNBS that permit to apply formulas for AVNBS to computation of NBR $2n$. At the end is produced the proof of the Goldbach's binary problem with help of ABR $2n$. For it apply method of a proof by contradiction in which we make an assumption that for any $2n$ not exist BR $2n$ in the VSTOP then make computations at this conditions then we come to contradiction. Hence our assumption is false and for all $2n > 2$ exist BR $2n$ in the VSTOP.

1 Introduction

On 7 June, 1742, the Prussian mathematician Christian Goldbach wrote a letter to Leonhard Euler in which he proposed the following conjecture: Every integer greater than 2 can be written as the sum of three primes. He considered 1 to be a prime number. In mathematics, a prime number is a natural number that has exactly two natural number divisors, which are 1 and the prime... , a convention subsequently abandoned. A modern version of Goldbach's original conjecture is: Every integer greater than 5 can be written as the sum of three primes. Euler, becoming interested in the problem, replied by noting that this conjecture is equivalent with another version: Every even integer greater than 2 can be written as the sum of two primes. Euler's version is the form in which the conjecture is usually expressed today. It is also known as the strong, even, or binary Goldbach's conjecture.

2 Method

Researching great number proposed solutions of this problem it can make conclusion that analytical methods exhausted their resources since neither one of them don't bring to purpose. And for solution of Goldbach' binary problem it needs to develop new method . Considered all representations of even integer $2n$, and not only binary representations in the view of a sum of two odd primes (BRVSTOP). It can make conclusion that to BRVSOTP no approach but it is for other binary representations of $2n$ for example in the view of a sum of two odd composite integers (BRVSTOCI) or in the view of a sum of two odd integers one is composite other is prime (BRVSTOICP). So came idea to develop the arithmetic of binary representations of even positive integer $2n$ (ABR $2n$). In which it was found the connection between types of binary representation of even integer $2n$ (BR $2n$). Which underlie of axioms of ABR $2n$. Thus the New method as follows : To ABR $2n$ is used standard procedure : the proof of contradiction i.e. don't break generality we make assumption that for any value of $2n$ it is not the BVRSTOP . Next we compute all corollaries from it into ABR $2n$. Specifically from it appears two identities which were not the early. From these identities it follows that to the each number of noncomposite ("1" + primes) $< 2n$ (p) correspond to infinite set of $2n$ that contradict by proposition proved in ABR $2n$ which assert that each p correspond to finite set of $2n$ included between neighboring values of p . Next it follows standard conclusion that our assumption that the BVRSTOP are not for

some value is false and it is true that the BVRSTOP are for all $2n > 2$. Hence $\forall 2n > 2$ exists at least one representation of even positive integer $2n$ in the view of a sum of two odd prime positive integers or "1" and odd prime positive integer.

3 The arithmetic of binary representations of even positive integer $2n$

3.1 General conception

Definition 3.1 *The binary representations of even positive integer $2n$ in $ABR2n$ are defined as bijective mappings :*

$$f : X \rightarrow Y \quad (1)$$

$$y = 2n - x \quad (2)$$

Where:

$$X = \{x | x \in N, 1 \leq x < n\} \quad (3)$$

$$Y = \{y | y \in N, n < y < 2n\} \quad (4)$$

n - positive integer .

$$|X| = |Y| \quad (5)$$

Definition 3.2 *The set of binary representations of even positive integer $2n$ ($SBR2n$) are defined as follows :*

$$SBR2n = \{1 + (2n - 1) = 2n ;$$

$$2 + (2n - 2) = 2n ;$$

...

$$n - 1 + (2n - (n - 1)) = 2n\}$$

The last is got by represent (2)

in the view of $x + y = 2n$.

Definition 3.3 *The set XUY consist of: even positive integers , odd positive integers inclusive odd composite positive integers and noncomposite positive integers (primes and "1").*

Remark 3.1 *Since n is not mapped into Y and it is mapped into itself (automorphism) therefore mapping $n \rightarrow n$ and corresponding binary representation $n + n = 2n$ do not enter into $SBR2n$ which is formed only from bijective mappings. By this reason n do not enter into XUY . But it is not*

denoted that as a result of the exception of binary representations: $2+2=4$ and $3+3=6$ from $SBR2n$, 4 and 6 have not binary representations in the view of a sum of two noncomposite positive integers. But it is not the case. In $SBR2n$ is in existence the binary representations $1+3=4$ and $1+5=6$. It is significant "1" with primes are related in $ABR2n$ to noncomposite positive integers. At that special status of "1" in N is ignored in $ABR2n$.

Definition 3.4 s_e - the number of even composite positive integers into $SBR2n < 2n$.

$$s_e - \text{positive integer} > 0 \quad \forall 2n > 6$$

$$s_e = 0 \quad \forall 2 < 2n < 8.$$

Definition 3.5 s_o - the number of odd composite positive integers into $SBR2n < 2n$.

$$s_o - \text{positive integer} > 0 \quad \forall 2n > 8$$

$$s_o = 0 \quad \forall 2 < 2n < 10.$$

Definition 3.6 p - the number of odd noncomposite integers (primes and "1") into $SBR2n < 2n$.

$$p - \text{positive integer} > 0 \quad \forall 2n > 2.$$

Corollary 3.1 By reason of exclusion n from the set XUY if n is even then n or s_e are decremented by two (since it is excluded $n + n = 2n$):
 $n^* = n - 2$; $s_e^* = s_e - 2$.

Corollary 3.2 By reason of exclusion n from the set $X U Y$ If n is odd then n or s_o are decremented by two:
 $n^* = n - 2$; $s_o^* = s_o - 2$.

Corollary 3.3 By reason of exclusion n from the set $X U Y$ If n is odd prime then n or p are decremented by two:
 $n^* = n - 2$; $p^* = p - 2$.

Remark 3.2 It needs to say that corollaries 3.1 , 3.2 , 3.3 are took into account automatically at forming $SBR2n$ since it makes until appearance of representation in the view of $n + n = 2n$ which throw-off . Thus if n is odd or odd prime then n, s_o automatically are decremented by two at the direct computation them into $SBR2n$. If n is even then n, s_o, p are not decremented since s_o, p by definitions related to odd composite and noncomposite integers.

Remark 3.3 *In the equations which include n is even n, s_o, p are not decremented (see rem 3.2).*

At the computations which related to s_e n is decremented by two.

Remark 3.4 *In the equations which include n is odd or odd prime then it is decremented by one since n enters in the equation with value which is required the correction.*

Proposition 3.1 $\forall n > 1$ always is fulfilled the condition $|X| = |Y|$.

Proof 3.1 *Taking into account that $2n$ do not enter into the set XUY and n is excluded from the set XUY (see remark 3.1) then we have :*

$$|XUY| = 2n - 2 = 2(n - 1) \quad (6)$$

Hence $|XUY|$ is even for any n and thus $\forall n > 1$ always is fulfilled the condition $|X| = |Y|$. \square

3.2 The types of binary representations even positive integer $2n$

In depend of that are x, y in the view of $x + y = 2n$ prime or composite it can be four types binary representations of even positive integer $2n$:

Definition 3.7 *There is Type "H" if x -odd prime positive integer or "1" and y - odd prime positive integer.*

$|H|$ - the number of binary representations of Type "H".

$|H|$ - positive integer $> 0 \forall 2n > 2$ (see theorem (10.1.)).

Definition 3.8 *There is Type "Q" if x -odd composite positive integer; and y -odd composite positive integer .*

$|Q|$ - the number of binary representations of Type "Q".

$|Q|$ - positive integer $> 0 \forall 2n > 22$

excepting $2n = 26; 28; 32; 38$; in which $|Q| = 0$

(see numerical solution $|Q| = 0$ subsection (12.2) .

$|Q| = 0 \forall 2 < 2n < 24$.

Definition 3.9 *There is Type "L" if x - odd prime positive integer or "1" and y -odd composite positive integer or x -odd composite positive integer and y - odd prime positive integer.*

$|L|$ - the number of binary representations of Type "L".

$|L|$ - positive integer $> 0 \forall 2n > 8$.

$|L| = 0 \forall 2 < 2n < 10$.

Definition 3.10 *There is Type "E" if x -even positive integer and y -even positive integer .*

$|E|$ - *the number of binary representations of Type"E".*

$|E|$ - *positive integer $> 0 \forall 2n > 4$.*

3.3 The axioms of ABR2n

Axiom 3.1 *The number of the binary representations type H (NBRH) is connected with The number of the binary representations type L (NBRL) as follows:*

$$2|H| + |L| = p \quad (7)$$

$\forall 2n > 2$.

The equation (7) asserts that odd noncomposite positive integers less than $2n$ are allotted to types "H", "L" in compliance with balance (7).

Remark 3.5 *From equations (7 and 8) it follows :*

$$|H| = (p - |L|)/2 \quad |Q| = (s_0 - |L|)/2 .$$

In these equations is division by '2'. From this it can conclude that equations are correct only for n is even. But it is not the case. Below we show that in ABR2n all equations where is division by '2' It always is possibility for any n 's and p 's parity.

Proposition 3.2 *The equation (7) is true for any n 's and p 's parity.*

Proof 3.2 *For n is even p is even*

By equation (7) we get: $|H| = (p - |L|)/2$.

By equation (26) $|L| = n - p - 2|Q|$ and $|L|$ is even

for n is even p is even then $(p - |L|)$ is even .

Hence the division by"2" is possibility then $|H|$

is integer .

Thus equation (7) is true for n is even p is even.

Next for n is even ; p is odd

By equation (7) we get: $|H| = (p - |L|)/2$.

By equation (26) $|L| = n - p - 2|Q|$ and $|L|$ is odd

for n is even and p is odd then $(p - |L|)$ is even .

Hence the division by"2" is possibility then $|H|$

is integer .

Thus equation (7) is true for n is even p is odd.

Next for n is odd or odd prime p is even
 By equation (7) we get: $|H| = (p - |L|)/2$.
 By equation (27) $|L| = n - p - 1 - 2|Q|$ and $|L|$ is even
 for n is odd or odd prime and p is even then $(p - |L|)$ is even .
 Hence the division by "2" is possibility then $|H|$
 is integer .
 Thus equation (7) is true for n is odd or odd prime p is even.

Next for n is odd or odd prime p is odd
 By equation (7) we get: $|H| = (p - |L|)/2$.
 By equation (27) $|L| = n - p - 1 - 2|Q|$ and $|L|$ is odd
 for n is odd or odd prime and p is odd then $(p - |L|)$ is even .
 Hence the division by "2" is possibility then $|H|$
 is integer .
 Thus equation (7) is true for n is odd or odd prime p is odd.

Finally equation (7) is true for any n 's and p 's parity. \square

Axiom 3.2 The number of the binary representations Type Q (NBRQ) is connected with the number of the binary representations Type "L" (NBRL) as follows:

$$2|Q| + |L| = s_0 \quad (8)$$

$\forall 2n > 2$.

The expression(8) asserts that odd composite positive integers less than $2n$ are allotted to types "Q", "L" in compliance with balance (8).

Proposition 3.3 The equation (8) is true for any n 's and p 's parity.

Proof 3.3 For n is even p is even
 By equation (8) we get: $|Q| = (s_0 - |L|)/2$.
 By equation (7) $|L| = p - 2|H|$ and $|L|$ is even
 for n is even and p is even .
 By equation (17) $s_0 = n - p$ is even then $(s_0 - |L|)$ is even .
 Hence the division by "2" is possibility then $|Q|$
 is integer .
 Thus equation (8) is true for n is even ; p is even.

Next for n is even p is odd
 By equation (8) we get: $|Q| = (s_0 - |L|)/2$.

By equation (7) $|L| = p - 2|H|$ and $|L|$ is odd
for n is even p is odd .

By equation (17) $s_0 = n - p$ is odd then $(s_0 - |L|)$ is even .

Hence the division by "2" is possibility then $|Q|$
is integer

Thus equation (8) is true for n is even p is odd

Next for n is odd or odd prime p is even

By equation (8) we get: $|Q| = (s_0 - |L|)/2$.

By equation (7) $|L| = p - 2|H|$ and $|L|$ is even
for p is even .

By equation (18) $s_0 = n - p - 1$ is even then $(s_0 - |L|)$ is even .

Hence the division by "2" is possibility then $|Q|$
is integer .

Thus equation (8) is true for n is odd or odd prime p is even.

Next for n is odd or odd prime p is odd

By equation (8) we get: $|Q| = (s_0 - |L|)/2$.

By equation (7) $|L| = p - 2|Q|$ and $|L|$ is odd
for p is odd .

By equation (18) $s_0 = n - p - 1$ is odd then $(s_0 - |L|)$ is even .

Hence the division by "2" is possibility then $|Q|$
is integer .

Thus equation (8) is true for n is odd or odd prime p is odd.

Finally equation (8) is true for any n 's and p 's parity. \square

Definition 3.11 G - the general number of binary representations in
 $ABR2n$.

G - positive integer $\forall 2n > 2$.

Axiom 3.3

$$|Q| + |L| + |H| + |E| = G \quad (9)$$

Definition 3.12 F - the general number of binary representations
with odd positive integers .

F - positive integer $\forall 2n > 2$.

Axiom 3.4

$$|Q| + |L| + |H| = F \quad (10)$$

3.4 The computation of $G, F, S_o, p, |Q|, |L|, |H|, |E|$

3.4.1 The computation of G

Proposition 3.4

$$G = n - 1 \quad (11)$$

$\forall n > 1$

Proof 3.4 *The general number of elements in the set XUY by equation (6) equals $2(n - 1)$. Taking into account that in forming of each binary representation participate with two elements from the set XUY then we have: $G = 1/2(2(n - 1)) = n - 1 \forall n > 1$. \square*

3.4.2 The computation of $|E|$

Proposition 3.5

$$|E| = [(n - 1)/2] \quad (12)$$

$\forall n > 1$

Proof 3.5 *By definition (3.3) by elements of the set XUY are even integer which consist of half of all integers*

Then the number of even integers in the set XUY equals:

$$1/2|XUY| = 1/2(2(n - 1)) = n - 1 .$$

Taking into account that in forming of each binary representation

participate by two elements from the set XUY we have: $|E| = (n - 1)/2$.

Taking into account that for n - even $|E|$ is not integer

that breaks the status of $|E|$ (see definition (3.10))

then $|E| = [(n - 1)/2]$ is aliquot of $(n - 1)/2$

then we get : $|E| = [(n - 1)/2] \forall n > 1$ \square

Proposition 3.6

$$|E| = n/2 - 1 \quad (13)$$

for n is even $n > 1$

Proof 3.6 *For $n = 2i$ where $i \in N$ by equation (12) we get: $|E| =$*

$$[(2i - 1)/2] = [i - 1/2] = [(i - 1) + 1/2] = i - 1$$

returning to n finally we get: $|E| = n/2 - 1 \ n > 1$. \square

Proposition 3.7

$$|E| = (n - 1)/2 \quad (14)$$

for n is odd or odd prime $n > 2$

Proof 3.7 For $n = 2i + 1$ where $i \in \mathbb{N}$ by equation (12) we get: $|E| = [(2i + 1 - 1)/2] = [i] = i$
returning to n finally we get: $|E| = (n - 1)/2$ $n > 2$. \square

3.4.3 The computation of F

Proposition 3.8

$$F = n/2 \quad (15)$$

for n is even $n > 1$

Proof 3.8 Subtracting equation (10) from equation (9) we get: $|E| = G - F$ whence $F = G - |E|$. Taking into account equations (11), (12) we get:

$F = (n - 1) - [(n - 1)/2]$ for $n = 2i$ we get:
 $F = 2i - 1 - [(2i - 1)/2] = 2i - 1 - [i - 1/2] =$
 $= 2i - 1 - [i - 1 + 1/2] = 2i - 1 - (i - 1) = i$
returning to n finally we get: $F = n/2$ $n > 1$

Remark 3.6 Since n is even then the division by "2" is possibility. Thus F is integer and equation (15) is true for n is even. \square

Proposition 3.9

$$F = (n - 1)/2 \quad (16)$$

for n is odd or odd prime $n > 2$

Proof 3.9 Subtracting equation (10) from equation (9) we get: $|E| = G - F$ whence $F = G - |E|$. Taking into account equations (11), (12) we get:

$F = (n - 1) - [(n - 1)/2]$ for $n = 2i + 1$ we get:
 $F = 2i - i = i$ returning to n finally we get: $F = (n - 1)/2$ $n > 2$. \square

Remark 3.7 Since n is odd or odd prime then $(n - 1)$ is even. Hence the division by "2" is possibility. Thus F is integer and equation (16) is true for n is odd or odd prime.

3.4.4 The computation of s_o

Proposition 3.10

$$s_o = n - p \quad (17)$$

.

for n is even, $n > 1$

Proof 3.10 By definition (3.3) for computation s_o it needs to subtract from $|XUY|$ the number of even composite positive integers $(n - 2)$ here (-2) takes into account corollary 3.1 $n^* = n - 2$ and also subtract the number of odd noncomposite positive integers p then we get:

$$s_o = 2(n - 1) - (n - 2) - p = n - p \text{ for } n \text{ is even } n > 1 . \quad \square$$

Proposition 3.11

$$s_o = n - p - 1 \quad (18)$$

.

for n is odd or odd prime $n > 1$

Proof 3.11 Making the change equation (17) by rem 3.4 we get: $s_o = n - p - 1$ for n is odd $n > 1$. \square

3.4.5 The computation of p

Definition 3.13 p_1 – the first approximation of p .

p_2 – the second approximation of p .

p_3 – the third approximation of p .

Proposition 3.12

$$p_1 = \text{round}(2n/\ln 2n) \quad (19)$$

Proposition 3.13

$$p_2 = \text{round}(2n/\ln 2n + 2n/(\ln 2n)^2) \quad (20)$$

Proposition 3.14

$$p_3 = \text{round}(2n/\ln 2n + 2n/\ln^2 2n + 4n/\ln^3 2n) \quad (21)$$

Proof 3.12 As everybody knows [1] that the number of the primes less than $2n$ is expressed as follows:

$$\pi(2n) = (2n/\ln 2n) \int_0^1 (1 - (\ln y/\ln 2n) + (\ln^2 y/\ln^2 2n) + \dots) dy$$

We are limited to three of the first terms of the series.

Integrating in parts then we get:

$$\pi(2n) = (2n/\ln 2n)(1 + 1/\ln 2n + 2/\ln^2 2n)$$

Whence taking into account definition (3.6) that p is (primes + 1) then we get :

$$p = \text{round}(2n/\ln 2n + 2n/\ln^2 2n + 4n/\ln^3 2n + 1)$$

But taking into account that "2" is not odd prime and

that by definition (3.6) p is the number of odd prime $< 2n$ then

$$p = \text{round}(2n/\ln 2n + 2n/\ln^2 2n + 4n/\ln^3 2n + 1 - 1)$$

Finally we get:

$$p = \text{round}(2n/\ln 2n + 2n/\ln^2 2n + 4n/\ln^3 2n)$$

Whence we get:

$$p_1 = \text{round}(2n/\ln 2n)$$

$$p_2 = \text{round}(2n/\ln 2n + 2n/\ln^2 2n)$$

$$p_3 = \text{round}(2n/\ln 2n + 2n/\ln^2 2n + 4n/\ln^3 2n) \quad \square$$

3.4.6 The computation of $|Q|$

Proposition 3.15

$$|Q| = (n - p - |L|)/2 \quad (22)$$

for n is even $n > 1$

Proof 3.13 By equation (8) we get: $|Q| = (s_o - |L|)/2$. Taking into account equation (17) we get for n is even

$$|Q| = (n - p - |L|)/2 . \text{ Now we show that for } n \text{ is even } (n - p - |L|) = (n - (p + |L|)) \text{ is even:}$$

By equation (28) $|L| = p - 2|H|$ whence $|L|$ has parity of p hance $(p + |L|)$ is even

for any p 's and $|L|$'s parity and for n is even then $(n - p - |L|)$ is even then the division by "2" is possibility

thus $|Q|$ is integer and equation (22) is true for n is even $n > 1$. \square

Proposition 3.16

$$|Q| = (n - p - 1 - |L|)/2 \quad (23)$$

.

for n is odd or odd prime $n > 1$

Proof 3.14 Changing equation (22) by rem (3.4) we get: $|Q| = (n - p - 1 - |L|)/2 = (n - 1 - (p + |L|)) / 2$ $n > 1$.

Now we show that $(n - p - 1 - |L|)$ is even:

By equation (28) $|L| = p - 2|H|$ whence $|L|$ has parity of p hence $(p + |L|)$ is even for any p 's and $|L|$'s parity

then $(n - 1 - (p + |L|))$ is even then the division by "2" is possibility thus $|Q|$ is integer and equation (23) is true for n is odd or odd prime $n > 1$.

□

Proposition 3.17

$$|Q| = (n - 2p + 2|H|)/2 \quad (24)$$

.

for n is even $n > 1$

Proof 3.15 subtracting equation (7) from equation (8) we get: $2|Q| - 2|H| = s_0 - p$ Whence we get: $|Q| = (2|H| + s_0 - p)/2$. Taking into account equation (17)

$|Q| = (2|H| + n - p - p)/2 = (n - 2p + 2|H|)/2$. and for n is even $(n - 2p + 2|H|)$ is even then the division by "2"

is possibility thus $|Q|$ is integer and equation (24) is true for n is even $n > 1$. □

Proposition 3.18

$$|Q| = (n - 2p + 2|H| - 1)/2 \quad (25)$$

for n is odd or odd prime $n > 1$

Proof 3.16 Changing equation (24) by rem (3.4) we get: $|Q| = (n - 2p + 2|H| - 1)/2$ $n > 1$.

For n is odd or odd prime $(n - 2p + 2|H| - 1)$ is even then the division by "2" is possibility thus $|Q|$ is integer and equation (25) is true for n is odd or odd prime $n > 1$. □

3.4.7 The computation of $|L|$

Proposition 3.19

$$|L| = n - p - 2|Q| \quad (26)$$

for n is even $n > 1$

Proof 3.17 By equation (8) we get: $|L| = (s_o - 2|Q|)$ Taking into account (17) we get for n is even

$|L| = (n - p - 2|Q|)$ For n is even $n > 1$. \square

Proposition 3.20

$$|L| = n - p - 1 - 2|Q| \quad (27)$$

for n is odd or odd prime $n > 1$

Proof 3.18 Changing equation (26) by rem (3.4) we get: $|L| = (n - p - 1 - 2|Q|)/2$ for n is odd or odd prime $n > 1$. \square

Proposition 3.21

$$|L| = p - 2|H| \quad (28)$$

$\forall n > 1$

Proof 3.19 By equation (7) and rem (3.3) we get: $|L| = p - 2|H|$

for n is even $n > 1$

For n is odd or odd prime by rem (3.4) we also get: $|L| = p - 2|H|$ $n > 1$

Thus $|L| = p - 2|H|$ $\forall n > 1$. \square

3.4.8 The computation of $|H|$

Proposition 3.22

$$|H| = (p - |L|)/2 \quad (29)$$

$\forall n > 1$

Proof 3.20 By equation(7) and rem (3.3) we get: $|H| = (p - |L|)/2$ for n is even $n > 1$.

For n is odd or odd prime by rem (3.4) we also get: $|H| = (p - |L|)/2$ $n > 1$.

Now we show that $\forall n$ $(p - |L|)$ is even

By equation (26) $|L| = n - p - 2|Q|$ whence $|L| + p$ is even for n is even .

By equation (27) $|L| = n - p - 1 - 2|Q|$ whence $|L| + p$ is even for n is odd or odd prime.

It is possibility if $|L|$ and p have equal parity then $(p - |L|)$ is even for any p 's and $|L|$'s parity.

Thus $(p - |L|)$ is even $\forall n$.

Hence the division by "2" is possibility then $|H|$ is integer and equation (29) is true $\forall n > 1$. \square

Proposition 3.23

$$|H| = (2p + 2|Q| - n)/2 \quad (30)$$

for n is even $n > 1$

Proof 3.21 Subtracting equation (7) from equation (8) we get:

$2|Q| - 2|H| = s_0 - p$ Whence we get: $|H| = (2|Q| - s_0 + p)/2$. Taking into account equation (17) we have for n is even:

$|H| = (2|Q| - (n - p) + p)/2 = (2p + 2|Q| - n)/2$. Now we can see that for n is even $(2p + 2|Q| - n)$ is even

then the division by "2" is possibility thus $|H|$ is integer and equation (30) is true for n is even $n > 1$ \square

Proposition 3.24

$$|H| = (2p + 2|Q| - n + 1)/2 \quad (31)$$

for n is odd or odd prime $n > 1$

Proof 3.22 Subtracting equation (7) from equation (8) we get:

$2|Q| - 2|H| = s_0 - p$ Whence we get: $|H| = (2|Q| - s_0 + p)/2$. Taking into account (18) we have for:

$|H| = (2|Q| - (n - p - 1) + p)/2 = (2p + 2|Q| - n + 1)/2$. Now we can see that for n is odd or odd prime

$(2p + 2|Q| - n + 1)$ is even then the division by "2" is possibility thus $|H|$ is integer and equation (31) is true

for n is odd or odd prime $n > 1$. \square

Definition 3.14 $|Q| - |H|$ - lower limit of possible range of $|Q| - |H|$.

$|Q| - |H|$ - positive integer $> 0 \quad \forall 2n > 120$.

$|Q| - |H|$ - negative integer $< 0 \quad \forall 2 < 2n < 120$.

excepting $2n = 94; 96; 100; 106; 118$; for which $|Q| - |H| = 0$ (see subsection 12.1)

.

Proposition 3.25

$$|Q| - |H| = (n - 2p)/2 \quad (32)$$

for n is even $n > 1$

Proof 3.23 Subtracting equation (7) from equation (8) we get: $2|Q| - 2|H| = s_0 - p$ Whence we get: $|Q| - |H| = (s_0 - p)/2 = .$ Taking into account equation (17) we :

$|Q| - |H| = (n - p - p)/2 = (n - 2p)/2$. Since for n is even $(n - 2p)$ is even then the division by "2"

is possibility thus $|Q| - |H|$ is integer and equation (32) is true for n is even $n > 1$ \square

Proposition 3.26

$$|Q| - |H| = (n - 2p - 1)/2 \quad (33)$$

for n is odd or odd prime $n > 1$

Proof 3.24 Subtracting equation (7) from equation (8) we get:

$2|Q| - 2|H| = s_0 - p$ Whence we get: $|Q| - |H| = (s_0 - p)/2 = .$ Taking into account equation (18)

we have :

$|Q| - |H| = (n - 1 - p - p)/2 = (n - 2p - 1)/2$. For n is odd or odd prime $(n - 2p - 1)$ is even then

the division by "2" is possibility thus $|Q| - |H|$ is integer

and equation (33) is true for n is odd or odd prime $n > 1$. \square

Remark 3.8 It was controlled ABR2n for some values of 2n. For each value of 2n it was computed by direct computation the following: $p, s_0, |Q|, |L|, |H|, F$ for values $< 2n$. Next we compare data direct computations with the same parameters which are computed by equations of ABR2n and we get full coincidence. The full text of the control ABR2n see subsection (12.3) of appendices.

4 The limited values of possible range of $|Q|, |L|, |H|$

Definition 4.1 $|H|_b$ -lower limit of possible range of $|H|$.

Axiom 4.1

$$|H|_b = 0 \quad (34)$$

Definition 4.2 $p(n)$ is the number of odd noncomposite integers

(primes and "1") $< n$

$p(n)$ is positive integer $> 0 \forall n > 1$

Definition 4.3 $p - p(n)$ - the number of odd noncomposite integers in Y

Definition 4.4 $|H|_c$ -upper limit of possible range of $|H|$.

Proposition 4.1

$$|H|_c = p - p(n) \quad (35)$$

Proof 4.1 *By Law of distribution of primes the number of odd noncomposite integers in X is greater than the number of odd noncomposite integers in $Y : p(n) > p - p(n)$.*

Since the number of primes decreases with increase of n . Hence maximal number of pair of odd noncomposite integers in the set XUY equals the number of odd noncomposite integers in $Y : p - p(n)$

then $|H|_c = p - p(n)$. \square

Corollary 4.1 *The number of unpaired odd noncomposite positive integers in X equals $2p(n) - p$ and are allotted to type "L". Then $|L| > 0 \forall 2n > 8$*

Proof 4.2 *The number of unpaired odd noncomposite positive integers in X by definitions 4.2, 4.3 equals :*

$$(p(n) - (p - p(n))) = 2p(n) - p$$

and are allotted to type "L". \square

Definition 4.5 $|L|_b$ - lower limit of possible range of $|L|$.

Proposition 4.2

$$|L|_b = 2p(n) - p \quad (36)$$

Proof 4.3 *Substituting upper limit of $|H|$ by (35) to (7) then we get: lower limit for $|L| : |L|_b = 2p(n) - p$ \square*

Definition 4.6 $|L|_c$ -upper limit of possible range of $|L|$.

Proposition 4.3

$$|L|_c = p \quad (37)$$

Proof 4.4 *Substituting lower limit of $|H|$ by (34) to (7) then we get upper limit for $|L| : |L|_c = p$ \square*

Definition 4.7 $|Q|_b$ - lower limit of possible range of $|Q|$.

$|Q|_b$ - positive integer $> 0 \forall 2n > 120$ see proposition (9.2).

$|Q|_b$ - negative integer $< 0 \forall 2 < 2n < 120$ see proposition (9.1).

excepting $2n = 94; 96; 100; 106; 118$ for which $|Q|_b = 0$ see subsection (12.1).

At that $2n = 120; (|Q|_b = 0)$ is border point.

Proposition 4.4

$$|Q|_b = (n - 2p)/2 \quad (38)$$

n is even $n > 1$

Proof 4.5 *Substituting upper limit of $|L|$ by equation (37) to equation (8) then we get*

lower limit for $|Q|$: $|Q|_b = (S_0 - p)/2$.

Substituting S_0 by equation (17), then we get:

$|Q|_b = (n - 2p)/2$ $n > 1$. \square

Proposition 4.5

$$|Q|_b = (n - 2p - 1)/2 \quad (39)$$

n is odd or odd prime $n > 1$

Proof 4.6 *Substituting upper limit of $|L|$ by equation (37) to equation (8) then we get*

lower limit for $|Q|$: $|Q|_b = (S_0 - p)/2$.

Substituting S_0 by equation (18) finally we get: $|Q|_b = (n - 2p - 1)/2$

for n is odd or odd prime $n > 1$. \square

Definition 4.8 $|Q|_c$ - upper limit of possible range of $|Q|$.

Proposition 4.6

$$|Q|_c = (n - 2p(n))/2 \quad (40)$$

for is even

Proof 4.7 *Substituting lower limit of $|L|$ by equation (36) to equation (8) then we get*

upper limit for $|Q|$: $|Q|_c = (S_0 - (2p(n) - p))/2$.

Substituting S_0 by equation (17) finally we get: $|Q|_c = (n - 2p(n))/2$;

for n is even \square

Proposition 4.7

$$|Q|_c = (n - 2p(n) - 1)/2 \quad (41)$$

for n is odd or odd prime

Proof 4.8 *Substituting lower limit of $|L|$ by equation (36) to equation (8) then we get*

upper limit for $|Q|$: $|Q|_c = (S_0 - (2p(n) - p))/2$

Substituting S_0 by equation (18) finally we get: $|Q|_c = (n - 2p(n) - 1)/2$;

for n is odd or odd prime . \square

Proposition 4.8

$$|Q| - |H| = |Q|_b \quad (42)$$

$\forall n > 1$

Proof 4.9 By equation (32) we have: $|Q| - |H| = (n - 2p)/2$ for n is even By equation (33) we have: $|Q| - |H| = (n - 2p - 1)/2$ for n is odd or odd prime.

By equation (38) we have: $|Q|_b = (n - 2p)/2$ for n is even. By equation (39) we have: $|Q|_b = (n - 2p - 1)/2$ for n is odd or odd prime.

Whence we get: $|Q| - |H| = |Q|_b \forall n > 1. \square$

5 Average value of the number of binary sums are formed from odd composite positive integers $< 2n$

Definition 5.1 S -ordered set of odd composite positive integers $< 2n$
 s - element of S

$|S| = s_o$ - power of S

s_i - vary over all s

s_j - vary over all s

Definition 5.2 $V = \{v_k | v_k \in N, v_k = s_i + s_j\}$

is a set by elements of which are every possible binary sums of odd composite

integers $< 2n$. (each with all the rest)

Since $s_i < 2n; s_j < 2n$ then $\max v_k < 4n$.

$|V|$ - the power of set V .

Definition 5.3 $W = \{w | w \in N, w = 2k, 1 \leq k \leq 2n\}$

is a set of even composite positive integers $< 4n + 2$

(inf $W = 2$; sup $W = 4n$).

$|W|$ - the power of set $W |W| = 2n$.

Definition 5.4 $|Q|_m$ - mean quantity of binary sums $v_k = s_i + s_j$ which can be formed of odd composite positive integers $< 2n$ and which are mapped into W by surjective mapping :
 $f : V \Rightarrow W$

$$|Q|_m = |V|/|W| \quad (43)$$

i.e. uniform mapping regardless of real.

$|Q|_m$ - positive rational number > 0

$\forall 2n > 4.$

Proposition 5.1

$$|V| = s_0^2 \quad (44)$$

Proof 5.1 *The number of every possible binary sums in the view of $v_k = s_i + s_j$ are formed of odd composite positive integers $< 2n$ is equal the power of Cartesian product: $S \times S$ [2] [3]. Then:
 $|V| = s_0 \cdot s_0 = s_0^2$ \square*

Proposition 5.2

$$|Q|_m = s_0^2/2n \quad (45)$$

Proof 5.2 *By equations (43 and 44) we have : $|Q|_m = |V|/|W| = s_0^2/2n$
 \square*

Proposition 5.3

$$|Q|_m = (n - p)^2/2n \quad (46)$$

For n is even $n > 1$

Proof 5.3 *Substituting s_0 by equation (17) to equation (45) then we get : $|Q|_m = (n - p)^2/2n$;
*For n is even $n > 1$ \square**

Proposition 5.4

$$|Q|_m = (n - p - 1)^2/2n \quad (47)$$

For n is odd or odd prime $n > 1$

Proof 5.4 *Substituting s_0 by equation (18) to equation (45) then we get :
 $|Q|_m = (n - p - 1)^2/2n$;
*For n is odd or odd prime $n > 1$ \square**

6 Average value of the number of binary sums are formed from odd noncomposite positive integers $< 2n$

Definition 6.1 P - ordered set of odd noncomposite positive integers $< 2n$

p - elements of P

$|P| = p$ - power of P

p_i - vary over all p

p_j - vary over all p

Definition 6.2 $T = \{t_k | t_k \in N, t_k = p_i + p_j\}$

is a set by elements of which are every possible binary sums of odd noncomposite integers $< 2n$. (each with all the rest)

Since $p_i < 2n; p_j < 2n$ then $\max t_k < 4n$.

$|T|$ - the power of set T .

Definition 6.3 $|H|_m$ is mean quantity of binary sums $p_i + p_j = 2k$ which can be formed of odd

noncomposite positive integers $< 2n$ and which are mapped into W by surjective mapping :

$f : T \Rightarrow W$

$$|H|_m = |T|/|W| \quad (48)$$

i.e. uniform mapping regardless of real.

$|H|_m$ -positive rational number $> 0 \quad \forall 2n > 4$.

Proposition 6.1

$$|T| = p^2 \quad (49)$$

Proof 6.1 The number of every possible binary sums

are formed of odd noncomposite positive integers $< 2n$

in the view of $p_i + p_j = 2k$ is equal the power of Cartesian product $P \times P$ [2] [3] .

Then $|T| = p \cdot p = p^2 \quad \square$

Proposition 6.2

$$|H|_m = p^2/2n \quad (50)$$

$\forall n > 1$

Proof 6.2 By equations (48 and 49) we have : $|H|_m = p^2/2n \forall n > 1 \quad \square$

7 The deviation of $|Q|, |H|$ from $|Q|_m, |H|_m$

Definition 7.1 The deviation of $|Q|$ from $|Q|_m$ is:

$$\begin{aligned} \Delta|Q| &= |Q|_m - |Q| \\ \Delta|Q| &> 0 \text{ if } |Q|_m > |Q| \\ \Delta|Q| &< 0 \text{ if } |Q|_m < |Q| \end{aligned}$$

Definition 7.2 The deviation of $|H|$ from $|H|_m$ is:

$$\begin{aligned} \Delta|H| &= |H|_m - |H| \\ \Delta|H| &> 0 \text{ if } |H|_m > |H| \\ \Delta|H| &< 0 \text{ if } |H|_m < |H| \end{aligned}$$

7.1 Relationship between deviations of $\Delta|Q|$ and $\Delta|H|$

Proposition 7.1

$$\Delta|Q| = \Delta|H| + |Q|_m - |H|_m - |Q|_b \quad (51)$$

$\forall n > 1$

Proof 7.1 By equation (42) $|Q| - |H| = |Q|_b$

By definition (7.1) $\Delta|Q| = |Q|_m - |Q|$

By definition (7.2) $\Delta|H| = |H|_m - |H|$

Whence : $|Q| = |Q|_m - \Delta|Q|$

$|H| = |H|_m - \Delta|H|$

Then by equation (42)

$$(|Q|_m - \Delta|Q|) - (|H|_m - \Delta|H|) = |Q|_b$$

$$|Q|_m - \Delta|Q| - |H|_m + \Delta|H| = |Q|_b$$

Whence:

$$\Delta|Q| = \Delta|H| + |Q|_m - |H|_m - |Q|_b . \quad \square$$

Remark 7.1 It was controlled the equation for at the some values of $2n$. For each value of $2n$ it was took data direct computations of $p, s_o, |Q|, |l|, |H|$ from section (12.3) . Next we compute by equation (51) using data direct computations. The compare of $\Delta|Q|$ computed by data of direct computations with computed value by equation (51) shows full coincidence of results. The full text of the control you can see section (13).

8 The relationship between n and p

8.1 The relationship between n and p for $n < 60$

Proposition 8.1

$$2p > n \quad (52)$$

for n is even $n < 60$

Proof 8.1 From the numerical solution (see.subsection 12.1) it follows that:

$$2p > n \text{ for } n \text{ is even } n < 60 \quad \square$$

Proposition 8.2

$$2p + 1 > n \quad (53)$$

for n is odd or odd prime $n < 59$

Proof 8.2 From the numerical solution (see.subsection 12.1) it follows that:

$$2p + 1 > n \text{ for } n \text{ is odd or odd prime } n < 59 \quad \square$$

8.2 The relationship between n and p for $n > 60$

Proposition 8.3

$$n > 2p \quad (54)$$

for n is even $n > 60$

Proposition 8.4

$$n > 2p + 1 \quad (55)$$

for n is odd or odd prime $n > 59$.

Proof 8.3 We need to prove that: $n > 2p$ for n is even, $n > 60$.
 $n > 2p + 1$ for n is odd or odd prime, $n > 59$.

For it we must find a dependence of differences :

$$f_1 = n - 2p; \text{ and } f_2 = n - (2p + 1); \text{ from } n$$

Substituting for p its the second order approximation by equation (20) we get:

$$f_1 = n - 2(2n/\ln 2n + 2n/\ln^2 2n)f_2 = n - (2(2n/\ln 2n + 2n/\ln^2 2n) + 1) .$$

$$f_1 = n - 4n/\ln 2n - 4n/\ln^2 2n f_2 = n - 4n/\ln 2n - 2n/\ln^2 2n) - 1 .$$

Next we computation the derivatives :

$$(f_1)' = (\ln^4 2n - 4\ln^3 2n + 8\ln 2n) / \ln^4 2n .$$

$$(f_2)' = (\ln^4 2n - 4\ln^3 2n + 8\ln 2n) / \ln^4 2n .$$

Whence $(f_1)' > 0$; $(f_2)' > 0$; $\forall 2n > 2$.

Then f_1, f_2 increase $\forall 2n > 2$.

Next we compute the points of intersection of f_1, f_2 with abscissa axis.

For it we need to test of fulfillment of conditions:

$$(f_1(2n) = 0 \ f_1(2n + 2) > 0) \text{ and } (f_2(2n) = 0 \ f_2(2n + 4) > 0).$$

As follows from numerical solution (see subsection 12.1).

The points of intersection are:

$2n = 120$ for n is even; and $2n = 118$ for n is odd or odd prime

Since conditions are fulfilled only for them.

Hence $f_1 > 0$ for n is even, $n > 60$.

And $f_2 > 0$ for n is odd or odd prime $n > 59$

Thus $n > 2p$; for n is even , $n > 60$.

And $n > 2p + 1$ for n is odd or odd prime $n > 59$. \square

8.2.1 The control of relationship $n > 2p$; $n > 2p + 1$

Remark 8.1 p^* is decremental value of p by corollary 3.3

From data direct computations exmp 12.5; examp 12.6 ; examp 12.7 we have :

$$n = 66 , p = 32 ; n = 67, p^* = 30 \ n = 69, p = 33$$

By proposition (8.3) $66 > 64$.

By proposition (8.4) $67 > 60 + 1$ $69 > 66 + 1$. Hence propositions (8.3 and 8.4) are correct for n stated above.

8.3 The correspondence between n and $p \ \forall n$

Definition 8.1 The correspondence of p to finite set of $2n$ it is when for several $2n$ it takes at the same p at computations by equations ABR $2n$.

Proposition 8.5 To each p is correspond finite set of $2n$ included between neighboring values of p .

Proof 8.4 Let for any $2n_1$ exists $p < 2n_1$ and for $2n_k$ exists $p + 1 < 2n_k$ then to each p is correspond to set of $2n = 2n_1 \dots 2n_k$ This set is restrictedly. Hence to each p is correspond finite set of $2n$ included between neighboring values of p . \square

Remark 8.2 *It was controlled the correspondence of finite set of $2n$ included between neighboring values of p to it. It was shown that each p correspond to finite set of $2n$ included between neighboring values of p . The full text of the control we can see the subsection (13.1).*

9 The properties of $|Q|_b$

Proposition 9.1 $|Q|_b$ is negative integer in the range of $2 < 2n < 120$.

Excepting $2n = 94, 96, 100, 118$.

Proof 9.1 *By proof (4.5) $|Q|_b$ is integer for n is even .*

By proof (4.6) $|Q|_b$ is integer for n is odd or odd prime .

Then $|Q|_b$ is integer $\forall n > 1$.

Since by propositions (8.1 and 8.2) $2p > n$ and $2p + 1 > n$ then by equation (38) $|Q|_b < 0$ for n is even < 60 .

And by equation (39) $|Q|_b < 0$ for n is odd or odd prime < 60 .

By direct computation with help of numerical solution of $|Q|_b$ (see subsection 12.1)

we find $2n$ for which $|Q|_b = 0$.

Thus In the range of $2 < 2n < 120$ $|Q|_b$ is negative integer excepting $2n = 94, 96, 100, 118$. (see subsection 12.1) \square

Proposition 9.2 $|Q|_b$ is positive integer $\forall 2n > 120$

Proof 9.2 *By proof (4.5) $|Q|_b$ is integer for n is even .*

By proof (4.6) $|Q|_b$ is integer for n is odd or odd prime .

Then $|Q|_b$ is integer $\forall n > 1$.

Since by propositions (8.3 and 8.4) $2p < n$ and $2p + 1 < n$ then by equation (38) $|Q|_b > 0$ for n is even > 60 .

And by equation (39) $|Q|_b > 0$ for n is odd or odd prime > 60 .

Thus $|Q|_b$ is positive integer $\forall 2n > 120$ \square

Corollary 9.1 *There is no less than $|Q|_b$ of representations of the type "Q" $\forall 2n > 120$.*

Remark 9.1 *The following proposition explains the cause by which $|Q| = 0$ in the range of $2 < 2n < 120$.*

Proposition 9.3 *In the range of $2 < 2n < 120$ if fulfilled the condition $|H| = ||Q|_b|$ then $|Q| = 0$.*

Proof 9.3 By equation (42) and proposition (9.1) we have for $2 < 2n < 120$

$$|H| = |Q| + ||Q|_b|.$$

Let $|Q| = 0$; then $|H| = ||Q|_b|$.

Thus if it is fulfilled the condition $|H| = ||Q|_b|$ then $|Q| = 0$

In the range of $2 < 2n < 120$. \square

10 The solution of the Goldbach's binary problem

Lemma 10.1 $\forall 2 < 2n < 120$ exists at least one representation of type "H".

Proof 10.1 Taking into account equation (42) we get: $|H| = |Q| - |Q|_b$. Since in the range of $2 < 2n < 120$ by proposition(9.1) $|Q|_b < 0$ then $|H| = |Q| + ||Q|_b|$

Whence $|H| > 0$ excepting $2n = 94; 96; 100; 106; 118$; in which $|Q|_b = 0$ (see subsection 12.1).

For this $2n$ the truth of lemma follows from that for $2n$ in which $|Q|_b = 0$ then $|Q| > 0$ in this points since the points of exclusion for $|Q|$ $2n = 26, 28, 32, 38$ (see subsection 12.2) don't coincidence with points of exclusion $|Q|_b$ $2n = 94, 96, 100, 106, 118$ (see subsection 12.1).

Thus $|H| > 0 \forall 2 < 2n < 120$ \square

Theorem 10.1 $\forall 2n > 2$ exists at least one representation of even positive integer $2n$ in the view of a sum of two odd prime positive integers or "1" and odd prime positive integer.

Proof 10.2 Earlier by Lemma we proved that $|H| > 0 \forall 2 < 2n < 120$. For the full proof of the theorem we need to prove that

$$|H| > 0 \forall 2n > 120$$

Let $|H| = 0$ for some value $n > 60$ then by definition (7.2) we get:

$$\Delta|H| = |H|m \tag{56}$$

Also from definition (7.1) we get:

$$|Q| = |Q|m - \Delta|Q| \tag{57}$$

Taking into account that at $|H| = 0$ from equation (42) it follows that:

$$|Q| = |Q|_b \tag{58}$$

we get:

$$|Q|_b = |Q|_m - \Delta|Q| \quad (59)$$

Taking into account proposition (7.1) and equation (56) we get:

$$|Q|_b = |Q|_m - (|Q|_m - |H|_m - |Q|_b + |H|_m) \quad (60)$$

Whence :

$$|Q|_b = |Q|_b \quad (61)$$

Taking into account equation (38) we get identity for n is even, $n > 1$:

$$(n - 2p)/2 = (n - 2p)/2 \quad (62)$$

Taking into account equation (39) we get identity for n is odd or odd prime, $n > 1$:

$$(n - 2p - 1)/2 = (n - 2p - 1)/2 \quad (63)$$

Thus in the result of assumption that $|H| = 0$ for some value $n > 60$ we come to identities: (61 and 62) next we transform these identities into as follows :

$$2n(n - 2p) = 2n(n - 2p); 2n^2 - 4pn = 2n^2 - 4pn; \text{ whence : } 4pn = 4pn; \text{ or } 2np = 2np .$$

$$2n(n - 2p - 1) = 2n(n - 2p - 1); 2n^2 - 4pn - 2n = 2n^2 - 4pn - 2n; \text{ whence : } 4pn = 4pn; \text{ or :}$$

$$2np = 2np \quad (64)$$

We fix value of p and will be substitute values of infinite series $2n$ to(63) and identity (63) is correct for all values of $2n$

then it is determine following relationship between n and p : each p correspond to infinite set of $2n$.

That contradict by proposition (8.5) which assert that each p correspond to finite set of $2n$ included between neighboring values of p .

Hence our assumption that $|H| = 0$ for some value $n > 60$ is false and it is true that $|H| > 0 \forall 2n > 120$. Thus we proved that $|H| > 0 \forall 2n > 2$.

Hence $\forall 2n > 2$ exists at least one representation of even positive integer $2n$ in the view of a sum of two odd prime positive integers or "1" and odd prime positive integer. \square

Corollary 10.1

$$|Q| > |Q|_b \quad (65)$$

$\forall n > 1$

Proof 10.3 From equation (42) we get:

$$|Q| = |Q|_b + |H|$$

Then for $|H| > 0$ we get:

$$|Q| > |Q|_b \forall n > 1. \quad \square$$

Corollary 10.2

$$|L| < p \tag{66}$$

$\forall n > 1$

Proof 10.4 From equation (2.7) we get:

$$|L| = p - 2|H|$$

Then for $|H| > 0$ we get:

$$|L| < p; \forall n > 1. \quad \square$$

11 The computation of the real values of $|Q|, |H|$

11.1 The relative accuracy of computation of $|Q|, |H|$

Definition 11.1 The relative accuracy of computation of $|Q|$ as follows below :

$$\delta_Q = (\Delta|Q|/|Q|_m)100\% \tag{67}$$

Definition 11.2 The relative accuracy of computation of $|H|$ as follows below:

$$\delta_H = (\Delta|H|/|H|_m)100\% \tag{68}$$

11.2 The estimation of δ_Q

Proposition 11.1

$$(100p^2/(n-p)^2) > \delta_Q \tag{69}$$

for n is even $n > 2$;

Proof 11.1 By definition (7.1) we have $|Q| = |Q|_m - \Delta|Q|$.

By definition. (11.1) we have:

$$\Delta|Q| = (\delta_Q|Q|_m)/100$$

then we get : $|Q| = |Q|_m - (\delta_Q|Q|_m)/100$.

By corollary (10.1) $(|Q|_m - (\delta_Q|Q|_m)/100) > |Q|_b$.

Whence it follows that $(100(|Q|_m - |Q|_b)/|Q|_m) > \delta_Q$.

Taking into account equations (46 and 38) .

Then we get:

$$(100((n-p)^2)/2n - (n-2p)/2)/((n-p)^2)/2n > \delta_Q.$$

Hence $(100(p^2)/(n-p)^2) > \delta_Q$ for n is even $n > 2$. \square

Proposition 11.2

$$((100(p^2 + 2p - n + 1)/(n - p - 1)^2)) > \delta_Q \quad (70)$$

for n is odd or oddprime $n > 4$

Proof 11.2 By definition (7.1) we have $|Q| = |Q|_m - \Delta|Q|$.

By definition. (11.1) we have:

$$\Delta|Q| = (\delta_Q|Q|_m)/100$$

then we get : $|Q| = |Q|_m - (\delta_Q|Q|_m)/100$.

By corollary (10.1) $|Q|_m - (\delta_Q|Q|_m)/100 > |Q|_b$.

Whence it follows that $100((|Q|_m - |Q|_b)/|Q|_m) > \delta_Q$.

Taking into account equations (47 and 39) then we get:

Then we get:

$$(100((n-p-1)^2)/2n - (n-2p-1)/2)/((n-p-1)^2)/2n > \delta_Q .$$

Hence $(100(p^2 + 2p - n + 1)/(n - p - 1)^2) > \delta_Q$.

for n is odd or odd prime $n > 4$. \square

11.3 The dependence of $\delta_Q(2n)$

Theorem 11.1 For n is even > 2 If $n \rightarrow \infty$ then $\delta_Q \rightarrow 0$.

Proof 11.3 We equate δ_Q with its estimation by equation(68)

then for n is even we have: $\delta_Q = 100(p^2)/(n-p)^2$.

We replace p with its first order approximation by equation (19) we get:

$$\delta_Q = (100 \cdot 4n^2/\ln^2 2n)/(n^2 \ln^2 2n - 4n^2 \ln 2n + 4n^2) \ln^2 2n = 400/\ln^2 2n(1 - 4/\ln 2n + 4/\ln^2 2n)$$

Whence follows that if $n \rightarrow \infty$ then $\delta_Q \rightarrow 0$

for n is even $n > 2$. \square

Theorem 11.2 For n is odd or odd prime > 4 If $n \rightarrow \infty$ then $\delta_Q \rightarrow 0$.

Proof 11.4 We equate δ_Q with its estimation by equation (69) then for n is odd or odd prime we have: $\delta_Q = 100(p^2 + 2p - n + 1)/(n - p - 1)^2$.

We replace p with its first order approximation by equation (19) we get:
 $\delta_Q = (4n^2 + 4n \ln 2n - n \ln^2 2n + \ln^2 2n)/(n^2 \ln^2 2n - 4n^2 \ln 2n + 4n^2 - 2n \ln^2 2n + 4n \ln 2n + \ln^2 2n)$;

$\delta_Q = 4n^2 \ln^2 2n (1/\ln^2 2n + 1/n \ln 2n - 1/4n + 1/4n^2) / 4n^2 \ln^2 2n (1/4 - 1/\ln 2n + 1/\ln^2 2n - 1/2n + 1/n \ln 2n + 1/4n^2)$

$(1/\ln^2 2n + 1/n \ln 2n - 1/4n + 1/4n^2) / (1/4 - 1/\ln 2n + 1/\ln^2 2n - 1/2n + 1/n \ln 2n + 1/4n^2)$

Whence follows that if $n \rightarrow \infty$ then $\delta_Q \rightarrow 0$.

for n is odd or odd prime $n > 4$.

Thus we proved that if $n \rightarrow \infty$ then $\delta_Q \rightarrow 0 \forall n > 4$ \square

11.4 The character of dependence of $|Q|(2n)$

Theorem 11.3 If $n \rightarrow \infty$ then $|Q| \rightarrow |Q|_m$.

Proof 11.5 By Theorems (11.1 and 11.2) $\delta_Q \rightarrow 0$ if $n \rightarrow \infty$ then by Definition 11.1 $\Delta|Q| \rightarrow 0$ since $\delta_Q \rightarrow 0$.

Whence by Definition 7.1 we have:

If $n \rightarrow \infty$ then $|Q| \rightarrow |Q|_m$. \square

11.5 The formulas for computation of the real values of $|Q|, |H|$

Remark 11.1 By theorem (11.3) we can replace the computation of $|Q|$ with $|Q|_m$. The mistake of it is decreased with increase of n .

$$|Q| = \text{round}((n - p)^2 / 2n) \quad (71)$$

for n is even $n > 16$

$$|Q| = \text{round}((n - p - 1)^2 / 2n) \quad (72)$$

for n is odd or odd prime $n > 17$

Where: $p = \text{round}(2n/\ln 2n + 2n \ln^2 2n)$. By (20)

The computed value of $|Q|$ it can be used for computation of $|H|$ by equations(30 and 31):

$|H| = (2|Q| + 2p - n)/2$ for n is even $n > 16$

$|H| = (2|Q| + 2p - n + 1)/2$ for n is odd or odd prime $n > 17$

12 The appendices

12.1 The numerical solution of $|Q|_b = 0$ in the range $2 < 2n < 138$

Remark 12.1 *If n is prime then value of p is decremented by one .
 $p^* = p - 1$ by corollary 3.3 .*

- $2n = 4; n = 2; p = 2; |Q|_b = -1$ by (38).
- $2n = 6; n = 3; p^* = 2; |Q|_b = -1$ by (39).
- $2n = 8; n = 4; p = 4; |Q|_b = -2$ by (38).
- $2n = 10; n = 5; p^* = 3; |Q|_b = -1$ by (39).
- $2n = 12; n = 6; p = 5; |Q|_b = -2$ by (38).
- $2n = 14; n = 7; p^* = 5; |Q|_b = -2$ by (39).
- $2n = 16; n = 8; p = 6; |Q|_b = -2$ by (38).
- $2n = 18; n = 9; p = 7; |Q|_b = -3$ by (39).
- $2n = 20; n = 10; p = 8; |Q|_b = -3$ by (38).
- $2n = 22; n = 11; p^* = 7; |Q|_b = -2$ by (39).
- $2n = 24; n = 12; p = 9; |Q|_b = -3$ by (38).
- $2n = 26; n = 13; p^* = 8; |Q|_b = -2$ by (39).
- $2n = 28; n = 14; p = 9; |Q|_b = -2$ by (38).
- $2n = 30; n = 15; p = 10; |Q|_b = -3$ by (39).
- $2n = 32; n = 16; p = 11; |Q|_b = -3$ by (38).
- $2n = 34; n = 17; p^* = 10; |Q|_b = -2$ by (39)
- $2n = 36; n = 18; p = 11; |Q|_b = -2$ by (38).
- $2n = 38; n = 19; p^* = 11; |Q|_b = -2$ by (39).
- $2n = 40; n = 20; p = 12; |Q|_b = -2$ by (38).
- $2n = 42; n = 21; p = 13; |Q|_b = -3$ by (39).
- $2n = 44; n = 22; p = 14; |Q|_b = -3$ by (38).
- $2n = 46; n = 23; p^* = 13; |Q|_b = -2$ by (39).
- $2n = 48; n = 24; p = 15; |Q|_b = -3$ by (38).
- $2n = 50; n = 25; p = 15; |Q|_b = -3$ by (39).
- $2n = 52; n = 26; p = 15; |Q|_b = -2$ by (38)
- $2n = 54; n = 27; p = 16; |Q|_b = -3$ by (39).
- $2n = 56; n = 28; p = 16; |Q|_b = -2$ by (38).
- $2n = 58; n = 29; p^* = 15; |Q|_b = -1$ by (39).
- $2n = 60; n = 30; p = 17; |Q|_b = -2$ by (38).
- $2n = 62; n = 31; p^* = 17; |Q|_b = -2$ by (39).
- $2n = 64; n = 32; p = 18; |Q|_b = -2$ by (38).
- $2n = 66; n = 33; p = 18; |Q|_b = -2$ by (39).
- $2n = 68; n = 34; p = 19; |Q|_b = -2$ by (38).

- $2n = 70; n = 35; p = 19; |Q|_b = -2$ by (39).
 $2n = 72; n = 36; p = 20; |Q|_b = -2$ by (38).
 $2n = 74; n = 37; p^* = 20; |Q|_b = -2$ by (39).
 $2n = 76; n = 38; p = 21; |Q|_b = -2$ by (38).
 $2n = 78; n = 39; p = 21; |Q|_b = -2$ by (39).
 $2n = 80; n = 40; p = 22; |Q|_b = -2$ by (38).
 $2n = 82; n = 41; p^* = 21; |Q|_b = -1$ by (39).
 $2n = 84; n = 42; p = 23; |Q|_b = -2$ by (38).
 $2n = 86; n = 43; p^* = 22; |Q|_b = -1$ by (39).
 $2n = 88; n = 44; p = 23; |Q|_b = -1$ by (38).
 $2n = 90; n = 45; p = 24; |Q|_b = -2$ by (39).
 $2n = 92; n = 46; p = 24; |Q|_b = -1$ by (38).
 $2n = 94; n = 47; p^* = 23; |Q|_b = 0$ by (39).
 $2n = 96; n = 48; p = 24; |Q|_b = 0$ by (38).
 $2n = 98; n = 49; p = 25; |Q|_b = -1$ by (39).
 $2n = 100; n = 50; p = 25; |Q|_b = 0$ by (38).
 $2n = 102; n = 51; p = 26; |Q|_b = -1$ by (39).
 $2n = 104; n = 52; p = 27; |Q|_b = -1$ by (38).
 $2n = 106; n = 53; p^* = 26; |Q|_b = 0$ by (39).
 $2n = 108; n = 54; p = 28; |Q|_b = -1$ by (38).
 $2n = 110; n = 55; p = 29; |Q|_b = -2$ by (39).
 $2n = 112; n = 56; p = 29; |Q|_b = -1$ by (38).
 $2n = 114; n = 57; p = 30; |Q|_b = -2$ by (39).
 $2n = 116; n = 58; p = 30; |Q|_b = -1$ by (38).
 $2n = 118; n = 59; p = 29; |Q|_b = 0$ by (39).
 $2n = 120; n = 60; p = 30; |Q|_b = 0$ by (38).
 $2n = 122; n = 61; p = 29; |Q|_b = 1$ by (39).
 $2n = 124; n = 62; p = 30; |Q|_b = 1$ by (38).
 $2n = 126; n = 63; p = 30; |Q|_b = 1$ by (39).
 $2n = 128; n = 64; p = 31; |Q|_b = 1$ by (38).
 $2n = 130; n = 65; p = 31; |Q|_b = 1$ by (39).
 $2n = 132; n = 66; p = 32; |Q|_b = 1$ by (38).
 $2n = 134; n = 67; p = 31; |Q|_b = 2$ by (39).
 $2n = 136; n = 68; p = 32; |Q|_b = 2$ by (38).
 $2n = 138; n = 69; p = 33; |Q|_b = 1$ by (39).

**12.2 The numerical solution of $|Q| = 0$
in the range $8 < 2n < 134$**

- $2n = 10; |Q| = 0; (10 - 9 = 1) .$
 $2n = 12; |Q| = 0; (12 - 9 = 3) .$
 $2n = 14; |Q| = 0; (14 - 9 = 5) .$
 $2n = 16; |Q| = 0; (16 - 9 = 7) .$
 $2n = 18; |Q| = 0; (18 - 9 = 9) .$
 $2n = 20; |Q| = 0; (20 - 9 = 11) .$
 $2n = 22; |Q| = 0; (22 - 9 = 13) .$
 $2n = 24; |Q| > 0; (24 - 9 = 15) .$
 $2n = 26; |Q| = 0; (26 - 9 = 17) .$
 $2n = 28; |Q| = 0; (28 - 9 = 19) .$
 $2n = 30; |Q| > 0; (30 - 9 = 21) .$
 $2n = 32; |Q| = 0; (32 - 9 = 23) .$
 $2n = 34; |Q| > 0; (34 - 9 = 25) .$
 $2n = 36; |Q| > 0; (30 - 9 = 21) .$
 $2n = 38; |Q| = 0; (38 - 9 = 29) .$
 $2n = 40; |Q| > 0; (40 - 15 = 25) .$
 $2n = 42; |Q| > 0; (42 - 9 = 33) .$
 $2n = 44; |Q| > 0; (44 - 9 = 35) .$
 $2n = 46; |Q| > 0; (46 - 21 = 25) .$
 $2n = 48; |Q| > 0; (48 - 9 = 39) .$
 $2n = 50; |Q| > 0; (50 - 15 = 35) .$
 $2n = 52; |Q| > 0; (52 - 25 = 27) .$
 $2n = 54; |Q| > 0; (54 - 9 = 45) .$
 $2n = 56; |Q| > 0; (56 - 21 = 35) .$
 $2n = 58; |Q| > 0; (58 - 9 = 49) .$
 $2n = 60; |Q| > 0; (60 - 9 = 51) .$
 $2n = 62; |Q| > 0; (62 - 9 = 21) .$
 $2n = 64; |Q| > 0; (64 - 9 = 55) .$
 $2n = 66; |Q| > 0; (66 - 9 = 57) .$
 $2n = 68; |Q| > 0; (68 - 33 = 35) .$
 $2n = 70; |Q| > 0; (70 - 15 = 55) .$
 $2n = 72; |Q| > 0; (72 - 9 = 63) .$
 $2n = 74; |Q| > 0; (74 - 9 = 21) .$
 $2n = 76; |Q| > 0; (76 - 21 = 55) .$
 $2n = 78; |Q| > 0; (78 - 9 = 69) .$
 $2n = 80; |Q| > 0; (80 - 15 = 65) .$
 $2n = 82; |Q| > 0; (82 - 25 = 57) .$
 $2n = 84; |Q| > 0; (84 - 9 = 75) .$

$$\begin{aligned}
2n &= 86; |Q| > 0; (86 - 9 = 77) . \\
2n &= 88; |Q| > 0; (88 - 25 = 63) . \\
2n &= 90; |Q| > 0; (90 - 9 = 81) . \\
2n &= 92; |Q| > 0; (92 - 15 = 77) . \\
2n &= 94; |Q| > 0; (94 - 9 = 85) . \\
2n &= 96; |Q| > 0; (96 - 9 = 87) . \\
2n &= 98; |Q| > 0; (98 - 21 = 77) . \\
2n &= 100; |Q| > 0; (100 - 15 = 85) . \\
2n &= 102; |Q| > 0; (102 - 9 = 93) . \\
2n &= 104; |Q| > 0; (104 - 9 = 95) . \\
2n &= 106; |Q| > 0; (106 - 15 = 91) . \\
2n &= 108; |Q| > 0; (108 - 9 = 99) . \\
2n &= 110; |Q| > 0; (110 - 15 = 95) . \\
2n &= 112; |Q| > 0; (112 - 21 = 91) . \\
2n &= 114; |Q| > 0; (114 - 9 = 105) . \\
2n &= 116; |Q| > 0; (116 - 21 = 95) . \\
2n &= 118; |Q| > 0; (118 - 25 = 93) . \\
2n &= 120; |Q| > 0; (120 - 9 = 111) . \\
2n &= 122; |Q| > 0; (122 - 35 = 87) . \\
2n &= 124; |Q| > 0; (124 - 9 = 115) . \\
2n &= 126; |Q| > 0; (126 - 9 = 117) . \\
2n &= 128; |Q| > 0; (128 - 9 = 119) . \\
2n &= 130; |Q| > 0; (130 - 9 = 121) . \\
2n &= 132; |Q| > 0; (132 - 9 = 123) .
\end{aligned}$$

12.3 The control of ABR2n

Remark 12.2 We take into account that : $p, s_o, |Q|, |L|, |H|, |Q|_b, |Q| - |H|$, are defined for values $< 2n$.

I.e. the direct computations of them are made \forall values $< 2n$, but the computations by equations are made for $n = 2n/2$.

Exemple 12.1 The computation of $|Q|, |L|, |H|, s_o, G, F$ for $2n = 4$

Construction SBR2n:

$$2n = 4 ; n = 2$$

\ominus - odd noncomposite; \oplus - odd composite , \otimes - even composite .

$$\ominus 1 + 3\ominus = 4(H)$$

Data of direct computations for even $n = 2$ at that it is took into account remark (3.2)

$$p = 2, s_o = 0, |Q| = 0, |L| = 0, |H| = 1, G = 1, F = 1 .$$

The computations of parameters of binary representations of the positive integer $2n = 4$ with help of arithmetic stated above

at that $p, s_o, |Q|, |L|, |H|$, are took from the data of direct computations.

$$n = 4/2 = 2 ;$$

$$G = n - 1 = 2 - 1 = 1; \text{ by equation (11)}$$

$$|E| = n/2 - 1 = 1 - 1 = 0; \text{ by equation (13)}$$

$$F = n/2 = 2/2 = 1; \text{ by equation (15)}$$

$$s_o = n - p = 2 - 2 = 0; \text{ by equation (17)}$$

$$|L| = n - p - 2|Q| = 2 - 2 - 0 = 0; \text{ by equation (26)}$$

$$|L| = p - 2|H| = 2 - 2 = 0; \text{ by equation (28)}$$

$$|H| = (2|Q| + 2p - n)/2 = (0 + 4 - 2)/2 = 1; \text{ by equation (30)}$$

$$|H| = (p - |L|)/2 = (2 - 0)/2 = 1; \text{ by equation (29)}$$

$$|Q| = (n - p - |L|)/2 = (2 - 2 - 0)/2 = 0; \text{ by equation (22)}$$

$$|Q| = (n - 2p + 2|H|)/2 = (2 - 4 + 2)/2 = 0; \text{ by equation (24)}$$

$$|Q| - |H| = 0 - 1 = -1;$$

$$|Q| - |H| = (n - 2p)/2 = (2 - 4)/2 = -1; \text{ by equation (32)}$$

$$p = 2|H| + |L| = 2 + 0 = 2; \text{ by equation (7)}$$

$$s_o = 2|Q| + |L| = 0 + 0 = 0; \text{ by equation (8)}$$

$$F = |Q| + |L| + |H| = 0 + 0 + 1 = 1; \text{ by equation (10)}$$

The compare of data of direct computations with computed values shows full coincidence of results.

Exemple 12.2 The computation of $|Q|, |L|, |H|, s_o, G, F$ for $2n = 6$

.

\circ - odd noncomposite; \oplus - odd composite ; \otimes - even composite .

Construction SBR_{2n} :

$$2n = 6 ; n = 3$$

$$\circ 1 + 5\circ = 6(H)$$

$$\otimes 2 + 4\otimes = 6(E)$$

Data of direct computations for odd $n = 3$ at that it is took into account remark (3.2)

$$p^* = 2; s_o = 0; |Q| = 0; |L| = 0; |H| = 1; G = 2; F = 1$$

The computations of parameters of binary representations of the positive integer $2n = 6$ with help of arithmetic stated above

at that $p, s_o, |Q|, |L|, |H|$, are took from the data of direct computations.

$$n = 6/2 = 3;$$

$$G = n - 1 = 3 - 1 = 2; \text{ by equation (11)}$$

$$|E| = (n - 1)/2 = 1; \text{ by equation (12)}$$

$F = (n - 1)/2 = (3 - 1)/2 = 1$; by equation (16)
 $s_o = n - p - 1 = 3 - 2 - 1 = 0$; by equation (18)
 $|L| = n - p - 2|Q| - 1 = 3 - 2 - 0 - 1 = 0$; by equation (27)
 $|L| = p - 2|H| = 2 - 2 = 0$; by equation (28)
 $|H| = (2|Q| + 2p - n + 1)/2 = (0 + 4 - 3 + 1)/2 = 1$; by equation (31)
 $|H| = (p - |L|)/2 = (2 - 0)/2 = 1$; by equation (29)
 $|Q| = (n - p - |L| - 1)/2 = (3 - 2 - 0 - 1)/2 = 0$; by equation (23)
 $|Q| = (n - 2p + 2|H| - 1)/2 = (3 - 4 + 2 - 1)/2 = 0$; by equation (25)
 $|Q| - |H| = 0 - 1 = -1$; $|Q| - |H| = (n - 2p - 1)/2 = (3 - 4 - 1)/2 = -1$;
 by equation (33)
 $p = 2|H| + |L| = 2 + 0 = 2$; by equation (7)
 $s_o = 2|Q| + |L| = 0 + 0 = 0$; by equation (8)
 $F = |Q| + |L| + |H| = 0 + 0 + 1 = 1$; by equation (10)
 The compare of data of direct computations with computed values shows
 full coincidence of results.

Exemple 12.3 The computation of $|Q|$, $|L|$, $|H|$, s_o , G , F for $2n = 12$

Construction SBR $2n$:

$$2n = 12 ; n = 6$$

\circ - odd noncomposite; \oplus - odd composite ; \otimes - even composite .

$$\circ 1 + 11\circ = 12(H)$$

$$\otimes 2 + 10\otimes = 12(E)$$

$$\circ 3 + 9\oplus = 12(L)$$

$$\otimes 4 + 8\otimes = 12(E)$$

$$\circ 5 + 7\circ = 12(H)$$

Data of direct computations for even $n = 2$, at that it is took into account
 remark (3.2)

$$p = 5; s_o = 1; |Q| = 0; |L| = 1; |H| = 2; G = 5; F = 3$$

The computations of parameters of binary representations of the positive
 integer $2n = 12$ with help of arithmetic stated above

$$n = 12/2 = 6 ;$$

$$G = n - 1 = 6 - 1 = 5; \text{ by equation (11)}$$

$$|E| = n/2 - 1 = 3 - 1 = 2; \text{ by equation (13)}$$

$$F = n/2 = 6/2 = 3; \text{ by equation (15)}$$

$$s_o = n - p = 6 - 5 = 1; \text{ by equation (17)}$$

$$|L| = n - p - 2|Q| = 6 - 5 - 0 = 1; \text{ by equation (26)}$$

$$|L| = p - 2|H| = 5 - 4 = 1; \text{ by equation (28)}$$

$$|H| = (2|Q| + 2p - n)/2 = (0 + 10 - 6)/2 = 2; \text{ by equation (30)}$$

$$|H| = (p - |L|)/2 = (5 - 1)/2 = 2; \text{ by equation (29)}$$

$$|Q| = (n - p - |L|)/2 = (6 - 5 - 1)/2 = 0; \text{ by equation (22)}$$

$$|Q| = (n - 2p + 2|H|)/2 = (6 - 10 + 4)/2 = 0; \text{ by equation (24)}$$

$$|Q| - |H| = 0 - 2 = -2;$$

$$|Q| - |H| = (n - 2p)/2 = (6 - 10)/2 = -2; \text{ by equation (32)}$$

$$p = 2|H| + |L| = 4 + 1 = 5; \text{ by equation (7)}$$

$$s_o = 2|Q| + |L| = 0 + 1 = 1; \text{ by equation (8)}$$

$$F = |Q| + |L| + |H| = 0 + 1 + 2 = 3; \text{ by equation (10)}$$

The compare of data of direct computations with computed values shows full coincidence of results.

Exemple 12.4 The computation of $|Q|$, $|L|$, $|H|$, s_o , G , F for $2n = 14$.

\circ - odd noncomposite; \oplus - odd composite ; \otimes - even composite .

Construction SBR2n:

$$2n = 14 ; n = 7$$

$$\circ 1 + 13\circ = 14(H)$$

$$\otimes 2 + 12\otimes = 14(E)$$

$$\circ 3 + 11\circ = 14(H)$$

$$\otimes 4 + 10\otimes = 14(E)$$

$$\circ 5 + 9\oplus = 14(L)$$

$$\otimes 6 + 8\otimes = 14(E)$$

Data of direct computations for odd $n = 7$, at that it is took into account remark (3.2).

$$p^* = 5; s_o = 1; |Q| = 0; |L| = 1; |H| = 2; G = 6; F = 3$$

The computations of parameters of binary representations of the positive integer $2n = 14$ with help of arithmetic stated above

$$n = 14/2 = 7;$$

$$G = n - 1 = 7 - 1 = 6; \text{ by equation (11)}$$

$$|E| = (n - 1)/2 = 3; \text{ by equation (12)}$$

$$F = (n - 1)/2 = (7 - 1)/2 = 3; \text{ by equation (16)}$$

$$s_o = n - p - 1 = 7 - 5 - 1 = 1; \text{ by equation (18)}$$

$$|L| = n - p - 2|Q| - 1 = 7 - 5 - 0 - 1 = 1; \text{ by equation (27)}$$

$$|L| = p - 2|H| = 5 - 4 = 1; \text{ by equation (28)}$$

$$|H| = (2|Q| + 2p - n + 1)/2 = (0 + 10 - 7 + 1)/2 = 2; \text{ by equation (31)}$$

$$|H| = (p - |L|)/2 = (5 - 1)/2 = 2; \text{ by equation (29)}$$

$$|Q| = (n - p - |L| - 1)/2 = (7 - 5 - 1 - 1)/2 = 0; \text{ by equation (23)}$$

$$|Q| = (n - 2p + 2|H| - 1)/2 = (7 - 10 + 4 - 1)/2 = 0; \text{ by equation (25)}$$

$$|Q| - |H| = 0 - 2 = -2; |Q| - |H| = (n - 2p - 1)/2 = (7 - 10 - 1)/2 = -2; \text{ by equation (33)}$$

$$p = 2|H| + |L| = 4 + 1 = 5; \text{ by equation (7)}$$

$s_o = 2|Q| + |L| = 0 + 1 = 1$; by equation (8)

$F = |Q| + |L| + |H| = 0 + 1 + 2 = 3$; by equation (10)

The compare of data of direct computations with computed values shows full coincidence of results.

Exemple 12.5 The computation of $|Q|$, $|L|$, $|H|$, s_o , G , F for $2n = 132$

Remark 12.3 The representations with x is even; y is even are excluded since

$|Q|$, $|L|$, $|H|$, s_o , p , by definitions related to odd composite and noncomposite integers.

\circledast - odd noncomposite; \oplus - odd composite ;

Construction SBR $2n$:

$2n = 132$; $n = 66$

$\circledast 1 + 131\circledast = 132(H)$

$\circledast 3 + 129\oplus = 132(L)$

$\circledast 5 + 127\circledast = 132(H)$

$\circledast 7 + 125\oplus = 132(L)$

$\oplus 9 + 123\oplus = 132 (Q)$

$\circledast 11 + 121\oplus = 132 (L)$

$\circledast 13 + 119\oplus = 132 (L)$

$\oplus 15 + 117\oplus = 132 (Q)$

$\circledast 17 + 115\oplus = 132 (L)$

$\circledast 19 + 113\circledast = 132 (H)$

$\oplus 21 + 111\oplus = 132 (Q)$

$\circledast 23 + 109\circledast = 132 (H)$

$\oplus 25 + 107\circledast = 132 (L)$

$\oplus 27 + 105\oplus = 132 (Q)$

$\circledast 29 + 103\circledast = 132 (H)$

$\circledast 31 + 101\circledast = 132 (H)$

$\oplus 33 + 99\oplus = 132 (Q)$

$\oplus 35 + 97\circledast = 132 (L)$

$\circledast 37 + 95\oplus = 132 (L)$

$\oplus 39 + 93\oplus = 132 (Q)$

$\circledast 41 + 91\oplus = 132 (L)$

$\circledast 43 + 89\circledast = 132 (H)$

$\oplus 45 + 87\oplus = 132 (Q)$

$\circledast 47 + 85\oplus = 132 (L)$

$$\oplus 49 + 83 \ominus = 132 \text{ (L)}$$

$$\oplus 51 + 81 \oplus = 132 \text{ (Q)}$$

$$\ominus 3 + 79 \ominus = 132 \text{ (H)}$$

$$\oplus 55 + 77 \oplus = 132 \text{ (Q)}$$

$$\oplus 57 + 75 \oplus = 132 \text{ (Q)}$$

$$\ominus 59 + 73 \ominus = 132 \text{ (H)}$$

$$\ominus 61 + 71 \ominus = 132 \text{ (H)}$$

$$\oplus 63 + 69 \oplus = 132 \text{ (Q)}$$

$$\oplus 65 + 67 \ominus = 132 \text{ (L)}$$

Data of direct computations for even $n = 66$, at the compare it is took into account remark (3.2).

$$p = 32; s_o = 34; |Q| = 11; |L| = 12; |H| = 10; F = 33$$

The computations of parameters of binary representations of the positive integer $2n = 132$ with help of arithmetic stated above

at that $p, s_o, |Q|, |L|, |H|$, are took from the data of direct computations.

$$n = 132/2 = 66;$$

$$G = n - 1 = 66 - 1 = 65; \text{ by equation (11)}$$

$$|E| = n/2 - 1 = 66/2 - 1 = 32; \text{ by equation (13)}$$

$$F = n/2 = 66/2 = 33; \text{ by equation (15)}$$

$$s_o = n - p = 66 - 32 = 34; \text{ by equation (17)}$$

$$|L| = n - p - 2|Q| = 66 - 32 - 22 = 12; \text{ by equation (26)}$$

$$|L| = p - 2|H| = 32 - 20 = 12; \text{ by equation (28)}$$

$$|H| = (2|Q| + 2p - n)/2 = (22 + 64 - 66)/2 = 10; \text{ by equation (30)}$$

$$|H| = (p - |L|)/2 = (32 - 12)/2 = 10; \text{ by equation (29)}$$

$$|Q| = (n - p - |L|)/2 = (66 - 32 - 12)/2 = 11; \text{ by equation (22)}$$

$$|Q| = (n - 2p + 2|H|)/2 = (66 - 64 + 20)/2 = 11; \text{ by equation (24)}$$

$$|Q| - |H| = 11 - 10 = 1; \text{ by equation ()}$$

$$|Q| - |H| = (n - 2p)/2 = (66 - 64)/2 = 1; \text{ by equation (32)}$$

$$p = 2|H| + |L| = 20 + 12 = 32; \text{ by equation (7)}$$

$$s_o = 2|Q| + |L| = 22 + 12 = 34; \text{ by equation (8)}$$

$$F = |Q| + |L| + |H| = 11 + 12 + 10 = 33; \text{ by equation (10)}$$

The compare of data of direct computations with computed values shows full coincidence of results.

Exemple 12.6 The computation of $|Q|, |L|, |H|, s_o, G, F$ for $2n = 138$.

Remark 12.4 The representations with x is even; y is even are excluded since

$|Q|, |L|, |H|, s_o, p$, by definitions related to odd composite and prime integers.

\circledast - odd noncomposite; \oplus - odd composite ;

Construction SBR2n:

$$2n = 138 ; n = 69$$

$$\circledast 1 + 137\circledast = 138(H)$$

$$\circledast 3 + 135\oplus = 138(L)$$

$$\circledast 5 + 133\oplus = 138(L)$$

$$\circledast 7 + 131\circledast = 138(H)$$

$$\oplus 9 + 129\oplus = 138(Q)$$

$$\circledast 11 + 127\circledast = 138(H)$$

$$\circledast 13 + 125\oplus = 138(L)$$

$$\oplus 15 + 123\oplus = 138(Q)$$

$$\circledast 17 + 121\oplus = 138(L)$$

$$\circledast 19 + 119\oplus = 138(L)$$

$$\oplus 21 + 117\oplus = 138(Q)$$

$$\circledast 23 + 115\oplus = 138(L)$$

$$\oplus 25 + 113\circledast = 138(L)$$

$$\oplus 27 + 111\oplus = 138(Q)$$

$$\circledast 29 + 109\circledast = 138(H)$$

$$\circledast 31 + 107\circledast = 138(H)$$

$$\oplus 33 + 105\oplus = 138(Q)$$

$$\oplus 35 + 103\circledast = 138(L)$$

$$\circledast 37 + 101\circledast = 138(H)$$

$$\oplus 39 + 99\oplus = 138(Q)$$

$$\circledast 41 + 97\circledast = 138(H)$$

$$\circledast 43 + 95\oplus = 138(L)$$

$$\oplus 45 + 93\oplus = 138(Q)$$

$$\circledast 47 + 91\oplus = 138(L)$$

$$\oplus 49 + 89\circledast = 138(L)$$

$$\oplus 51 + 87\oplus = 138(Q)$$

$$\circledast 53 + 85\oplus = 138(L)$$

$$\oplus 55 + 83\circledast = 138(L)$$

$$\oplus 57 + 81\oplus = 138(Q)$$

$$\circledast 59 + 79\circledast = 138(H)$$

$$\circledast 61 + 77\oplus = 138(L)$$

$$\oplus 63 + 75\oplus = 138(Q)$$

$$\oplus 65 + 73\circledast = 138(L)$$

$$\circledast 67 + 71\circledast = 138(H)$$

Data of direct computations for odd $n = 69$, at that it is took into account remark (3.2).

$$p = 33; s_o = 35; |Q| = 10; |L| = 15; |H| = 9; F = 34$$

The computations of parameters of binary representations of the positive integer $2n = 132$ with help of arithmetic stated above

at that $p, s_o, |Q|, |L|, |H|$, are took from the data of direct computations.

$$n = 138/2 = 69;$$

$$G = n - 1 = 69 - 1 = 68; \text{ by equation (11)}$$

$$|E| = (n - 1)/2 = 34; \text{ by equation (12)}$$

$$F = (n - 1)/2 = (69 - 1)/2 = 34; \text{ by equation (16)}$$

$$s_o = n - p - 1 = 69 - 33 - 1 = 35; \text{ by equation (18)}$$

$$|L| = n - p - 2|Q| - 1 = 69 - 33 - 20 - 1 = 15; \text{ by equation (27)}$$

$$|L| = p - 2|H| = 33 - 18 = 15; \text{ by equation (28)}$$

$$|H| = (2|Q| + 2p - n + 1)/2 = (20 + 66 - 69 + 1)/2 = 9; \text{ by equation (31)}$$

$$|H| = (p - |L|)/2 = (33 - 15)/2 = 9; \text{ by equation (29)}$$

$$|Q| = (n - p - |L| - 1)/2 = (69 - 33 - 15 - 1)/2 = 10; \text{ by equation (23)}$$

$$|Q| = (n - 2p + 2|H| - 1)/2 = (69 - 66 + 18 - 1)/2 = 10; \text{ by equation (25)}$$

$$|Q| - |H| = 10 - 9 = 1; |Q| - |H| = (n - 2p - 1)/2 = (69 - 66 - 1)/2 = 1; \text{ by equation (33)}$$

$$p = 2|H| + |L| = 18 + 15 = 33; \text{ by equation (7)}$$

$$s_o = 2|Q| + |L| = 20 + 15 = 35; \text{ by equation (8)}$$

$$F = |Q| + |L| + |H| = 10 + 15 + 9 = 34; \text{ by equation (10)}$$

The compare of data of direct computations with computed values shows full coincidence of results.

Exemple 12.7 The computation of $|Q|, |L|, |H|, s_o, G, F$ for $2n = 134$

Remark 12.5 The representations with x is even; y is even are excluded since

$|Q|, |L|, |H|, s_o, p$, by definitions related to odd composite and prime integers.

\oslash - odd noncomposite; \oplus - odd composite ;

Construction $SBR_{2n}::$

$$2n = 134 \quad n = 67$$

$$\oslash 1 + 133\oplus = 134(L)$$

$$\oslash 3 + 131\oslash = 134(H)$$

$$\oslash 5 + 129\oplus = 134(L)$$

$$\oslash 7 + 127\oslash = 134(H)$$

$$\oplus 9 + 125\oplus = 134(Q)$$

$$\oslash 11 + 123\oplus = 134(L)$$

$$\oslash 13 + 121\oplus = 134(L)$$

$$\begin{aligned}
\oplus 15 + 119 \oplus &= 134(Q) \\
\circlearrowleft 17 + 117 \oplus &= 134(L) \\
\circlearrowleft 19 + 115 \oplus &= 134(L) \\
\oplus 21 + 113 \circlearrowleft &= 134(L) \\
\circlearrowleft 23 + 111 \oplus &= 134(L) \\
\oplus 25 + 109 \circlearrowleft &= 134(L) \\
\oplus 27 + 107 \circlearrowleft &= 134(L) \\
\circlearrowleft 29 + 105 \oplus &= 134(L) \\
\circlearrowleft 31 + 103 \circlearrowleft &= 134(H) \\
\oplus 33 + 101 \circlearrowleft &= 134(L) \\
\oplus 35 + 99 \oplus &= 134(Q) \\
\circlearrowleft 37 + 97 \circlearrowleft &= 134(H) \\
\oplus 39 + 95 \oplus &= 134(Q) \\
\circlearrowleft 41 + 93 \oplus &= 134(L) \\
\circlearrowleft 43 + 91 \oplus &= 134(L) \\
\oplus 45 + 89 \oplus &= 134(Q) \\
\circlearrowleft 47 + 87 \oplus &= 134(L) \\
\oplus 49 + 85 \oplus &= 134(Q) \\
\oplus 51 + 83 \circlearrowleft &= 134(L) \\
\circlearrowleft 53 + 81 \oplus &= 134(L) \\
\oplus 55 + 79 \circlearrowleft &= 134(L) \\
\oplus 57 + 77 \oplus &= 134(Q) \\
\circlearrowleft 59 + 75 \oplus &= 134(L) \\
\circlearrowleft 61 + 73 \oplus &= 134(H) \\
\oplus 63 + 71 \circlearrowleft &= 134(L) \\
\oplus 65 + 69 \circlearrowleft &= 134(Q)
\end{aligned}$$

Data of direct computations for odd prime $n = 67$, at that it is took into account remark (3.2)

$$p^* = 30; s_o = 36; |Q| = 8; |L| = 20; |H| = 5; F = 33$$

The computations of parameters of binary representations of the positive integer $2n = 134$

with help of arithmetic stated above

at that $p, s_o, |Q|, |L|, |H|$, are took from the data of direct computations.

$$n = 134/2 = 67;$$

$$G = n - 1 = 67 - 1 = 66; \text{ by equation (11)}$$

$$|E| = (n - 1)/2 = 33; \text{ by equation (12)}$$

$$F = (n - 1)/2 = (67 - 1)/2 = 33; \text{ by equation (16)}$$

$$s_o = n - p - 1 = 67 - 30 - 1 = 36; \text{ by equation (18)}$$

$$|L| = n - p - 2|Q| - 1 = 67 - 30 - 16 - 1 = 20; \text{ by equation (27)}$$

$$|L| = p - 2|H| = 30 - 10 = 20; \text{ by equation (28)}$$

$$|H| = (2|Q| + 2p - n + 1)/2 = (16 + 60 - 67 + 1)/2 = 5; \text{ by equation (31)}$$

$|H| = (p - |L|)/2 = (30 - 20)/2 = 5$; by equation (29) $|Q| = (n - p - |L| - 1)/2 = (67 - 30 - 20 - 1)/2 = 8$; by equation (23)

$|Q| = (n - 2p + 2|H| - 1)/2 = (67 - 60 + 10 - 1)/2 = 8$; by equation (25)

$|Q| - |H| = 8 - 5 = 3$;

$|Q| - |H| = (n - 2p - 1)/2 = (67 - 60 - 1)/2 = 3$; by equation (33)

$p = 2|H| + |L| = 10 + 20 = 30$; by equation (7)

$s_o = 2|Q| + |L| = 16 + 20 = 36$; by equation (8)

$F = |Q| + |L| + |H| = 8 + 20 + 5 = 33$; by equation (10)

The compare of data of direct computations with computed values shows full coincidence of results.

13 The control of equation for $\Delta|Q|$

Exemple 13.1 The computation $\Delta|Q|$ for n is even: $n = 2, 2n = 4$.

Data of direct computations we take from exmp 12.1

$p = 2; s_o = 0; |Q| = 0; |L| = 0; |H| = 1; G = 1; F = 1$

$|Q|_m = ((n - p)^2/2n) = ((2 - 2)^2/4) = 0$. by equation (46)

$|H|_m = (p^2)/2n = (2^2)/4 = 1$; by equation (50)

$\Delta|Q| = |Q|_m - |Q| = 0 - 0 = 0$; by definition (7.1)

$\Delta|H| = |H|_m - |H| = 1 - 1 = 0$; by definition (7.2)

$|Q|_b = (n - 2p)/2 = (2 - 4)/2 = -1$; by equation (38)

The computation of $\Delta|Q|$ by equation (51).

$\Delta|Q| = |Q|_m - |H|_m - |Q|_b + \Delta|H| = 0 - 1 + 1 - 0 = 0$.

The compare of $\Delta|Q|$ computed by data of direct computations with computed value

by equation (51) shows full coincidence of results.

The computation $\Delta|Q|$ for n is odd prime: $n = 3; 2n = 6$.

Data of direct computations for odd prime $n = 3$ we take from exmp 12.2

$p^* = 2; s_o = 0; |Q| = 0; |L| = 0; |H| = 1; G = 2; F = 1$

$|Q|_b = |Q| - |H| = 0 - 1 = -1$. by equation (42)

$|Q|_m = ((n - p - 1)^2)/2n = ((3 - 2 - 1)^2/6) = 0$. by equation (47)

$|H|_m = (p^2)/2n = (2^2)/6 = 2/3$; by equation (50)

$\Delta|Q| = |Q|_m - |Q| = 0 - 0 = 0$; by definition (7.1)

$\Delta|H| = |H|_m - |H| = 2/3 - 1 = -1/3$; by definition (7.2)

$|Q|_b = (n - 2p - 1)/2 = (3 - 4 - 1)/2 = -1$; by equation (39)

The computation of $\Delta|Q|$ by equation (51).

$\Delta|Q| = |Q|_m - |H|_m - |Q|_b + \Delta|H| = 0 - 2/3 + 1 - 1/3 = 0$.

The compare of $\Delta|Q|$ computed by data of direct computations with computed value

by equation (51) shows full coincidence of results.

Exemple 13.2 The computation $\Delta|Q|$ for n is even: $n = 6, 2n = 12$.

Data of direct computations we take from exmp 3.3

$p = 5; s_o = 1; |Q| = 0; |L| = 1; |H| = 2; G = 5; F = 3$

$|Q|_m = ((n - p)^2/2n) = ((6 - 5)^2/12) = 1/12$. by equation (46)

$|H|_m = (p^2)/2n = (5^2)/12 = 25/12$; by equation (50)

$\Delta|Q| = |Q|_m - |Q| = 1/12 - 0 = 1/12$; by definition (7.1)

$\Delta|H| = |H|_m - |H| = 25/12 - 2 = 1/12$; by definition (7.2)

$|Q|_b = (n - 2p)/2 = (6 - 10)/2 = -2$; by equation (38)

The computation of $\Delta|Q|$ by equation (51) .

$\Delta|Q| = |Q|_m - |H|_m - |Q|_b + \Delta|H| = 1/12 - 25/12 + 2 + 1/12 = 1/12$.

The compare of $\Delta|Q|$ computed by data of direct computations with computed value

by equation (51) shows full coincidence of results.

Exemple 13.3 The computation $\Delta|Q|$ for n is odd prime: $n = 7; 2n = 14$

Data of direct computations for odd prime $n = 67$ we take from exmp 12.4

$p^* = 5; s_o = 1; |Q| = 0; |L| = 1; |H| = 2; G = 6; F = 3$

$|Q|_b = |Q| - |H| = 0 - 2 = -2$. by equation (42)

$|Q|_m = ((n - p - 1)^2)/2n = ((7 - 5 - 1)^2/14) = 1/14$. by equation (47)

$|H|_m = (p^2)/2n = (5^2)/14 = 25/14$; by equation (50)

$\Delta|Q| = |Q|_m - |Q| = 1/14 - 0 = 1/14$; by definition(7.1)

$\Delta|H| = |H|_m - |H| = 25/14 - 2 = -3/14$; by definition (7.2)

$|Q|_b = (n - 2p - 1)/2 = (7 - 10 - 1)/2 = -2$; by equation (39)

The computation of $\Delta|Q|$ by equation (51) .

$\Delta|Q| = |Q|_m - |H|_m - |Q|_b + \Delta|H| = 1/14 - 25/14 + 2 - 3/14 = 1/14$.

The compare of $\Delta|Q|$ computed by data of direct computations with computed value

by equation (51) shows full coincidence of results.

Exemple 13.4 The computation $\Delta|Q|$ for n is even: $n = 66, 2n = 132$.

Data of direct computations we take from exmp 12.5

$p = 32; s_o = 34; |Q| = 11; |L| = 12; |H| = 10; F = 33$

$|Q|_m = ((n - p)^2/2n) = ((66 - 32)^2/132) = 8,7575757$. by equation (46)

$|H|_m = (p^2)/2n = (32^2)/132 = 7,7575757$; by equation (50)

$\Delta|Q| = |Q|_m - |Q| = 8,7575757 - 11 = -2,2424243$; by definition (7.1)

$\Delta|H| = |H|_m - |H| = 7,7575757 - 10 = -2,2424243$; by definition (7.2)

$|Q|_b = (n - 2p)/2 = (66 - 64)/2 = 1$; by equation (38)

The computation of $\Delta|Q|$ by equation (51) .

$\Delta|Q| = |Q|_m - |H|_m - |Q|_b + \Delta|H| = 8,7575757 - 7,7575757 - 1 - 2,2424243 = -2,2424243$.

The compare of $\Delta|Q|$ computed by data of direct computations with computed value

by equation (51) shows full coincidence of results.

Exemple 13.5 The computation $\Delta|Q|$ for n is odd : $n = 69; 2n = 138$.
Data of direct computations we take from expm 12.6

$$p = 33; s_o = 35; |Q| = 10; |L| = 15; |H| = 9;$$

$$|Q|_m = ((n - p - 1)^2 / 2n) = ((69 - 33 - 1)^2 / 138) = 8,8768115. \text{ by equation (47)}$$

$$|H|_m = (p^2) / 2n = (33^2) / 138 = 7,8913043; \text{ by equation (50)}$$

$$\Delta|Q| = |Q|_m - |Q| = 8,8768115 - 10 = -1,123189; \text{ by definition (7.1)}$$

$$\Delta|H| = |H|_m - |H| = 7,8913043 - 9 = -1,1086957; \text{ by definition (7.2)}$$

$$|Q|_b = (n - 2p - 1) / 2 = (69 - 66 - 1) / 2 = 1 ; \text{ by equation (39)}$$

The computation of $\Delta|Q|$ by equation (51) .

$$\Delta|Q| = |Q|_m - |H|_m - |Q|)b + \Delta|H| = 8,8768115 - 7,8913043 - 1 - 1,1086957 = -1,123189 .$$

The compare of $\Delta|Q|$ computed by data of direct computations with computed value

by equation (51) shows full coincidence of results.

Exemple 13.6 The computation $\Delta|Q|$ for n is odd prime: $n = 67; 2n = 134$.

Data of direct computations for odd prime $n = 67$ we take from expm 12.7

$$p = 30; s_o = 36; |Q| = 8; |L| = 20; |H| = 5;$$

$$|Q|_b = |Q| - |H| = 8 - 5 = 3 . \text{ by equation (42)}$$

$$|Q|_m = ((n - p - 1)^2) / 2n = ((67 - 30 - 1)^2 / 134) = 9,6716417. \text{ by equation (47)}$$

$$|H|_m = (p^2) / 2n = (30^2) / 134 = 6,7164179; \text{ by equation (50)}$$

$$\Delta|Q| = |Q|_m - |Q| = 9,6716417 - 8 = 1,6716417; \text{ by definition (7.1)}$$

$$\Delta|H| = |H|_m - |H| = 6,7164179 - 5 = 1,7164179; \text{ by definition (7.2)}$$

$$|Q|_b = (n - 2p - 1) / 2 = (67 - 60 - 1) / 2 = 3 ; \text{ by equation (39)}$$

The computation of $\Delta|Q|$ by equation (51) .

$$\Delta|Q| = |Q|_m - |H|_m - |Q|)b + \Delta|H| = 2,9552238 - 3 + 1,7164179 = 1,6716417 .$$

The compare of $\Delta|Q|$ computed by data of direct computations with computed value

by equation (51) shows full coincidence of results.

Hence equation (51) is true $\forall n > 1$.

13.1 The examples of the correspondence of $2n$ included between neighboring values of p to it

For example we consider the set of XUY for $2n = 32$. We can see that the set of XUY consist of several sets. To each set of subsets is corresponded at the same value of p . Next for example of computation s_o we show that use at the same value of p for corresponding set gives corrected result which is controlled by direct computation s_o for each subsets of corresponding set.

Exemple 13.7 *Let we have the sat of numbers*

$X U Y \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, \}$ U

$\{17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31\}$

at that $n = 16$ is excluded by remark (3.1).

Corollaries (3.1 and 3.2 and 3.3) are changed following way:

$s_e^* = s_e - 1; s_o^* = s_o - 1 : p^* = p - 1$

since in this case is excluded one n .

We can see that :

$p = 2$ correspond to subset for $2n = 4$

$p = 3$ correspond to subset for $2n = 6$

$p = 4$ correspond to subsets for $2n = 8, 10$

$p = 5$ correspond to subset for $2n = 12$

$p = 6$ correspond to subsets for $2n = 14, 16$

$p = 7$ correspond to subset for $2n = 18$

$p = 8$ correspond to subsets for $2n = 20, 22$

$p = 9$ correspond to subsets for $2n = 24, 26, 28$

$p = 10$ correspond to subset for $2n = 30$

We control the following correspondences: $p = 4$ correspond $2n = 8, 10$

subsets: $X U Y \{1, 2, 3\} U \{5, 6, 7\}$

$n = 4$ is excluded by rem.(3.1).

$X U Y \{1, 2, 3, 4\} U \{6, 7, 8, 9\}$ $n = 5$ is excluded (rem3.1).

$p = 6$ correspond $2n = 14, 16$ subsets: $X U Y \{1, 2, 3, 4, 5, 6\} U$

$\{8, 9, 10, 11, 12, 13\}$ $n = 7$ is excluded (rem3.1).

$X U Y \{1, 2, 3, 4, 5, 6, 7\} U \{9, 10, 11, 12, 13, 14, 15\}$ $n = 8$ is excluded (rem3.1).

$p = 8$ correspond $2n = 20, 22$ subsets: $X U Y \{1, 2, 3, 4, 5, 6, 7, 8, 9\} U$

$\{11, 12, 13, 14, 15, 16, 17, 18, 19\}$ $n = 10$ is excluded (rem3.1). ;

$X U Y \{1, 2, 3, 4, 5, 6, 7, 9, 10\} U$

$\{12, 13, 14, 15, 16, 17, 18, 19, 20, 21\}$ $n = 11$ is excluded (rem3.1).

$p = 9$ correspond $2n = 24, 26, 28$ subsets: $X U Y \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11\} U$

$\{13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23\}$ $n = 12$ is excluded (rem3.1).

$X U Y \{1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12\} U$

$\{14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25\}$ $n = 13$ is excluded (rem3.1).

$X U Y \{1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13\}U$

$\{15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27\}$ $n = 14$ is excluded (rem3.1).

By equation (17) we get :

for $2n - 8$; $n = 4$; $p = 4$.

$s_o = n - p = 4 - 4 = 0$.

By direct computation for $2n = 8$, $s_o = 0$.

for $2n - 10$; $n = 5$; $p^* = 3$ (corrolary (3.3) with real changes).

By equation (18) we get :

$s_o = n - p - 1 = 5 - 3 - 1 = 1$

By direct computation for $2n = 10$, $s_o = 1$.

for $2n - 14$; $n = 7$; $p^* = 5$ (corrolary (3.3) with real changes) .

By equation (18) we get :

$s_o = n - p - 1 = 7 - 5 - 1 = 1$

By direct computation for $2n = 14$, $s_o = 1$.

for $2n - 16$; $n = 8$; $p = 6$.

By equation (17) we get :

$s_o = n - p = 8 - 6 = 2$

By direct computation for $2n = 16$, $s_o = 2$.

for $2n - 20$; $n = 10$; $p = 8$.

By equation (17) we get :

$s_o = n - p = 10 - 8 = 2$

By direct computation for $2n = 20$, $s_o = 2$.

for $2n - 22$; $n = 11$; $p^* = 7$ (corrolary (3.3) with real changes) .

By equation (18) we get :

$s_o = n - p - 1 = 11 - 7 - 1 = 3$

By direct computation for $2n = 22$, $s_o = 3$.

for $2n - 24$; $n = 12$; $p = 9$.

By equation (17) we get :

$s_o = n - p = 12 - 9 = 3$

By direct computation for $2n = 24$, $s_o = 3$.

for $2n - 26$; $n = 13$; $p^* = 8$ (corrolary (3.3) with real changes) .

By equation (18) we get :

$s_o = n - p - 1 = 13 - 8 - 1 = 4$

By direct computation for $2n = 26$, $s_o = 4$.

for $2n - 28$; $n = 14$; $p = 9$.

By equation (17) we get :

$s_o = n - p = 14 - 9 = 5$

By direct computation for $2n = 28$, $s_o = 5$.

Thus for several $2n$ it takes at the same p at computations by equations $ABR2n$ and these computations are correct.

Hence for each p correspond to finite set of $2n$ included between neighboring values of p

References

- [1] J.B.Zeldovich , A.D.Myshkis: *The elements of applied mathematics*, pgs: 562-567, publishers "Nauka",Moscow (1965)
- [2] Warner, S: *Modern Algebra*, page 6. Dover Press, 1990.
- [3] Singh, S. (2009, August 27). Cartesian product. Retrieved from the Connexions Web site: <http://cnx.org/content/m15207/1.5/>