

# Quantum Impedances: Background Independent Relations between Gravity and Electromagnetism

Peter Cameron\*  
Strongarm Studios  
Mattituck, NY USA

(Dated: February 3, 2013)

As every circuit designer knows, the flow of energy is governed by impedance matching. Classical or quantum impedances, mechanical or electromagnetic, fermionic or bosonic, topological,... To understand the flow of energy it is essential to understand the relations between the associated impedances. The connection between electromagnetism and gravitation can be made explicit by examining the impedance mismatch between the electrically charged Planck particle and the electron and photon. This mismatch is shown to be the ratio of the gravitational and electromagnetic forces.

## INTRODUCTION

This letter presents a preliminary exploration of the role of background independent quantum impedances[1–3] in gravitation.

In the earlier work on impedances, the main players were the photon and the electron. That earlier work presented a model in which the unstable particles were seen as resonant excitations of the network of electron impedances, the ‘scattering matrix’, by the photon.

The present work looks at the extreme high energy limit of the interaction of the photon and the electron with a third player, the Planck particle[4]. In what follows the Planck particle is presented, and its gravitational and Coulomb interactions with the electron are introduced.

The impedance mismatch between the Planck particle and photon and electron is then shown to be equal to the ratio of the gravitational and electromagnetic forces at the level of a few parts per billion (the empirically determined Newtonian gravitational constant  $G$  is measured with an accuracy of about a part in ten thousand).

This suggests that the gravitational force might be taken to be electromagnetic in origin. The enormous difference in strengths can be understood in terms of the equally enormous impedance mismatch to the Planck particle.

## THE PLANCK PARTICLES

We have two ways to define Planck particles, one each for massive and massless particles.

For **massive particles** we equate the reduced Compton wavelength and the Schwarzschild radius

$$\frac{\hbar}{mc} = \frac{mG}{c^2} \quad (1)$$

Solving for the mass  $m$  gives the reduced Planck mass[5]

$$m_{Pl} = \sqrt{\frac{\hbar c}{G}} \simeq 2.1765 * 10^{-8} kg \simeq 10^{19} GeV \quad (2)$$

and the reduced Planck length

$$L_{Pl} = \frac{\hbar}{m_{Pl}c} = \sqrt{\frac{\hbar G}{c^3}} \simeq 1.6162 * 10^{-35} m \quad (3)$$

The Planck particle as thus defined is strictly mechanical. It has no electromagnetic properties.

The Planck particle may be similarly defined in terms of the **massless photon**, again as one whose wavelength and Schwarzschild radius are equal. For the photon it is the simple matter of the equivalence  $E = mc^2$  of energy and mass. From that and  $E = h\nu$  we have the photon wavelength

$$\lambda = \frac{c}{\nu} = \frac{hc}{E} = \frac{hc}{mc^2} = \frac{h}{mc} \quad (4)$$

and, excepting the factor of  $2\pi$ , can proceed as for the massive particles with equation (1).

The Planck particle as thus defined is electromechanical, an electromagnetic black hole[6].

## THE INTERACTIONS

The **gravitational force** between the Planck particle and the electron can be written as

$$F_{grav} = G \frac{m_e m_{Pl}}{\lambda_e^2} = 8.873 419 056 * 10^{-24} N \quad (5)$$

where  $m_e$  is the mass and  $\lambda_e$  the reduced Compton wavelength of the electron.

The **Coulomb force** between the electron and a Planck particle carrying the charge of a positron is

$$F_{Coul} = \frac{1}{4\pi\epsilon_0} \frac{e^2}{\lambda_e^2} = 1.547 138 513 * 10^{-3} N \quad (6)$$

The **ratio** of these two forces is

$$ratio_F = \frac{F_{grav}}{F_{Coul}} = 1.743 565 251 * 10^{-20} \quad (7)$$




FIG. 1. Photon and quantum Hall impedances as a function of spatial scale as defined by photon energy. The role of the fine structure constant  $\alpha$  is prominent in the figure

## THE IMPEDANCES

Near and far field photon impedances[7] and the scale invariant quantum Hall impedance[8] are shown in the figure. The wavelength of the photon is the Compton wavelength of the electron. The energy of such a photon is 0.511MeV, the mass of the electron.

Similarly, we defined the Planck particle in two ways, one each for massive and massless particles, both with the same wavelength and energy. This is true in general. If the wavelength of a photon is the same as the Compton wavelength of a particle, then particle and photon have the same energy.

There are two possibilities for calculating the impedance mismatch between the electron and the Planck particle, either by matching them directly or via the intermediary of the photon. The simpler approach, and the one in greater harmony with QED, utilizes the photon. There are at least two ways to accomplish this:

- match the near field impedance of a .511MeV photon to the scale invariant quantum Hall impedance of the Planck particle at the Planck length
- match the dipole mode of the electron to a  $10^{19}$ GeV photon at the Planck length.

The first approach assumes that one of the Planck particle impedances can be taken to be the quantum Hall impedance<sup>1</sup>, which is well documented in the literature. It requires that the Planck particle be given the attribute of electric charge, in this case the charge of the positron.

The second approach requires introduction of the quantum dipole impedance of the electron[2]. Both produce the same result. In the interest of simplicity and brevity, only the first will be presented here.

<sup>1</sup> the equality between the quantum Hall impedance and an inertial impedance associated with the centripetal force was presented in earlier work [2, 3]

The near and far field amplitudes of the electric and magnetic components of the dipole impedance seen by the photon can be written as [7]

$$Z_E = Z_0 \left| \frac{1 + \frac{\lambda}{ir} + \left(\frac{\lambda}{ir}\right)^2}{1 + \frac{\lambda}{ir}} \right| \quad (8)$$

$$Z_M = Z_0 \left| \frac{1 + \frac{\lambda}{ir}}{1 + \frac{\lambda}{ir} + \left(\frac{\lambda}{ir}\right)^2} \right| \quad (9)$$

where  $\lambda$  is the photon wavelength,  $r$  is the length scale of interest, and  $Z_0 \simeq 377\Omega$  is the free space impedance seen by the photon in the far field. These are the photon impedances plotted in the figure. On the scale of this page the Planck particle sits slightly less than a full page width to the left of the electron Compton wavelength. The singularity is at infinity, totally mismatched, decoupled from the photon.

Looking first at the **electric component** of the photon near field impedance, and taking  $\lambda$  to be the electron Compton wavelength and  $r$  the Planck length gives

$$Z_E = 9.001\ 262\ 858 * 10^{24}\Omega \quad (10)$$

and

$$ratio_{Z_E} = \frac{Z_0}{\alpha Z_E} = 5.735\ 580\ 908 * 10^{-21} \quad (11)$$

At the few parts per billion level this is the ratio of the gravitational to Coulomb force calculated in equation (7), so that

$$\frac{ratio_F}{ratio_{Z_E}} = 1.000\ 000\ 004 \quad (12)$$

The accuracy of this result might be surprising at first glance, given the much larger experimental uncertainty in the gravitational constant  $G$ . However  $G$  is present in both numerator and denominator of this ratio of ratios and hence cancels out. *The result is independent of the value of  $G$ .*

One wonders whether it would be reasonable to suggest that the quantum impedance model is gravitationally gauge invariant, and how this might be related to background independence.

The same calculation for the **magnetic component** of the photon near field impedance gives

$$Z_M = 1.576\ 731\ 302 * 10^{-20}\Omega \quad (13)$$

and

$$ratio_{Z_M} = \frac{Z_M}{\alpha Z_0} = 5.735\ 580\ 908 * 10^{-21} \quad (14)$$

so that, as for the electric component of the impedance

$$\frac{ratio_F}{ratio_{Z_M}} = 1.000\ 000\ 004 \quad (15)$$

## DISCUSSION

The reader might object that the Planck particle exists only in theory, that if such a particle could somehow be produced it would not be stable, would immediately radiate its energy away. However, the possibility of interaction with the virtual Planck particle remains, just as interaction with the vacuum permits renormalization of QED in theory, or the measurement of the Casimir force in practice.

The reasoning presented in the previous sections was adopted in the interest of making the simplest possible presentation of the role of quantum impedances in gravitation. It therefore employed only those impedances found in the commonly accepted body of physics knowledge, namely the photon and quantum Hall impedances.

The step to generalized quantum impedances requires the introduction of a model[2, 3] that is not needed for the present purpose. The reader is encouraged to explore that model, in the hope that the logical foundation of the calculations presented here will become more transparent and additional new possibilities will reveal themselves.

## CONCLUSION

Classical or quantum, mechanical or electromagnetic, fermionic or bosonic, topological,... The flow of energy is governed by impedance matching.

This understanding is common to electrical engineers, some mechanical engineers, most accelerator physicists, and a subset of the physics community, working almost exclusively in condensed matter. It requires a sophistication of network analysis now accessible to electrical engineers and complex systems specialists.

The Planck particle provides a much needed additional anchor point for the iterative software that might eventually sort out the mode structures and coupling mechanisms, and perhaps solve the localization problem[9–13].

If the gravitational mass of the electron follows from mismatched electromagnetic interaction with the virtual Planck particle, then this will be true for all massive particles, each via its own routes through the impedance networks.

It is interesting to speculate upon possible relationships between string and loop theory and quantum impedances in the vicinity of the Planck scale[14].

## ACKNOWLEDGEMENTS

The author thanks Michael Suisse for discussions, literature searches, and critical readings of the many drafts of this letter, and Corky Maul for helpful conversations that focused attention on the relations between quantum impedances and gravity.

Thanks are also due to the referee who pointed out an exact relationship between the ratio of the forces and that of the electron and Planck particle masses, shown here in terms of their electromagnetic impedances as well.

$$\frac{F_{grav}}{F_{Coul}} = \frac{m_e}{\alpha m_{Pl}} = \frac{Z_0}{\alpha Z_E} = \frac{Z_M}{\alpha Z_0}$$

The role of the fine structure constant in these equations will be explored in a note to follow[14].

---

\* petethepop@aol.com

- [1] Cameron, P., "The Two Body Problem and Mach's Principle", submitted to Am. Jour. Phys (1975), in revision. The unrevised version of this note was published as an appendix to the Electron Impedances note [2].
- [2] Cameron, P., "Electron Impedances", *Apeiron*, vol.18, no.2, p.222-253 (2011). <http://redshift.vif.com/JournalFiles/V18N02PDF/V18N2CAM.pdf>
- [3] Cameron, P., "Generalized Quantum Impedances: A Background Independent Model for the Unstable Particles" (July 2012). <http://vixra.org/abs/1207.0022>
- [4] [http://en.wikipedia.org/wiki/Planck\\_particle](http://en.wikipedia.org/wiki/Planck_particle)
- [5] SI units are used throughout. All constants are CODATA2006 except alpha, which is taken from G. Gabrielse, "New Measurement of the Electron Magnetic Moment and the Fine Structure Constant", available at <http://arxiv.org/pdf/0801.1134.pdf>
- [6] Wheeler, J., "Geons". *Phys. Rev.* 97 (2): 511 (1957).
- [7] Capps, C., "Near Field or Far Field?", *Electronic Design News*, p.95 (16 Aug 2001). <http://edn.com/design/communications-networking/4340588/Near-field-or-far-field->
- [8] von Klitzing, K. et al, "New method for high-accuracy determination of the fine-structure constant based on quantized Hall resistance", *PRL*, vol.45, no.6, p.494-497 (1980). <http://140.120.11.121/~wanwan/Reference/QHE/Klitzing%20QHE%201980.pdf>
- [9] Lehnert, B., "On the Self-Confinement of Electromagnetic Radiation", *TRITA-EPP-79-13* (1979) <http://cds.cern.ch/record/1068895/>
- [10] Williamson, J., and van der Mark, M. "Is the Electron a Photon with Toroidal Topology?", *Annales de la Fondation Lous deBroglie*, vol.22, no.2, 133 (1997). <http://www.cybsoc.org/electron.pdf>  
This paper has a good extensive bibliography.
- [11] Vaga, T., "Particles at Cutoffs in the Electromagnetic Spectrum", *Phys. Essays*, vol.14, no.3, p.203-207 (2001). <http://physicsessays.org/doi/abs/10.4006/1.3025484?journalCode=phes>
- [12] Ranada, A., "On Topology and Electromagnetism", *Ann. Phys. (Berlin)* 524, No. 2, A35A37 (2012). <http://onlinelibrary.wiley.com/doi/10.1002/andp.201100710/pdf>
- [13] Arrayas, M. and Trueba, J. "Exchange of helicity in a knotted electromagnetic field", *Ann. Phys. (Berlin)* 524, No. 2, 7175 (2012). <http://onlinelibrary.wiley.com/doi/10.1002/andp.201100119/pdf>
- [14] Cameron, P. and Suisse, M., "Background Independent Quantum Impedances in String and Loop Theory", to be published.