

The relation of colour charge to electric charge

Dirac has shown how the Klein-Gordon equation can be factored into two linear parts using 4x4 Dirac gamma matrices.

[Dirac, P.A.M., *The Principles of Quantum Mechanics*, 4th edition (Oxford University Press) ISBN 0-19-852011-5]

$$(\partial_t^2 - \partial_x^2 - \partial_y^2 - \partial_z^2 + m^2) I = (-i[s\gamma^0\partial_t + r\gamma^1\partial_x + g\gamma^2\partial_y + b\gamma^3\partial_z] - wmI) (i[s\gamma^0\partial_t + r\gamma^1\partial_x + g\gamma^2\partial_y + b\gamma^3\partial_z] - wmI)$$

where r,g,b and s,w equal +1 or -1.

For leptons r,g,b all equal -1 and for quarks two of r,g,b are equal to +1 and the third equals -1. The signs are all negated for anti-particles as in the equation above.

When s = +1, count the number of plus signs (say) for r,g,b which is 0 for leptons and 2 for quarks.

When s = -1, count the number of minus signs (say) for r,g,b which is 3 for leptons and 1 for quarks.

For material particles r,g,b all equal -1 which is always true for leptons and true for three distinct quarks with r,g,b equal to -1 separately or a quark and an appropriate anti-quark.

Let $\hat{Y} = i\gamma_0\gamma_1\gamma_2\gamma_3\gamma_4$ where $\gamma_0, \gamma_1, \gamma_2, \gamma_3, \gamma_4$ are vectors which anti-commute and where:

$$\gamma_1^2 = \gamma_2^2 = \gamma_3^2 = -I \qquad \gamma_0^2 = \gamma_4^2 = I$$

$$\text{Then: } \gamma_0\hat{Y} = \hat{Y}\gamma_0 \quad \gamma_1\hat{Y} = \hat{Y}\gamma_1 \quad \gamma_2\hat{Y} = \hat{Y}\gamma_2 \quad \gamma_3\hat{Y} = \hat{Y}\gamma_3 \quad \gamma_4\hat{Y} = \hat{Y}\gamma_4 \qquad \hat{Y}^2 = I$$

$$\text{Let: } \hat{s} = \frac{1}{2}(I + s\hat{Y}) \quad \hat{r} = \frac{1}{2}(I + r\hat{Y}) \quad \hat{g} = \frac{1}{2}(I + g\hat{Y}) \quad \hat{b} = \frac{1}{2}(I + b\hat{Y}) \quad \hat{w} = \frac{1}{2}(I + w\hat{Y})$$

$$\hat{s}^2 = \hat{s} \quad \hat{r}^2 = \hat{r} \quad \hat{g}^2 = \hat{g} \quad \hat{b}^2 = \hat{b} \quad \hat{w}^2 = \hat{w}$$

A charged particle moving in an electro-magnetic-weak field will have $\partial_t, \partial_x, \partial_y, \partial_z, \partial_m$ modified by the scalar-vector-scalar potential of the field, where $\partial_t, \partial_x, \partial_y, \partial_z, \partial_m$ do not commute with each other. Thus:

$$\begin{aligned} & (\hat{s}\gamma_0\partial_t + \hat{r}r\gamma_1\partial_x + \hat{g}g\gamma_2\partial_y + \hat{b}b\gamma_3\partial_z + \hat{w}w\gamma_4\partial_m) (\hat{s}\gamma_0\partial_t + \hat{r}r\gamma_1\partial_x + \hat{g}g\gamma_2\partial_y + \hat{b}b\gamma_3\partial_z + \hat{w}w\gamma_4\partial_m) \\ &= \hat{s}\partial_t^2 - \hat{r}\partial_x^2 - \hat{g}\partial_y^2 - \hat{b}\partial_z^2 + \hat{w}\partial_m^2 \\ & \quad + \hat{s}\hat{w}w\gamma_0\gamma_4(\partial_t\partial_m - \partial_m\partial_t) \\ & \quad + \hat{s}\gamma_0[\hat{r}r\gamma_1(\partial_t\partial_x - \partial_x\partial_t) + \hat{g}g\gamma_2(\partial_t\partial_y - \partial_y\partial_t) + \hat{b}b\gamma_3(\partial_t\partial_z - \partial_z\partial_t)] \quad (= 0 \text{ for a neutrino}) \\ & \quad + \hat{w}w\gamma_4[\hat{r}r\gamma_1(\partial_m\partial_x - \partial_x\partial_m) + \hat{g}g\gamma_2(\partial_m\partial_y - \partial_y\partial_m) + \hat{b}b\gamma_3(\partial_m\partial_z - \partial_z\partial_m)] \\ & \quad + \hat{r}r\hat{g}g\gamma_1\gamma_2(\partial_x\partial_y - \partial_y\partial_x) + \hat{g}g\hat{b}b\gamma_2\gamma_3(\partial_y\partial_z - \partial_z\partial_y) + \hat{b}b\hat{r}r\gamma_3\gamma_1(\partial_z\partial_x - \partial_x\partial_z) \\ &= \hat{s}\partial_t^2 - \hat{r}\partial_x^2 - \hat{g}\partial_y^2 - \hat{b}\partial_z^2 + \hat{w}\partial_m^2 \\ & \quad + \hat{s}\hat{w}w\gamma_0\gamma_4(\partial_t\partial_m - \partial_m\partial_t) \\ & \quad + \hat{s}\gamma_0[\hat{r}r\gamma_1(\partial_t\partial_x - \partial_x\partial_t) + \hat{g}g\gamma_2(\partial_t\partial_y - \partial_y\partial_t) + \hat{b}b\gamma_3(\partial_t\partial_z - \partial_z\partial_t)] \quad (= 0 \text{ for a neutrino}) \\ & \quad + \hat{w}w\gamma_4[\hat{r}r\gamma_1(\partial_m\partial_x - \partial_x\partial_m) + \hat{g}g\gamma_2(\partial_m\partial_y - \partial_y\partial_m) + \hat{b}b\gamma_3(\partial_m\partial_z - \partial_z\partial_m)] \\ & \quad - \hat{r}\hat{g}b\gamma_1\gamma_2(\partial_x\partial_y - \partial_y\partial_x) - r\hat{g}\hat{b}\gamma_2\gamma_3(\partial_y\partial_z - \partial_z\partial_y) - \hat{r}g\hat{b}\gamma_3\gamma_1(\partial_z\partial_x - \partial_x\partial_z) \end{aligned}$$