

Florentin SMARANDACHE



INTRODUCTION TO
NEUTROSOPHIC MEASURE,
NEUTROSOPHIC INTEGRAL,
AND NEUTROSOPHIC PROBABILITY

Florentin Smarandache

**INTRODUCTION TO
NEUTROSOPHIC MEASURE,
NEUTROSOPHIC INTEGRAL,
AND NEUTROSOPHIC PROBABILITY**

Sitech - Education

*Sitech, 200402 Craiova, Romania, Aleea teatrului, Nr. 2, Bloc T1, parter, Tel/Fax
0251/414003, sitech.ro*

and

*Education Publishing, 1313 Chesapeake Avenue, Columbus, Ohio 43212, USA, Tel.
614/4850721*

*@ All rights reserved. This book is protected by copyright. No part of this book may
be reproduced in any form or by any means, including photocopying or using any
information storage and retrieval system without written permission from the
copyright owner.*

Peer-Reviewers:

Prof. Valeri Kroumov, Okayama University of Science, Japan.

Dr. Ștefan Vlăduțescu, University of Craiova, Romania.

Marius Coman, researcher, Bucharest, Romania.

Dr. Linfan Mao, Chinese Academy of Sciences, Beijing, China.

Many books can be downloaded from the following

Digital Library of Science:

<http://fs.gallup.unm.edu/eBooks-otherformats.htm>

EAN: 9781599732534

ISBN: 978-1-59973-253-4

CONTENTS

Neutrosophic Science (preface): 9

Chapter 1.

Introduction to Neutrosophic Measure: 11

- 1.1. Introduction: 12
- 1.2. Definition of Neutrosophic Measure: 12
- 1.3. Neutrosophic Measure Space: 13
- 1.4. Normalized Neutrosophic Measure: 13
- 1.5. Finite Neutrosophic Measure Space: 14
- 1.6. σ -Finite Neutrosophic Measure: 14
- 1.7. Neutrosophic Axiom of Non-Negativity: 14
- 1.8. Measurable Neutrosophic Set and Measurable Neutrosophic Space: 14
- 1.9. Neutrosophic Measurable Function: 15
- 1.10. Neutrosophic Probability Measure: 15
- 1.11. Neutrosophic Category Theory: 15
- 1.12. Properties of Neutrosophic Measure: 17
- 1.13. Neutrosophic Measure Continuous from Below or Above: 18
- 1.14. Generalizations: 18
- 1.15. Examples: 19

Chapter 2.

Introduction to Neutrosophic Integral: 22

- 2.1. Definition of Neutrosophic Integral: 23
- 2.2. First Example of Neutrosophic Integral:
Indeterminacy Related to Function's Values: 23
- 2.3. Second Example of Neutrosophic Integral:
Indeterminacy Related to Lower Limit: 25

Chapter 3.

Introduction to Neutrosophic Probability: 27

- 3.1. First Example of Indeterminacy: 28
- 3.2. Second Example of Indeterminacy: 30
- 3.3. Third Example of Indeterminacy: 30
- 3.4. Fourth Example of Indeterminacy: 31
- 3.5. Fifth Example of Indeterminacy: 32
- 3.6. Sixth Example of Indeterminacy: 32
- 3.7. The Seventh Example of Indeterminacy: 32
- 3.8. The Eighth Example of Indeterminacy: 33
- 3.9. Ninth Example of Indeterminacy: 33
- 3.10. Tenth Example of Indeterminacy: 33
- 3.11. Eleventh Natural Example of
Indeterminacy: 33
- 3.12. Example of a Neutrosophic Continuous
Random Variable: 34
- 3.13. First Types of Indeterminacies: 35
- 3.14. Second Types of Indeterminacies: 35
- 3.15. Distinction between Indeterminacy and
Randomness: 38
- 3.16. Neutrosophic Random Variables: 39
- 3.17. Many Possible Neutrosophic Measures and
Probabilities: 40

- 3.18. Definition of Neutrosophic Probability: 41
- 3.19. Neutrosophic Probability vs. Imprecise Probability: 42
- 3.20. Sigma-Algebra of Events: 45
- 3.21. Definition of Classical Probability: 45
- 3.22. Neutrosophic Sigma-Algebra of Events: 45
- 3.23. Neutrosophic Probability Measure: 46
- 3.24. Neutrosophic Probability Mass Function: 47
- 3.25. Neutrosophic Probability Axioms: 48
- 3.26. Consequences of Neutrosophic Probability Axioms: 50
- 3.27. Interpretations of the Neutrosophic Probability: 52
- 3.28. Neutrosophic Notions: 52
- 3.29. Example with Neutrosophic Frequentist Probability: 53
- 3.30. Example with Neutrosophic Frequentist Probability on a Neutrosophic Product Space: 55
- 3.31. Example with Double Indeterminacy: 57
- 3.32. Neutrosophic Example with Tossing a Coin Multiple Times: 58
- 3.33. Example with Sum of Chances of an Event: 62
- 3.34. Paraconsistent Neutrosophic Probability: 63
- 3.35. Incomplete Neutrosophic Probability: 64

- 3.36. Neutrosophic Mutually Exclusive Events: 65
- 3.37. Neutrosophic Experimental Probability: 67
- 3.38. Neutrosophic Survey: 68
- 3.39. Neutrosophic Conditional Probability for Independent Events: 68
- 3.40. Neutrosophic Probability of an Impossible Event: 69
- 3.41. Neutrosophic Probability of a Sure Event: 69
- 3.42. Neutrosophic Bayesian Rule: 69
- 3.43. Neutrosophic Multiplicative Rule: 71
- 3.44. Neutrosophic Negation (or Neutrosophic Probability of Complement Events): 74
- 3.45. De Morgan's Neutrosophic Laws: 74
- 3.46. Neutrosophic Double Negation: 75
- 3.47. Neutrosophic Expected Value: 76
- 3.48. Neutrosophic Probability and Neutrosophic Logic Used in The Soccer Games: 77
- 3.49. A Neutrosophic Question: 81
- 3.50. Neutrosophic Discrete Probability Spaces: 84
- 3.51. Classification of Neutrosophic Probabilities: 87
- 3.52. The Fundamental Neutrosophic Counting Principle: 88

- 3.53. A Formula for the Fusion of Subjective Neutrosophic Probabilities: 89
- 3.54. Numerical Example of Fusion of Subjective Neutrosophic Probabilities: 93
- 3.55. General Formula for Fusing Classical Subjective Probabilities Provided by Two Sources: 95
- 3.56. Different Ways of Combining Neutrosophic Subjective Probabilities Provided by Two Sources: 95
- 3.57. Neutrosophic Logic Inference type in Fusing Subjective Neutrosophic Probabilities: 98
- 3.58. Neutrosophic Logic vs. Subjective Neutrosophic Probability: 101
- 3.59. Removing Indeterminacy: 101
- 3.60. n-Valued Refined Neutrosophic Probability Space and Neutrosophic Probability: 102
- 3.61. Neutrosophic Markov Chain: 106
- 3.62. Applications of Neutrosophics: 114

Chapter 4.

Neutrosophic Subjects for Future Research: 115

Neutrosophic Subjects: 116
References: 118

ADDENDA: 121

- Books on Neutrosophics: 122
- More Articles on Neutrosophics: 125
- Seminars on Neutrosophics: 138
- International Conferences on
Neutrosophics: 139
- Ph. D. Dissertations on Neutrosophics:
139-140

Neutrosophic Science

(Preface)

Since the world is full of indeterminacy, the neutrosophics found their place into contemporary research.

*We now introduce for the first time the notions of **neutrosophic measure** and **neutrosophic integral**. We develop the 1995 notion of **neutrosophic probability** and give many practical examples.*

Neutrosophic Science means development and applications of neutrosophic logic/set/measure/integral/probability etc. and their applications in any field.

It is possible to define the neutrosophic measure and consequently the neutrosophic integral and neutrosophic probability in many ways, because there are various types of indeterminacies, depending on the problem we need to solve.

Indeterminacy is different from randomness. Indeterminacy can be caused by physical space materials and type of construction, by items involved in the space, or by other factors.



*Fig. 1. An example of indeterminacy.
What is tossed, 1, 3 or 5?*

Neutrosophic measure is a generalization of the classical measure for the case when the space contains some indeterminacy.

*Neutrosophic probability is a generalization of the classical and imprecise probabilities.
Several classical probability rules are adjusted in the form of neutrosophic probability rules.*

Finally, the neutrosophic probability is extended to n -valued refined neutrosophic probability.

The author

Chapter 1.
Introduction to Neutrosophic Measure

1.1. Introduction.

Let $\langle A \rangle$ be an item. $\langle A \rangle$ can be a notion, an attribute, an idea, a proposition, a theorem, a theory, etc. And let $\langle \text{anti}A \rangle$ be the opposite of $\langle A \rangle$; while $\langle \text{neut}A \rangle$ be neither $\langle A \rangle$ nor $\langle \text{anti}A \rangle$ but the neutral (or indeterminacy, unknown) related to $\langle A \rangle$.

For example, if $\langle A \rangle = \text{victory}$, then $\langle \text{anti}A \rangle = \text{defeat}$, while $\langle \text{neut}A \rangle = \text{tie game}$.

If $\langle A \rangle$ is the degree of truth value of a proposition, then $\langle \text{anti}A \rangle$ is the degree of falsehood of the proposition, while $\langle \text{neut}A \rangle$ is the degree of indeterminacy (i.e. neither true nor false) of the proposition.

Also, if $\langle A \rangle = \text{voting for a candidate}$, $\langle \text{anti}A \rangle = \text{voting against that candidate}$, while $\langle \text{neut}A \rangle = \text{not voting at all, or casting a blank vote, or casting a black vote}$.

In the case when $\langle \text{anti}A \rangle$ does not exist, we consider its measure be null $\{ m(\text{anti}A) = 0 \}$. And similarly when $\langle \text{neut}A \rangle$ does not exist, its measure is null $\{ m(\text{neut}A) = 0 \}$.

1.2. Definition of Neutrosophic Measure.

We introduce for the first time the scientific notion of neutrosophic measure.

Let X be a neutrosophic space, and Σ a σ -neutrosophic algebra over X . A *neutrosophic measure* ν is defined by for neutrosophic set $A \in \Sigma$ by

$$\begin{aligned} \nu : X &\rightarrow R^3, \\ \nu(A) &= (m(A), m(\text{neut}A), m(\text{anti}A)), \end{aligned} \quad (1)$$

with $antiA$ = the opposite of A , and $neutA$ = the neutral (indeterminacy) neither A nor $antiA$ (as defined above);

for any $A \subseteq X$ and $A \in \Sigma$,

$m(A)$ means *measure of the determinate part of A* ;

$m(neutA)$ means *measure of indeterminate part of A* ;

and $m(antiA)$ means *measure of the determinate part of $antiA$* ;

where ν is a function that satisfies the following two properties:

a) Null empty set: $\nu(\Phi) = (0, 0, 0)$.

b) Countable additivity (or σ -additivity): For all countable collections $\{A_n\}_{n \in L}$ of disjoint neutrosophic sets in Σ , one has:

$$\nu\left(\bigcup_{n \in L} A_n\right) = \left(\sum_{n \in L} m(A_n), \sum_{n \in L} m(neutA_n), \sum_{n \in L} m(antiA_n) - (n-1)m(X)\right)$$

where X is the whole neutrosophic space, and

$$\sum_{n \in L} m(antiA_n) - (n-1)m(X) = m(X) - \sum_{n \in L} m(A_n) = m\left(\bigcap_{n \in L} antiA_n\right).$$

(2)

1.3. Neutrosophic Measure Space.

A neutrosophic measure space is a triplet (X, Σ, ν) .

1.4. Normalized Neutrosophic Measure.

A neutrosophic measure is called normalized if

$$\nu(X) = (m(X), m(neutX), m(antiX)) = (x_1, x_2, x_3),$$

$$\text{with } x_1 + x_2 + x_3 = 1,$$

$$\text{and } x_1 \geq 0, x_2 \geq 0, x_3 \geq 0. \quad (3)$$

Where, of course, X is the whole neutrosophic measure space.

1.5. Finite Neutrosophic Measure Space.

Let $A \subset X$. We say that $\nu(A) = (a_1, a_2, a_3)$ is finite if all a_1 , a_2 , and a_3 are finite real numbers.

A *neutrosophic measure space* (X, Σ, ν) is called *finite* if $\nu(X) = (a, b, c)$ such that all a , b , and c are finite (rather than infinite).

1.6. σ -Finite Neutrosophic Measure.

A neutrosophic measure is called σ -finite if X can be decomposed into a countable union of neutrosophically measurable sets of finite neutrosophic measure.

Analogously, a set A in X is said to have a *σ -finite neutrosophic measure* if it is a countable union of sets with finite neutrosophic measure.

1.7. Neutrosophic Axiom of Non-Negativity.

We say that the neutrosophic measure ν *satisfies the axiom of non-negativity*, if:

$$\forall A \in \Sigma, \nu(A) = (a_1, a_2, a_3) \geq 0 \text{ if } a_1 \geq 0, a_2 \geq 0, \text{ and } a_3 \geq 0. \quad (4)$$

While a neutrosophic measure ν , that satisfies only the null empty set and countable additivity axioms (hence not the non-negativity axiom), takes on at most one of the $\pm\infty$ values.

1.8. Measurable Neutrosophic Set and Measurable Neutrosophic Space.

The members of Σ are called measurable neutrosophic sets, while (X, Σ) is called a measurable neutrosophic space.

1.9. Neutrosophic Measurable Function.

A function $f : (X, \Sigma_X) \rightarrow (Y, \Sigma_Y)$, mapping two measurable neutrosophic spaces, is called *neutrosophic measurable function* if $\forall B \in \Sigma_Y, f^{-1}(B) \in \Sigma_X$ (the inverse image of a neutrosophic Y -measurable set is a neutrosophic X -measurable set).

1.10. Neutrosophic Probability Measure.

As a particular case of neutrosophic measure ν is the neutrosophic probability measure, i.e. a neutrosophic measure that measures probable/possible propositions

$$-0 \leq \nu(X) \leq 3^+, \quad (5)$$

where X is the whole neutrosophic probability sample space.

We use nonstandard numbers, such I^+ for example, to denominate the absolute measure (measure in all possible worlds), and standard numbers such as I to denominate the relative measure (measure in at least one world). Etc.

We denote the neutrosophic probability measure by \mathcal{NP} for a closer connection with the classical probability \mathcal{P} .

1.11. Neutrosophic Category Theory.

The neutrosophic measurable functions and their neutrosophic measurable spaces form a neutrosophic

category, where the functions are arrows and the spaces objects.

We introduce the neutrosophic category theory, which means the study of the neutrosophic structures and of the neutrosophic mappings that preserve these structures.

The classical category theory was introduced about 1940 by Eilenberg and Mac Lane.

A neutrosophic category is formed by a class of neutrosophic objects X, Y, Z, \dots and a class of neutrosophic morphisms (arrows) ν, ξ, ω, \dots such that:

- a) If $Hom(X, Y)$ represent the neutrosophic morphisms from X to Y , then $Hom(X, Y)$ and $Hom(X', Y')$ are disjoint, except when $X = X'$ and $Y = Y'$;
- b) The composition of the neutrosophic morphisms verify the axioms of
 - i) Associativity: $(\nu \circ \xi) \circ \omega = \nu \circ (\xi \circ \omega)$
 - ii) Identity unit: for each neutrosophic object X there exists a neutrosophic morphism denoted id_X , called *neutrosophic identity* of X such that $id_X \circ \nu = \nu$ and $\xi \circ id_X = \xi$




Fig. 2

1.12. Properties of Neutrosophic Measure.

a) *Monotonicity.*

If A_1 and A_2 are neutrosophically measurable, with $A_1 \subseteq A_2$, where

$$\nu(A_1) = (m(A_1), m(\text{neut}A_1), m(\text{anti}A_1)),$$

and
$$\nu(A_2) = (m(A_2), m(\text{neut}A_2), m(\text{anti}A_2)),$$

then

$$m(A_1) \leq m(A_2), m(\text{neut}A_1) \leq m(\text{neut}A_2), m(\text{anti}A_1) \geq m(\text{anti}A_2). \quad (6)$$

Let $\nu(X) = (x_1, x_2, x_3)$ and $\nu(Y) = (y_1, y_2, y_3)$. We say that $\nu(X) \leq \nu(Y)$, if $x_1 \leq y_1, x_2 \leq y_2$, and $x_3 \geq y_3$.

b) *Additivity.*

$$\text{If } A_1 \cap A_2 = \Phi, \text{ then } \nu(A_1 \cup A_2) = \nu(A_1) + \nu(A_2), \quad (7)$$

where we define

$$(a_1, b_1, c_1) + (a_2, b_2, c_2) = (a_1 + a_2, b_1 + b_2, a_3 + b_3 - m(X)), \quad (8)$$

where X is the whole neutrosophic space, and

$$\begin{aligned}
a_3 + b_3 - m(X) &= m(X) - m(A) - m(B) = m(X) - a_1 - a_2 \\
&= m(\text{anti}A \cap \text{anti}B).
\end{aligned}
\tag{9}$$

1.13. Neutrosophic Measure Continuous from Below or Above.

A neutrosophic measure ν is continuous from below if, for A_1, A_2, \dots neutrosophically measurable sets with $A_n \subseteq A_{n+1}$ for all n , the union of the sets A_n is neutrosophically measurable, and

$$\nu\left(\bigcup_{n=1}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \nu(A_n)
\tag{10}$$

And a neutrosophic measure ν is continuous from above if for A_1, A_2, \dots neutrosophically measurable sets, with $A_n \supseteq A_{n+1}$ for all n , and at least one A_n has finite neutrosophic measure, the intersection of the sets A_n and neutrosophically measurable, and

$$\nu\left(\bigcap_{n=1}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \nu(A_n).
\tag{11}$$

1.14. Generalizations.

1.14.a. Neutrosophic measure is a generalization of the *fuzzy measure*, because when $m(\text{neut}A) = 0$ and $m(\text{anti}A)$ is ignored, we get

$$\nu(A) = (m(A), 0, 0) \equiv m(A),
\tag{12}$$

and the two fuzzy measure axioms are verified:

a) If $A = \Phi$, then $\nu(A) = (0, 0, 0) \equiv 0$

b) If $A \subseteq B$, then $\nu(A) \leq \nu(B)$.

1.14.b. The neutrosophic measure is practically a triple classical measure: a classical measure of the determinate part of a neutrosophic object, a classical part of the indeterminate part of the neutrosophic object, and another classical measure of the determinate part of the opposite neutrosophic object. Of course, if the indeterminate part does not exist (its measure is zero) and the measure of the opposite object is ignored, the neutrosophic measure is reduced to the classical measure.

1.15. Examples.

Let's see some examples of neutrosophic objects and neutrosophic measures.

a) If a book of 100 sheets (covers included) has 3 missing sheets, then

$$\nu(\text{book}) = (97, 3, 0), \quad (13)$$

where ν is the neutrosophic measure of the book number of pages.

b) If a surface of 5×5 square meters has cracks of 0.1×0.2 square meters, then

$$\nu(\text{surface}) = (24.98, 0.02, 0), \quad (14)$$

where ν is the neutrosophic measure of the surface.

c) If a die has two erased faces then

$$v(\text{die}) = (4, 2, 0),$$

where v is the neutrosophic measure of the die's number of correct faces.

d) An approximate number N can be interpreted as a neutrosophic measure $N = \underline{d} + \underline{i}$, where \underline{d} is its determinate part, and \underline{i} its indeterminate part. Its anti part is considered 0 .

For example if we don't know exactly a quantity q , but only that it is between let's say $q \in [0.8, 0.9]$, then $q = 0.8 + i$, where 0.8 is the determinate part of q , and its indeterminate part $i \in [0, 0.1]$.

We get a negative neutrosophic measure if we approximate a quantity measured in an inverse direction on the x -axis to an equivalent positive quantity.

For example, if $r \in [-6, -4]$, then $r = -6 + i$, where -6 is the determinate part of r , and $i \in [0, 2]$ is its indeterminate part. Its anti part is also 0 .

e) Let's measure the truth-value of the proposition

$G =$ "through a point exterior to a line one can draw only one parallel to the given line".

The proposition is incomplete, since it does not specify the type of geometrical space it belongs to. In an *Euclidean geometric space* the proposition G is true; in a *Riemannian geometric space* the proposition G is false (since there is no parallel passing through an

exterior point to a given line); in a *Smarandache geometric space* (constructed from mixed spaces, for example from a part of Euclidean subspace together with another part of Riemannian space) the proposition G is indeterminate (true and false in the same time).

$$\nu(G) = (1,1,1). \quad (15)$$

- f) In general, not well determined objects, notions, ideas, etc. can become subject to the neutrosophic theory.

Chapter 2:
Introduction to Neutrosophic Integral

2.1. Definition of Neutrosophic Integral

Using the neutrosophic measure, we can define a neutrosophic integral.

The neutrosophic integral of a function f is written as:

$$\int_X f d\nu \quad (16)$$

where X is the a neutrosophic measure space, and the integral is taken with respect to the neutrosophic measure ν .

Indeterminacy related to integration can occur in multiple ways: with respect to value of the function to be integrated, or with respect to the lower or upper limit of integration, or with respect to the space and its measure.

2.2. First Example of Neutrosophic Integral: Indeterminacy Related to Function's Values

Let

$$f_N: [a, b] \rightarrow R \quad (17)$$

where the neutrosophic function is defined as:

$$f_N(x) = g(x) + i(x) \quad (18)$$

with $g(x)$ the determinate part of $f_N(x)$, and $i(x)$ the indeterminate part of $f_N(x)$, where for all x in $[a, b]$ one has:

$$i(x) \in [0, h(x)], h(x) \geq 0. \quad (19)$$

Therefore the values of the function $f_N(x)$ are approximate, i.e.

$$f_N(x) \in [g(x), g(x) + h(x)]. \quad (20)$$




Fig. 3

Similarly, the neutrosophic integral is an approximation:

$$\int_a^b f_N(x)dv = \int_a^b g(x)dx + \int_a^b i(x)dx \quad (21)$$

2.3. Second Example of Neutrosophic Integral: Indeterminacy Related to the Lower Limit

Suppose we need to integrate the function

$$f: X \rightarrow R \quad (22)$$

on the interval $[a, b]$ from X , but we are unsure about the lower limit a . Let's suppose that the lower limit "a" has a determinant part " a_1 " and an indeterminate part ε , i.e.

$$a = a_1 + \varepsilon \quad (23)$$

where

$$\varepsilon \in [0, 0.1]. \quad (24)$$




Fig. 4

Therefore

$$\int_a^b f dv = \int_{a_1}^b f(x) dx - i_1 \quad (25)$$

where the indeterminacy i_1 belongs to the interval:

$$i_1 \in [0, \int_{a_1}^{a_1+0.1} f(x) dx]. \quad (26)$$

Or, in a different way:

$$\int_a^b f dv = \int_{a_1+0.1}^b f(x) dx + i_2 \quad (27)$$

where similarly the indeterminacy i_2 belongs to the interval:

$$i_2 \in [0, \int_{a_1}^{a_1+0.1} f(x) dx]. \quad (28)$$

Chapter 3:
Introduction to Neutrosophic Probability

3.1. First Example of Indeterminacy.

The idea of extending the neutrosophic principle, which is based on indeterminacy, to probability, came to my mind when I tossed a die outside, on my stairs made of concrete, but the concrete was broken, had small cracks, and the die got stuck on an edge in a crack. There was no clear face to see, hence it was an indeterminacy.



Fig. 5. Indeterminate die state

Thus tossing a die on a cracked surface one can get:

$\{1, 2, 3, 4, 5, 6, \textit{indeterminacy}\}$.

This is its sample space.

A cubic die (with 12 edges and 8 vertices) tossed on an irregular surface has the chance to fall on a vertex or on an edge in a small slit or crack (not on one of its faces). Therefore, tossing the die can turn on an indeterminate outcome.

Whence, the neutrosophic probability \mathcal{NP}_t of tossing, for example $\{1\}$ is less than $\frac{1}{6}$, since there are seven possible outcomes $\mathcal{NP}_t(1) < \frac{1}{6}$, not like in classical probability where $\mathcal{P}(1) = \frac{1}{6}$.

The more irregularities on the surface (as below), the more indeterminacy occurs:



Fig. 6

In the classical probability the die and the surface it rolls on are considered perfect, hence there is no indeterminacy due to the materials. There is only randomness.

In neutrosophic probability one has, besides randomness, indeterminacy due to construction materials and shapes of the die and of the surface.

If the die is not regular, and the faces have different areas, or the die's center of mass is not in the geometrical die's center, then the probability will be proportional to the face's surface, and the closer is the center of mass to a face the higher the probability for that face.

The die's mass of inhomogeneous density will influence the probability outcome.

3.2. Second Example of Indeterminacy.

Let's consider a regular die (with six faces), having two faces whose print is erased (let's say faces 5 and 6). Then:

$$\mathcal{NP}(1) = \mathcal{NP}(2) = \mathcal{NP}(3) = \mathcal{NP}(4) = \frac{1}{6},$$

$$\mathcal{NP}(5) = \mathcal{NP}(6) = 0,$$

$$\text{while } \mathcal{NP}(\text{indeterm}) = \frac{2}{6}, \quad (29)$$

when the die is tossed on a regular surface.

3.3. Third Example of Indeterminacy.

On a surface with cracks there is a chance that flipping a coin, the coin falls into a crack and gets stuck on its edge; then we have again indeterminacy.

$$\mathcal{NP}(\text{Head}) = \mathcal{NP}(\text{Tale}) < \frac{1}{2} \quad (30)$$

and the sample space is $\{\text{Head}, \text{Tale}, \text{indeterminacy}\}$.



Fig. 7. Indeterminacy related to tossing a coin

3.4. Fourth Example of Indeterminacy.

An urn with two types of votes: A -ballots and B -ballots, but some votes are deteriorated, and we can't determine if it's written A or B . Therefore, we have indeterminate votes.

In many practical applications we may not even know the exact number of indeterminate votes, or of A -ballots, or of B -ballots. Therefore, the indeterminacy is even bigger.

3.5. Fifth Example of Indeterminacy.

If there are two candidates A and B for presidency, and the probability that A wins is 0.46 , it doesn't mean that the probability that B wins is 0.54 , since there may be blank votes (from the voters not choosing any candidate) or black votes (from the voters that reject both candidates).

For example, the probability that B wins could be 0.45 , while the difference $1-0.46-0.45 = 0.09$ would be the probability of blank and black votes together. Therefore we have a neutrosophic probability:
 $\mathcal{NP}(A) = (0.46, 0.09, 0.045)$

3.6. Sixth Example of Indeterminacy.

If a meteorology center reports that the chance of rain tomorrow is 60%, it does not mean that the chance of not raining is 40%, since there might be hidden parameters (weather factors) that the meteorology center is not aware of.

There might be an unclear weather, for example, cloudy and humid day, that some people can interpret as rainy day and others as non-rainy day. The ambiguity arouses indeterminacy.

3.7. The Seventh Example of Indeterminacy.

If some drug tests are 95% reliable, it doesn't mean that 5% they are unreliable, because there might be some unknown effects of the drugs that we are not sure they are beneficial or harmful.

3.8. The Eighth Example of Indeterminacy.

A roulette wheel has 38 numbers. But, having been used too much, several of its numbers have been erased, and one cannot read.

Therefore, we get again indeterminacy.

3.9. Ninth Example of Indeterminacy.

A deck of 52 cards has 3 damaged cards that we are unable to read. Then we have indeterminacy. If the damaged cards are visibly broken, we don't have equiprobability.

3.10. Tenth Example of Indeterminacy.

Probability in a soccer game.

Classical probability is incomplete, because it computes for a team the chance of winning, or the chance of not winning, but not all three chances as in neutrosophic probability: winning, having tie game, or losing.

3.11. Eleventh Natural Example of Indeterminacy.

Indeterminacy occurs (yet rarely) whether a series of newborns will be girls or boys, since some transgender

children can have undetermined (ambiguous) sex, i.e. partially male and partially female).

3.12. Example of a Neutrosophic Continuous Random Variable.

The previous examples used neutrosophic discrete random variables.

Let's now consider a spinner as bellow:




Fig. 8

The continuous sample space is $\Omega = [0, 360]$. Let's say that spinner's table is erased between $270^\circ - 360^\circ$, so if the spinner gets in this area (*IV*th quadrant) we are not able to read a number, we consider it indeterminacy

zone. Therefore $\mathcal{NP}(indeterm) = \frac{1}{4}$. We have a continuous random variable.

$$\begin{aligned} \mathcal{NP}([90,100]) &= \left(ch([90,100]), ch(indeterm), ch(\overline{[90,100]}) \right) \\ &= \left(\frac{10}{360}, \frac{90}{360}, \frac{260}{360} \right) \end{aligned} \tag{31}$$

3.13. First Types of Indeterminacies.

One has at least two types of indeterminacies:

- a) The indeterminacy due to the space (for example the surface on which the dices are tossed on, the urn on which the votes are introduced, etc.).
- b) The indeterminacy due to the items contained into the physical space (for example the defect dice, the unclear ballots, etc.).

3.14. Second Types of Indeterminacies.

- a) We have indeterminacy not related to a particular event, which is a *constant indeterminacy*. For example, tossing a regular die on a irregular surface which has cracks. No matter what outcome we look for 1, 2, ..., or 6,

the indeterminacy (chance that the die falls in a crack and has an unclear reading) is the same.

b) But we may have *indeterminacy related to each event*. For example, let the sample space be:

$\{\textit{sunny day, rainy day, snowfall day}\}$

as a weather forecast for one week from today.

A meteorologist approximately computes the chance of each event, using various parameters, such as: statistics of past weather, today's weather, etc. and gives the following (imprecise) probabilities:

$$\{[0.1, 0.2], [0.5, 0.7], [0.3, 0.6]\}, \quad (32)$$

where $[0.1, 0.2]$ means the probability of sunny day,

$[0.5, 0.7]$ probability of rainy day,

and $[0.3, 0.6]$ probability of snowfall day.

Thus, we have different indeterminacies which are related to the occurrence of each event.

Neutrosophically, we can write it as:

$$NP(\textit{sunny day}) = 0.1 + i_1, \quad \text{where } i_1 \in [0.0, 0.1],$$

$$NP(\textit{rainy day}) = 0.5 + i_2, \quad \text{where } i_2 \in [0.0, 0.2],$$

and

$$NP(\textit{snowfall day}) = 0.3 + i_3,$$

$$\text{where } i_3 \in [0.0, 0.3]$$

with i_1, i_2, i_3 indeterminacies.

Let's compute the union of events

$NP(\text{sunny day } \mathbf{or} \text{ snowfall day})$.

$$\begin{aligned} NP(\text{sunny day}) + NP(\text{snowfall day}) \\ &= (0.1 + i_1) + (0.3 + i_3) \\ &= (0.1 + 0.3) + (i_1 + i_3) \\ &= 0.4 + i_4, \end{aligned}$$

where $i_4 \in [0.0, 0.1] + [0.0, 0.3] = [0.0, 0.4]$.

This could also be computed simply as in classical imprecise probability:

$$\begin{aligned} P(\text{sunny day } \mathbf{or} \text{ snowfall day}) = \\ [0.1, 0.2] + [0.3, 0.6] = [0.4, 0.8] = 0.4 + i_4, \end{aligned}$$

where $i_4 \in [0.0, 0.4]$.

Similarly for intersection of events:

$$\begin{aligned} NP(\text{sunny } \mathbf{and} \text{ snowfall day}) \\ &= (0.1 + i_1) \cdot (0.3 + i_3) \\ &= (0.1)(0.3) \\ &+ \{0.3i_1 + 0.1i_3 + i_1i_3\} \\ &= 0.03 \\ &+ \{[0.0, 0.3] + [0.0, 0.3] \\ &+ [0.0, 0.3]\} = 0.03 + i_5, \end{aligned}$$

where $i_5 \in [0.0, 0.9]$.

This is because:

$$\{\text{Indeterminacy}\} \cdot \{\text{number}\} = \{\text{indeterminacy}\}$$

and

$$\begin{aligned} \{\text{indeterminacy}\} \cdot \{\text{indeterminacy}\} \\ = \{\text{indeterminacy}\}. \end{aligned}$$

Classically:

$$P(\text{sunny day and snowfall day}) = [0.1, 0.2] \cdot [0.3, 0.6] = [0.03, 0.12] = 0.03 + i_5, \\ \text{where } i_5 \in [0.0, 0.09].$$

Similarly for negation of events:

$$NP(\text{not a sunny day}) = 1 - (0.1 + i_1) \\ = 1 - 0.1 - i_1 = 0.9 - i_1 = \\ = 0.8 + i_6, \text{ where } i_6 \in [0.0, 0.1].$$

Classically:

$$P(\text{not a sunny day}) = 1 - [0.1, 0.2] \\ = [0.8, 0.9] = 0.8 - i_6,$$

where $i_6 \in [0.0, 0.1]$.

- c) Or *mixt indeterminacies*: to some events there is a chance of indeterminacy > 0 , while to other events there is not.

A similar example as the previous, but we change the data:

$$\{[0.1, 0.2], [0.5, 0.7], 0.3\}. \quad (33)$$

Therefore, there is indeterminacy related to the first and second events, but not to the third.

3.15. Distinction between Indeterminacy and Randomness.

Indeterminacy is different from randomness. Indeterminacy is due to the defects of the construction of the physical space (where an event can occur), and/or to the imperfect construction of the physical objects involved in the event, etc.

Therefore, neutrosophic probability analyses both: the random phenomena, and the indeterminacy related to these phenomena.

In consequence, neutrosophic probability deals with two types of variables: random variables and indeterminacy variables, and two types of processes: stochastic process and respectively indeterminate process.

3.16. Neutrosophic Random Variables.

A *classical random (stochastic) variable* is subject to change due to randomness, while the *neutrosophic random (stochastic) variable* is subject to change due to both randomness and indeterminacy.

A neutrosophic random variable's values represent the possible outcomes and possible indeterminacies. The randomness and indeterminacy can be objective or subjective.

Alike classical random variables, the neutrosophic random variables can be classified as:

- *discrete*, that is it can take a value in a specified list of exact values and a finite number of indeterminacies;
- *continuous*, that is it can take a value or an indeterminacy in an interval, or in a collection of intervals;

- *mixt*, that is it can take a value or indeterminacy either in a specified list of exact values, or in an interval or in a collection of intervals (mixture of discrete and continuous).

Another classification, alike classical random variables, for neutrosophic random variables is

- *finite*; having of course a finite number of possible outcomes and possible indeterminacies;
- *infinite*; having an infinite number of possible outcomes or indeterminacies.

An infinite neutrosophic random variable can be

- *countably*;
- or *uncountably*.

A *neutrosophic random variable* X is *admissible* if it is possible to compute the chance that the value of X is less than any particular number, together with its corresponding indeterminacy and its nonchance. Which is equivalent to the possibility of computing the chance that the value of X is in any range, range that must be mapped to a subset of the neutrosophic sample space $\nu\Omega$.

3.17. Many Possible Neutrosophic Measures and Probabilities.

We may be able to define the neutrosophic measure and neutrosophic probability in many ways, since we work with approximations and indeterminacies. Their definitions may depend on each particular application.

3.18. Definition of Neutrosophic Probability

Neutrosophic probability (or likelihood) is a particular case of the neutrosophic measure. It is an estimation of an event (different from indeterminacy) to occur, together with an estimation that some indeterminacy may occur, and the estimation that the event does not occur.

Neutrosophic Probability and Neutrosophic Statistics started in *1995*, but was not developed and applied as much as neutrosophic logic and neutrosophic set that are widely used.

A **neutrosophic random variable** is a variable that may have an indeterminate (unclear, ambiguous) outcome.

A **neutrosophic random (stochastic) process** represents the evolution over time of some neutrosophic random values. It is a collection of neutrosophic random variables.

The classical probability deals with fair dice, coins, roulettes, spinners, decks of cards, random walks, while neutrosophic probability deals with unfair, imperfect such objects, variables and processes.

The neutrosophic probability is a generalization of the classical probability because, when the chance of indeterminacy of a stochastic process is zero, these two probabilities coincide.

3.19. Neutrosophic Probability vs. Imprecise Probability.

In Imprecise Probability (IP), the probability of an event A ,

$$IP(A) = (a, b) \subseteq [0, 1] \quad (34)$$

is an interval included into $[0, 1]$, not a crisp number.

The Neutrosophic Probability that an event A occurs is

$$\mathcal{NP}(A) = (ch(A), ch(neutA), ch(antiA)) = (T, I, F), \quad (35)$$

but sometimes instead of “neutA” we say “indeterminacy related to A” and we denote it by “indeterm_A”; also we note “antiA” by \bar{A} ;

where T, I, F are standard or nonstandard subsets of the nonstandard unitary interval $]0, 1^+[$, and T is the chance that A occurs, denoted $ch(A)$; I is the indeterminate chance related to A, $ch(indeterm_A)$; and F is the chance that A does not occur, $ch(\bar{A})$.

So, \mathcal{NP} is a generalization of the Imprecise Probability as well.

Therefore, using other notations we have:

$$\mathcal{NP}(A) = (ch(A), ch(indeterm_A), ch(\bar{A})). \quad (36)$$

We used the notations T (truth), I (indeterminate), and F (falsehood) in order to be consistent with those from neutrosophic logic and neutrosophic set, widely spread.

In the most general case, T, I, F are standard or non-standard subsets of the unitary non-standard interval $]^{-0, 1^+}$, in order to be able to make distinction between *absolute sure event* (sure event in all possible worlds -- whose probability value is 1^+), and *relative sure event* (i.e. sure event in at least one world, but not in all words -- whose probability is I , where $I < 1^+$).

Similarly, for *absolute impossible event* (impossible event in all possible worlds -- whose probability is $^{-0}$), and *relative impossible event* (i.e. impossible event in at least one world, but not in all words -- whose probability is $^{-0}$, where $^{-0} < 0$).

$$1^+ = 1 + \varepsilon \text{ and } ^{-0} = 0 - \varepsilon, \quad (37)$$

where ε is a very tiny positive number.

For technical applications we'll use only standard sets and the standard unit interval $[0, 1]$. And throughout this book, with few exceptions.

Let's note by majuscules the subsets T, I, F and by lower-case letters the crisp numbers t, i, f . For the crisp neutrosophic probability, when T, I, F are just standard or non-standard numbers in $]^{-0, 1^+}$, in the most general case one has:

$$^{-0} \leq t + i + f \leq 3^+, \quad (38)$$

considering that the tree components t, i, f are independent (as in neutrosophic logic and in neutrosophic set).

If only two components are dependent, while the third one is independent from them, then

$$-0 \leq t+i+f \leq 2^+ . \quad (39)$$

If all three components are dependent two by two, then

$$-0 \leq t+i+f \leq 1^+ . \quad (40)$$

Let's consider the standard case.

- 1) If $t+i+f=1$ one has *complete probability* (the most common application), or *normalized probability*.
- 2) If $t+i+f < 1$ one has *incomplete probability* (because the source of information or the stochastic process is incomplete, i.e. not well known).
- 3) If $t+i+f > 1$ one has *paraconsistent probability* (because of conflicting sources of information that transmit us contradictory information; for example one source may compute the chance that an event occurs using some criteria (parameters influencing the event), but it is not able to compute the chance that the event does not occur, while another independent source of information may compute the chance that the event does not occur using different criteria (different parameters), but not able to compute the chance that the event occur).

Similarly, for computing the chance of indeterminacy of the stochastic process by a third independent source of information. Therefore it is possible to get the sum $t+i+f \neq 1$. (41)

3.20. Sigma-Algebra of Events.

A sigma-algebra or σ -algebra of X , in the measure theory, is a collection of subsets of the set X such that

- 1) $\Phi \in \Sigma$;
- 2) $X \in \Sigma$;
- 3) If $A \in \Sigma$ then the complement of A , $C(A) \in \Sigma$;
- 4) If $A_1, A_2, \dots, A_n \in \Sigma$, then the countable union $A_1 \cup A_2 \cup \dots \cup A_n \in \Sigma$.

3.21. Definition of Classical Probability.

The classical probability measure is a mapping:

$$\mathcal{P} : X \rightarrow [0,1] \quad (42)$$

where X is a sample space, such that $\mathcal{P}(X)=1$ and \mathcal{P} is additive for the union

$$\mathcal{P}(A \cup B) = \mathcal{P}(A) + \mathcal{P}(B) \text{ for } A \cap B = \phi, \quad (43)$$

even for infinite unions:

$$\mathcal{P}\left(\bigcup_{n \geq 0} A_n\right) = \sum_{n \geq 0} \mathcal{P}(A_n) \quad (44)$$

for A_i disjoint two by two, that lie in the sigma-algebra of events Σ of X .

3.22. Neutrosophic Sigma-Algebra of Events.

The neutrosophic sigma-algebra of events $\nu\Sigma$ will be defined in the same way, with the distinction that the set X contains some indeterminacy. Therefore there are some subjects of X that are indeterminate parts.

3.23. Neutrosophic Probability Measure.

The neutrosophic probability measure is a mapping:

$$\mathcal{NP} : X \rightarrow [0,1]^3 \quad (45)$$

where X is a neutrosophic sample space (i.e. X contains some indeterminacy),

$$\mathcal{NP}(A) = \left(ch(A), ch(indeterm_A), ch(\bar{A}) \right), \quad (46)$$

or, using other notations, we have:

$$\mathcal{NP}(A) = \left(ch(A), ch(neutA), ch(antiA) \right) \quad (47)$$

where $indeterm_A$ means the indeterminacy that may occur when trying to have event A occurs, such that the neutrosophic probability of the whole space X has the property that:

$$\mathcal{NP}(X) = (\alpha, \beta, \gamma),$$

where $0 \leq \alpha, \beta, \gamma \leq 1$, and

$$0 \leq \alpha + \beta + \gamma \leq 3. \quad (48)$$

Therefore, the sum of the three components of the neutrosophic probability of the whole sample space is not required to be equal to 1 as in classical probability, since there cases where it is strictly less than 1 , or strictly greater than 1 .

We also have:

$$\mathcal{NP}(A \cup B) = \left(ch(A) + ch(B), ch(indeterm_{A \cup B}), ch(\overline{A \cup B}) \right) \quad (49)$$

for $A \cap B = \phi$, and for infinite unions:

$$\mathcal{NP}\left(\bigcup_{n \geq 0} A_n\right) = \left(\sum_{n \geq 0} ch(A_n), ch(indeterm_{\bigcup_{n \geq 0} A_n}), ch\left(\overline{\bigcup_{n \geq 0} A_n}\right) \right) \quad (50)$$

for A_n disjoint two by two that lie in the neutrosophic sigma algebra of events.

Remark. Although in most cases the sum of the three components is 1 (in normalized probability):

$$ch(A) + ch(neutA) + ch(antiA) = 1 \quad (51)$$

or using similar notations

$$ch(A) + ch(indeterm_A) + ch(\overline{A}) = 1, \quad (52)$$

we still recommend to computing all three components because it arises cases when the probability is not normalized.

3.24. Neutrosophic Probability Mass Function.

A *Neutrosophic Probability Mass Function*

(*npmf*) is a function

$$f: \nu\Omega \rightarrow [0, 1]^3$$

$$f(x) \in [0, 1]^3 \text{ for all } x \in \nu\Omega,$$

$$f(x) = (ch(x), ch(indeterm_x), ch(\bar{x})). \quad (53)$$

A *Neutrosophic Event* is any subset E of the neutrosophic sample space $\nu\Omega$. The neutrosophic probability of any event E is defined as:

$$NP(E) = (\sum_{x \in E} ch(x), ch(indeterm_E), \sum_{y \in \bar{E}} ch(y)). \quad (54)$$

3.25. Neutrosophic Probability Axioms.

They are extensions of Kolmogorov axioms from classical probability.

$(\nu\Omega, NF, NP)$ is a neutrosophic probability space, where $\nu\Omega$ is a neutrosophic sample space, NF is a neutrosophic event space, and NP is a neutrosophic probability measure.

First Axiom.

The neutrosophic probability of an event A

$$NP(A) = (ch(A), ch(indeterm_A), ch(\bar{A})), \quad (55)$$

where $ch(A) \geq 0$,

$$ch(indeterm_A) \geq 0,$$

$$ch(\bar{A}) \geq 0, \text{ for any } A \in NF;$$

with the notations that " $indeterm_A$ " means indeterminacy related to event A , and \bar{A} is the opposite event of A (the *antiA* event).

Second Axiom.

The neutrosophic probability of the sample space is between -0 and 3^+ .

$$NP(v\Omega) = (\sum_{x \in v\Omega} ch(x), ch(indeterm_{v\Omega}), ch(anti v\Omega)), \quad (56)$$

where

$$-0 \leq \sum_{x \in v\Omega} ch(x) + ch(indeterm_{v\Omega}) + ch(anti v\Omega) \leq 3^+, \quad (57)$$

with the notation $indeterm_{v\Omega}$ means total indeterminacy that may occur in the neutrosophic sample space.

For the classical complete (normalized) sample space,

$ch(anti v\Omega) = 0$, but for incomplete sample space

$$ch(anti v\Omega) > 0. \quad (58)$$

Third Axiom.

This axiom is concerned with *neutrosophic σ -additivity*:

$$NP(A_1 \cup A_2 \cup \dots) =$$

$$\left(\sum_{j=1}^{\infty} ch(A_j), ch(indeterm_{A_1 \cup A_2 \cup \dots}), ch(\overline{A_1 \cup A_2 \cup \dots}) \right), \quad (59)$$

where A_1, A_2, \dots is a countable sequence of disjoint (or mutually exclusive) neutrosophic events.

If we relax the third axiom we get a *neutrosophic quasiprobability distribution*.

3.26. Consequences of Neutrosophic Probability Axioms.

a) *Monotonicity.*

If A and B are two neutrosophic events, with $A \subseteq B$, with

$$NP(A) = (ch(A), ch(indeterm_A), ch(\bar{A}))$$

$$NP(B) = (ch(B), ch(indeterm_B), ch(\bar{B})),$$

then $ch(A) \leq ch(B), \quad (60)$

$$ch(indeterm_A) \leq ch(indeterm_B), \quad (61)$$

$$ch(\bar{A}) \geq ch(\bar{B}). \quad (62)$$

b) *Neutrosophic Probability of the Empty Set.*

$$NP(\emptyset) = (0, 0, 0). \quad (63)$$

c) *Bounding the Neutrosophic Probability.*

$$NP(A) = (ch(A), ch(indeterm_A), ch(\bar{A}))$$

$$\text{where } 0 \leq ch(A) \leq 1, \quad (64)$$

$$0 \leq ch(indeterm_A) \leq 1, \quad (65)$$

$$0 \leq ch(\bar{A}) \leq 1. \quad (66)$$

d) *Neutrosophic Addition Law (or Neutrosophic Sum Rule):*

For any two neutrosophic events A and B we have:

$$\begin{aligned} NP(A \cup B) = \\ (ch(A) + ch(B) - ch(A \cap B), ch(indeterm_{A \cup B}), ch(\overline{A \cup B})). \end{aligned} \quad (67)$$

If $A \cap B = \emptyset$, then

$$NP(A \cup B) = (ch(A) + ch(B), ch(indeterm_{A \cup B}), ch(\overline{A \cup B})). \quad (68)$$

e) *Neutrosophic Inclusion-Exclusion Principle.*

$$NP(v\Omega \setminus A) = (ch(v\Omega) - ch(A), ch(indeterm_{v\Omega \setminus A}), ch(A)). \quad (69)$$

Also, if $A \subseteq B$, then:

$$NP(B \setminus A) = (ch(B) - ch(A), ch(indeterm_{B \setminus A}), ch(\overline{B \setminus A})).$$

(70)

3.27. Interpretations of the Neutrosophic Probability.

Neutrosophic Probability can also have two interpretations, as the classical probability:

- a) *Objective form*, or describing objective state of affairs, whose most popular version is the neutrosophic frequentist probability; and
- b) *Subjective form*, or a degree of belief in an event to occur.

3.28. Neutrosophic Notions.

If an experiment produces indeterminacy, that is called a *neutrosophic experiment*. Collecting all results, including the indeterminacy, we get the *neutrosophic sample space* (or the *neutrosophic probability space*) of the experiment.

The *neutrosophic power set* of the neutrosophic sample space is formed by all different collections (that may or may not include the indeterminacy) of possible results. These collections are called *neutrosophic events*.

3.29. Example with Neutrosophic Frequentist Probability.

Let's consider a more concrete example.

Using the Frequentist Neutrosophic Probability we can (approximately of course) determine what is the chance that the die tosses as indeterminate. Similarly as in classical probability, we can use a computer simulation, based upon connections between *neutrosophic mathematical model* (i.e. models involving indeterminacy) and our everyday life. Neutrosophic statisticians can use simulations to approximate the probability of die uncertainty tossed on a specific irregular surface. With computers a large number of trials can be simulated in short time.

Suppose we obtain that the chance of getting indeterminacy $ch(indeterm)=0.10$ for tossing a regular die on an irregular surface. The neutrosophic sample space is then:

$$\nu\Omega = \{1, 2, 3, 4, 5, 6, indeterm\}. \quad (71)$$

Then, the neutrosophic probability of tossing event A is

$$\mathcal{NP}(A) = \left(ch(A), ch(indeterm_A), ch(\bar{A}) \right) \quad (72)$$

where $ch(\cdot)$ mans “chance”, and \bar{A} is the opposite event of A (chance that *anti* A occurs).

For example:

$$\begin{aligned} \mathcal{NP}(1) &= \left(ch(\{1\}), ch(indeterm_{\{1\}}), ch(\{\bar{1}\}) \right) \\ &= \left(\frac{1-0.10}{6}, 0.10, 5 \cdot \frac{1-0.10}{6} \right) = (0.15, 0.10, 0.75) \\ &= \mathcal{NP}(2) = \dots = \mathcal{NP}(6). \end{aligned} \quad (73)$$

In general:

$$\mathcal{NP}(\bar{A}) = \left(ch(\bar{A}), ch\left(indeterm_{(\bar{A})} \right), ch(A) \right). \quad (74)$$

Hence

$$\begin{aligned} \mathcal{NP}(\bar{1}) &= \left(ch(\bar{1}), ch\left(indeterm_{(\bar{1})} \right), ch(1) \right) \\ &= \left(ch(\{2,3,4,5,6\}), ch\left(indeterm_{\{2,3,4,5,6\}} \right), ch(1) \right) \\ &= (5(0.15), 0.10, 0.15) = 0.75, 0.10, 0.15. \end{aligned} \quad (75)$$

Also, for

$$\begin{aligned} \mathcal{NP}(1 \text{ or } 2) &= \left(ch(1 \text{ or } 2), ch\left(indeterm_{(1 \text{ or } 2)} \right), ch(\overline{1 \text{ or } 2}) \right) \\ &= \left(ch(1) + ch(2), ch\left(indeterm_{(1 \text{ or } 2)} \right), ch(\overline{1 \text{ and } 2}) \right) \\ &= (0.15 + 0.15, 0.10, ch(\{3,4,5,6\})) = (0.30, 0.10, 4 \cdot (0.15)) \\ &= (0.30, 0.10, 0.60). \end{aligned} \quad (76)$$

In general:

$$\mathcal{NP}(A \text{ or } B) = \left(ch(A) + ch(B), ch\left(indeterm_{(A \text{ or } B)} \right), ch(\overline{A \text{ or } B}) \right) \quad (77)$$

for $A \cap B = \phi$.

For *neutrosophic non-exclusive events* in general one has:

$$\begin{aligned}
\mathcal{NP}(A \text{ or } B) &= \left(ch(A \text{ or } B), ch\left(\underset{A \text{ or } B}{\text{indeterm}}\right), ch\left(\overline{A \text{ or } B}\right) \right) \\
&= \left(ch(A) + ch(B) - ch(A \cap B), ch(\text{indeterm}), ch(\overline{A \text{ and } B}) \right).
\end{aligned}
\tag{78}$$

Whence, if $A = \{1, 2, 3\}$, $B = \{2, 3, 4, 5\}$, then:

$$\begin{aligned}
\mathcal{NP}(\{1, 2, 3\} \text{ or } \{2, 3, 4, 5\}) &= \\
&= \left(3(0.15) + 4(0.15) - 2(0.15), 0.10, ch(\{4, 5, 6\} \text{ and } \{1, 6\}) \right) \\
&= \left(0.75, 0.10, ch(\{6\}) \right) = (0.75, 0.10, 0.15).
\end{aligned}$$

In general, for independent events, one has:

$$\begin{aligned}
\mathcal{NP}(A \text{ and } B) &= \left(ch(A \text{ and } B), ch\left(\underset{A \text{ and } B}{\text{indeterm}}\right), ch\left(\overline{A \text{ and } B}\right) \right) \\
&= \left(ch(A) \cdot ch(B), ch\left(\underset{A \text{ and } B}{\text{indeterm}}\right), ch\left(\overline{A \text{ and } B}\right) \right).
\end{aligned}
\tag{79}$$

3.30. Example with Neutrosophic Frequentist Probability on a Neutrosophic Product Space.

Let suppose we toss the previous regular die on an irregular surface twice. Therefore we have two independent events. What is the neutrosophic probability

of getting {3} on the first tossing and {4} on the second tossing?

The first neutrosophic space with corresponding chances:

$$\nu\Omega_1 = \left\{ \begin{array}{ccccccc} 1, & 2, & 3, & 4, & 5, & 6, & \text{indeterm} \\ \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} \\ 0.15 & 0.15 & 0.15 & 0.15 & 0.15 & 0.15 & 0.10 \end{array} \right\}, \quad (80)$$

and the second neutrosophic space with corresponding chances:

$$\nu\Omega_2 = \left\{ \begin{array}{ccccccc} 1, & 2, & 3, & 4, & 5, & 6, & \text{indeterm} \\ \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} & \underbrace{\quad} \\ 0.15 & 0.15 & 0.15 & 0.15 & 0.15 & 0.15 & 0.10 \end{array} \right\} \quad (81)$$

Whence we construct their *neutrosophic product space*:

$$\left\{ \begin{array}{ccc} (1,1), (1,2), \dots, (1,6) & (1,I), (2,I), \dots, (6,I) & (I,I) \\ (1,1), (1,2), \dots, (1,6) & (I,1), (I,2), \dots, (I,6) & \\ \dots & & \\ (6,1), (6,2), \dots, (6,6) & & \end{array} \right\} \quad (82)$$

where I= *indeterminacy* ,

with corresponding chances:

$$\left\{ \begin{array}{ccc} 0.0225, 0.0225, \dots, 0.0225 & 0.0150, 0.0150, \dots, 0.0150 & 0.0100 \\ 0.0225, 0.0225, \dots, 0.0225 & 0.0150, 0.0150, \dots, 0.0150 & \\ \dots & & \\ 0.0225, 0.0225, \dots, 0.0225 & & \end{array} \right\} \quad (83)$$

Hence,

$$ch(\{3\} \text{ and } \{4\}) = 0.15(0.15) = 0.00225;$$

$$ch\left(\begin{array}{c} \text{indeterm} \\ \{3\} \text{ or } \{4\} \end{array}\right) = 12(0.0150) + 0.0100 = 0.1800 + 0.0100 \\ = 0.1900;$$

$$ch(\overline{\{3\}} \text{ and } \overline{\{4\}}) = \\ = \left(ch(\overline{\{3\}} \wedge \nu\Omega_2), \nu\Omega_1 \wedge \overline{\{4\}}, \{3\} \wedge \{1,2,3,5,6\}, \{1,2,4,5,6\} \wedge \{4\} \right) \\ = 35(0.0225) = 0.7875.$$

$$\mathcal{NP}(\{3\} \text{ and } \{4\}) = (0.0225, 0.1900, 0.7875). \quad (84)$$

We have considered that (1,I),..., (6,I), (I,1),..., (I,6) are indeterminacies, while (I,I) obviously is a double indeterminacy.

3.31. Example with Double Indeterminacy.

We change again the theoretical equipment.

Instead of a fair die, we consider now a defect die in the sense that two of its faces have the print erased, for example the erased faces are {5} and {6}.

The new neutrosophic probability space is:

$$\nu\Omega = \{1,2,3,4, \text{indeterm}_d, \text{indeterm}_s\} \quad (85)$$

with two types of indeterminacies: one due to the physical die, denoted by indeterm_d , and the second due to the physical space, denoted by indeterm_s .

We consider that chance of $indeterm_s$ is the same as in the previous frequentist examples: $ch(indeterm_s) = 0.10$, and $ch(1) = \dots = ch(4) = 0.15$ as before.

But from two erased prints we get $ch(indeterm_d) = 2(0.15) = 0.30$.

Thus

$$\begin{aligned} ch\left(\text{total } indeterm\right) &= ch(indeterm_s) + ch(indeterm_d) \\ &= 0.10 + 0.30 = 0.40, \end{aligned}$$

whence

$$\mathcal{NP}(1) = \dots = \mathcal{NP}(4) = (0.15, 0.40, 0.45). \quad (86)$$

This neutrosophic experiment is equivalent to experiment of having a perfect die with four faces (a tetrahedron), which is tossed on an irregular surface where the chance of indeterminacy (for the die to get stuck on one of its six edges or on one of its four vertices) is 0.40.

Therefore

$$\nu\Omega = \{1, 2, 3, 4, indeterm\}. \quad (87)$$

3.32. Neutrosophic Example with Tossing a Coin Multiple Times.

Let's consider a regular coin [with two faces: H (head) and T (tale)] flipped on an irregular surface. By

neutrosophic frequentist probability let's suppose the chance that the coin gets stuck on its edge into a surface crack is:

$$ch(\text{indeterm}) = 0.02 \quad (88)$$

Because the coin is fair, the chances of head or tale are equal:

$$ch(H) = ch(T) = \frac{1-0.02}{2} = 0.49. \quad (89)$$

The neutrosophic probability space is:

$$\nu\Omega = \{H, T, I\}, \quad (90)$$

where “ T ” stands for indeterm(inacy).

Therefore:

$$NP(H) = NP(T) = (0.49, 0.02, 0.49). \quad (91)$$

We flip the coin three times. What is the (neutrosophic) probability of getting HTT ?

The neutrosophic product space is:

$$\left\{ \begin{array}{ccc} H & T & I \\ 0.49 & 0.49 & 0.02 \end{array} \right\} \times$$

$$\left\{ \begin{array}{ccc} H & T & I \\ 0.49 & 0.49 & 0.02 \end{array} \right\} \times$$

$$\left\{ \begin{array}{ccc} H & T & I \\ 0.49 & 0.49 & 0.02 \end{array} \right\}$$

$$= \{ HHH, HHT, HTH, HTT, THH, THT, TTH, TTT;$$

and *indeterminacy of first order*:

$$IHH, IHT, ITH, ITT; HIH, HIT, TIH, TIT; HHI, HTI, THI, TTI;$$

also *indeterminacy of second order*:

$$IIH, IIT; IHI, ITI; HII, TII;$$

and *indeterminacy of third order*:

$$III \}, \tag{92}$$

which has $3^3 = 27$ elements.

Computing the chances:

$$ch(HHH) = ch(HHT) = \dots = ch(TTT) = (0.49)^3 = 0.117649;$$

$$ch(IHH) = ch(IHT) = \dots = ch(TTI) = (0.49)^2(0.02) = 0.004802,$$

for each first order indeterminacy;

$$ch(IIH) = ch(IIT) = \dots = ch(TII) = 0.49(0.02)^2 = 0.000196,$$

for each second order indeterminacy;

$$ch(III) = (0.02)^3 = 0.000008.$$

Therefore indeterminacy propagates.

The sum of total indeterminate chances is:

$$ch(\text{total indeterm}) = 12(0.004802) + 6(0.000196) + 1(0.000008) = 0.058808.$$

The chance that HTT occurs is

$$ch(HTT) = (0.49)^3 = 0.117649,$$

while the chance that HTT does not occur is:

$$ch(\overline{HTT}) = 7(0.117649) = 0.823543.$$

Finally,

$$NP(HTT) = (0.117649, 0.058808, 0.823543).$$

In the classical probability, where $ch(\text{indeterm}) = 0$,

we get $P(HTT) = 0.5^3 = 0.125$,

and, transcribed into neutrosophic form, we get:

$$NP(HTT) = (0.5^3, 0, 7(0.5)^3) = (0.125, 0, 0.875). \quad (93)$$

The chance of flipping three times in a row and getting HTT is smaller in the neutrosophic probability space than in the classical probability space, because of the strictly positive chance of having indeterminacy:

$$0.117649 < 0.125000. \quad (94)$$

3.33. Example with Sum of Chances of an Event.

In *classical probability*: If A is an event, then $P(A)$ is the sum of probabilities of all outcomes in the set A .

In *neutrosophic probability*, it is similar if

$$A = \{a_1, a_2, \dots, a_n\}. \quad (95)$$

$NP(A) = (\text{sum of chances of all outcomes in the set } A, ch(indeterm_A), ch(\bar{A})) =$

$$(\sum_{j=1}^n ch(a_j), ch(indeterm_A), ch(\bar{A})). \quad (96)$$

For example, if we retake one of the previous experiments of a regular die tossed on an irregular surface, where the chance of indeterminacy is 0.10 , then

$$\begin{aligned} & NP(\{1, 2, 3\}) \\ &= (ch\{1, 2, 3\}, ch(indeterm_{\{1,2,3\}}), ch(\overline{\{1, 2, 3\}})) \\ &= (ch(1) + ch(2) + ch(3), ch(indeterm_{\{1,2,3\}}), ch(\{4, 5, 6\})) \\ &= (0.15 + 0.15 + 0.15, 0.10, ch(4) + ch(5) + ch(6)) \\ &= (0.45, 0.10, 0.45), \end{aligned} \quad (97)$$

since $NP(1) = NP(2) = NP(3) = (0.15, 0.10, 0.75)$.

3.34. Paraconsistent Neutrosophic Probability.

The paraconsistent neutrosophic probability has the property that the sum of its components is strictly greater than I :

$$t + i + f \in (1, 3^+], \quad (98)$$

therefore one has contradictions between chances.

Forecasting an event from different criteria, we may obtain different chances of occurrence.

For example, suppose two handball teams G and H will compete in a game next week.

- a) According to the *history* of their previous disputes, team G is 60% favorable to win.
- b) But, according to their *last games in the actual season vs. other handball teams*, H is showing a better performance than G , and the experts conclude upon this criterion that H has 70% chance to win.
- c) Others believe that since G was often better than H , but in this season H contrarily played better than G , as a compensation it is 10% chance that their game be undecided (tie).

Therefore, $NP(G \text{ wins over } H) = (0.6, 0.1, 0.7)$,

with $0.6 + 0.1 + 0.7 > 1$. (99)

3.35. Incomplete Neutrosophic Probability.

The incomplete neutrosophic probability has the property that the sum of its components is strictly less than 1:

$$t + i + f \in]^-, 0, 1), \quad (100)$$

therefore one has incomplete (missing) information.

Lets' reconsider the previous example about two handball teams H and G that will compete in a game next week.

- a) If both teams have a weak performance in the present season and of almost equal values, then each one will have a slim chance to win on 20%.
- b) Studying the low number of their previous games when the results were tie, the handball experts conclude that it is a slim chance of 30% of having a tie game.

Therefore, $NP(G \text{ wins over } H) = (0.2, 0.3, 0.2)$,

with $0.2 + 0.3 + 0.2 < 1$. (101)

3.36. Neutrosophic Mutually Exclusive Events.

In *classical probability*, if A and B are *mutually exclusive (independent) events*, then

$$P(A \text{ or } B) = P(A) + P(B). \quad (102)$$

In *neutrosophic probability*, we have similar property for mutually exclusive events:

$$NP(A \text{ or } B) = (ch(A) + ch(B), ch(indeterm_{A \text{ or } B}), ch(\overline{A \text{ or } B})). \quad (103)$$

In *classical probability* for *non mutually exclusive events* A and B one has:

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B). \quad (104)$$

In *neutrosophic probability* for *non mutually exclusive neutrosophic events* one similarly has:

$$NP(A \text{ or } B) = (ch(A) + ch(B) - ch(A \text{ and } B), ch(indeterm_{A \text{ or } B}), ch(\overline{A \text{ or } B})). \quad (105)$$

For example, let's consider a deck of 52 cards, but such that 2 of them are deteriorated and one cannot read them. Let's draw at random a single card. What is the

neutrosophic probability of getting a face card (event A) or a heart card (event B)? We know that none of the face and heart cards were deteriorated. There are 12 face cards (four types of each of J , Q , and K), 13 heart cards, and 3 cards that are both face and heart.

$$\begin{aligned}
 NP(A \text{ or } B) &= \\
 (ch(A \text{ or } B), ch(indeterm_{A \text{ or } B}), ch(\overline{A \text{ or } B})) &= \\
 (ch(A) + ch(B) - \\
 ch(A \text{ and } B), ch(indeterm_{A \text{ or } B}), ch(\overline{A \text{ and } B})) &= \\
 \left(\frac{12}{52} + \frac{13}{52} - \frac{3}{52}, \frac{2}{52}, \frac{52-12-13+3-2}{52}\right) &= \left(\frac{22}{52}, \frac{2}{52}, \frac{28}{52}\right). \quad (106)
 \end{aligned}$$

Of course,

$$\begin{aligned}
 NP(A) &= \left(\frac{12}{52}, \frac{2}{52}, \frac{38}{52}\right), \\
 NP(B) &= \left(\frac{13}{52}, \frac{2}{52}, \frac{37}{52}\right), \\
 NP(A \text{ and } B) &= \left(\frac{3}{52}, \frac{2}{52}, \frac{47}{52}\right). \quad (107)
 \end{aligned}$$

We do not simplify the fractions because we can better compare these neutrosophic probabilities if we leave the same denominators for them all.

But let's say we don't know if any of the two erased cards are among the face or heart cards. Then:

$$NP(A) = \left(\left[\frac{10}{52}, \frac{12}{52}\right], \frac{2}{52}, \left[\frac{38}{52}, \frac{40}{52}\right]\right), \quad (108)$$

$$\text{and } NP(B) = \left(\left[\frac{11}{52}, \frac{13}{52} \right], \frac{2}{52}, \left[\frac{37}{52}, \frac{39}{52} \right] \right), \quad (109)$$

$$\text{and } NP(A \text{ and } B) = \left(\left[\frac{1}{52}, \frac{3}{52} \right], \frac{2}{52}, \left[\frac{47}{52}, \frac{49}{52} \right] \right), \quad (110)$$

$$\text{whence } NP(A \text{ or } B) = \left(\left[\frac{18}{52}, \frac{24}{52} \right], \frac{2}{52}, \left[\frac{26}{52}, \frac{32}{52} \right] \right), \quad (111)$$

because $ch(A \text{ or } B) =$

$$\begin{aligned} &= \left[\frac{10}{52}, \frac{12}{52} \right] + \left[\frac{11}{52}, \frac{13}{52} \right] - \left[\frac{1}{52}, \frac{3}{52} \right] \\ &= \left[\frac{21}{52}, \frac{25}{52} \right] - \left[\frac{1}{52}, \frac{3}{52} \right] \\ &= \left[\frac{21-3}{52}, \frac{25-1}{52} \right] = \left[\frac{18}{52}, \frac{24}{52} \right], \end{aligned}$$

and $ch(\overline{A \text{ or } B}) =$

$$\begin{aligned} &= ch(\text{whole neutrosophic probability space}) \\ &\quad - ch(\text{indeterm}) - ch(A \text{ or } B) \\ &= 1 - \frac{2}{52} - \left[\frac{16}{52}, \frac{24}{52} \right] = \frac{50}{52} - \left[\frac{18}{52}, \frac{24}{52} \right] = \left[\frac{26}{52}, \frac{32}{52} \right]. \end{aligned}$$

3.37. Neutrosophic Experimental Probability.

In *classical experimental probability* is

$$\frac{\text{number of times event } A \text{ occurs}}{\text{total number of trials}}. \quad (112)$$

Similarly, *Neutrosophic Experimental Probability* is:

$$\left(\frac{\text{number of times event } A \text{ occurs}}{\text{total number of trials}}, \frac{\text{number of times indeterminacy occurs}}{\text{total number of trials}}, \frac{\text{number of times event } A \text{ does not occur}}{\text{total number of trials}} \right). \quad (113)$$

3.38. Neutrosophic Survey.

A *Neutrosophic Survey* is a way to obtain neutrosophic experimental probability.

Example. Let's say that we toss a regular die five times on an irregular surface, and we get: 2, 5, 1, *indeterminacy*, 4.

3.39. Neutrosophic Conditional Probability for Independent Events.

In *classical probability*, if A and B are *independent events*, then $P(A \text{ given } B) = P(A)$. (114)

Similarly for *neutrosophic independent events*:

$$NP(A \text{ given } B) = NP(A), \quad (115)$$

because

$$\begin{aligned} ch(A \text{ given } B) &= ch(A), \\ ch(\text{indeterm}_A \text{ given } B) &= ch(\text{indeterm}_A), \end{aligned} \quad (116)$$

and

$$ch(\bar{A} \text{ given } B) = ch(\bar{A}). \quad (117)$$

3.40. Neutrosophic Probability of an Impossible Event (Φ) on the neutrosophic probability space $\nu\Omega$ is:

$$NP(\Phi) = \left(0, ch(\text{indeterm}), ch(\nu\Omega) - ch(\text{indeterm}) \right). \quad (118)$$

3.41. Neutrosophic Probability of a Sure Event ($\nu\Omega$) on the neutrosophic probability space $\nu\Omega$ is:

$$NP(\nu\Omega) = (1 - ch(\text{indeterm}), ch(\text{indeterm}), 0). \quad (119)$$

3.42. Neutrosophic Bayesian Rule.

In *classical probability*, the Bayesian Rule is:

$$P(A|B) = P(B|A) \frac{P(A)}{P(B)}. \quad (120)$$

Let's examine the neutrosophic version of this rule.

Suppose we have an urn with 5 A -votes, 2 indeterminate (unclear, erased) votes, and 3 B -votes.

If A is the event of extracting an A -vote from the urn, and B the event of extracting a B -vote from the urn, then:

$$NP(A) = \left(\frac{5}{10}, \frac{2}{10}, \frac{3}{10} \right), NP(B) = \left(\frac{3}{10}, \frac{2}{10}, \frac{5}{10} \right). \quad (121)$$

If one B -vote has been taken from the urn, then

$$NP(A|B) = \left(\frac{5}{9}, \frac{2}{9}, \frac{2}{9} \right). \quad (122)$$

But if one A -votes has been taken from the urn, then

$$NP(B|A) = \left(\frac{3}{9}, \frac{2}{9}, \frac{4}{9} \right). \quad (123)$$

In general, the *Neutrosophic Bayesian Rule* is:

$$NP(A|B) = (ch(A|B), ch(indeterm_A | B), ch(\bar{A}|B))$$

$$= \left(ch(B|A) \frac{ch(A)}{ch(B)}, ch(indeterm_A | B), ch(\bar{A}|B) \right). \quad (124)$$

Therefore, as in classical probability

$$ch(A|B) = ch(B|A) \frac{ch(A)}{ch(B)}. \quad (125)$$

For our particular example, we get:

$$\begin{aligned} NP(A|B) &= \\ &= \left(ch(B|A) \frac{ch(A)}{ch(B)}, ch(indeterm_A|B), ch(\bar{A}|B) \right) \\ &= \left(\frac{3}{9} \times \frac{\frac{5}{10}}{\frac{2}{3}}, \frac{2}{9}, ch(B|B) \right) \\ &= \left(\frac{5}{9}, \frac{2}{9}, \frac{2}{9} \right). \end{aligned} \quad (126)$$

3.43. Neutrosophic Multiplicative Rule.

In *classical probability*, the *Multiplication Rule for Probabilities* (equivalent with the *Conditional Probability*) is:

$$P(A \text{ and } B) = P(A) \cdot P(B \text{ given } A). \quad (127)$$

The *Multiplication Rule for Neutrosophic Probabilities* is:

$$\begin{aligned}
 NP(A \text{ and } B) = & (ch(A) \cdot ch(B \text{ given } A), \\
 & ch(indeterm_{A \text{ and } B}) + ch(indeterm_{A \text{ or } B} | A) - \\
 & ch(indeterm_{A \text{ and } B}) \cdot ch(indeterm_{A \text{ and } B} | A), \\
 & ch(A) \cdot ch(A \text{ given } A) + ch(B) \cdot ch(A \text{ given } B) + \\
 & ch(B) \cdot ch(B \text{ given } B)), \quad (128)
 \end{aligned}$$

because:

$$\begin{aligned}
 ch(indeterm \text{ for } (A \text{ and } B)) = & \\
 & = ch(indeterm) \cdot ch(indeterm|A) \\
 & + ch(indeterm) \cdot ch(A|A) \\
 & + ch(indeterm) \cdot ch(B|A) \\
 & + ch(indeterm|A) \cdot ch(A) \\
 & + ch(indeterm|A) \cdot ch(B) \\
 & = ch(indeterm|A) \\
 & \cdot \left[\underbrace{ch(indeterm) + ch(A) + ch(B)} \right] \\
 & + ch(indeterm) \\
 & \cdot \left[\underbrace{ch(A|A) + ch(B|A) + ch(indeterm|A)} \right] \\
 & - ch(indeterm|A) \Big] \\
 & = ch(indeterm) + ch(indeterm|A) \\
 & - ch(indeterm) \cdot ch(indeterm|A),
 \end{aligned}$$

due to the facts that

$$\left[\underbrace{ch(indeterm) + ch(A) + ch(B)} \right] = 1 \quad (129)$$

$$\text{and } \left[\underbrace{ch(A|A) + ch(B|A) + ch(indeterm|A)} \right] = 1. \quad (130)$$

Let's consider the previous neutrosophic example:

$$\begin{array}{ccc} 5 & 2 & 3 \\ A - \text{votes} & \text{indeterm} - \text{votes} & B - \text{votes} \end{array}$$

We pick two votes in succession without replacement.

Suppose A is the event that the first is an A-vote, and B is a B-vote.

We have:

$$ch(A) = \frac{5}{10}, ch(indeterm) = \frac{2}{10},$$

$$ch(B) = \frac{3}{10}, ch(A|A) = \frac{4}{9},$$

$$ch(indeterm|A) = \frac{2}{9}, ch(B|A) = \frac{3}{9},$$

$$ch(A|B) = \frac{5}{4}, ch(B|B) = \frac{2}{9},$$

whence:

$$NP(A \text{ and } B) = \left(\frac{5}{10} \cdot \frac{3}{9}, \frac{2}{10} + \frac{2}{9} - \frac{2}{10} \cdot \frac{2}{9}, \frac{5}{10} \cdot \frac{4}{9} + \frac{3}{10} \cdot \frac{5}{9} + \frac{3}{10} \cdot \frac{2}{9} \right) = \left(\frac{15}{90}, \frac{34}{90}, \frac{41}{90} \right). \quad (131)$$

3.44. Neutrosophic Negation (or Neutrosophic Probability of Complementary Events).

For any event A different from indeterminacy, from the sample space X , one has:

$$NP(A) = (ch(A), ch(indeterm_A), ch(antiA)), \quad (132)$$

whence the neutrosophic probability of the complement of A , noted as $\mathcal{L}(A)$ (or as $antiA$) is:

$$NP(\mathcal{L}(A)) = NP(antiA) = (ch(antiA), ch(indeterm_{antiA}), ch(anti(antiA))) = (ch(X)-ch(A), ch(indeterm_{antiA}), ch(A)). \quad (133)$$

3.45. De Morgan's Neutrosophic Laws.

$$\begin{aligned} NP(\mathcal{L}(A \cup B)) &= (ch(\mathcal{L}(A \cup B)), ch(indeterm_{\mathcal{L}(A \cup B)}), \\ &\quad ch(\mathcal{L}(\mathcal{L}(A \cup B)))) \\ &= (ch(\mathcal{L}(A) \cap \mathcal{L}(B)), ch(indeterm_{\mathcal{L}(A) \cap \mathcal{L}(B)}), \\ &\quad ch(\mathcal{L}(\mathcal{L}(A) \cap \mathcal{L}(B)))) \\ &= NP(\mathcal{L}(A) \cap \mathcal{L}(B)). \end{aligned} \quad (134)$$

Similarly,

$$\begin{aligned}
 NP(\mathcal{L}(A \cap B)) &= (\text{ch}(\mathcal{L}(A \cap B)), \text{ch}(\text{indeterm}_{\mathcal{L}(A \cap B)}), \\
 &\quad \text{ch}(\mathcal{L}(\mathcal{L}(A \cap B)))) \\
 &= (\text{ch}(\mathcal{L}(A) \cup \mathcal{L}(B)), \text{ch}(\text{indeterm}_{\mathcal{L}(A) \cup \mathcal{L}(B)}), \\
 &\quad \text{ch}(\mathcal{L}(\mathcal{L}(A) \cup \mathcal{L}(B)))) \\
 &= NP(\mathcal{L}(A) \cup \mathcal{L}(B)). \tag{135}
 \end{aligned}$$

3.46. Neutrosophic Double Negation.

In classical probability,

$$P(\text{anti}(\text{anti}A)) = P(A). \tag{136}$$

In neutrosophic probability, for A an event different from indeterminacy:

$$NP(A) = (\text{ch}(A), \text{ch}(\text{indeterm}_A), \text{ch}(\text{anti}A)), \tag{137}$$

then:

$$\begin{aligned}
 NP(\text{anti}A) &= (\text{ch}(\text{anti}A), \text{ch}(\text{indeterm}_{\text{anti}A}), \text{ch}(\text{anti}(\text{anti}A))) \\
 &= (\text{ch}(\text{anti}A), \text{ch}(\text{indeterm}_{\text{anti}A}), \text{ch}(A)) \tag{138}
 \end{aligned}$$

whence

$$\begin{aligned}
 NP(\text{anti}(\text{anti}A)) &= (\text{ch}(\text{anti}(\text{anti}A)), \text{ch}(\text{indeterm}_{\text{anti}(\text{anti}A)}), \\
 \text{ch}(\text{anti}A)) &= (\text{ch}(A), \text{ch}(\text{indeterm}_A), \text{ch}(\text{anti}A)) = NP(A). \tag{139}
 \end{aligned}$$

Let's reconsider the previous example about a urn with:

$$\begin{array}{ccc} 5 & 2 & 3 \\ A - \text{votes} & \text{indeterm} - \text{votes} & B - \text{votes} \end{array}$$

$NP(A) = (5/10, 2/10, 3/10)$,

then $NP(\text{anti}A) = (3/10, 2/10, 5/10)$,

and it follows that $NP(\text{anti}(\text{anti}A)) = (5/10, 2/10, 3/10) = NP(A)$.

3.47. Neutrosophic Expected Value.

Let's consider a neutrosophic discrete probability space X with the determined outcomes x_1, x_2, \dots, x_r and their respective chances to occur p_1, p_2, \dots, p_r , and with indeterminacies $\text{indeterm}_1, \text{indeterm}_2, \dots, \text{indeterm}_k$, then the Neutrosophic Expected Value (NE) is:

$$NE = \sum_{j=1}^r n_j p_j + \sum_{k=1}^s (m_k \cdot \text{ch}(\text{indeterm}_k)) \quad (140)$$

where n_j is the possible numerical outcome for the corresponding chance p_j , for all j , and m_k is the possible numerical outcome for the corresponding chance that indeterm_k occurs, for all k .

If we reconsider the previous neutrosophic example:

$$\begin{array}{ccc}
 5 & 2 & 3 \\
 A - \text{votes} & \text{indeterm} - \text{votes} & B - \text{votes}
 \end{array}$$

And the numerical outcomes for extracting an *A*-vote is loosing \$2.00, for extracting a *B*-vote is gaining \$3.00, while for extracting an indeterminate vote is loosing \$1.00. What is the neutrosophic expected value?

$$NE = -2 \times (5/10) + 3 \times (3/10) - 1 \times (2/10) = -\$0.30. \quad (141)$$

3.48. Neutrosophic Probability and Neutrosophic Logic Used in The Soccer Games.

For all games where there are three possible results (winning, loosing, or tie), neutrosophic probability works perfectly, but the classical or imprecise probabilities do not apply, since they can describe one result only.

Let's say: What is the probability that a soccer team wins in a soccer game? Neutrosophic probability gives all three chances: chance to win, chance to get a tie game, and chance to loose.

Suppose two soccer games will play: teams *Alpha* (α) vs. *Beta* (β), and *Gamma* (γ) vs. *Delta* (δ).

$$NP(\text{Alpha to win}) = (0.7, 0.2, 0.1), \quad (142)$$

which means that *Alpha* has 0.6 chance to win, 0.2 chance to tie game, and 0.1 chance to loose,

$$NP(\textit{Gamma to win}) = (0.3, 0.5, 0.2), \quad (143)$$

then what is the neutrosophic probability that both teams *Alpha* and *Gamma* win in their soccer games?

We make the product of the neutrosophic probability spaces:

$$\begin{aligned} & \left\{ \begin{array}{ccc} W_\alpha, & I_{\alpha\beta}, & L_\alpha \\ 0.7 & 0.2 & 0.1 \end{array} \right\} \times \\ & \left\{ \begin{array}{ccc} W_\gamma, & I_{\gamma\delta}, & L_\gamma \\ 0.3 & 0.5 & 0.2 \end{array} \right\} \\ & \hspace{15em} (144) \end{aligned}$$

where $W_\alpha = \alpha$ winning, $I_{\alpha\beta} =$ indeterminacy (tie games between α and β), $L_\alpha = \alpha$ loosing; similarly for $W_\gamma, I_{\gamma\delta}, L_\gamma$, which is

$$\left\{ \begin{array}{ccccccccc} W_\alpha W_\gamma, & W_\alpha I_{\gamma\delta}, & W_\alpha L_\gamma, & I_{\alpha\beta} W_\gamma, & I_{\alpha\beta} I_{\gamma\delta}, & I_{\alpha\beta} L_\gamma, & L_\alpha W_\gamma, & L_\alpha I_{\gamma\delta}, & L_\alpha L_\gamma \end{array} \right\}, \quad (145)$$

and the numbers below each possible outcome represent their corresponding chances to occur.

We can re-arrange the final result in many ways.

a) In the classical probability, we can say:

$$P((Alpha\ winning)\&(Gamma\ winning)) = 0.7(0.3) = 0.21, \quad (146)$$

while $1 - 0.21 = 0.79$ is the probability of the opposite event, negation of

$((Alpha\ winning)\&(Gamma\ winning))$, i.e. in the two soccer games either there is at least a tie game, or at least one of the teams *Alpha* or *Gamma* loses.

b) In the neutrosophic probability, the outcome is more refined.

$$\begin{aligned} b1) \quad NP((Alpha\ winning)\&(Gamma\ winning)) \\ = \{ ch(Alpha\ Winning \& Gamma\ winning), \\ \quad ch(at\ least\ one\ tie\ game), \\ ch(referring\ to\ Alpha\ and\ Gamma: \text{one losing,} \\ \quad \text{the other winning,} \\ \quad \text{or both losing}) \} = \\ = (0.21, 0.35 + 0.06 + 0.10 + 0.04 + \\ 0.05, 0.14 + 0.03 + 0.02) = \\ (0.21, 0.60, 0.19). \end{aligned} \quad (147)$$

$$\begin{aligned} b2) \quad NP((Alpha\ winning)\&(Gamma\ winning)) = \\ \{ (Alpha\ winning)\&(Gamma\ winning), \\ \quad ch(at\ least\ one\ of\ Alpha \\ and\ Gamma\ has\ a\ tie\ game, \text{and none loses}), \end{aligned}$$

$$\begin{aligned}
& ch(\text{at least one} \\
& \text{of Alpha or Gamma loses}) = \\
& = (0.21, 0.35 + 0.06 + 0.10, 0.14 + 0.04 + \\
& \quad 0.03 + 0.05 + 0.02) = (0.21, 0.51, 0.28).
\end{aligned}
\tag{148}$$

c) Another solution to this soccer game would be to use neutrosophic logic. Let's consider

$$P_1 = \{\text{Team Alpha will win}\},$$

and $P_2 = \{\text{Team Gamma will win}\}$

as two neutrosophic logical propositions whose values are $(0, 0.7, 0.2, 0.1)$, and respectively $(0.3, 0.5, 0.2)$.

Then one uses the neutrosophic operator “and” (Λ_N) as part of the N – norm:

$$P_1 \Lambda_N P_2 = (0.7 \Lambda_F 0.3, 0.2 \vee_F 0.5, 0.1 \vee_F 0.2),$$

(149)

where Λ_F is the fuzzy “and” operator (t-norm),

and \vee_F is the fuzzy “or” operator (t-conorm).

c1. If we take the fuzzy and/or operators min/max, we get:

$$\begin{aligned}
P_1 \Lambda_N P_2 = \\
& (\min(0.7, 0.3), \max(0.2, 0.5), \max(0.1, 0.2)) = \\
& (0.3, 0.5, 0.2).
\end{aligned}
\tag{150}$$

c2. If we take the fuzzy *and/or* operators
as $x \cdot y / x + y - x \cdot y$, (151)

we get

$$P_1 \Lambda_N P_2 = (0.7(0.3), 0.2 + 0.5 - 0.2(0.5), 0.1 + 0.2 - 0.1(0.2)) = (0.21, 0.60, 0.28). \quad (152)$$

(In neutrosophic logic, the sum of its three components may be different from 1.)

Similarly for other particular t-norms/t-conorms.

3.49. A Neutrosophic Question.

Rolling two regular dice on an irregular surface, what is the neutrosophic probability of getting a sum of 6?



Fig. 9. Double Indeterminacy

The five favorable cases will be:

$$1 + 5, 2 + 4, 3 + 3, 4 + 2, 5 + 1. \quad (153)$$

But what about:

$$6 + \textit{indeterm}, \textit{and indeterm} + 6?$$

Should we consider that

$$6 + \textit{indeterm} = 6 \quad (154)$$

(and consequently: $indeterm + 6 = 6$) ?

or should we say that

$$6 + indeterm = indeterm ?$$

Of course,

$$indeterm + indeterm = indeterm. \quad (155)$$

Surely, in a game the players can make conventions among themselves, for example that a number plus indeterminacy is equal to that number, but this would mean that indeterminacy is taken for zero, which is not quite true.

Let's compute $NP(sum = 6)$.

Neutrosophic Probability Spaces are:

$$\nu\Omega_1 = \{1, 2, 3, 4, 5, 6, indeterm_1\} \text{ for die \#1;}$$

$$\nu\Omega_2 = \{1, 2, 3, 4, 5, 6, indeterm_2\} \text{ for die \#2.}$$

The neutrosophic probability product space is

$$\begin{aligned} & \Omega_1 \times \nu\Omega_2 \\ & = \{ (1, 1), (1, 2), \dots, (1, 6), \dots (6, 1), (6, 2), \dots, (6, 6), \\ & \quad (indeterm_1, 1), (indeterm_1, 2), \dots, (indeterm_1, 6), \\ & \quad (1, indeterm_2), (2, indeterm_2), \dots, (6, indeterm_2), \end{aligned}$$

$$(\text{indeterm}_1, \text{indeterm}_2) \}. \quad (156)$$

Considering that through frequentist neutrosophic experiment for a single die we have found the chance of getting indeterminacy is 0.10, whence

$$ch(1) = \dots = ch(6) = \frac{1-0.1}{6} = 0.15, \text{ we get:}$$

$$\begin{aligned} NP(\text{sum} = 6) == & (ch(\text{sum} = 6), ch(\text{indeterm}_{\text{sum}=6}), ch(\text{sum} \neq \\ & 6 \text{ and no indeterm})) = (5 \cdot \\ & (0.15)(0.15), 12(0.10)(0.15) + \\ & 0.10(0.10), 31(0.15)(0.15)) = \\ & (0.1125, 0.1900, 0.6975). \end{aligned} \quad (157)$$

In classical probability, where there is no indeterminacy,

$$\begin{aligned} P(\text{sum} = 6) &= \frac{5}{36} \approx 0.1389 > 0.1125 \\ &= ch(\text{sum} = 6) \end{aligned} \quad (158)$$

from neutrosophic probability.

3.50. Neutrosophic Discrete Probability Spaces.

In general, if we have two neutrosophic discrete probability spaces, with the chances of occurring of each event listed below that event

$$\nu\Omega_1 = \begin{array}{cccc} \{A_1, & A_2, & \dots, & A_n, & \text{indeterm}_1\} \\ p_1, & p_2, \dots, & & p_n, & \text{ch(indeterm}_1) \end{array}$$

and

$$\nu\Omega_2 = \begin{array}{cccc} \{B_1, & B_2, & \dots, & B_n, & \text{indeterm}_2\} \\ q_1, & q_2, \dots, & & q_n, & \text{ch(indeterm}_2), \end{array}$$

then the neutrosophic probability of having event A_j and event B_k to occur is:

$$NP(A_j \text{ and } B_k) = \left(p_j \cdot q_k, I_1 + I_1 - I, I_2, \sum_{v=1, n} p_u q_v - p_j \cdot q_k \right). \quad (159)$$

This can be further generalized to the neutrosophic discrete probability product of s spaces:

$$\begin{aligned} \nu\Omega_1 \times \nu\Omega_2 \times \dots \times \nu\Omega_s \\ = \prod_{r=1}^s \{A_{r,1}, A_{r,2}, \dots, A_{r,n_r}, \text{indeterm}_r\} \end{aligned}$$

with corresponding neutrosophic probabilities

$$P_{r,1} \ P_{r,2} \ \dots \ P_{r,n_r} \ I_r.$$

$$\text{Then } NP(A_{1,j_1} \text{ and } A_{2,j_2} \text{ and } \dots \text{ and } A_{s,j_s}) =$$

$$\left(\prod_{r=1}^s p_{r,j_r}, \sum_{t=1}^s (-1)^{t-1} S_t, \sum_{\substack{k_1=1,n_1 \\ \dots \\ k_s=1,n_s}} (\prod_{r=1}^s p_{r,k_r}) - \right.$$

$$\left. \prod_{r=1}^s p_{r,j_r} \right), \quad (160)$$

where

$$S_1 = I_1 + I_2 + \dots + I_s \quad (s \text{ terms} = C_s^1)$$

$$S_2 = I_1 I_2 + I_1 I_3 + \dots + I_1 I_s + \dots$$

$$+ I_{s-1} I_s \quad \left(\frac{s(s-1)}{2} \text{ terms} = C_s^2 \right)$$

.....

$$S_t =$$

$$\sum_{(j_1, j_2, \dots, j_t) \in \mathcal{C}_{\{1, 2, \dots, s\}}^t} I_{j_1} I_{j_2} \dots I_{j_t} \quad \left(\frac{s!}{t!(s-t)!} \text{ terms} = C_s^t \right)$$

$$(161)$$

where $\mathcal{C}_{\{1, 2, \dots, s\}}^t$ is the family of all subsets of $\{1, 2, \dots, s\}$, such that the cardinal of each subset is t , for $1 \leq t \leq s$, t integer.

3.51. Classification of Neutrosophic Probabilities.

1. There is an *objective neutrosophic probability* when the chances of all events, including the chance of indeterminacy, can be computed objectively.

For example: Tossing a cubic die, which has two faces that are unreadable, on a regular surface. Let's consider that the numbers 5 and 6 are erased. We can exactly compute

$$ch(\text{indeterm}) = \frac{2}{6},$$

$$ch(1) = ch(2) = ch(3) = ch(4) = \frac{1}{6}.$$

2. The *frequentist neutrosophic probability* when at least the chance of one event, or the chance of some indeterminacy, cannot be computed objectively (exactly), but one can make experiments in order to compute frequentist chances.

For example: Tossing a regular die on an irregular surface having many cracks. We are not able to exactly compute the chance of indeterminacy (i.e when the die gets stuck in a crack on a vertex or on an edge). We may

experiment, tossing the die a number of times in order to compute chance of indeterminacy as number of favorable cases over total number of cases. But this is just an approximation. And if we repeat the experiment, we get a different result.

3. *Subjective Neutrosophic Probability* is neither possible to compute it objectively (exactly), nor to experiment it and compute it as frequentist chance.

For example, an aircraft is detected in the sky. A source estimates it as to be 60% friend, 30% hostile, and 10% neutral. The estimation is subjective. Another source could give us a different estimation.

3.52. The Fundamental Neutrosophic Counting Principle.

Let's consider a neutrosophic event E that can occur in e ways and e_1 indeterminacies. After E has occurred, a neutrosophic event F can occur in f ways and f_1 indeterminacies. Then, the neutrosophic event E followed by the neutrosophic event F can occur in $e \cdot f$ ways, and in $e_1 \cdot f + e \cdot f_1$ indeterminacies of first order, and in $e_1 \cdot f_1$ indeterminacies of second order.

Taking the previous example about tossing a cubic die, which has two faces 5 and 6 that are unreadable, but

on an irregular surface. And then another cubic die with all readable faces, tossed on an irregular surface. We have the following neutrosophic sample spaces:

$$\nu\Omega_1 = \{1, 2, 3, 4, \text{indeterm}_{die}, \text{indeterm}_{space}\} \quad (162)$$

$$\nu\Omega_2 = \{1, 2, 3, 4, 5, 6, \text{indeterm}_{space}\} \quad (163)$$

Then an event E followed by an event F can occur in

$$4 \cdot 6 = 24 \text{ ways,}$$

$$2 \cdot 6 + 4 \cdot 1 = 16 \text{ indeterminacies of first order,}$$

and

$$2 \cdot 1 = 2 \text{ indeterminacies of second order.}$$

3.53. A Formula for the Fusion of Subjective Neutrosophic Probabilities.

For subjective neutrosophic probability, in order for us to having a better aproximation, we'd need more sources of information relating us about *the same event*. We then combine all subjective chances given by them.

Suppose a satellite is detected by radar in the sky, which can be friendly (\underline{t}), neutral (\underline{i}), or hostile (\underline{f}). We have two observers that give us the following subjective neutrosophic probabilities:

$$NP_1(\text{satellite}) = (t_1, i_1, f_1), \quad (164)$$

where $t_1 = ch(\text{satellite is freindly})$,

$$i_1 = ch(\text{satellite is neutral}),$$

$$f_1 = ch(\text{satellite is hostile}),$$

and

$$NP_2(\text{satellite}) = (t_2, i_2, f_2). \quad (165)$$

We consider the following normalized probabilities:

$$t_1 + i_1 + f_1 = t_2 + i_2 + f_2 = 1, \quad (166)$$

but in case if they are non-normalized, the problem is solved in the same way. Note that \underline{t} stands for truth, \underline{i} stands for indeterminacy, and \underline{f} stands for falsehood.

$$(NP_1 \cap NP_2(t) = t_1 t_2 + \left(\frac{t_1^2 i_2}{t_1 + i_2} + \frac{t_2^2 i_1}{t_2 + i_1} \right) + \left(\frac{t_1^2 f_2}{t_1 + f_2} + \frac{t_2^2 f_1}{t_2 + f_1} \right)). \quad (167)$$

Because:

$t_1 \cdot i_2$ is redistributed back to the truth (t) and indeterminacy (i), proportionally with respect to t_1 , and respectively to i_2 :

$$\frac{x_1}{t_1} = \frac{y_1}{i_2} = \frac{t_1 i_2}{t_1 + i_2}, \quad (168)$$

where x_1 and y_1 are the parts from $t_1 \cdot i_2$ that are redistributed back to t (chance of friendly) and respectively to i (chance of neutral),

whence

$$x_1 = \frac{t_1^2 i_2}{t_1 + i_2}, \quad y_1 = \frac{t_1 i_2^2}{t_1 + i_2}. \quad (169)$$

Similarly $t_2 \cdot i_1$ is redistributed back to the truth (t) and indeterminacy (i), proportionally with respect to t_2 , and respectively to i_1 :

$$\frac{x_2}{t_2} = \frac{y_2}{i_1} = \frac{t_2 i_1}{t_2 + i_1}, \quad (170)$$

where x_2 and y_2 are the parts from $t_2 \cdot i_1$ that are redistributed back to t (chance of friendly) and respectively to i (chance of neutral),

whence

$$x_2 = \frac{t_2^2 i_1}{t_2 + i_1}, \quad y_2 = \frac{t_2 i_1^2}{t_2 + i_1}. \quad (171)$$

Again, $t_1 \cdot f_2$ is redistributed back to t and f (falsehood) proportionally with respect to t_1 and respectively f_2 :

$$\frac{x_3}{t_1} = \frac{z_1}{f_2} = \frac{t_1 f_2}{t_1 + f_2}, \quad (172)$$

whence

$$x_3 = \frac{t_1^2 f_2}{t_1 + f_2}, \quad z_1 = \frac{t_1 f_2^2}{t_1 + f_2}. \quad (173)$$

And, similarly $t_2 \cdot f_1$ is redistributed back to t and f proportionally with respect to t_2 and respectively f_1 :

$$\frac{x_4}{t_2} = \frac{z_2}{f_1} = \frac{t_2 f_1}{t_2 + f_1}, \quad (174)$$

whence

$$x_4 = \frac{t_2^2 f_1}{t_2 + f_1}, \quad z_2 = \frac{t_2 f_1^2}{t_2 + f_1}. \quad (175)$$

In the same way, $i_1 \cdot f_2$ is redistributed back to i and f proportionally with respect to i_1 and respectively f_2 :

$$\frac{y_3}{i_1} = \frac{z_3}{f_2} = \frac{i_1 f_2}{i_1 + f_2}, \quad (176)$$

whence

$$y_3 = \frac{i_1^2 f_2}{i_1 + f_2}, \quad z_3 = \frac{i_1 f_2^2}{i_1 + f_2}. \quad (177)$$

While $i_2 \cdot f_1$ is redistributed back to i and f proportionally with respect to i_2 and respectively f_1 :

$$\frac{y_4}{i_2} = \frac{z_4}{f_1} = \frac{i_2 f_1}{i_2 + f_1}, \quad (178)$$

whence

$$y_4 = \frac{i_2^2 f_1}{i_2 + f_1}, \quad z_4 = \frac{i_2 f_1^2}{i_2 + f_1}. \quad (179)$$

Then,

$$(NP_1 \cap NP_2)(i) = i_1 i_2 + \left(\frac{i_1^2 t_2}{i_1 + t_2} + \frac{i_2^2 t_1}{i_2 + t_1} \right) + \left(\frac{i_1^2 f_2}{i_1 + f_2} + \frac{i_2^2 f_1}{i_2 + f_1} \right) \quad (180)$$

and

$$(NP_1 \cap NP_2)(f) = f_1 f_2 + \left(\frac{f_1^2 t_2}{f_1 + t_2} + \frac{f_2^2 t_1}{f_2 + t_1} \right) + \left(\frac{f_1^2 i_2}{f_1 + i_2} + \frac{f_2^2 i_1}{f_2 + i_1} \right). \quad (181)$$

3.54. Numerical Example of Fusion of Subjective Neutrosophic Probabilities:

$$(0.6, 0.1, 0.3) \wedge_A (0.2, 0.3, 0.5) = (0.44097, 0.15000, 0.40903), \quad (182)$$

because

$$\left. \begin{aligned} t_1 = 0.6, i_1 = 0.1, f_1 = 0.3 \\ t_2 = 0.2, i_2 = 0.3, f_2 = 0.5 \end{aligned} \right\} \quad (183)$$

are replaced into the three previous neutrosophic formulas:

$$\begin{aligned}
(NP_1 \cap NP_2)(t) &= \\
&= 0.6(0.2) + \left[\frac{0.6^2(0.3)}{0.6 + 0.3} + \frac{0.2^2(0.1)}{0.2 + 0.1} \right] \\
&+ \left[\frac{0.6^2(0.5)}{0.6 + 0.5} + \frac{0.2^2(0.3)}{0.2 + 0.3} \right] \approx 0.44097;
\end{aligned}$$

$$\begin{aligned}
(NP_1 \cap NP_2)(i) &= \\
&= 0.1(0.3) + \left[\frac{0.1^2(0.2)}{0.1 + 0.2} + \frac{0.3^2(0.6)}{0.3 + 0.6} \right] \\
&+ \left[\frac{0.1^2(0.5)}{0.1 + 0.5} + \frac{0.3^2(0.3)}{0.3 + 0.3} \right] \approx 0.15000;
\end{aligned}$$

$$\begin{aligned}
(NP_1 \cap NP_2)(f) &= \\
&= 0.3(0.5) + \left[\frac{0.3^2(0.2)}{0.3 + 0.2} + \frac{0.5^2(0.6)}{0.5 + 0.6} \right] \\
&+ \left[\frac{0.3^2(0.3)}{0.3 + 0.3} + \frac{0.5^2(0.1)}{0.5 + 0.1} \right] \approx 0.40903.
\end{aligned}$$

(184)

Therefore, there is a higher chance that the satellite is friendly, because:

$$0.44097 > 0.40903 > 0.15000. \quad (185)$$

3.55. General Formula for Fusing Classical Subjective Probabilities Provided by Two Sources.

The principle of redistributing the conflicting chances, for example $t_1 i_2$, back to \underline{t} and \underline{i} , is the same as in PCR5 rule (Proportional Conflict Redistribution rule #5 from *The Dezert-Smarandache Theory of Paradoxist and Plausible Reasoning (DSmT)*) used in information fusion:

if two sources of information S_1 , and S_2 give the subjective probabilities P_1 and P_2 about the same event E to occur,

then combining both of them using PCR5 we get

for any event E in the subjective probability space Ω ,

$$(P_1 \wedge P_2)(E) = P_1(E) \cdot P_2(E) + \sum_{\substack{x \in \Omega \\ x \cap E = \emptyset}} \left[\frac{P_1(E)^2 \cdot P_2(x)}{P_1(E) + P_2(x)} + \frac{P_2(E)^2 \cdot P_1(x)}{P_2(E) + P_1(x)} \right] \quad (186)$$

3.56. Different Ways of Combining Neutrosophic Subjective Probabilities Provided by Two Sources.

Let's set the problem in a different way and use different notations. By combining subjective probabilities we don't get a single result, but many, since we do approximations.

A military vehicle is moving in a warzone. Two experts report the chances of the vehicle to be friendly (F), neutral (N), or hostile (H):

$$NP_1(\text{vehicle}) = (F_1, N_1, H_1) \quad (187)$$

and

$$NP_2(\text{vehicle}) = (F_2, N_2, H_2), \quad (188)$$

where all $F_1, N_1, H_1, F_2, N_2, H_2$ are chances (numbers in $[0,1]$), such that $F_1 + N_1 + H_1 = F_2 + N_2 + H_2 = 1$ (normalized neutrosophic probabilities).

Suppose $NP_1 \wedge NP_2 = (F, N, H)$, with similarly F, N, H in $[0,1]$ and $F + N + H = 1$.

Let's multiply $(F_1 + N_1 + H_1)(F_2 + N_2 + H_2) = 1 \times 1 = 1$.

We get 9 terms in the left side:

$$F_1F_2 + F_1N_2 + F_1H_2 + N_1F_2 + N_1N_2 + N_1H_2 + H_1F_2 + H_1N_2 + H_1H_2 = 1. \quad (189)$$

These 9 terms are distributed to F, N, H .

Of course F_1F_2 will go to F , N_1N_2 will go to N , and H_1H_2 will go to H .

The other 6 terms have to be distributed to F , N , H too.

We pay attention to the symmetry of distribution of these 6 terms to F and H .

a) *Pessimistic case:*

F_1N_1 and F_2N_1 to N .

Similarly H_1N_2 and H_2N_1 to N .

There are left F_1H_2 and F_2H_1 .

a1) We can either distribute both of them to N (in a *very pessimistic case*);

a2) or we can use, for example PCR5, to redistribute them back to F and H proportionally (in a *less pessimistic way*).

b) *Optimistic case:*

F_1N_1 and F_2N_1 to F .

Similarly H_1N_2 and H_2N_1 to H .

There are left F_1H_2 and F_2H_1 ,

b1) We can either distribute both of them to N (in a *less optimistic case*),

b2) or we can use for example PCR5 to redistribute them back to F and H proportionally (in a *very optimistic way*).

No normalization is needed, since the sum $F + N + H$ will be 1.

3.57. Neutrosophic Logic Inference type in Fusing Subjective Neutrosophic Probabilities.

Let the neutrosophic probability space be $\Phi = \{F, N, H\}$, where F = friend, N = neutral, H = hostile. If we consider that all intersections of events are empty:

$$F \cap N = F \cap H = N \cap H = \emptyset, \quad (190)$$

we can use the neutrosophic logic inference.

Suppose an aircraft is detected. We need to find out if it's a friend, or neutral, or hostile.

We have two sources that give the subjective chances:

	F	N	H
NP_1	a_1	a_2	a_3
NP_2	b_1	b_2	b_3

where all $a_1, a_2, a_3, b_1, b_2, b_3$ are positive and $a_1 + a_2 + a_3 = b_1 + b_2 + b_3 = 1$.

Then

$$NP_1 \wedge_p NP_2 = (a_1, a_2, a_3) \wedge_p (b_1, b_2, b_3) = (a_1 \wedge b_1, a_2 \vee b_2, a_3 \vee b_3) \quad (191)$$

in a *pessimistic way*,

or

$$NP_1 \wedge_o NP_2 = (a_1, a_2, a_3) \wedge_o (b_1, b_2, b_3) = (a_1 \wedge b_1, a_2 \wedge b_2, a_3 \vee b_3) \quad (192)$$

in an *optimistic way*.

" \wedge " is a t-norm operation, and " \vee " is a t-conorm operation. For example, \wedge/\vee can be as in fuzzy logic respectively:

min/max,

$xy / x + y - xy,$

$$\max\{0, x + y - 1\} / \min\{1, x + y\}, \text{ etc.} \quad (193)$$

Then we normalize each way if needed.

The N-norm and N-conorm were defined above with the help of t-norm and t-conorm.

Numerical Example:

	<i>F</i>	<i>N</i>	<i>H</i>
NP_1	0.4	0.1	0.5
NP_2	0.3	0.5	0.2

$$\begin{aligned}
 NP_1 \wedge_p NP_2 &= (0.4, 0.1, 0.5) \wedge_p (0.3, 0.5, 0.2) = \\
 &= (0.4 \wedge 0.3, 0.1 \vee 0.5, 0.5 \vee 0.2) = \\
 &= (\min\{0.4, 0.3\}, \max\{0.1, 0.5\}, \max\{0.5, 0.2\}) = \\
 &\quad (\text{using } \min/\max \text{ operators}) \\
 &= (0.3, 0.5, 0.5) \text{ normalized to } \left(\frac{3}{15}, \frac{5}{15}, \frac{5}{15}\right).
 \end{aligned}$$

$$\begin{aligned}
 NP_1 \wedge_0 NP_2 &= \\
 &(\min\{0.4, 0.3\}, \min\{0.1, 0.5\}, \max\{0.5, 0.2\}) = \\
 &(0.3, 0.1, 0.5) \text{ normalized to } \left(\frac{3}{9}, \frac{1}{9}, \frac{5}{9}\right). \quad (194)
 \end{aligned}$$

If we combine both pessimist and optimist results we get:

<i>F</i>	<i>N</i>	<i>H</i>
$\left[\frac{3}{15}, \frac{3}{9}\right]$	$\left[\frac{1}{9}, \frac{5}{15}\right]$	$\left[\frac{5}{15}, \frac{5}{9}\right]$

It would be interesting of computing using the other \wedge/\vee operators and compare or combine (for example making the average of) the results.

3.58. Neutrosophic Logic vs. Subjective Neutrosophic Probability.

In *neutrosophic logic*, the operator AND computes the conjunction of the two or more different logical propositions.

In *subjective neutrosophic probability*, there is a single neutrosophic probability space, and two or more sources of information that provide subjective chances about the events to occur. Then we use various procedures to aggregate the subjective probabilities provided by all sources in order to get the best estimation.

3.59. Removing Indeterminacy.

We can remove indeterminacy from the sample space, but then the second axiom of Kolmogorov is invalidated, because the neutrosophic probability of the whole sample space is strictly less than 1.

Let's suppose having a cubic die whose three faces 4, 5, 6 are erased and unreadable. We then have

$$ch(1) = ch(2) = ch(3) = \frac{1}{6} \quad (195)$$

and

$$ch(\text{indeterm}) = 3 \left(\frac{1}{6} \right) = \frac{3}{6} = \frac{1}{2}. \quad (196)$$

So, if we remove the indeterminacy, our neutrosophic sample space becomes:

$$\nu\Omega = \{1, 2, 3\} \quad (197)$$

and

$$ch(\nu\Omega) = \frac{1}{2} < 1. \quad (198)$$

$\nu\Omega$ is an *incomplete classical sample space*. The first and third axioms remain valid, but the second axiom is invalidated.

3.60. n-Valued Refined Neutrosophic Probability Space and Neutrosophic Probability.

Let's consider a handball game between two teams, Romania and Bulgaria. What is the neutrosophic probability that Romania wins?

The simplest sample space with respect to Romania is:

$$\nu\Omega = \{\text{victory, equality, defeat}\}. \quad (199)$$

Suppose that the neutrosophic probability measure NP (Romania wins) = (0.7, 0.1, 0.2). But we go further and refine the sample space and, implicitly, the probability measure.

The n -Valued Refined Neutrosophic Sample Space is:

$$\nu\Omega = \left\{ \{V_1, V_2, \dots, V_p\}, \{E_1, E_2, \dots, E_h\}, \{D_1, D_2, \dots, D_s\} \right\} \quad (200)$$

where $p, r, s \geq 1$ and $p + r + s = n$;

also:

V_1 = Romania wins with 1 goal difference (i.e. 1-0, 2-1, 3-2, etc.);

.....

V_{p-1} = Romania wins with $p-1$ goals difference;

V_p = Romania wins with p or more goals difference;

E_1 = tie game (equality), with score 0-0, 1-1;

E_2 = tie game, with score 2-2;

E_{r-1} = tie game, with score (r-1) to (r-1);

E_r = tie game, with score x to x , where $x \geq r$.

Similarly:

D_1 = Romania is defeated with 1 goal difference (i.e. 0-1, 1-2, 2-3, etc.);

.....

D_{s-1} = Romania is defeated with $s - 1$ goals difference;

D_s = Romania is defeated with s or more goals difference.

Consequently, the *n-Valued Refined Neutrosophic Probability Measure* could be:

$$\begin{aligned}
 NP' \text{ (Romania wins)} &= \\
 &= \left(\left(\underbrace{(0.4, 0.2, 0.5, 0.05, 0, \dots, 0)}_p \right), \left(\underbrace{(0.03, 0.05, 0.02, 0, \dots, 0)}_r \right), \left(\underbrace{(0.1, 0.08, 0.02, 0, \dots, 0)}_s \right) \right).
 \end{aligned}
 \tag{201}$$

In general, let's consider a neutrosophic probability space, and a neutrosophic event A .

$$NP(A) = (ch(A), ch(indeterm_A), ch(\bar{A})) =$$

by notation to (T, I, F) ,

where for simplicity $T = \text{truth}$, $I = \text{indeterminacy}$, and $F = \text{falsity}$ as in neutrosophic logic.

Then the *n-Valued Refined Neutrosophic Probability* is:

$$NP_h(A) = \left((T_1, T_2, \dots, T_p), (I_1, I_2, \dots, I_r), (F_1, F_2, \dots, F_s) \right) \quad (202)$$

with $p, r, s \geq 1$ and $p + r + s = n$;

and $T_j = \text{the chance that event } A \text{ occurs and the occurrence has the property } P_j$;

$I_k = \text{the indeterminacy related to the occurrence of event } A, \text{ such that the indeterminacy has the property } Q_k$;

$F_l = \text{the chance that event } A \text{ does not occur and the non-occurrence has the property } R_l$;

where $1 \leq j \leq p, 1 \leq k \leq r, \text{ and } 1 \leq l \leq s$.

Remarks.

- a) Such n-Valued refinement is not possible for all applications.

a1) For example, if we consider a cubic die with four erased faces 1, 2, 3, and 4, tossed on a regular surface, then $NP(5) = \left(\frac{1}{6}, \frac{4}{6}, \frac{1}{6}\right)$; but *we are not able to refine* the occurrence of {5}, neither the non-occurrence of {5}, nor the indeterminacy that might be related to this event.

a2) But if we consider a regular die tossed on an irregular surface with several small cracks and other deep cracks, *we may refine the indeterminacy* for each event, since the die may get stucked in a small crack with a vertex of faces, we can read (for example 4 & 5 & 6), or the die can fall in a deep crack that we are not able to see it at all. Yet, we are not able to refine the occurrence or non-occurrence of an event in this case.

b) The refinements can be done in multiple ways, depending on the properties P_j, Q_k, R_l we choose, for all j, k, l .

3.61. Neutrosophic Markov Chain.

It is a straight neutrosophic generalization of the classical Markov chain, i.e. some indeterminacy is taken into consideration in the classical probability space.

The neutrosophic Markov chain is a sequence of neutrosophic random variables X_1, X_2, \dots , with the property that the next neutrosophic state depends on the current neutrosophic state only:

$$NP(X_n = x | X_1 = x_1, X_2 = x_2, \dots, X_{n-1} = x_{n-1}) = NP(X_n = x | X_{n-1} = x_{n-1}). \quad (203)$$

It is a *neutrosophic mathematical system* that is characterized as memoryless.

A *neutrosophic transition* is a change of the state of a system with indeterminacy.

A *neutrosophic Markov chain of order m* , where $1 \leq m < \infty$, or neutro-sophic Markov chain with memory m , is:

$$\begin{aligned} NP(X_n = x | X_{n-1} = x_{n-1}, X_{n-2} = x_{n-2}, \dots, X_1 = x_1) = \\ = NP(X_n = x | X_{n-1} = x_{n-1}, X_{n-2} = x_{n-2}, \dots, X_{n-m} = x_{n-m}). \end{aligned} \quad (204)$$

We defined above the neutrosophic Markov chain for *discrete time*. For a *continuous time*, we use a continuous index:

$$NP(X_n = x | X_{n-1} = y) = NP(X_{n-1} = x | X_{n-2} = y), \text{ for all } n. \quad (205)$$

To illustrate an example about a discrete neutrosophic Markov chain, we use a *neutrosophic probability graph* (this should be distinguished from the *neutrosophic graph* and *neutrosophic fuzzy graph*, both introduced by W.B. Vasantha Kandasamy & F. Smarandache in our algebraic structure books since year 2003).

Let's consider the world economy, and its states: economic prosperity (P), economic recession (R), and economic depression (D).

Suppose we have the following neutrosophic probability graph, during a year:




Fig. 10

Upon the figure, an economic prosperity year is followed by another economic prosperity year 40% of the time while 10% of the time it is unknown, an economic recession year 20% of the time, while 10% of the time it is not followed by an economic recession, and an economic depression year 10% of the time and 5% of the time it is unknown while 5% of the time it is not followed by an economic recession year.

The neutrosophic transition matrix of this graph is:

$$\begin{array}{c}
 \begin{array}{ccc}
 & P & R & D \\
 P & (0.40, 0.10, 0.00) & (0.20, 0.00, 0.10) & (0.10, 0.05, 0.05) \\
 R & (0.19, 0.03, 0.08) & (0.35, 0.05, 0.00) & (0.24, 0.03, 0.04) \\
 D & (0.07, 0.03, 0.10) & (0.27, 0.09, 0.04) & (0.32, 0.06, 0.02)
 \end{array} \\
 NP = R \left[\begin{array}{ccc}
 (0.40, 0.10, 0.00) & (0.20, 0.00, 0.10) & (0.10, 0.05, 0.05) \\
 (0.19, 0.03, 0.08) & (0.35, 0.05, 0.00) & (0.24, 0.03, 0.04) \\
 (0.07, 0.03, 0.10) & (0.27, 0.09, 0.04) & (0.32, 0.06, 0.02)
 \end{array} \right]
 \end{array}$$

The state space is $\{P, R, D\}$.

The stochastic row vectors are:

$$\begin{array}{l}
 P = [1 \quad 0 \quad 0], \\
 R = [0 \quad 1 \quad 0], \\
 D = [0 \quad 0 \quad 1].
 \end{array}$$

Let X be any of these stochastic row vectors,

with the neutrosophic relation

$$X^{(n+1)} = X^{(n)}NP \tag{206}$$

for any time n .

Whence

$$\begin{aligned}
 X^{(n+2)} &= X^{(n+1)}NP \\
 &= [X^{(n)}NP]NP \\
 &= X^{(n)}(NP)^2 \tag{207}
 \end{aligned}$$

and more general:

$$X^{(n+m)} = X^{(n)}(NP)^m. \tag{208}$$

At the end we normalize the rows of matrix $(NP)^m$.

We define the multiplication of neutrosophic probabilities as

$$\begin{aligned}
 (a_1, b_1, c_1) \cdot (a_2, b_2, c_2) &= \\
 (a_1 a_2, \max\{b_1, b_2\}, \max\{c_1, c_2\}), \tag{209}
 \end{aligned}$$

and the addition of neutrosophic probabilities as:

$$\begin{aligned}
 (a_1, b_1, c_1) + (a_2, b_2, c_2) &= (a_1 + \\
 a_2, \min\{b_1, b_2\}, \min\{c_1, c_2\}). \tag{210}
 \end{aligned}$$

In case when the neutrosophic probability is reduced to classical probability (i.e. $b_1 = b_2 = c_1 = c_2 = 0$), we get the same result for neutrosophical probability matrix NP^n as for the classical probability matrix P^n .

Other multiplication and addition operators of neutrosophic probabilities can be defined as well. For example:

$$(a_1, b_1, c_1) \cdot (a_2, b_2, c_2) = (a_1 a_2, \min\{b_1, b_2\}, \max\{c_1, c_2\}), \quad (211)$$

or

$$(a_1, b_1, c_1) \cdot (a_2, b_2, c_2) = \left(a_1 a_2, \frac{b_1 + b_2}{2}, \max\{c_1, c_2\}\right), \text{ etc.} \quad (212)$$

and similarly for the addition of neutrosophic probabilities changing the middle component to $\max\{b_1, b_2\}$, or average $\frac{b_1 + b_2}{2}$, etc.

$$\text{Let's note } (NP)^2 = \{c_{ij}\}_{i,j}; \quad (213)$$

$$\begin{aligned} c_{11} = & [(0.40, 0.10, 0.00)(0.20, 0.00, 0.10)(0.10, 0.05, 0.05)] \cdot \\ & \left[\begin{array}{l} (0.40, 0.10, 0.00) \\ (0.19, 0.03, 0.08) \\ (0.07, 0.03, 0.10) \end{array} \right] = \begin{array}{l} (0.40, 0.10, 0.00) \\ P \rightarrow P \end{array} . \\ & \begin{array}{l} (0.40, 0.10, 0.00) \\ P \rightarrow P \end{array} + \\ & \begin{array}{l} (0.20, 0.00, 0.10) \\ P \rightarrow R \end{array} \begin{array}{l} (0.19, 0.03, 0.08) \\ R \rightarrow P \end{array} + \\ & \begin{array}{l} (0.10, 0.05, 0.05) \\ P \rightarrow D \end{array} \begin{array}{l} (0.07, 0.03, 0.10) \\ D \rightarrow P \end{array} = \end{aligned}$$

$$(0.16, 0.10, 0.00) + (0.038, 0.03, 0.10) + (0.007, 0.05, 0.10) = (0.205, 0.050, 0.000). \quad (214)$$

c_{11} means the neutrosophic probability of having an economic prosperity year (P), after 2 years, starting from prosperity:

$$[(P \rightarrow P) \wedge (P \rightarrow P)] \text{ or } [(P \rightarrow R) \wedge (R \rightarrow P)] \text{ or } [(P \rightarrow D) \wedge (D \rightarrow P)], \quad (215)$$

where, for example, $P \rightarrow R$ means “from prosperity to recession”, and so on.

$$\begin{aligned} c_{12} &= (0.080, 0.10, 0.10) + (0.070, 0.05, 0.10) \\ &\quad + (0.027, 0.09, 0.05) \\ &= (0.277, 0.050, 0.050) \end{aligned}$$

$$\begin{aligned} c_{12} &= (0.004, 0.10, 0.05) + (0.048, 0.02, 0.10) \\ &\quad + (0.032, 0.06, 0.05) \\ &= (0.084, 0.020, 0.050) \end{aligned}$$

$$\begin{aligned} c_{21} &= (0.076, 0.10, 0.08) + (0.0665, 0.05, 0.08) \\ &\quad + (0.0168, 0.03, 0.10) \\ &= (0.1593, 0.003, 0.080) \end{aligned}$$

$$\begin{aligned} c_{22} &= (0.038, 0.03, 0.10) + (0.1225, 0.05, 0.00) \\ &\quad + (0.0648, 0.09, 0.004) \\ &= (0.253, 0.003, 0.000) \end{aligned}$$

$$\begin{aligned}
c_{23} &= (0.019, 0.05, 0.08) + (0.0840, 0.05, 0.04) \\
&\quad + (0.0768, 0.06, 0.04) \\
&= (0.1798, 0.005, 0.004)
\end{aligned}$$

$$\begin{aligned}
c_{31} &= (0.028, 0.100, 0.100) \\
&\quad + (0.0513, 0.09, 0.08) \\
&\quad + (0.024, 0.060, 0.100) \\
&= (0.1017, 0.06, 0.08)
\end{aligned}$$

$$\begin{aligned}
c_{32} &= (0.014, 0.03, 0.10) + (0.0945, 0.09, 0.04) \\
&\quad + (0.0864, 0.09, 0.04) \\
&= (0.1949, 0.03, 0.04)
\end{aligned}$$

$$\begin{aligned}
c_{33} &= (0.007, 0.05, 0.10) + (0.0648, 0.09, 0.04) \\
&\quad + (0.1024, 0.06, 0.02) \\
&= (0.1742, 0.05, 0.02).
\end{aligned}$$

Thus

$$\begin{aligned}
&(NP)^2 \\
&= \begin{bmatrix} (0.205, 0.05, 0.0) & (0.277, 0.05, 0.05) & (0.084, 0.02, 0.05) \\ (0.1593, 0.03, 0.08) & (0.2253, 0.03, 0.0) & (0.1798, 0.05, 0.04) \\ (0.1017, 0.06, 0.08) & (0.1949, 0.03, 0.04) & (0.1742, 0.05, 0.02) \end{bmatrix}.
\end{aligned}
\tag{216}$$

We normalize the rows by dividing each of the nine components by their sum. We get with three decimal approximation:

$$\begin{aligned}
& (NP)_{norm}^2 \\
& = \begin{bmatrix} (0.261, 0.064, 0.000) & (0.352, 0.064, 0.064) & (0.106, 0.025, 0.064) \\ (0.201, 0.038, 0.101) & (0.284, 0.038, 0.000) & (0.226, 0.062, 0.050) \\ (0.135, 0.080, 0.107) & (0.260, 0.040, 0.053) & (0.232, 0.066, 0.027) \end{bmatrix}.
\end{aligned}
\tag{217}$$

According to this *neutrosophic transition matrix*, after two years the largest chance of the economy to be is in the state of recession.

3.62. Applications of Neutrosophics.

Once could use the neutrosophics in statistical physics, financial markets, risk management, mathematical biology, quantum theory, and in almost any humanistic or scientific field where indeterminacy, unknown, and in general where <neutA> (neutrality with respect to an item <A>) occur.

Chapter 4.
Neutrosophic Subjects for Future Research

Neutrosophic Subjects

1. Neutrosophic topologies, including neutrosophic metric spaces and smooth topological spaces.
2. Neutrosophic numbers ($a+bI$, where $I =$ indeterminacy and $I^2 = I$, $mI+nI = (m+n)I$, $0I = 0$, and a, b are real or complex numbers), and arithmetical operations, including ranking procedures for neutrosophic numbers.
3. Neutrosophic rough sets.
4. Neutrosophic relational structures, including neutrosophic relational equations, neutrosophic similarity relations, and neutrosophic ordering.
5. Neutrosophic geometry (Smarandache geometries).
6. Neutrosophic probability.
7. Neutrosophic logical operations, including n-norms, n-conorms, neutrosophic implicators, neutrosophic quantifiers.
8. Measures of neutrosophication.
9. Deneutrosophication techniques.
10. Neutrosophic multivalued mappings.
 11. Develop the neutrosophic measure (defined in this book).
 12. Develop the neutrosophic integral (defined in this book).
13. Neutrosophic differential calculus.
14. Neutrosophic mathematical morphology.
15. Neutrosophic algebraic structures.
16. Neutrosophic models.
17. Neutrosophic cognitive maps.

18. Neutrosophic relational maps.
19. Neutrosophic matrix, bimatrix, ..., n -matrix.
20. Neutrosophic graph, which is a graph that has at least one indeterminate edge or one indeterminate node.
21. Neutrosophic tree, which is a tree that has at least one indeterminate edge or one indeterminate node.
22. Neutrosophic fusion rules for information fusion.
23. Applications: neutrosophic relational databases, neutrosophic image (thresholding, denoising, segmentation) processing, neutrosophic linguistic variables, neutrosophic decision making and preference structures, neutrosophic expert systems, neutrosophic reliability theory, neutrosophic soft computing techniques in e-commerce and e-learning, image segmentation, etc.

References

1. Florentin Smarandache, *An Introduction to Neutrosophic Probability Applied in Quantum Physics*, SAO/NASA ADS Physics Abstract Service, <http://adsabs.harvard.edu/abs/2009APS..APR.E1078S>
2. Florentin Smarandache, *An Introduction to Neutrosophic Probability Applied in Quantum Physics*, Bulletin of the American Physical Society, 2009 APS April Meeting, Volume 54, Number 4, Saturday–Tuesday, May 2–5, 2009; Denver, Colorado, USA, <http://meetings.aps.org/Meeting/APR09/Event/102001>
3. Florentin Smarandache, *An Introduction to Neutrosophic Probability Applied in Quantum Physics*, Bulletin of Pure and Applied Sciences, Physics, 13-25, Vol. 22D, No. 1, 2003.
4. Florentin Smarandache, "Neutrosophy. / Neutrosophic Probability, Set, and Logic", American Research Press, Rehoboth, USA, 105 p., 1998.
5. Florentin Smarandache, *A Unifying Field in Logics: Neutrosophic Logic, Neutrosophic Set, Neutrosophic Probability and Statistics*, 1999, 2000, 2003, 2005.
6. Florentin Smarandache, *Neutrosophic Physics as a new field of research*, Bulletin of the American Physical Society, APS March Meeting 2012, Volume 57, Number 1, Monday–Friday, February 27–March 2

2012; Boston, Massachusetts,
<http://meetings.aps.org/Meeting/MAR12/Event/160317>

7. Florentin Smarandache, V. Christianto, A Neutrosophic Logic View to Schrodinger's Cat Paradox, Bulletin of the American Physical Society 2008 Joint Fall Meeting of the Texas and Four Corners Sections of APS, AAPT, and Zones 13 and 16 of SPS, and the Societies of Hispanic & Black Physicists Volume 53, Number 11. Friday–Saturday, October 17–18, 2008; El Paso, Texas,
<http://meetings.aps.org/link/BAPS.2008.TS4CF.E4.8>
8. Florentin Smarandache, Vic Christianto, The Neutrosophic Logic View to Schrodinger Cat Paradox, Revisited, Bulletin of the American Physical Society APS March Meeting 2010 Volume 55, Number 2. Monday–Friday, March 15–19, 2010; Portland, Oregon,
<http://meetings.aps.org/link/BAPS.2010.MAR.S1.15>
9. Florentin Smarandache, Neutrosophic Degree of Paradoxicity of a Scientific Statement, Bulletin of the American Physical Society 2011 Annual Meeting of the Four Corners Section of the APS Volume 56, Number 11. Friday–Saturday, October 21–22, 2011; Tuscon, Arizona,
<http://meetings.aps.org/link/BAPS.2011.4CF.F1.37>
10. Florentin Smarandache, n-Valued Refined Neutrosophic Logic and Its Applications to Physics, Bulletin of the American Physical Society 2013 Annual Fall Meeting of the APS Ohio-Region Section Volume 58, Number 9. Friday–Saturday, October 4–5, 2013; Cincinnati, Ohio,

<http://meetings.aps.org/Meeting/OSF13/Event/20564>

1

11. Florentin Smarandache, Neutrosophic Diagram and Classes of Neutrosophic Paradoxes, or To the Outer-Limits of Science, Bulletin of the American Physical Society, 17th Biennial International Conference of the APS Topical Group on Shock Compression of Condensed Matter Volume 56, Number 6. Sunday–Friday, June 26–July 1 2011; Chicago, Illinois, <http://meetings.aps.org/link/BAPS.2011.SHOCK.F1.1>
67

ADDENDA

Books on Neutrosophics

1. *Fuzzy Neutrosophic Models for Social Scientists*, by W. B. Vasantha Kandasamy, Florentin Smarandache, Education Publisher, Columbus, OH, 167 pp., 2013.
2. *Neutrosophic Super Matrices and Quasi Super Matrices*, by W. B. Vasantha Kandasamy, Florentin Smarandache, 200 p., Educational Publisher, Columbus, 2012.
3. *Neutrosafia ca reflectarea a realității neconvenționale*, de Florentin Smarandache, Tudor Păroiu, Ed. Sitech, Craiova, Romania, 130 p., 2012.
4. *Neutrosophic Interpretation of The Analects of Confucius (弗羅仁汀·司馬仁達齊, 傅昱華 論語的中智學解讀和擴充—正反及中智論語)*, *English-Chinese Bilingual (英漢雙語)*, by Florentin Smarandache, Fu Yuhua, Zip Publisher, Columbus, 268 p., 2011.
5. *Neutrosophic Interval Bialgebraic Structures*, by W. B. Vasantha Kandasamy, Florentin Smarandache, Zip Publishing, Columbus, 195 p., 2011.
6. *Finite Neutrosophic Complex Numbers*, by W. B. Vasantha Kandasamy, Florentin Smarandache, Zip Publisher, Columbus, Ohio, USA, 220 p., 2011.
7. *Neutrosophic Interpretation of Tao Te Ching (English-Chinese bilingual)*, by Florentin Smarandache & Fu Yuhua, Translation by Fu Yuhua, Chinese Branch Kappa, Beijing, 208 p., 2011.
8. *Neutrosophic Bilinear Algebras and Their Generalization*, by W.B. Vasantha Kandasamy, Florentin Smarandache, Svenska Fysikarkivet, Stockholm, Sweden, 402 p., 2010.
9. *Multispace & Multistructure. Neutrosophic Transdisciplinarity (100 Collected Papers of Sciences)*, Vol. IV, by Florentin Smarandache, North-European Scientific

- Publishers, Hanko, Finland, 800 p., 2010.
10. New Classes of Neutrosophic Linear Algebras, by W.B. Vasantha Kandasamy, F. Smarandache, K. Ilanthenral, CuArt, Slatina, 286 p., 2010.
 11. Neutrosophic Physics: More Problems, More Solutions (Collected Papers), editor Florentin Smarandache, North-European Scientific Publishers, Hanko, Finland, 94 p., 2010.
 12. Neutrosophic Logic, Wave Mechanics, and Other Stories (Selected Works: 2005-2008), by F. Smarandache, V. Christianto, Kogaion Editions, Bucharest, 129 p., 2009.
 13. Chinese Neutrosophy and Taoist Natural Philosophy [Chinese language], by F. Smarandache and Jiang Zhengjie, Xiquan Chinese Hse., Beijing, 150 p., 2008.
 14. Neutrality and Multi-Valued Logics, A R Press, Rehoboth, 119 p., 2007, by Andrew Schumann, Florentin Smarandache.
 15. Neutrosophy in Arabic Philosophy [English version], Renaissance High Press (Ann Arbor), 291 pp., 2007, by F. Smarandache & Salah Osman;
 - Translated into Arabic language by Dr. Osman Salah, 418 pp., *Munsha't al-Ma'arif* Publ. Hse., Jalal Huzie & Partners, Alexandria, Egypt, 2007.
 16. Multi-Valued Logic, Neutrosophy, and Schrödinger Equation, 107 p., Hexis, 2006, by F. Smarandache & V. Christianto.
 17. Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures, 219 p., Hexis, 2006, by W. B. Vasantha Kandasamy and F. Smarandache.
 18. Introduction to N-Adaptive Fuzzy Models to Analyze Public Opinion on AIDS, 235 p., Hexis, 2006, by W. B. Vasantha Kandasamy and F. Smarandache.
 19. Neutrosophic Rings, 154 p., Hexis, 2006, by W. B. Vasantha Kandasamy & F. Smarandache.
 20. Fuzzy Interval Matrices, neutrosophic Interval Matrices and Their Applications, 304 p., Hexis, 2006, by W. B. Vasantha Kandasamy & F. Smarandache.

21. Vedic Mathematics, 'Vedic' or 'Mathematics': A Fuzzy & Neutrosophic Analysis, Automaton, Los Angeles, 220 p., 2006, by W. B. Vasantha Kandasamy and F. Smarandache.
22. Neutrosophic Methods in General Relativity, 78 p., Hexis, 2005, by D. Rabounski, F. Smarandache, L. Borissova).
 - Russian translation by D. Rabounski, Нейтрософские методы в Общей Теории Относительности, Hexis, 105 p., 2006.
23. Interval Neutrosophic Sets and Logic: Theory and Applications in Computing, Hexis, 87 p., 2005, by H. Wang, F. Smarandache, Y.-Q. Zhang, R. Sunderraman.
24. Introduction to Bimatrices, Hexis, 181 p., 2005, by W. B. Vasantha Kandasamy, F. Smarandache, K. Ilanthenral.
25. Fuzzy and Neutrosophic Analysis of Women with HIV / AIDS (With Specific Reference to Rural Tamil Nadu in India), Hexis, 316 p., 2005; by W. B. Vasantha Kandasamy, F. Smarandache; translation of the Tamil interviews by Meena Kandasamy).
26. Applications of Bimatrices to some Fuzzy and Neutrosophic Models, Hexis, 273 pp., 2005, by W. B. Vasantha Kandasamy, F. Smarandache, K. Ilanthenral.
27. Analysis of Social Aspect of Migrant Labourers Living with HIV/AIDS Using Fuzzy Theory and Neutrosophic Cognitive Maps / With specific reference to Rural Tamilnadu in India, Xiquan, Phoenix, 471 p., 2004, by W. B. Vasantha Kandasamy, F. Smarandache.
28. Neutrosophic Dialogues, Xiquan, Phoenix, 97 p., 2004, by F. Smarandache and Feng Liu.
29. Fuzzy Relational Equations & Neutrosophic Relational Equations, Hexis, 301 pp., 2004, by W. B. Vasantha Kandasamy, F. Smarandache.
30. Basic Neutrosophic Algebraic Structures and their Applications to Fuzzy and Neutrosophic Models, Hexis, 149 pp., 2004, by W. B. Vasantha Kandasamy, F. Smarandache.
31. Fuzzy Cognitive Maps and Neutrosophic Cognitive Maps, Xiquan, Phoenix, 211 p., 2003, W. B. Vasantha Kandasamy,

F. Smarandache.

32. Proceedings of the First International Conference on Neutrosophy, Neutrosophic Logic, Neutrosophic Set, Neutrosophic Probability and Statistics, editor F. Smarandache, University of New Mexico, Gallup Campus, Xiquan, Phoenix, 147 pp., 2002.
33. Collected Papers, Vol. III, by F. Smarandache, Abaddaba Publ. Hse., Oradea, 160 p., 2000.
34. Neutrosophy. / Neutrosophic Probability, Set, and Logic, by F. Smarandache, Amer. Res. Press, Rehoboth, USA, 105 p., 1998;
 - Republished in 2000, 2003, 2005, A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability and Statistics (fourth edition), American Research Press, 156 p.;
 - Chinese translation by F. Liu, “A Unifying Field in Logics: Neutrosophic Logic. / Neutrosophy, Neutrosophic Set, Neutrosophic Probability and statistics”, Xiquan Chinese Branch, 121 p., 2003;
 - Russian partial translation by D. Rabounski: Hexis, **Сущность нейтрософии**, 32 p., 2006.

More Articles on Neutrosophics

1. N-norm and N-conorm in Neutrosophic Logic and Set, and the Neutrosophic Topologies (2005), by F. Smarandache, in Critical Review, Creighton University, Vol. III, 73-83, 2009.
2. n-ary Fuzzy Logic and Neutrosophic Logic Operators, by F. Smarandache, V. Christianto, in arXiv.org, and in <Studies in Logic Grammar and Rhetoric>, Belarus, 17 (30), 1-16, 2009.
3. Neutrosophic Logic and Set, and Paradoxes chapters by F. Smarandache, V. Christianto, F. Liu, Haibin Wang, Yanqing Zhang, Rajshekhar Sunderraman, André Rogatko, Andrew

Schumann, in *Multispace & Multistructure. Neutrosophic Transdisciplinarity*, NESP, Finland, pp. 395-548 and respectively 604-631, 2010.

4. The Fifth Function of University: “Neutrosophic E-function” of Communication-Collaboration-Integration of University in the Information Age, by F. Smarandache, S. Vlăduțescu.
5. The Neutrosophic Research Method in Scientific and Humanistic Fields, by Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 732-733, 2010.
6. Single Valued Neutrosophic Sets, by Haibin Wang, Florentin Smarandache, Yanqing Zhang, Rajshekhar Sunderraman, in *Multispace and Multistructure*, Vol. 4, 410-413, 2010.
7. Neutrosophic Soft Set, by Pabitra Kumar Maji, *Annals of Fuzzy Mathematics and Informatics*, Vol. 5, No. 1, 157-168, January 2013.
8. A Neutrosophic Soft Set Approach to A Decision Making Problem, by Pabitra Kumar Maji, *Annals of Fuzzy Mathematics and Informatics*, Vol. 3, No. 2, 313-319, April 2012.
9. Correlation Coefficients of Neutrosophic Sets by Centroid Method, by I. M. Hanafy, A. A. Salama, K. M. Mahfouz, *International Journal of Probability and Statistics* 2013, 2(1): 9-12.
10. Análisis de textos de José Martí utilizando mapas cognitivos neutrosóficos, por Maikel Leyva-Vazquez, Karina Perez-Teruel, F. Smarandache, 2013, <http://vixra.org/abs/1303.0216>
11. Correlation of Neutrosophic Data, by I. M. Hanafy, A.A.Salama and K. Mahfouz, *International Refereed Journal of Engineering and Science (IRJES)*, Vol. 1, Issue 2, 39-43, 2012.
12. Neutrosophic Filters, by A. A. Salama & H. Alagamy, *International Journal of Computer Science Engineering and Information Technology Research (IJCSEITR)*, Vol. 3, Issue 1, Mar 2013, 307-312.
13. Neutrosophic Masses & Indeterminate Models. Applications to Information Fusion, by Florentin Smarandache, *Proceedings*

of the 15th International Conference on Information Fusion, Singapore, 9-12 July 2012.

14. A Geometric Interpretation of the Neutrosophic Set – A Generalization of the Intuitionistic Fuzzy Set, 2011 IEEE International Conference on Granular Computing, edited by Tzung-Pei Hong, Yasuo Kudo, Mineichi Kudo, Tsau-Young Lin, Been-Chian Chien, Shyue-Liang Wang, Masahiro Inuiguchi, GuiLong Liu, IEEE Computer Society, National University of Kaohsiung, Taiwan, 602-606, 8-10 November 2011.
15. Applications of Neutrosophic Logic to Robotics / An Introduction, by Florentin Smarandache, Luige Vladareanu, 2011 IEEE International Conference on Granular Computing, edited by Tzung-Pei Hong, Yasuo Kudo, Mineichi Kudo, Tsau-Young Lin, Been-Chian Chien, Shyue-Liang Wang, Masahiro Inuiguchi, GuiLong Liu, IEEE Computer Society, National University of Kaohsiung, Taiwan, 607-612, 8-10 November 2011.
16. Color Image Segmentation Based on Neutrosophic Method, by Ling Zhang, Ming Zhang, H. D. Cheng, *Optical Engineering*, 51(3), 037009, 2012.
17. Intuitionistic Neutrosophic Soft Set, by Said Broumi, F. Smarandache, *Journal of Information and Computing Science*, Vol. 8, No. 2, 2013, *pp.* 130-140.
18. A Novel Neutrosophic Logic SVM (N-SVM) and its Application to Image Categorization, by Wen Ju and H. D. Cheng, *New Mathematics and Natural Computation (World Scientific)*, Vol. 9, No. 1, 27-42, 2013.
19. Activism and Nations Building in Pervasive Social Computing Using Neutrosophic Cognitive Maps (NCMs), by A. Victor Devadoss, M. Clement Joe Anand, *International Journal of Computing Algorithm*, Volume: 02, Pages: 257-262, October 2013.
20. A Study of Quality in Primary Education Combined Disjoint Block Neutrosophic Cognitive Maps (CDBNCM), by A. Victor Devadoss, M. Clement Joe Anand, A. Joseph Bellarmin, Indo-

Bhutan International Conference On Gross National Happiness
Vol. 02, Pages: 256-261, October 2013.

21. Segmentation of Breast Ultrasound Images Based on Neutrosophic Method, by Ming Zhang, Ling Zhang, H. D. Cheng, *Optical Engineering*, 49(11), 117001-117012, 2010.
22. A Neutrosophic Approach to Image Segmentation Based on Watershed Approach, by Ming Zhang, Ling Zhang, H. D. Cheng, *Signal Processing*, 90(5), 1510-1517, 2010.
23. "On neutrosophic paraconsistent topology", by F .G. Lupiáñez, *Kybernetes* 39 (2010), 598-601.
24. Strategy on T, I, F Operators. A Kernel Infrastructure in Neutrosophic Logic, by Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 414-419, 2010.
25. On Similarity and Entropy of Neutrosophic Sets, by Pinaki Majumdar & S. K. Samanta. M.U.C Women College, Burdwan (W. B.), India, 2013.
26. An Effective Neutrosophic Set-Based Preprocessing Method for Face Recognition, by Mohammad Reza Faraji and Xiaojun Qi, Utah State University, Logan, 2013.
27. Toward Dialectic Matter Element of Extenics Model, by Liu Feng, Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 420-429, 2010.
28. Self-Knowledge and Knowledge Communication, by Liu Feng and Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 430-435, 2010.
29. A Neutrosophic Description Logic, by Haibin Wang, Andre Rogatko, Florentin Smarandache, Rajshekhar Sunderraman, Proceedings of 2006 IEEE International Conference on Granular Computing, edited by Yan-Qing Zhang and Tsau Young Lin, Georgia State University, Atlanta, 305-308, 2006.
30. Neutrosophic Relational Data Model, by Haibin Wang, Rajshekhar Sunderraman, Florentin Smarandache, André Rogatko, in <Critical Review> (Society for Mathematics of Uncertainty, Creighton University), Vol. II, 19-35, 2008.

31. Short Definitions of Neutrosophic Notions [in Russian], by F. Smarandache, translated by A. Schumann, Philosophical Lexicon, Minsk-Moscow, Econompres, Belarus-Russia, 2008.
32. Neutrosophic Logic Based Semantic Web Services Agent, by Haibin Wang, Yan-Qing Zhang, Rajshekhar Sunderraman, Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 505-519, 2010.
33. Neutrosophic Transdisciplinarity (Multi-Space & Multi-Structure), by Florentin Smarandache, Arhivele Statului, Filiala Vâlcea, Rm. Vâlcea, 1969; presented at Scoala de Vara Internationala, Interdisciplinara si Academica, Romanian Academy, Bucharest, 6-10 July 2009.
34. Neutrosophic Logic as a Theory of Everything in Logics, by Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 525-527, 2010.
35. Blogs on Applications of Neutrosophics and Multispace in Sciences, by Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 528-548, 2010.
36. A Neutrosophic Multicriteria Decision Making Method, by Athar Kharal, National University of Science and Technology, Islamabad, Pakistan.
37. A multicriteria decision-making method using aggregation operators for simplified neutrosophic sets, by J. Ye, *Journal of Intelligent and Fuzzy Systems* (2013) doi: 10.3233/IFS-130916.
38. Single valued neutrosophic cross-entropy for multicriteria decision making problems, by Jun Ye, *Applied Mathematical Modelling* (2013) doi: 10.1016/j.apm.2013.07.020.
39. Multicriteria decision-making method using the correlation coefficient under single-valued neutrosophic environment, by Jun Ye, *International Journal of General Systems*, Vol. 42, No. 4, 386-394, 2013.
40. Similarity Measures between Interval Neutrosophic Sets and their Multicriteria Decision Method, by Jun Ye, Shaoxing Univ., China.

41. Neutrosophic Degree of a Paradoxicity, by Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 605-607, 2010.
42. Neutrosophic Diagram and Classes of Neutrosophic Paradoxes, or To The Outer-Limits of Science, by Florentin Smarandache, *Prog. Physics*, Vol. 4, 18-23, 2010.
43. S-denying a Theory, by Florentin Smarandache, in *Multispace and Multistructure*, Vol. 4, 622-629, 2010.
44. Five Paradoxes and a General Question on Time Traveling, by Florentin Smarandache, *Prog. Physics*, Vol. 4, 24, 2010.
45. H. D. Cheng, Yanhui Guo and Yingtao Zhang, *A Novel Image Segmentation Approach Based on Neutrosophic Set and Improved Fuzzy C-means Algorithm*, *New Mathematics and Natural Computation*, Vol. 7, No. 1 (2011) 155-171.
46. Degree of Negation of an Axiom, by F. Smarandache, to appear in the *Journal of Approximate Reasoning*, arXiv:0905.0719.
47. *Remedy for Effective Cure of Diseases using Combined Neutrosophic Relational Maps*, by M R Bivin, N.Saivaraju and K S Ravichandran, *International Journal of Computer Applications*, 12(12):18?23, January 2011. Published by Foundation of Computer Science.
48. Neutrosophic Research Method, by F. Smarandache, in *Multispace & Multistructure. Neutrosophic Transdisciplinarity*, NESP, Finland, pp. 395-548 and respectively 732-733, 2010.
49. *A Tool for Qualitative Causal Reasoning On Complex Systems*, by Tahar Guerram, Ramdane Maamri, and Zaidi Sahnoun, *IJCSI International Journal of Computer Science Issues*, Vol. 7, Issue 6, November 2010.
50. *A Study on Suicide problem using Combined Overlap Block Neutrosophic Cognitive Maps*, by P. Thiruppathi, N.Saivaraju, K.S. Ravichandran, *International Journal of Algorithms, Computing and Mathematics*, Vol. 3, Number 4, November 2010.

51. *On various neutrosophic topologies*, Francisco Gallego Lupiáñez, “Recent advances in Fuzzy Systems”, WSEAS (Athens, 2009), 59-62.
52. *Interval neutrosophic sets and Topology*, by F .G. Lupiáñez, *Kybernetes* 38 (2009), 621-624.
53. *On various neutrosophic topologies*, by F .G. Lupiáñez, *Kybernetes* 38 (2009), 1009-1013.
54. *Interval neutrosophic sets and topology*, Francisco Gallego Lupiáñez, *Kybernetes: The International Journal of Systems & Cybernetics*, Volume 38, Numbers 3-4, 2009 , pp. 621-624(4).
55. *Neutrosophic logics on Non-Archimedean Structures*, by Andrew Schumann, Critical Review, Creighton University, USA, Vol. III, 36-58, 2009.
56. *Positive, Negative and Neutral Law of Universal Gravitation*, by Fu Yuhua, Fu Anjie, Zhao Ge, *New Science and Technology*, 2009 (12), 30-32.
57. *Intuitionistic Neutrosophic Set*, by Monoranjan Bhowmik and Madhumangal Pal, *Journal of Information and Computing Science*, England, Vol. 4, No. 2, 2009, pp. 142-152.
58. *Discrimination of Outer Membrane Proteins using Reformulated Support Vector Machine based on Neutrosophic Set*, by Wen Ju and H. D. Cheng, *Proceedings of the 11th Joint Conference on Information Sciences (2008)*, Published by Atlantis Press.
59. *Adaptive fuzzy cognitive maps vs neutrosophic cognitive maps: decision support tool for knowledge based institution*, by Goutam Bernajee, *Journal of Scientific and Industrial Research*, 665-673, Vol. 67, 2008.
60. *A Method of Imprecise Query Solving*, by Smita Rajpal, M.N. Doja, Ranjit Biswas, *International Journal of Computer Science and Network Security*, Vol. 8 No. 6, pp. 133-139, June 2008, http://paper.ijcsns.org/07_book/200806/20080618.pdf
61. *On Neutrosophic Topology*, by F .G. Lupiáñez, *Kybernetes* 37 (2008), 797-800.

62. *Interval neutrosophic sets and Topology*, by F .G. Lupiáñez, “*Applied and Computational Mathematics*”, WSEAS (Athens , 2008), 110-112.
63. *A Method of Neutrosophic Logic to Answer Queries in Relational Database*, by Smita Rajpal, M.N. Doja and Ranjit Biswas, *Journal of Computer Science* 4 (4): 309-314, 2008.
64. *Ensemble Neural Networks Using Interval Neutrosophic Sets and Bagging*, by Pawalai Kraipeerapun, Chun Che Fung, Kok Wai Wong, Third International Conference on Natural Computation (ICNC 2007), Haikou, Hainan, China, August 24-August 27, 2007.
65. *Lithofacies Classification from Well Log Data using Neural Networks, Interval Neutrosophic Sets and Quantification of Uncertainty*, by Pawalai Kraipeerapun, Chun Che Fung, and Kok Wai Wong, *World Academy of Science, Engineering and Technology*, 23, 2006.
66. *Redesigning Decision Matrix Method with an indeterminacy-based inference process*, by Jose L. Salmeron, Florentin Smarandache, *Advances in Fuzzy Sets and Systems*, Vol. 1(2), 263-271, 2006.
67. *Neural network ensembles using interval neutrosophic sets and bagging for mineral prospectivity prediction and quantification of uncertainty*, by P. Kraipeerapun, C. C. Fung, W. Brown and K. W. Wong, 2006 IEEE Conference on Cybernetics and Intelligent Systems, 7-9 June 2006, Bangkok, Thailand.
68. *Processing Uncertainty and Indeterminacy in Information Systems success mapping*, by Jose L. Salmeron, Florentin Smarandache, arXiv:cs/0512047v2.
69. *The Combination of Paradoxical, Uncertain, and Imprecise Sources of Information based on DSmT and Neutro-Fuzzy Inference*, by Florentin Smarandache, Jean Dezert; in arXiv:cs/0412091v1. A version of this paper published in *Proceedings of 10th International Conference on Fuzzy Theory*

and Technology (FT&T 2005), Salt Lake City, Utah, USA, July 21-26, 2005.

70. *A simplification of the neutrosophic sets. Neutrosophic logic and intuitionistic fuzzy sets*, by Kalin Georgiev, Conference proceedings, Notes on IFS, Volume 11 (2005) Number 2, pages 28-31; Presented at: 9th ICIFS, Sofia, Bulgaria, 7-8 May 2005.
71. *Book Review by Milan Mares: W. B. Vasantha Kandasamy and Florentin Smarandache, Fuzzy Cognitive Maps and Neutrosophic Cognitive Maps*, Kybernetika, Vol. 40 (2004), No. 1, [151]-15.
72. *Set-Theoretic Operators on Degenerated Neutrosophic Set*, by H. Wang, Y. Zhang, R. Sunderraman, F. Song, Georgia State University, Atlanta, 2004.
73. *Neutrosophy in situation analysis*, Anne-Laure Joussetme, Patrick Maupin, Proc. of Fusion 2004 Int. Conf. on Information Fusion, pp. 400-406, Stockholm, Sweden, June 28-July 1, 2004 (<http://www.fusion2004.org>).
74. *Preamble to Neutrosophic Logic*, by C. Lee, Multiple-Valued Logic / An International Journal, Vol. 8, No. 3, 285-296, June 2002.
75. *Neutrosophy, a New Branch of Philosophy*, by Florentin Smarandache, Multiple-Valued Logic / An International Journal, Vol. 8, No. 3, 297-384, June 2002.
76. *A Unifying Field in Logics: Neutrosophic Field*, by Florentin Smarandache, Multiple-Valued Logic / An International Journal, Vol. 8, No. 3, 385-438, June 2002.
77. *Open Questions to Neutrosophic Inferences*, by Jean Dezert, Multiple-Valued Logic / An International Journal, Vol. 8, No. 3, 439-472, June 2002.
78. *Logic: A Misleading Concept. A Contradiction Study toward Agent's Logic*, by Feng Liu, Florentin Smarandache, Proceedings of the First International Conference on Neutrosophy, Neutrosophic Logic, Neutrosophic Set, Neutrosophic Probability and Statistics, University of New Mexico, Gallup Campus, 2001.

79. *Six Neutral Fundamental Reactions Between Four Fundamental Reactions*, by Fu Yuhua, Fu Anjie, Zhao Ge, <http://wbabin.net/physics/yuhua2.pdf>.
80. *On Rugina's System of Thought*, by Florentin Smarandache, International Journal of Social Economics, Vol. 28, No. 8, 623-647, 2001.
81. *Intentionally and Unintentionally. On Both, A and Non-A, in Neutrosophy*, by Feng Liu, Florentin Smarandache, Presented to the First International Conference on Neutrosophy, Neutrosophic Logic, Set, and Probability, University of New Mexico, Gallup, December 1-3, 2001.
82. *Neutrosophic Transdisciplinarity*, by F. Smarandache, 1969.
83. Online English Dictionary, *Definition of Neutrosophy*.
84. *Deployment of neutrosophic technology to retrieve answer for queries posed in natural language*, by Arora, M. ; Biswas, R.; Computer Science and Information Technology (ICCSIT), 2010 3rd IEEE International Conference on, Vol. 3, DOI: 10.1109/ICCSIT.2010.5564125, 2010, 435 – 439.
85. *Neutrosophic modeling and control*, Aggarwal, S. ; Biswas, R. ; Ansari, A.Q. Computer and Communication Technology (ICCCT), 2010 International Conference on, DOI: 10.1109/ICCCT.2010.5640435, 2010, 718 – 723.
86. *Truth-value based interval neutrosophic sets*, Wang, H. ; Yan-Qing Zhang ; Sunderraman, R., Granular Computing, 2005 IEEE International Conference on, Vol. 1, DOI: 10.1109/GRC.2005.1547284, 2005, 274 – 277;
87. *A geometric interpretation of the neutrosophic set — A generalization of the intuitionistic fuzzy set*, Smarandache, F., Granular Computing (GrC), 2011 IEEE International Conference on, DOI: 10.1109/GRC.2011.6122665, 2011, 602 – 606.

88. *MRI denoising based on neutrosophic wiener filtering*, Mohan, J. ; Yanhui Guo ; Krishnaveni, V.; Jeganathan, K., Imaging Systems and Techniques (IST), 2012 IEEE International Conference on, DOI: 10.1109/IST.2012.6295518, 2012, 327 – 331.

89. *Applications of neutrosophic logic to robotics: An introduction*, Smarandache, F. ; Vladareanu, L., Granular Computing (GrC), 2011 IEEE International Conference on, DOI: 10.1109/GRC.2011.6122666, 2011, 607 – 612;

90. *A Neutrosophic approach of MRI denoising*, Mohan, J. ; Krishnaveni, V. ; Guo, Yanhui Image Information Processing (ICIIP), 2011 International Conference on, DOI: 10.1109/ICIIP.2011.6108880, 2011, 1 – 6.

91. *Neural Network Ensembles using Interval Neutrosophic Sets and Bagging for Mineral Prospectivity Prediction and Quantification of Uncertainty*, Kraipeerapun, P. ; Chun Che Fung ; Brown, W. ; Kok-Wai Wong, Cybernetics and Intelligent Systems, 2006 IEEE Conference on, DOI: 10.1109/ICCIS.2006.252249, 2006, 1 – 6;

92. *Neutrosophic masses & indeterminate models: Applications to information fusion*, Smarandache, F., Information Fusion (FUSION), 2012 15th International Conference on, 2012, 1051 – 1057.

93. *Red Teaming military intelligence - a new approach based on Neutrosophic Cognitive Mapping*, Rao, S.; Intelligent Systems and Knowledge Engineering (ISKE), 2010 International Conference on, DOI: 10.1109/ISKE.2010.5680765, 2010, 622 – 627.

94. *Neutrosophic masses & indeterminate models. Applications to information fusion*, Smarandache, F., Advanced Mechatronic

Systems (ICAMechS), 2012 International Conference on, 2012, 674 – 679.

95. *Validating the Neutrosophic approach of MRI denoising based on structural similarity*, Mohan, J. ; Krishnaveni, V. ; Guo, Yanhui; Image Processing (IPR 2012), IET Conference on, DOI: 10.1049/cp.2012.0419, 2012, 1 – 6.

96. *Ensemble Neural Networks Using Interval Neutrosophic Sets and Bagging*, Kraipeerapun, P. ; Chun Che Fung ; Kok Wai Wong; Natural Computation, 2007. ICNC 2007. Third International Conference on, Vol. 1, DOI: 10.1109/ICNC.2007.359, 2007, 386 – 390.

97. *Comparing performance of interval neutrosophic sets and neural networks with support vector machines for binary classification problems*, Kraipeerapun, P. ; Chun Che Fung, Digital Ecosystems and Technologies, 2008. DEST 2008. 2nd IEEE International Conference on, DOI: 10.1109/DEST.2008.4635138, 2008, 34 – 37.

98. *Quantification of Uncertainty in Mineral Prospectivity Prediction Using Neural Network Ensembles and Interval Neutrosophic Sets*, Kraipeerapun, P. ; Kok Wai Wong ; Chun Che Fung ; Brown, W.; Neural Networks, 2006. IJCNN '06. International Joint Conference on, DOI: 10.1109/IJCNN.2006.247262, 2006, 3034 – 3039.

99. *A neutrosophic description logic*, Haibin Wang ; Rogatko, A. ; Smarandache, F.; Sunderraman, R.; Granular Computing, 2006 IEEE International Conference on DOI: 10.1109/GRC.2006.1635801, 2006, 305 – 308.

100. *Neutrosophic information fusion applied to financial market*, Khoshnevisan, M. ; Bhattacharya, S.; Information Fusion, 2003.

Proceedings of the Sixth International Conference of
Vol. 2, DOI: 10.1109/ICIF.2003.177381, 2003, 1252 – 1257.

101. *Neutrosophic set - a generalization of the intuitionistic fuzzy set*,
Smarandache, F.
Granular Computing, 2006 IEEE International Conference on,
DOI: 10.1109/GRC.2006.1635754, 2006, 38 – 42.

102. *From Fuzzification to Neutrosophication: A Better Interface between Logic and Human Reasoning*, Aggarwal, S. ; Biswas, R. ; Ansari, A.Q. Emerging Trends in Engineering and Technology (ICETET), 2010 3rd International Conference on, DOI: 10.1109/ICETET.2010.26, 2010, 21 – 26.

103. *A novel image enhancement approach for Phalanx and Epiphyseal/metaphyseal segmentation based on hand radiographs*, Chih-Yen Chen ; Tai-Shan Liao ; Chi-Wen Hsieh ; Tzu-Chiang Liu ; Hung-Chun Chien; Instrumentation and Measurement Technology Conference (I2MTC), 2012 IEEE International, DOI: 10.1109/I2MTC.2012.6229651, 2012, 220–224.

104. *Quantification of Vagueness in Multiclass Classification Based on Multiple Binary Neural Networks*, Kraipeerapun, P. ; Chun Che Fung ; Kok Wai Wong
Machine Learning and Cybernetics, 2007 International Conference on, Vol. 1, DOI: 10.1109/ICMLC.2007.4370129, 2007 140 – 144.

105. *Automatic Tuning of MST Segmentation of Mammograms for Registration and Mass Detection Algorithms*, Bajger, M. ; Fei Ma ; Bottema, M.J.; Digital Image Computing: Techniques and Applications, 2009. DICTA '09. DOI: 10.1109/DICTA.2009.72, 2009. 400 – 407,

106. *Externalizing Tacit knowledge to discern unhealthy nuclear intentions of nation states*, Rao, S., Intelligent System and Knowledge Engineering, 2008. ISKE 2008. 3rd International Conference on, Vol. 1, DOI: 10.1109/ISKE.2008.4730959, 2008, 378 – 383.

107. *Vagueness, a multifacet concept - a case study on Ambrosia artemisiifolia predictive cartography*, Maupin, P. ; Joussemme, A.-L. Geoscience and Remote Sensing Symposium, 2004. IGARSS '04. Proceedings. 2004 IEEE International, Vol. 1, DOI: 10.1109/IGARSS.2004.1369036, 2004.

108. *Analysis of information fusion combining rules under the dsm theory using ESM inputs*, Djiknavorian, P. ; Grenier, D. ; Valin, P. ; Information Fusion, 2007 10th International Conference on, DOI: 10.1109/ICIF.2007.4408128, 2007, 1 – 8.

Seminars on Neutrosophics

1. *An Introduction to Information Fusion Level 1 and to Neutrosophic Logic/Set with Applications*, by F. Smarandache, ENSIETA (National Superior School of Engineers and the Study of Armament), Brest, France, 2 July 2010.
2. *An Introduction to Fusion Level 1 and to Neutrosophic Logic/Set with Applications*, by F. Smarandache at Air Force Research Laboratory, in Rome, NY, USA, July 29, 2009.
3. *An Introduction to Neutrosophic Logic in Arabic Philosophy*, by F. Smarandache & Salah Osman, Minufiyya University, Shebin Elkom, Egypt, 17 December 2007.

International Conferences on Neutrosophics

1. International Conference on Applications of Plausible, Paradoxical, and Neutrosophic Reasoning for Information Fusion, Cairns, Queensland, Australia, 8-11 July 2003.
2. First International Conference on Neutrosophy, Neutrosophic Logic, Set, Probability and Statistics, University of New Mexico, Gallup Campus, 1-3 December 2001.

Ph. D. Dissertations on Neutrosophics

1. Eng. Ionel Alexandru Gal, *Contributions to the Development of Hybrid Force-Position Control Strategies for Mobile Robots Control*, advisers Dr. Luige Vlădăreanu & Dr. Florentin Smarandache, Institute of Solid Mechanics, Romanian Academy, Bucharest, October 14, 2013.
2. Smita Rajpal, *Intelligent Searching Techniques to Answer Queries in RDBMS*, *Ph D Dissertation in progress*, under the supervision of Prof. M. N. Doja, Department of Computer Engineering Faculty of Engineering, Jamia Millia Islamia, New Delhi, India, 2011.
3. Ming Zhang, *Novel Approaches to Image Segmentation Based on Neutrosophic Logic*, *Ph D Dissertation*, Utah State University, Logan, Utah, USA, All Graduate Theses and Dissertations, Paper 795, <http://digitalcommons.usu.edu/etd/795>, 12-1-201, 2010.

4. Haibin Wang, *Study on Interval Neutrosophic Set and Logic*, Georgia State University, Atlanta, USA, 2005.
5. Sukanto Bhattacharya, *Utility, Rationality and Beyond - From Finance to Informational Finance* [using Neutrosophic Probability], Bond University, Queensland, Australia, 2004.



*In this book, we introduce for the first time the notions of **neutrosophic measure** and **neutrosophic integral**, and we develop the 1995 notion of **neutrosophic probability**. We present many practical examples.*

It is possible to define the neutrosophic measure and consequently the neutrosophic integral and neutrosophic probability in many ways, because there are various types of indeterminacies, depending on the problem we need to solve. Neutrosophics study the indeterminacy. Indeterminacy is different from randomness. It can be caused by physical space materials and type of construction, by items involved in the space, etc.

