

# Examples where the Conjunctive and Dempster's Rules are Insensitive

Florentin Smarandache  
Mathematics and Science Department  
University of New Mexico  
Gallup, NM 87301, USA  
smarand@unm.edu

Jean Dezert  
The French Aerospace Lab.  
ONERA/DTIM/EVF  
91761 Palaiseau Cedex, France  
jean.dezert@onera.fr

Valeri Kroumov  
Faculty of Engineering  
Okayama University of Science  
Okayama 700-0005, Japan  
val@ee.ous.ac.jp

**Abstract**—In this paper we present several counter-examples to the Conjunctive rule and to Dempster rule of combinations in information fusion.

**Keywords**— conjunctive rule, Dempster rule, DSMT, counter-examples to Conjunctive rule, counter-examples to Dempster rule, information fusion

## I. INTRODUCTION

In *Counter-Examples to Dempster's Rule of Combination* {Ch. 5 of *Advances and Applications to DSMT on Information Fusion*, Vol. I, pp. 105-121, 2004} [1], J. Dezert, F. Smarandache, and M. Khoshnevisan have presented several classes of fusion problems which could not be directly approached by the classical mathematical theory of evidence, also known as Dempster-Shafer Theory (DST), either because Shafer's model for the frame of discernment was impossible to obtain, or just because Dempster's rule of combination failed to provide coherent results (or no result at all). We have showed and discussed the potentiality of the DSMT combined with its classical (or hybrid) rule of combination to attack these infinite classes of fusion problems.

We have given general and concrete counter-examples for Bayesian and non-Bayesian cases.

In this article we construct new classes where both the conjunctive and Dempster's rule are insensitive.

## II. DEZERT-TCHAMOVA COUNTER-EXAMPLE

In [2], J. Dezert and A. Tchamova have introduced for the first time the following counter-example with some generalizations. This first type of example has then been discussed in details in [3,4] to question the validity of foundations of Dempster-Shafer Theory (DST). In the next sections of this short paper, we provide more counter-examples extending this idea. Let the frame of discernment  $\Theta = \{A, B, C\}$ , under Shafer's model (i.e. all intersections are empty), and  $m_1(\cdot)$  and  $m_2(\cdot)$  be two independent sources of information that give the below masses:

--	--	--	--	--

Focal Elements	A	C	$A \cup B$	$A \cup B \cup C$
$m_1$	a	0	1-a	0
$m_2$	0	1-b <sub>1</sub> -b <sub>2</sub>	b <sub>1</sub>	b <sub>2</sub>

Table 1

where the parameters  $a, b_1, b_2 \in [0,1]$ , and  $b_1+b_2 \leq 1$ .

Applying the conjunctive rule, in order to combine  $m_1 \oplus m_2 = m_{12}$ , one gets:

$$m_{12}(A) = a(b_1+b_2) \quad (1)$$

$$m_{12}(C) = 0 \quad (2)$$

$$m_{12}(A \cup B) = (1-a)(b_1+b_2) \quad (3)$$

$$m_{12}(A \cup B \cup C) = 0 \quad (4)$$

and the conflicting mass

$$m_{12}(\phi) = 1-b_1-b_2 = K_{12}. \quad (5)$$

After normalizing by dividing by  $1-K_{12} = b_1+b_2$  one gets Dempster's rule result  $m_{DS}(\cdot)$ :

$$m_{DS}(A) = \frac{m_{12}(A)}{1-K_{12}} = \frac{a(b_1+b_2)}{b_1+b_2} = a = m_1(A)$$

$$m_{DS}(A \cup B) = \frac{m_{12}(A \cup B)}{1-K_{12}} = \frac{(1-a)(b_1+b_2)}{b_1+b_2} = 1-a = m_1(A \cup B) \quad (6)$$

Counter-intuitively after combining two sources of information,  $m_1(\cdot)$  and  $m_2(\cdot)$ , with Dempster's rule, the result does not depend at all on  $m_2(\cdot)$ . Therefore Dempster's rule is insensitive to  $m_2(\cdot)$  no matter what the parameters  $a, b_1, b_2$  are equal to.



For example let  $\Theta = \{A, B, C\}$ , in Shafer's model. We show that the conflicts between sources are not correctly reflected by the conjunctive rule, and that a certain non-vacuous non-uniform source is ignored by Dempster's rule.

Let's consider the masses:

	A	B	C	$A \cup B \cup C$	
$m_1$	1	0	0	0	(the most specific mass)
$m_2$	1/3	1/3	1/3	0	(very unspecific mass)
$m_3$	0.6	0.4	0	0	(mass between the very unspecific and the most specific masses)
$m_0$	0.2	0.2	0.2	0.4	(not vacuous mass, not uniform mass)

Then the conflict  $K_{10} = 0.4$  between  $m_1(\cdot)$  and  $m_0(\cdot)$  is the same as the conflict  $K_{20}$  between  $m_2(\cdot)$  and  $m_0(\cdot)$ , and similarly the same as the conflict  $K_{30}$  between  $m_3(\cdot)$  and  $m_0(\cdot)$ , which is not normal, since  $m_1(\cdot)$  is the most specific mass while  $m_2(\cdot)$  is the most unspecific mass.

Let's check other thing combining two sources using Dempster's rule:

$$m_1 \oplus m_0 = m_1, m_2 \oplus m_0 = m_2, m_3 \oplus m_0 = m_3,$$

which is not normal.

In order to get the "normal behavior" we combine  $m_1(\cdot)$  and  $m_0(\cdot)$  with PCR5, and similarly for others:  $m_2(\cdot)$  combined with  $m_0(\cdot)$ , and  $m_3(\cdot)$  combined with  $m_0(\cdot)$ .

In order to know what should have been the "normal behavior" for the conflict (the initial conflict was  $K_{10} = 0.4$ ), let's make a small change to  $m_0(\cdot)$  as below:

	A	B	C	$A \cup B \cup C$	
$m_1$	1	0	0	0	(the most specific mass)
$m_2$	1/3	1/3	1/3	0	(very unspecific mass)
$m_3$	0.6	0.4	0	0	(mass between the very unspecific and the most specific masses)
$m_0$	0.3	0.2	0.1	0.4	(not vacuous mass, not uniform mass)

$$K_{10} = 0.30$$

$$K_{20} = 0.40$$

$$K_{30} = 0.34$$

Now, the conflicts are different.

## VIII. CONCLUSION

We showed in this paper that: first the conflict was the same, no matter what was one of the sources (and it is abnormal that a non-vacuous non-uniform source has no impact on the conflict), and second that the result using Dempster's rule is not all affected by a non-vacuous non-uniform source of information.

Normally, the most specific mass (*bba*) should dominate the fusion result.

Therefore, the conflicts between sources are not correctly reflected by the conjunctive rule, and certain non-vacuous non-uniform sources are ignored by Dempster's rule in the fusion process.

## REFERENCES

- [1] F. Smarandache, J. Dezert, "Advances and Applications of DSMT for Information Fusion," Vols. I, II, III, American Res. Press, Rehoboth, 2004, 2006, respectively 2009; <http://fs.gallup.unm.edu/DSmT.htm>.
- [2] J. Dezert, A. Tchamova, "On the behavior of Dempster's rule of combination," Presented at the spring school on Belief Functions Theory and Applications (BFTA), Autrans, France, 4–8 April 2011 (<http://hal.archives-ouvertes.fr/hal-00577983/>).
- [3] J. Dezert., P. Wang, A. Tchamova, "On The Validity of Dempster-Shafer Theory." Proceedings of the International Conference of Information Fusion, Singapore, July, 2012.
- [4] A. Tchamova., J. Dezert, "On the Behavior of Dempster's Rule of Combination and the Foundations of Dempster-Shafer Theory," (Best paper award), in Proc. of IEEE IS'2012, Sofia, Bulgaria, Sept. 6-8, 2012.