

Derivation of the Born rule from many-worlds interpretation and probability theory

[Home](#) > [Quantum mechanics](#) > [Derivation of Born rule of quantum mechanics](#)

2016/07/09
Published 2012/09/26
K. Sugiyama¹

The Born rule is a rule that a probability we observe a small particle like an electron is proportional to the square of the absolute value of the wave function. In this paper, we try to derive the Born rule from the many-worlds interpretation.




Figure 4-25: The number of elementary events is the square of the number of elementary states.

Many researchers have tried to derive the Born rule (also called Born's rule, Born's law, or probability interpretation) from Many-Worlds Interpretation (MWI). However, no one succeeds. Thus, the derivation of Born's rule had become an important issue for MWI. We try to derive Born's rule by introducing an elementary event of probability theory to the quantum theory as a new method.

We interpret the wave function as a manifold like a three-dimensional sphere, and interpret the absolute value of the wave function as the surface area of the manifold. We suppose that the manifold exists in the discrete space that has lattice points. We interpret a point on the surface of the manifold as a state that we

cannot divide any more, an elementary state. We draw an arrow from any point to any point. We interpret an arrow as an event that we cannot divide any more, an elementary event.

Probability is proportional to the number of elementary events, and the number of elementary events is the square of the number of elementary states. The number of elementary states is proportional to the surface area of the manifold, and the surface area of the manifold is the absolute value of the wave function. Therefore, the probability is proportional to the square of the absolute value of the wave function.

CONTENTS

1	Introduction.....	3
1.1	Subject.....	3
1.2	The importance of the subject.....	3
1.3	Past derivation method.....	3
1.4	New derivation method of this paper.....	4
2	Traditional method of deriving and the problem.....	4
2.1	The Born rule.....	4
2.2	Everett's many-worlds interpretation.....	4
2.2.1	The basis problem of many-worlds interpretation.....	5
2.2.2	The probability problem of many-worlds interpretation.....	6
3	Review of existing ideas.....	6
3.1	Universal Wave function of Wheeler and DeWitt.....	6
3.2	Barbour's many-particle wave function of the universe.....	6
3.3	Laplace's Probability Theory.....	7
3.4	Penrose's spin networks.....	8
3.5	Feynman's path integral.....	9
3.6	Dirac's quantum field theory.....	13
3.7	Kaluza-Klein theory.....	14
3.8	Euler's formula.....	14
3.9	Cauchy-Riemann equation.....	15
3.10	Hamilton's Quaternion.....	15
3.11	Cauchy-Riemann-Fueter equation.....	16
3.12	Cartan's differential form.....	17
4	A new method of deriving.....	19
4.1	Universe of Two-dimensional space-time.....	19
4.1.1	Closed path of Two-dimensional space-time.....	19
4.1.2	Introduction of the absolute value of wave function.....	20
4.1.3	Introduction of the phase of wave function.....	22
4.1.4	Hierarchical universe.....	24
4.1.5	Equations.....	24
4.2	Universe of four-dimensional space-time.....	26
4.2.1	Closed path of four-dimensional space-time.....	26
4.2.2	Introduction of the absolute value of wave function.....	27
4.2.3	Introduction of the phase of wave function.....	29
4.2.4	Hierarchical universe.....	31
4.2.5	Equations.....	31
4.3	Normal space.....	33
4.4	Wave space-time.....	35
4.5	Elementary event of many-worlds interpretation.....	36
4.6	Elementary state of many-worlds interpretation.....	37
4.7	Introduction of an elementary state to the quantum theory.....	38
4.8	Application of path integral to the field.....	39
4.9	Introduction of elementary event to the quantum theory.....	39

4.10	Derivation of Born's rule	41
5	Conclusion	43
6	Supplement	43
6.1	Supplement of the many-particle wave function	43
6.2	Supplement of the method of deriving Born's rule	44
6.3	Supplement of basis problem in many-worlds interpretation	45
6.4	Interpretation of time in many-worlds interpretation.....	46
6.5	Supplement of Long distance transition	46
6.6	The solution to the basis problem of MWI.....	47
7	Future Issues	47
8	Consideration of the future issues.....	48
8.1	Consideration of principles	48
8.1.1	Hierarchical principle.....	48
8.1.2	Event principle	48
8.2	Consideration of formulation for the quantum field theory.....	48
8.3	Consideration of discrete space	49
8.3.1	Discrete space from 24-hypercube.....	49
8.3.2	Discrete space Consideration from finite group	50
8.3.3	Consideration of the uncertain status	50
8.3.4	Time in the hierarchical universe.....	52
9	Appendix.....	53
9.1	Definition of Terms.....	53
9.2	Arrangement of Terms.....	53
10	Acknowledgment.....	54
11	References.....	54

1 Introduction

1.1 Subject

According to Born's rule, an observed probability of a particle is proportional to the square of the absolute value of the wave function. On the other hand, according to the many-worlds interpretation, we observe the particle of the various places in the various events. It is the subject of this paper to derive Born's rule by counting the number of the events.

1.2 The importance of the subject

Wave function collapse and Born's rule are the principle of the quantum mechanics. We can eliminate the wave function collapse from the quantum mechanics by Many-Worlds Interpretation (MWI), but we cannot eliminate Born's rule.

For this reason, many researchers have tried to derive Born's rule from MWI. However, it has not resulted in the success. Therefore, it has become an important subject to derive Born's rule.

1.3 Past derivation method

Hugh Everett III² claimed that he derived Born's rule from Many-Worlds Interpretation (MWI) in 1957. After that, many researchers claimed that they derived Born's rule from the method that is different from the method of Everett. James Hartle³ claimed in 1968, Bryce DeWitt⁴ claimed in 1970 and Neil Graham⁵ claimed in 1973 that they derived Born's rule.

However, Adrian Kent pointed out that their method of deriving Born's rule was insufficient⁶ in 1990. David Deutsch⁷ tried to derive Born's rule in 1999. Sumio Wada⁸ also tried it in 2007. However, many researchers do not agree on the methods of deriving Born's rule.

1.4 New derivation method of this paper

In the probability theory, we explain the probability by the concept of the elementary event. Therefore, we might be able to explain the probability of the quantum theory by the same concept. We try to derive the probability of the quantum theory by introducing a concept of the elementary event to the quantum theory as the new method of this paper.

2 Traditional method of deriving and the problem

2.1 The Born rule

Max Born⁹ proposed the Born rule in 1926. It is also called Born's rule, the Born law, the Born's law, or the probability interpretation. Born's rule is a principle of quantum mechanics. We express the state of the particle by the wave function $\psi(x)$ in quantum mechanics. We show an example of the wave function in the following figure.




Figure 2-1: An example of a wave function

The observed probability of a particle is proportional to the square of the absolute value of the wave function. We express the observed probability $P(x)$ of a particle at the position x as follows.

$$P(x) = |\psi(x)|^2 \quad (2.1)$$

According to the Copenhagen interpretation that is a general interpretation of quantum mechanics, we cannot mention the state of the particle before the observation because the wave function does not exist physically. However, the wave function might exist physically. One of the interpretations based on the existence of a wave function is a many-worlds interpretation.

2.2 Everett's many-worlds interpretation

Everett proposed Many-Worlds Interpretation (MWI) in order to deal with the universal wave function. He tried to derive Born's rule from the measure theory.

For example, we consider the Stern-Gerlach experiment.

We express the state vector of the spin of an electron by the eigenstate vector of the spin along the z -axis direction as follows.

$$a|\psi\rangle = a_1|z+\rangle + a_2|z-\rangle \quad (2.2)$$

The coefficients a and a_k are complex number.

Here, we have normalized $|\psi\rangle$, $|z+\rangle$ and, $|z-\rangle$.

$$\langle\psi|\psi\rangle = 1 \quad (2.3)$$

$$\langle z+|z+\rangle = 1 \quad (2.4)$$

$$\langle z-|z-\rangle = 1 \quad (2.5)$$

In order to derive the probability, Everett introduced a new concept, measure. He expressed the measure by a positive function $m(a)$. He requested the following equation for the measure.

$$m(a) = m(a_1) + m(a_2) \quad (2.6)$$

He adduced a probability conservation law to justify the request. We write the function $m(a)$ satisfying the above equation by using a positive constant c as follows.

$$m(a) = c|a|^2 \quad (2.7)$$

$$m(a_1) = c|a_1|^2 \quad (2.8)$$

$$m(a_2) = c|a_2|^2 \quad (2.9)$$

Andrew Gleason¹⁰ generally proved the above equation in 1957. His proof is called the Gleason's theorem. Everett considered the infinite time measurement, and concluded that the measure behaves like the probability. However, MWI of Everett has the **basis problem** and the **probability problem**. I will explain them in the following sections.

2.2.1 The basis problem of many-worlds interpretation

If we define the measure by using a particular basis, we need to show how to select a particular basis. However, Everett did not show how to select a particular basis in his paper.

For example, we consider the Stern-Gerlach experiment. We express a spin state of any direction by superposition of the upward spin state along z -axis and the downward spin state along z -axis. Therefore, we express the wave function of an electron by the basis of the spin eigenstate along z -axis as follows.

$$a|\psi\rangle = a_1|z+\rangle + a_2|z-\rangle \quad (2.10)$$

We express the measure of the spin eigenstate along z -axis as follows.

$$m(a_1) = |a_1|^2 \quad (2.11)$$

$$m(a_2) = |a_2|^2 \quad (2.12)$$

On the other hand, we also express the wave function of an electron by the basis of the spin eigenstate along x -axis as follows.

$$a|\psi\rangle = b_1|x+\rangle + b_2|x-\rangle \quad (2.13)$$

We express the measure of the spin eigenstate along x -axis as follows.

$$m(b_1) = |b_1|^2 \quad (2.14)$$

$$m(b_2) = |b_2|^2 \quad (2.15)$$

If the measure of Everett is a quantity that has a physical meaning, it should not change by choice of a basis of eigenstate. Therefore, we need a method to choose a specific basis. Everett did not show the method.

2.2.2 The probability problem of many-worlds interpretation

Everett tried to derive Born's rule from the measure theory. Then, Everett did not give the physical meaning to the measure. However, to request the conservation law of the probability for the equation of the measure is equivalent to defining the measure as the probability. Therefore, it is circular reasoning to show that measure acts like a probability for infinite time measurement.

If the number of worlds is proportional to the measure, it is necessary to clarify the mechanism by which the number is proportional to the measure of a world. If the number of worlds is not proportional to the measure, it is necessary to explain how the probability of occurrence of a world is proportional to the measure.

3 Review of existing ideas

3.1 Universal Wave function of Wheeler and DeWitt

John Wheeler and Bryce DeWitt¹¹ proposed the universal wave function in 1967. We have the wave function by the Hamiltonian operator H and the ket vector $|\psi\rangle$ as follows.

$$H|\psi\rangle = 0 \quad (3.1)$$

This ket vector $|\psi\rangle$ is not a normal function but a functional.

A functional is mathematically almost equivalent to a function of many variables. Since the discussion based on the functional is difficult, we use a function of many variables for discussion in this paper. The following sections describe the many-particle wave function, which is a function of many variables.

3.2 Barbour's many-particle wave function of the universe

Julian Barbour¹² expressed the universe by using the many-particle wave function in his book *The End of Time* in 1999.

We suppose that the number of the particles in the universe is n , and the k -th particle's position is $r_k = (x_k, y_k, z_k)$. Then we express the many-particle wave function ψ as follows.

$$\psi = \psi(r_1, r_2, r_3, \dots, r_n) \quad (3.2)$$

The many dimensional space expressing the positions of all the particles is called configuration space.




Figure 3-1: Many-particle wave function

The configuration space expresses all the possible worlds those exist physically in the past, the present, and the future, because a point in the configuration expresses the positions of all the particles. In other words, many-particle wave function expresses all the possible worlds in many-worlds interpretation.

If the combination of the positions of all particles of a world is decided, the state of the clock of the world will be decided. If the state of the clock of the world is decided, the time of the clock of the world is decided. Therefore, many-particle wave function does not need time as the argument of the function.

It is possible to choose a position or a momentum as a basis of a wave function. This paper chooses the position as a basic basis, since we always observe a position finally by an experiment.

The number of particles changes in the quantum field theory. Therefore, it is impossible to express the quantum field by the many-particle wave function. We need a functional in order to express the quantum field. On the other hand, it is possible to express the functional by many-variable function approximately. Then, we use many-variable function, many-particle wave function in order to argue easily in this paper.

We express the probability P that we observe a world in the configuration space as follows.

$$P = |\psi(r_1, r_2, r_3, \dots, r_n)|^2 \quad (3.3)$$

In order to consider the reason why we express the probability by this equation, we will review the probability theory in the following section.

3.3 Laplace's Probability Theory

Pierre-Simon Laplace¹³ summarized the classical probability theory in 1814. He described the following calculation method of the probability.

The probability of an event is the ratio of the number of cases favorable to it, to the number of all cases possible when nothing leads us to expect that any one of these cases should occur more than any other, which renders them, for us, equally possible.

This "equally possible" case is an elementary event in probability theory. All elementary events have a same probability of occurrence.

An elementary event is also called an atomic event. In this paper, we call "equally possible" case an **elementary event**.

We suppose that the number of all elementary events is N_a , and the number of elementary events of an event is N . Then, we express the probability P of occurrence of the event as follows.

$$P = \frac{N}{N_a} \propto N \quad (3.4)$$

$$N \ll N_A \quad (3.5)$$

For example, we suppose that the five balls are in the bag. Three of five balls are red and two balls are blue. We suppose that the probability of the event that we take out the red ball is P . Then, the probability is $3/5$.




Figure 3-2: Event is a set of elementary events

We explain the reason by the concept of an elementary event. According to the probability theory, we interpret the event that we take out a ball as an elementary event. We interpret an event as a set of elementary events.

In order to derive Born's rule, we need to find elementary events of quantum theory. An elementary event of probability theory generally we cannot divide anymore, so it is expected that an elementary event of quantum theory also cannot be divided anymore.

3.4 Penrose's spin networks

Roger Penrose¹⁴ proposed spin networks in 1971. According to the spin networks, we express the space as a graph with a line that connects a point and the other point. This graph is called spin network. Since the space-time is discrete, the space-time has a minimum length and minimum time.




Figure 3-3: Penrose's spin network

In this paper, though we do not use a spin network, we assume that space-time is discrete as well as by this theory and the space is a graph that connects the points. In this paper, we assume that the minimum length is Planck length ℓ_P and the minimum time is Planck time t_P .

$$\ell_P = \sqrt{\frac{\hbar G}{c^3}} \approx 1.6 \times 10^{-35} [m] \quad (3.6)$$

$$t_P = \sqrt{\frac{\hbar G}{c^5}} \approx 5.4 \times 10^{-44} [s] \quad (3.7)$$

We call the minimum domain that is constructed by the Planck length elementary domain.

If the space-time is discrete, we need to review the theory that has been constructed based on the continuous space-time. Therefore, in the next section, we review what happens in the path integral in the case of discrete space-time.

3.5 Feynman's path integral

Richard Feynman¹⁵ proposed path integral in 1948. It provides a new quantization method. In the path integral, we need to take the sum of all the possible paths of the particle.

We express the probability amplitude $K(b, a)$ from the position a to the position b as follows.

$$K(b, a) = \int_a^b Dx(t) \exp\left(\frac{i}{\hbar} S[b, a]\right) \quad (3.8)$$

The probability amplitude $K(b, a)$ is called propagator. The symbol $Dx(t)$ represents the sum of the probability amplitudes for all paths. We express the wave functions by the propagator as follows.

$$\psi(b, t_b) = K(b, a)\psi(a, t_a) \quad (3.9)$$

In the path integral, an event that a particle moves from a position a to the other position b is made to correspond to the propagator $K(b, a)$. We get the wave function of time t_b by multiplying the propagator $K(b, a)$ to a wave function of time t_a .

As shown in the following figure, there is not only a normal path of α but also the other path of β to travel long distance in a short period of time. Such path might have a speed that is greater than the speed of light. Since the path is contrary to the special relativity, the path is not allowed. In this paper, we call such a movement of the path **long distance transition**.




Figure 3-4: Feynman's path integral

Generally, the textbook of a path integral explains as follows.

The sum of a minutely different path near a path α becomes large. On the other hand, the sum of a minutely different path near a path β becomes small. For this reason, long distance transition is suppressed and the path β does not remain.

Then, what happens after the minimum time t_P ? We express the propagator from a position a to a position b after a minimum time t_P as follows.

$$K(b, a) = \exp\left(\frac{i}{\hbar} S[b, a]\right) \quad (3.10)$$

In this paper, we assume the discrete time. Since we cannot divide minimum time any more, when the departure point and the point of arrival are decided, it cannot take a minutely different path near a path β . For this reason, we cannot suppress long distance transition and the path β remains.

Therefore, if we apply the path integral to the discrete space-time and the position of a particle is determined like a delta function of the Dirac, long distance transition occurs after the minimum time t_P .

$$\psi(x', t_b) = K(x', x) \delta(x - a) \quad (3.11)$$




Figure 3-5: Long distance transition in the path integral

However, we do not observe the long distance transition. We deduce the reason is that the position of the particle is **distributed with a normal distribution** like the following figure.




Figure 3-6: A wave function of a localized state

Therefore, position x is distributed with deviation Δx , momentum p is also distributed with deviation Δp . According to the Uncertainty Principle, the product of Δx and Δp is close to the Planck constant $\hbar/2$.

$$\Delta x \Delta p \approx \frac{\hbar}{2} \quad (3.12)$$

We call the state of the wave function with a normal distribution **localized state**.

We express a wave function of a particle with momentum p as follows.

$$\psi(x) = \exp\left(\frac{i}{\hbar} px\right) \quad (3.13)$$

We suppose that this particle has a mass m and the velocity v . The momentum is shown below.

$$p = mv \quad (3.14)$$

We get the following formula by substituting this formula to the wave function.

$$\psi(x) = \exp\left(\frac{i}{\hbar} mvx\right) \quad (3.15)$$

We express the velocity v by the moving distance x and the Planck length t_P .

$$v = \frac{x}{t_P} \quad (3.16)$$

We get the following formula by substituting this formula to the wave function.

$$\psi(x) = \exp\left(i \frac{m}{\hbar t_P} x^2\right) \quad (3.17)$$

From the above formula, the wavelength of the wave function is long at the short range. On the other hand, the wavelength of the wave function is short at the long range.

In the short distance, the sum of the path integral of localized state becomes large. On the other hand, in long distance, the sum of the path integral of localized state becomes small. We call this phenomenon “**suppression of long distance transition due to localized states.**”

If the state is localized state, the long distance transition does not occur after the minimum time t_P . Therefore, the localized state is localized near the place after the time t_P . For this reason, we deduce that network structure of the path integral is realized, as shown in the following figure.




Figure 3-7: Network structure of the path integral

In this paper, we call the network structure of “**path network structure of the path integral.**”

We suppose that there is an event AB that is a transition from a state A to a state B . If the state A has three positions and the state B has three positions, the event AB has $3 \times 3 = 9$ paths.

In "network structure of the integral path," the number of paths is the square of the number of positions. On the other hand, according to Born's rule, the probability becomes the square of the absolute value of the wave function. In this paper, we discuss the similarities of these squares.

3.6 Dirac's quantum field theory

Paul Dirac¹⁶ proposed the quantum field theory to explain the emission and absorption of electromagnetic waves in 1927. We express the fundamental commutation relation¹⁷ of the quantum field theory in the case of one-dimensional space as follows.

$$[\psi(x), \pi(y)] = i\hbar\delta(x - y) \quad (3.18)$$

Then ψ is the field and π is the conjugate operator of the field ψ . The variable x and y are positions. The function δ is Dirac's delta function.

This commutation relation is similar to the following commutation relation between position x and momentum p .

$$[x, p] = i\hbar \quad (3.19)$$

This indicates that field ψ is a physical quantity that has a property similar to the position x . In this paper, we call the physical quantity “**positional physical quantity.**”

We got a field $\psi(x)$ by the first quantization for the position x . On the other hand, the field $\psi(x)$ is "positional physical quantity" like the position x . Therefore, we get a new field $\Psi(x, \psi(x))$ by the second

quantization for the field ψ . We call the field $\Psi(x, \psi(x))$ **the second wave function**. We express the second wave function $\Psi(x, \psi(x))$ in the following figure.




Figure 3-8: The second wave function

It is possible to interpret the second wave function $\Psi(x, \psi(x))$ as a functional $\Phi[\psi(x)]$. We express the functional $\Phi[\psi(x)]$ by the many-particle wave function $\psi(x_1, x_2, x_3, \dots, x_n)$ approximately. To argue a point easily, we use many-particle wave functions by this paper.

3.7 Kaluza-Klein theory

Theodor Kaluza¹⁸ proposed in 1921 and Oskar Klein¹⁹ proposed in 1926 the extra space like a one-dimensional circle, in order to unify the electromagnetic field and gravity. This theory is called Kaluza-Klein theory.

We express a new space $M^4 \times S^1$ by using a normal four-dimensional space-time M^4 and an extra space S^1 like a one-dimensional circle as follows.

$$M \times S^1 \tag{3.20}$$




Figure 3-9: Kaluza-Klein theory

3.8 Euler's formula

Euler published the following formula in 1748.
(Euler's formula)

$$\exp(i\phi) = \cos \phi + i \sin \phi \quad (3.21)$$

Imaginary number i satisfies the following equation.

$$i^2 = -1 \quad (3.22)$$

We express the complex number as follows.

$$s = t + ix \in \mathbb{C} \quad (3.23)$$

$$t, x \in \mathbb{R} \quad (3.24)$$

We express the complex conjugate as follows.

$$\bar{s} = t - ix \in \mathbb{C} \quad (3.25)$$

We express the complex function as follows.

$$f(s) \in \mathbb{C} \quad (3.26)$$

We express the square of the absolute value of the complex number as follows.

$$|s|^2 = s\bar{s} \quad (3.27)$$

We use the following symbols as follows.

$$\operatorname{Re}(s) = \frac{1}{2}(s + \bar{s}) = t \quad (3.28)$$

$$\operatorname{Im}(s) = \frac{1}{2}(s - \bar{s}) = ix \quad (3.29)$$

3.9 Cauchy-Riemann equation

Augustin Louis Cauchy²⁰ introduced the following equation in 1814 for complex analysis. Riemann²¹ used the following equation in 1851.

(Cauchy-Riemann equation)

$$\frac{\partial f}{\partial t} + i \frac{\partial f}{\partial x} = 0 \quad (3.30)$$

We express the above equation shortly as follows.

(Cauchy-Riemann equation)

$$\frac{\partial f}{\partial \bar{s}} = 0 \quad (3.31)$$

Cauchy introduced the following formula.

(Cauchy's integral formula)

$$f(s) = \oint_S \frac{dt}{2\pi i} \frac{f(t)}{(t-s)} \quad (3.32)$$

S is the contour path.

3.10 Hamilton's Quaternion

William Rowan Hamilton²² proposed the quaternion in 1843.

$$i^2 = j^2 = k^2 = ijk = -1 \quad (3.33)$$

We express the quaternion as follows.

$$s = t + ix + jy + kz \in \mathbb{H} \quad (3.34)$$

$$t, x, y, z \in \mathbb{R} \quad (3.35)$$

We express the quaternion conjugate as follows.

$$\bar{s} = t - ix - jy - kz \in \mathbb{H} \quad (3.36)$$

We express the quaternion function as follows.

$$f(s) \in \mathbb{H} \quad (3.37)$$

We express the square of the absolute value of the quaternion as follows.

$$|s|^2 = s\bar{s} \quad (3.38)$$

We use the following symbols as follows.

$$\operatorname{Re}(s) = \frac{1}{2}(s + \bar{s}) = t \quad (3.39)$$

$$\operatorname{Im}(s) = \frac{1}{2}(s - \bar{s}) = ix + jy + kz \quad (3.40)$$

3.11 Cauchy-Riemann-Fueter equation

Fueter²³ introduced the following equation in 1934 for quaternionic analysis.
(Cauchy-Riemann-Fueter equation)

$$\frac{\partial f}{\partial \tau} + i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z} = 0 \quad (3.41)$$

We express the above equation shortly as follows.
(Cauchy-Riemann-Fueter equation)

$$\frac{\partial f}{\partial \bar{s}} = 0 \quad (3.42)$$

$$\frac{\partial}{\partial \bar{s}} = \frac{\partial}{\partial t} + i \frac{\partial}{\partial x} + j \frac{\partial}{\partial y} + k \frac{\partial}{\partial z} \quad (3.43)$$

Fueter introduced the following formula.
(Cauchy-Fueter integral formula)

$$f(s) = \frac{1}{2\pi^2} \oint_{S^3} \frac{(q-s)^{-1}}{|q-s|^2} Dq f(q) \quad (3.44)$$

Here, S^3 is a three-dimensional closed surface. The detail of the quaternionic analysis was described in the Anthony Sudbery's paper²⁴ in 1979.

We use the following formula.
(Integral formula of quaternion)

$$f(s) = \oint_S \frac{dt}{2\pi u} \frac{f(t)}{(t-s)} \quad (3.45)$$

Here, u is a unit quaternion.

Please refer to the following paper for the integral formula of quaternion.

•Derivation of the reflection integral equation of the zeta function by the quaternionic analysis [2014/5/18]

http://www.geocities.jp/x_seek/Quaternion_e.htm

3.12 Cartan's differential form

Elie Cartan²⁵ defined differential form in 1899 in order to describe manifold by the method that is independent to the coordinates.

Though the differential form $d\omega$ is infinitesimal, we use **difference form** $\delta\omega$ of finitesimal

We express the surface area A of the manifold S as follows.

$$A = \int_S h(\omega) |d\omega| \quad (3.46)$$

We express the difference form δS of the surface area of the manifold S as follows.

$$\delta S(\omega) = h(\omega) \delta\omega \quad (3.47)$$




Figure 3-10: Manifold

Here, we express the difference form δS_1 of the surface area of the manifold S_1 as follows.

$$\delta S_1(\omega) = h_1(\omega) \delta\omega \quad (3.48)$$

Then, we express the difference form δS_2 of the surface area of the manifold S_2 as follows.

$$\delta S_2(\omega) = h_2(\omega) \delta\omega \quad (3.49)$$

We get the following manifold S as the superposition of the manifold S_1 and S_2 .

$$S = S_1 + S_2 \quad (3.50)$$

We sum the complex numbers of wave functions every position for the superposition of a wave function. Therefore, we deduce that we sum the surface areas of manifolds at every solid angle for the superposition of manifolds.

Then, we express the difference form δS of the manifold S as follows.

$$\delta S(\omega) = \delta S_1(\omega) + \delta S_2(\omega) \quad (3.51)$$

Therefore, we have the following formula for the spherical harmonics.

$$h(\omega) = h_1(\omega) + h_2(\omega) \quad (3.52)$$

We define the superposition of the manifolds by the above formula.

4 A new method of deriving

4.1 Universe of Two-dimensional space-time

4.1.1 Closed path of Two-dimensional space-time

We consider the universe U of two-dimensional space-time.

We express the world line s of the particle by a complex number.

$$s = t + ix \quad (4.1)$$

$$i^2 = -1 \quad (4.2)$$

We express the wave function of this particle as follows.

$$\psi(t, x) \in \mathbb{C} \quad (4.3)$$

We suppose that particles are generated by pair production and destroyed by pair annihilation.




Figure 4-1: Pair production and pair annihilation

We express the closed path C by the circle C of radius R as follows.




Figure 4-2: Closed path C

We express this closed path as follows.

$$s = r \exp(i\theta) \quad (4.4)$$

We express the circumference a of this circle C as follows.

$$a = \int_0^{2\pi} r |i \exp(i\theta)| d\theta \quad (4.5)$$

Here we introduce the **complex solid angle** ω .

$$\omega = \exp(i\theta) \quad (4.6)$$

$$d\omega = i \exp(i\theta) d\theta \quad (4.7)$$

We express the closed path as follows.

$$s = t + ix = r\omega \quad (4.8)$$

Then we express the circumference a of this circle C as follows.

$$a = \int_C r |d\omega| \quad (4.9)$$

We express the difference form of the circumference a as follows.

$$\delta s(\omega) = r \delta\omega \quad (4.10)$$

4.1.2 Introduction of the absolute value of wave function

We introduce a circle S as an extra space like Kaluza-Klein theory.

We express the point on the circle C by matrix representation of a complex number as follows.

$$C = ET + IX \quad (4.11)$$

$$I^2 = -E \quad (4.12)$$

$$E = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (4.13)$$

$$I = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad (4.14)$$

We call the circle an amplitude circle or an **amplitude 1-sphere**.




Figure 4-3: Amplitude 1-sphere S

We express the circumference A of the amplitude 1-sphere S by the radius R and the solid angle Ω as follows.

$$A = \int_S R |d\Omega| \quad (4.15)$$

If the radius R is the function of r and ω , the circumference A becomes the function of r and ω .

$$A(r, \omega) = \int_S R(r, \omega) |d\Omega| \quad (4.16)$$

We express the difference form δS of the sphere S .

$$\delta S(r, \omega, \Omega) = R(r, \omega) \delta\Omega \quad (4.17)$$

We interpret the circumference A as the absolute value of the wave function.

$$A(r, \omega) = |\psi(t, x)| \quad (4.18)$$

4.1.3 Introduction of the phase of wave function

In order to introduce the phase of wave function, we rotate the sphere S by the rotational transform angle Φ . We transform the sphere S to the new sphere S' by the rotational transform which depends on ω .

$$P(\omega): S(r, \omega) \rightarrow S'(r, \omega) \quad (4.19)$$

We define the rotational transform of the rotational transform angle Φ that depends on ω as follows.

$$P(\omega) = \exp(I\Phi(\omega)) \quad (4.20)$$

We express the function $\Phi(\omega)$ by the natural number n as follows.

$$\Phi(\omega) = n\omega \quad (4.21)$$

We interpret the rotational angle Φ as the phase of the wave function.

$$\Phi(\omega) = \arg(\psi(t, x)) \quad (4.22)$$

We transform the sphere S by the rotational transform as follows.

$$\delta S'(r, \omega, \Omega') = \exp(I\Phi(\omega)) \delta S(r, \omega, \Omega) \quad (4.23)$$




Figure 4-4: Rotation of the amplitude 1-sphere S

We define the superposition of a sphere S_1 and a sphere S_2 as follows.

$$\delta S(r, \omega, \Omega) = \delta S_1(r, \omega, \Omega) + \delta S_2(r, \omega, \Omega) \quad (4.24)$$

The superposition of the sphere and the sphere that is rotated by the angle 180 degrees is zero.

$$\delta S'(r, \omega, \Omega) = \exp(I\pi) \delta S(r, \omega, \Omega) \quad (4.25)$$

$$0 = \delta S(r, \omega, \Omega) + \delta S'(r, \omega, \Omega) \quad (4.26)$$

The direct product of the closed path C of the particle and the sphere S becomes a torus T .

$$T = C \times S \quad (4.27)$$

$$\delta T(\omega, \Omega) = \delta C(\omega) \times \delta S(r, \omega, \Omega) \quad (4.28)$$

$$\delta T(\omega, \Omega) = r(\omega) \delta \omega R(r, \omega, \Omega) \delta \Omega \quad (4.29)$$

We call this torus T a **torus world-sheet**.




Figure 4-5: The torus world-sheet

Here, we introduce the following new solid angle ν .

$$\nu = (\omega, \Omega) \quad (4.30)$$

Here we introduce the following new solid radius ρ .

$$\rho = (r, R) \quad (4.31)$$

Here we introduce the following new function $f(\rho, \nu)$.

$$f(\rho, \nu) = r(\omega) R(\omega, \Omega) \quad (4.32)$$

We express the torus world-sheet T by the function $f(\rho, \nu)$ as follows.

$$\delta T(\rho, \nu) = f(\rho, \nu) \delta \nu \quad (4.33)$$

The torus world-sheet is twisted like a helical torus as the following figure.




Figure 4-6: The torus world-sheet is twisted like a helical torus.

This dimension of the torus world-sheet is same as the dimension of the universe because the universe is two-dimensional space-time in this section.

Here we use a surprising idea.

We interpret the torus world-sheet as a new space-time. We call the space-time **toric space-time**.

We interpret the toric space-time an independent universe. We call the universe **the second universe**.

It is possible to construct the third and the forth universe in the same way that we construct the second universe. We construct many universes by repeating the same way. We call these universes **hierarchical universe**.

4.1.4 Hierarchical universe

We show the hierarchical universe as follows.




Figure 4-7: Hierarchical universe

We express the above hierarchical universe by the following symbol.

$$\dots \rightarrow U_1 \rightarrow U_2 \rightarrow U_3 \rightarrow \dots \quad (4.34)$$

4.1.5 Equations

We express the position s by the complex number as follows.

$$s = \tau + ix \in \mathbb{C} \quad (4.35)$$

Then the wave function becomes a complex function.

$$f(s) \in \mathbb{C} \quad (4.36)$$

We assume that the complex function is an analytic function.

Analytic functions satisfy the Cauchy-Riemann equation.

(Cauchy-Riemann equation)

$$\frac{\partial f}{\partial \tau} + i \frac{\partial f}{\partial x} = 0 \quad (4.37)$$

We call the equation **path differential equation**.

We define the complex conjugate as follows.

$$\bar{s} = \tau - ix \in \mathbb{C} \quad (4.38)$$

Then we express the path differential equation shortly as follows.

$$\frac{\partial}{\partial \bar{s}} f(s) = 0 \quad (4.39)$$

Analytic function satisfy the Cauchy's integral formula.
(Cauchy's integral formula)

$$f(s) = \oint_S \frac{dt}{2\pi i} \frac{f(t)}{(t-s)} \quad (4.40)$$

S is the contour path.

We interpret the Cauchy's integral formula as the **path integral equation** of Feynman's path integral.




Figure 4-8: The path integral equation of Feynman's path integral

We interpret that the particle on the circle S transit from the position t to the position s for the long distance directly.

We call the new interpretation the **path integral of space-time view** that is different from the traditional Feynman's path integral.

It is possible to use these equations for wave functions of each universe because the wave functions of each universe are complex functions.

4.2 Universe of four-dimensional space-time

4.2.1 Closed path of four-dimensional space-time

We consider the universe U of four-dimensional space-time.

We express the world line s of the particle by quaternion.

$$s = t + ix + jy + kz \quad (4.41)$$

$$i^2 = j^2 = k^2 = ijk = -1 \quad (4.42)$$

We express the wave function of this particle as follows.

$$\psi(t, x, y, z) \in \mathbb{C} \quad (4.43)$$

We suppose that particles are generated by pair production and destroyed by pair annihilation.




Figure 4-9: Pair production and pair annihilation

We express the closed path s by the circle C of radius r as follows.




Figure 4-10: Closed path C

We express this closed path as follows.

$$s = r \exp(i\theta) \quad (4.44)$$

We express the circumference a of this circle C as follows.

$$a = \int_0^{2\pi} r |i \exp(i\theta)| d\theta \quad (4.45)$$

Here we introduce the **quaternionic solid angle** ω .

$$\omega = \exp(i\theta) \quad (4.46)$$

$$d\omega = i \exp(i\theta) d\theta \quad (4.47)$$

We express the closed path as follows.

$$s = t + ix + jy + kz = r\omega \quad (4.48)$$

Then we express the circumference a of this circle C as follows.

$$a = \int_C r |d\omega| \quad (4.49)$$

We express the difference form of the circumference a as follows.

$$\delta s(\omega) = r \delta\omega \quad (4.50)$$

4.2.2 Introduction of the absolute value of wave function

We introduce a three-dimensional sphere (3-sphere) S as an extra space like Kaluza-Klein theory.

We express the point S on the 3-sphere by matrix representation of quaternion.

$$S = ET + IX + JY + KZ \quad (4.51)$$

$$I^2 = J^2 = K^2 = IJK = -E \quad (4.52)$$

$$E = \begin{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \\ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \end{pmatrix} \quad (4.53)$$

$$I = \begin{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \\ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} & -\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \end{pmatrix} \quad (4.54)$$

$$J = \begin{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ -\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \end{pmatrix} \quad (4.55)$$

$$K = \begin{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} & \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \\ \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \end{pmatrix} \quad (4.56)$$

We call the circle **amplitude 3-sphere**.




Figure 4-11: Amplitude 3-sphere S

We express the circumference A of the amplitude 3-sphere S by the radius R and the solid angle Ω as follows.

$$A = \int_S R^3 |d\Omega| \quad (4.57)$$

If the radius R is the function of r and ω , the circumference A becomes the function of r and ω .

$$A(r, \omega) = \int_S R^3(r, \omega) |d\Omega| \quad (4.58)$$

We express the difference form δS of the sphere S .

$$\delta S(r, \omega, \Omega) = R^3(r, \omega) \delta\Omega \quad (4.59)$$

We interpret the circumference A as the absolute value of the wave function.

$$A(r, \omega) = |\psi(t, x, y, z)| \quad (4.60)$$

4.2.3 Introduction of the phase of wave function

In order to introduce the phase of wave function, we rotate the sphere S by the rotational transform angle Φ . We transform the sphere S to the new sphere S' by the rotational transform which depends on ω .

$$P(\omega): S(r, \omega) \rightarrow S'(r, \omega) \quad (4.61)$$

We define the rotational transform of the rotational transform angle Φ that depends on ω as follows.

$$P(\omega) = \exp(I\Phi(\omega)) \quad (4.62)$$

We express the function $\Phi(\omega)$ by the natural number n as follows.

$$\Phi(\omega) = n\omega \quad (4.63)$$

We interpret the rotational angle Φ as the phase of the wave function.

$$\Phi(\omega) = \arg(\psi(t, x, y, z)) \quad (4.64)$$

We transform the sphere S by the rotational transform as follows.

$$\delta S'(r, \omega, \Omega') = \exp(I\Phi(\omega)) \delta S(r, \omega, \Omega) \quad (4.65)$$




Figure 4-12: Rotation of the amplitude 3-sphere S

We define the superposition of a sphere S_1 and a sphere S_2 as follows.

$$\delta S(r, \omega, \Omega) = \delta S_1(r, \omega, \Omega) + \delta S_2(r, \omega, \Omega) \quad (4.66)$$

The superposition of the sphere and the sphere that is rotated by the angle 180 degrees is zero.

$$\delta S'(r, \omega, \Omega) = \exp(I\pi) \delta S(r, \omega, \Omega) \quad (4.67)$$

$$0 = \delta S(r, \omega, \Omega) + \delta S'(r, \omega, \Omega) \quad (4.68)$$

The direct product of the closed path C of the particle and the sphere S becomes a manifold like a torus T .

$$T = C \times S \quad (4.69)$$

$$\delta T(\omega, \Omega) = \delta C(\omega) \times \delta S(r, \omega, \Omega) \quad (4.70)$$

$$\delta T(\omega, \Omega) = r(\omega) \delta \omega R(r, \omega, \Omega) \delta \Omega \quad (4.71)$$

We call this manifold T a **torus world-sheet**.




Figure 4-13: The torus world-sheet

Here we introduce the following new solid angle ν .

$$\nu = (\omega, \Omega) \quad (4.72)$$

Here we introduce the following new solid radius ρ .

$$\rho = (r, R) \quad (4.73)$$

Here we introduce the following new function $f(\rho, \nu)$.

$$f(\rho, \nu) = r(\omega) R(\omega, \Omega) \quad (4.74)$$

We express the torus T by the function $f(\rho, \nu)$ as follows.

$$\delta T(\rho, \nu) = f(\rho, \nu) \delta \nu \quad (4.75)$$

The torus world-sheet is twisted like a helical torus as the following figure.




Figure 4-14: The torus world-sheet is twisted like a helical torus.

This dimension of the torus world-sheet is same as the dimension of the universe because the universe is four-dimensional space-time in this section.

Here we use a surprising idea.

We interpret the torus world-sheet as a new space-time. We call the space-time **toric space-time**.

We interpret the toric space-time an independent universe. We call the universe **the second universe**.

It is possible to construct the third and the fourth universe in the same way that we construct the second universe. We construct many universes by repeating the same way. We call these universes **hierarchical universe**.

4.2.4 Hierarchical universe

We show the hierarchical universe as follows.




Figure 4-15: Hierarchical universe

We express the above hierarchical universe by the following symbol.

$$\dots \rightarrow U_1 \rightarrow U_2 \rightarrow U_3 \rightarrow \dots \quad (4.76)$$

4.2.5 Equations

We express the position s by the quaternion number as follows.

$$s = \tau + ix + jy + kz \in \mathbb{H} \quad (4.77)$$

Then, the wave function becomes a quaternionic function.

$$f(s) \in \mathbb{H} \quad (4.78)$$

We assume that quaternionic functions are analytic functions. Analytic functions satisfy the Cauchy-Riemann-Fueter equation. (Cauchy-Riemann-Fueter equation)

$$\frac{\partial f}{\partial \tau} + i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z} = 0 \quad (4.79)$$

We call the above equation **path differential equation** in this paper.

We define the quaternionic conjugate as follows.

$$\bar{s} = \tau - ix - jy - kz \in \mathbb{H} \quad (4.80)$$

Then we express the path differential equation shortly as follows.

$$\frac{\partial}{\partial \bar{s}} f(s) = 0 \quad (4.81)$$

Analytic functions satisfy the integral formula of quaternion. (Integral formula of quaternion)

$$f(s) = \oint_S \frac{dt}{2\pi u} \frac{f(t)}{(t-s)} \quad (4.82)$$

Here, u is a unit quaternion.

We interpret the integral formula of quaternion as the **path integral equation** of Feynman's path integral.




Figure 4-16: Path integral equation of Feynman's path integral

We interpret that the particle on the circle S^1 transit from the position t to the position s for the long distance directly.

We call the new interpretation a **space-time view path integral** that is different from the traditional Feynman's path integral.

It is possible to use these equations for the wave function of each universe because the wave functions of each universe are quaternionic functions.

4.3 Normal space

We express the surface area a of the normal space U as follows.

$$a = \int_U r^3 d\omega \tag{4.83}$$

We express the difference form of the normal space U as follows.

$$\delta s(\omega) = r^3 \delta\omega \tag{4.84}$$

Here we replace the r^3 to the function $f(\omega)$.

$$f(\omega) = r^3 \tag{4.85}$$

Then we express the following formula.

$$\delta s(\omega) = f(\omega)\delta\omega \tag{4.86}$$

We interpret the above formula like the following figure.
 We call the interpretation **manifold view**.




Figure 4-17: Manifold view of the normal space U

Here we rewrite the formula as follows by a new function $f(r, \omega)$.

$$\delta s(r, \omega) = f(r, \omega)\delta\omega \tag{4.87}$$

We interpret the above formula like the following figure.
 We call the interpretation **spherical harmonics view**.




Figure 4-18: Spherical harmonics view of the normal space U

In the spherical harmonics view, we interpret the function $f(r, \omega)$ as the spherical harmonics. The spherical harmonics $f(r, \omega)$ is the solution of the Laplace equation of the spheric polar coordinates. Therefore, spherical harmonics $f(r, \omega)$ satisfies the following Laplace equation.

$$(\partial_s^2 + \partial_s^2)f = 0 \quad (4.88)$$

Here, we used the following symbols.

$$\partial_{\bar{s}} = \frac{\partial}{\partial t} + i \frac{\partial}{\partial x} + j \frac{\partial}{\partial y} + k \frac{\partial}{\partial z} \quad (4.89)$$

$$\partial_s = \frac{\partial}{\partial t} - i \frac{\partial}{\partial x} - j \frac{\partial}{\partial y} - k \frac{\partial}{\partial z} \quad (4.90)$$

4.4 Wave space-time

We express the normal space-time U by the radius r and the solid angle ω as follows.

$$\delta S(r, \omega) = f(r, \omega) \delta \omega \quad (4.91)$$

We express the amplitude 3-sphere S by the radius R and the solid angle Ω as follows.

$$\delta S(R, \Omega) = F(R, \Omega) \delta \Omega \quad (4.92)$$

We define the wave space-time W as the direct product of the normal space-time U and the amplitude 3-sphere S as follows.

$$\delta W(r, \omega, R, \Omega) = \delta S(r, \omega) \times \delta S(r, \omega, R, \Omega) \quad (4.93)$$

$$\delta W(r, \omega, R, \Omega) = f(r, \omega) \delta \omega F(r, \omega, R, \Omega) \delta \Omega \quad (4.94)$$

Here we introduce the new solid angle.

$$\nu = (\omega, \Omega) \quad (4.95)$$

Here we introduce the new radius.

$$\rho = (r, R) \quad (4.96)$$

Here we introduce the new function.

$$g(\rho, \nu) = f(r, \omega) F(r, \omega, R, \Omega) \quad (4.97)$$

Then we express the wave space-time shortly.

$$\delta W(\rho, \nu) = g(\rho, \nu) \delta \nu \quad (4.98)$$




Figure 4-19: Wave space-time

The spherical harmonics $g(\rho, \nu)$ is the solution of Laplace equation of the spheric polar coordinates. Therefore, the spherical harmonics $g(\rho, \nu)$ satisfies the following harmonic equation.

$$\left(\partial_{\bar{s}}^2 + \partial_s^2 + \partial_{\bar{S}}^2 + \partial_S^2\right)g = 0 \quad (4.99)$$

Here, we used the following symbols.

$$\partial_{\bar{s}} = \frac{\partial}{\partial t} + i \frac{\partial}{\partial x} + j \frac{\partial}{\partial y} + k \frac{\partial}{\partial z} \quad (4.100)$$

$$\partial_s = \frac{\partial}{\partial t} - i \frac{\partial}{\partial x} - j \frac{\partial}{\partial y} - k \frac{\partial}{\partial z} \quad (4.101)$$

$$\partial_{\bar{S}} = \frac{\partial}{\partial t} + I \frac{\partial}{\partial x} + J \frac{\partial}{\partial y} + K \frac{\partial}{\partial Z} \quad (4.102)$$

$$\partial_S = \frac{\partial}{\partial T} - I \frac{\partial}{\partial X} - J \frac{\partial}{\partial Y} - K \frac{\partial}{\partial Z} \quad (4.103)$$

4.5 Elementary event of many-worlds interpretation

In the case of the Copenhagen interpretation, we cannot introduce an elementary event to the quantum theory, because we always observe one event at one observation.

Therefore, we introduce elementary events to quantum mechanics by embracing the Many-Worlds Interpretation (MWI) in this paper. In MWI, all the events those occur in one observation occur. However, one observer cannot observe all the events at the same time, because the observer itself is involved in an event.

If we interpret an event of quantum theory as a set of elementary events, we can derive the probability that an event occurs from the number of elementary events. If the event R or event B occurs in some observations, a world branched to the world that event R occurs and the other world that event B occurs in the MWI.

For example, if the number of elementary events of the event R is three, and B is two, the probability of occurrence of the event R is $3/5$. We call a world that an elementary event occurs the **elementary world**. We interpret a world as a set of elementary worlds.




Figure 4-20: A world is a set of elementary worlds in many-worlds interpretation

The concrete implementation method of elementary events is described in the following sections.

4.6 Elementary state of many-worlds interpretation

In the wave function of a many-particle system in configuration space (many-particle wave function), we call the position certain state that positions of all particles are decided **a position certain state**.

However, in the actual experiment, a particle spreads in the narrow range. Therefore, actual state diffuses in the narrow range in configuration space. We can regard the state as the set of the position certain states. We call the state **a localized state**.




Figure 4-21: Elementary state in configuration space

In addition, we interpret a wave function of the position certain state (position eigenstate) as a manifold. We interpret an absolute value of the wave function as the surface area of the manifold. We put a point on

the surface of the manifold at a fixed interval. We interpret the point as **an elementary state**. The number of elementary events is proportional to an absolute value of the wave function, because the number of elementary events is proportional to the surface area of the manifold.

In the discussion of this paper, there is no difference between the discussion using the many-particle wave function and the discussion using the wave function of one particle. Therefore, in the discussion of this paper, we do not use the many-particle wave function but the wave function of one particle.

4.7 Introduction of an elementary state to the quantum theory

We express the wave function $\psi(x, t)$ by Dirac delta function as follows.

$$\psi(x, t) = \int \psi(y, t) \delta(x - y) dy \tag{4.104}$$

We interpret the state $\psi(y, t)$ as the state that the position y of the particle is fixed, "position certain state." Then we compose the **elementary state** that cannot be separated any more by dividing the "position certain state." K. Sugiyama²⁶ introduced the elementary in 1999.

We divided the virtual high-dimensional Euclidean space by using "elementary domain" and we suppose that a lattice point is an elementary state. The position certain state is 1-sphere or 3-sphere and the lattice point on the sphere is an elementary states of the position certain state.




Figure 4-22: Elementary state of many-worlds interpretation

Since the surface area S of the manifold is the absolute value $|\psi(y, t)|$ of the wave function, we describe the number $M(y, t)$ of the elementary state by using Planck length ℓ_P as follows.

$$M(y, t) = \frac{S}{\ell_P^3} = \frac{|\psi(y, t)|}{\ell_P^3} \tag{4.105}$$

4.8 Application of path integral to the field

In quantum field theory, we quantize the field itself.

We interpret the field is an independent universe according to the hierarchical principle of this paper. Therefore, we apply the path integral to the field.

We were able to apply the path integral to the position x that is the "positional physical quantity." Therefore, we deduce that we can apply the path integral to the field ψ that is "positional physical quantity." Then, we apply the path integral to the following new function.

$$\Psi(x, t, \psi(x, t)) \tag{4.106}$$

There was a network structure of the path integral for the position x that is "positional physical quantity." Therefore, we apply a network structure of the path integral for the field ψ that is "positional physical quantity" like the following figure.




Figure 4-23: Application of network structure of path integral to field itself

We call the space-time that the new wave function Ψ exists is the second universe.

In the above figure, we apply the "network structure of the path integral" to the region that is smaller than $\Delta\psi$.

4.9 Introduction of elementary event to the quantum theory

We introduce a new concept, elementary event to the quantum theory in this paper.

We express an event as a transition from one state to the other state in quantum theory. Therefore, we express an elementary event as a transition from one elementary state to the other elementary state.

We interpret an elementary state as a point. We interpret an elementary event as an arrow from a point to the other point. Since we can draw a line from any point to any point, we deduce that an elementary event from any elementary state to any elementary state exists.

If the arrow from the point A to the point B exists, the arrow from the point B to the point A also exists conversely. If the number of points is M , the number of arrows becomes M^2 . In other words, if the number of elementary states is M , the number of elementary events becomes M^2 .

Though there is no clear evidence of the existence of an elementary event, we deduce it by the following reasons.




Figure 4-24: Elementary event and elementary state of many-worlds interpretation

We assume that an elementary event of quantum theory has the same properties as elementary events of probability theory. In other words, the probability of occurrence of an event is proportional to the number of elementary events those are included in the event.

In addition, we define the event that is a transition from any position certain state to any position certain state "path certain event." The path certain event is a set of elementary events.

Actual states are localized by the uncertainty principle. We call the states localized states. We call the event from any localized state to any localized state a "localized event."

If we apply the path integral to the discrete space-time, the long distance transition from position certain state occurs. However, "long distance transition" is suppressed due to the localized states. It means that the number of elementary events of localized event that is Long distance transition is very rare.

The existence of an elementary state and an elementary event suggests that an existence probability and a probability of occurrence are different concepts. If the number of elementary states of a state is m , the state's existence probability is proportional to m . If the number of elementary events of an event is n , the event's existence probability is proportional to n .

4.10 Derivation of Born's rule

This section describes how to derive this probability.

We express the observation probability $P(x, t)$ of the particle by the wave function $\psi(x, t)$ as follows.

$$P(x, t) = |\psi(x, t)|^2 \quad (4.107)$$

On the other hand, we express the probability P based on the Laplace's calculation method of the probability as follows.

$$P = \frac{N}{N_a} \quad (4.108)$$

Here N_a is the number of all elementary events and N is the number of elementary events those are expected. If N_a is sufficiently larger than N , the probability P is proportional to N .

$$P \propto N \quad (4.109)$$

Actual state is localized state. We apply the "network structure of path integral" to the localized state. Since "long distance transition" does not occur for localized state, the length of transition is small after a minimum time t_P .

The number $M(x', t')$ of elementary events of the localized state $\psi(x', t')$ Δx is proportional to the surface area of the manifold. We apply the "network structure of path integral" to the position on the surface area of the manifold.

Since the manifold after the minimum time almost same as the original manifold, we approximate it by the same manifold. We express the number $M(x, t)$ of elementary states on the surface area of the manifold as follows.

$$M(x, t) = \frac{S \Delta x}{\ell_P^3 \ell_P} = \frac{|\psi| \Delta x}{\ell_P^4} \quad (4.110)$$

Therefore, the number of the elementary state is proportional to the absolute value of the wave function if the Δx is almost constant.

$$M(x, t) \propto |\psi| \quad (4.111)$$

According to the uncertainty principle, the deviation Δp of momentum is almost constant if Δx is almost constant. Therefore, the number M of the elementary state approximately does not change.

$$M(x, t) \sim M(x, t + t_p) \quad (4.112)$$

The number N of elementary events is the number of transitions from all elementary states at time t' to all elementary states at time t'' . Therefore, the number N of elementary events is the square of the number M of elementary states.

$$N = M^2 \quad (4.113)$$

We express those elementary events in the following figure.




Figure 4-25: The number of elementary events is the square of the number of elementary states.

The probability P is proportional to the number of the elementary events N .

$$P \propto N \quad (4.114)$$

On the other hand, the number N of the elementary events is equal to the square of the number M of elementary states.

$$N = M^2 \quad (4.115)$$

The number M of elementary events is proportional to the absolute value of the wave function.

$$M \propto |\psi| \quad (4.116)$$

Therefore, the probability P is proportional to the square of the absolute value of the wave function.

$$P \propto |\psi|^2 \quad (4.117)$$

We derived the Born's rule by the above method.

5 Conclusion

We explained the method to derive Born's rule from many-worlds interpretation and probability theory.

Probability is proportional to the number of elementary events. The number of elementary events is the square of the number of elementary states because we apply the "network structure of path integral" to the elementary state. The number of elementary states is proportional to the absolute value of the wave function. Therefore, the probability is proportional to the absolute value of the wave function.

6 Supplement

6.1 Supplement of the many-particle wave function

We call an elementary state, a position certain state and a localized state for the universe "elementary world," "position certain world" and "localized world" respectively.

In addition, we call an elementary event, a path certain event and a localized event for the universe "elementary history", "path certain history" and "localized history" respectively.




Figure 6-1: Elementary history and elementary world of many-worlds interpretation

We interpret one point of the configuration space of the many-particle wave function as the state that the positions of all particles are determined. The state is "position certain world."

In the view of classical mechanics, the point is our world. In the view of the quantum mechanics, localized world is our world.

I guess that the absolute value of the many-particle wave function of the universe is most nearly zero in the almost area. The domain that the absolute value is large is localized like a network structure.

6.2 Supplement of the method of deriving Born's rule

The simplest way to derive Born's rule from Many-Worlds Interpretation (MWI) is that we connect the number of worlds to the probability.

If the probability of occurrence of event A is higher than the probability of occurrence of event B , we deduce that the number of worlds those event A occurred is greater than the number of worlds those event B occurred.

For example, we suppose that we make the 100 planets those are exactly same as Earth. If the event A occurred on 80 planets and the event B occurred on 20 planets, then we interpret that the probability of the occurrence of the event A is 80%.

However, it is not clear how to count the world. Therefore, we count the number of elementary worlds of the localized world that event A occurred.

We express the number M of elementary worlds of the localized world by the wave function ψ that event A occurred as follows.

$$M = \frac{|\psi| (\Delta x)^{3n}}{\ell_p^3 \ell_p^{3n}} = \frac{|\psi| (\Delta x)^{3n}}{\ell_p^{3+3n}} \quad (6.1)$$

Δx is the position deviation, and n is the number of all particles. The number of elementary worlds is proportional to the absolute value of the wave function.

$$M \propto |\psi| \quad (6.2)$$

On the other hand, the probability is proportional to the square of the absolute value of the wave function. Therefore, we cannot explain the probability by using the number of the elementary worlds.

To solve this problem, we explain the probability by using the number of histories. We express the number N of elementary histories of the localized history that event A occurred as follows.

$$N = M^2 \quad (6.3)$$

The probability is proportional to the number of elementary histories.

$$P \propto N \quad (6.4)$$

The number of histories is the square of the number of elementary worlds.

$$N = M^2 \quad (6.5)$$

On the other hand, the number of the elementary worlds is proportional to the absolute values of the wave function.

$$M \propto |\psi| \quad (6.6)$$

Therefore, the probability is proportional to the square of the absolute value of the wave function.

$$P \propto |\psi|^2 \quad (6.7)$$

6.3 Supplement of basis problem in many-worlds interpretation

In many-worlds interpretation, the absence of a particular basis of the wave function is a problem.

For example, we consider the Stern-Gerlach experiment of the spin of electrons. In this experiment, we measure the spin by using a magnetic field gradient. Since the basis of the spin is determined by the direction of the gradient magnetic field, there is no particular basis for the spin.

In this paper, we chose position as the particular basis. We could also choose the momentum as the particular basis, but we did not do so, because we express the basis of the momentum by using a set of elementary states those the positions are basis.

For spin, there is no way to select a particular basis. In this paper, we are considering the manifold of a particle of spin 1/2. We might be able to express the spin by using the manifold.

6.4 Interpretation of time in many-worlds interpretation

The position of all particles is different for a point in the configuration space of many-particle wave function. Therefore, we define the time for a point in the configuration space. Since a point corresponds to a position certain world, we interpret the time as a parameter to classify the position certain worlds.

A position certain world transits the minimum length continuously in the configuration space. I guess that we feel the transition as a time.




Figure 6-2: Many-worlds interpretation and arrow of time

If a transition of a direction exists, the transition of the opposite direction also exists. However, since there are many "elementary worlds" of future more overwhelmingly than the number of elementary worlds of past, we feel that our elementary world always transits to elementary world of the future. In this way, many-worlds interpretation explains the arrow of time by.

6.5 Supplement of Long distance transition

In this paper, we have been thinking about one particle is localized in one place. Here we consider the wave function of one particle that was localized in one place at a time. We suppose that the wave function was separated and localized in two places. We call the state "many localized states." In this case, what would happen?

Elementary event exists between any two elementary states. The world does not become disorder because long distance transition is suppressed due to the "localized state." We determine the number of elementary events between two localized states only by the number of elementary states of the two localized states.

Therefore, if there are "many localized states," the transition between the states those are localized in two places will occur.




Figure 6-3: Long distance transition between localized states

I call the phenomenon "localized long distance transition" or "localized shift."

Then, will localized shift between localized states those have different time occur?

In this case, since the elementary event exists between any two elementary states, the localized shift occurs, too.

I do not deduce that the localized teleport send information, because we cannot send any information by using EPR correlation.

6.6 The solution to the basis problem of MWI

The basis problem is a problem that the status of a world changes by choosing the basis of spin. The cause of the problem is to count a world that the spin has one direction as one world.

I showed that we can derive the spin from the rotation of 3-sphere in the following paper.

- Derivation of two-valuedness and angular momentum of spin-1/2 from rotation of 3-sphere [2013/5]
http://www.geocities.jp/x_seek/Spin_e.htm

This means that a world that the spin has one direction is constructed from many worlds. Choosing the basis of spin is equivalent to changing the rotational axis of the 3-sphere.

Therefore, we can solve the basis problem of MWI by the rotation of 3-sphere.

7 Future Issues

Future issues are shown as follows.

- (1) Consideration of the principle
- (2) Formulation for the quantum field theory
- (3) Consideration of the discrete space
- (4) Formulation for the relativistic mechanics
- (5) Formulation for the gravity theory

We consider some of these issues in the following chapters.

8 Consideration of the future issues

8.1 Consideration of principles

We consider the hierarchical principle and the event principle.

8.1.1 Hierarchical principle

I propose the following **hierarchical principle**.

- (1) Wave functions are quaternionic functions.
- (2) The direct product of the closed path of a particle and the wave function is the other universe.
- (3) Wave functions in the other universe are also quaternionic functions.

We call the theory based on the hierarchy principle the **hierarchy theory**.

8.1.2 Event principle

I propose the following **event principle**.

- (1) An elementary event is the transition from an elementary state to the other elementary state.
- (2) Event probability of an event is proportional to the number of elementary events those the event includes.

We call the theory based on the event principle the **event theory**.

8.2 Consideration of formulation for the quantum field theory

A position certain state has a phase and an absolute value of the wave function. Therefore, it is possible to use “suppression of long distance transition due to localized states” for the position certain state. On the other hand, an elementary state does not have a phase and an absolute value of the wave function. Therefore, it is impossible to use “suppression of long distance transition due to localized states” for the elementary state. In order to solve the problem, we consider the quantum field theory.

In the quantum mechanics, the position and the momentum of a particle have a commutation relation. It means that the position of the particle is distributed. On the other hand, in the quantum field theory the amplitude and the general momentum of the wave function have a commutation relation. It means that the amplitude of the wave function is distributed.

Then I propose the following new function.

$$\Psi(x, \psi(x)) \tag{8.1}$$

We call the function **the second wave function** because we get the wave function by the second quantization of the field. The second wave function exists in the second universe. The elementary state of the first universe is the position certain state of the second universe. Therefore, it is possible to use “suppression of long distance transition due to localized states” for the elementary state.

8.3 Consideration of discrete space

8.3.1 Discrete space from 24-hypercube

In this paper, we call the discrete space of the elementary state **an event space**.

For example, we consider the 24-hypercube as the model of the event space. 24-hypercube γ_{24} is the direct product of the 24 one-dimensional cube γ_1 .

$$\gamma_{24} = \gamma_1 \times \gamma_1 \times \gamma_1 \times \cdots \times \gamma_1 \quad (8.2)$$

We show the vertices, edges, faces, and cells of the 24-hypercube as below.

Table 8-1: The number of vertices, edges, faces, and cells of the 24-hypercube

Event space	Faces	Number
E_0	Vertices	$\binom{24}{0} 2^{24-0}$
E_1	Edges	$\binom{24}{1} 2^{24-1}$
E_2	Faces	$\binom{24}{2} 2^{24-2}$
E_3	3-faces (Cells)	$\binom{24}{3} 2^{24-3}$
E_k	k -faces	$\binom{24}{k} 2^{24-k}$
E_{23}	23-faces	$\binom{24}{23} 2^{24-23}$
E_{24}	24-faces	$\binom{24}{24} 2^{24-24}$

We use the following abbreviation.

E-state: Elementary state

E-event: Elementary event

(3) 0-event space: Vertices is e-state. Edge is e-event and vertices in the 1-event space.

(4) 1-event space: Vertices is e-state. Edge is e-event and vertices in the 2-event space.

(5) 2-event space: Vertices is e-state. Edge is e-event and vertices in the 3-event space.

We construct the following sequence of the event space by repeating the above process.

$$E_0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow \cdots \rightarrow E_{24} \quad (8.3)$$

8.3.2 Discrete space Consideration from finite group

We construct the discrete space from the finite group.

The representation on the vector space V of the group G is the map from the group G to the general linear matrix $GL(V)$.

$$\rho: G \rightarrow GL(V) \quad (8.4)$$

$$\rho(gh) = \rho(g)\rho(h) \quad (8.5)$$

$$g, h \in G \quad (8.6)$$

$$\rho(g), \rho(h) \in \rho(G) \quad (8.7)$$

We use the symbol g as the abbreviation for the representation $\rho(g)$ of the element of the group in this paper. We use the symbol G as the abbreviation for the set $\rho(G)$ of the representation of the element of the group.

We consider one fixed vector v_1 in the vector space V . We transform the vector v_1 to the vector v by the element g of the finite group G . Then vector v and element g have one-to-one onto mapping. Therefore, we interpret the element g as the vector v .

$$v_1 \in V \quad (8.8)$$

$$g \in G \quad (8.9)$$

$$v = gv_1 \quad (8.10)$$

We interpret an element g as an elementary state. We call the group **world group**.

We consider the element of the direct product of the two world groups.

$$(g_1, g_2) \in G \times G \quad (8.11)$$

We interpret an element as an elementary event.

We construct a new group as the direct product of two world groups.

$$H = G \times G \quad (8.12)$$

We call the group **history group**.

8.3.3 Consideration of the uncertain status

We consider the uncertain status.

We define **certain information entropy** H for the observed information as follows.
(Certain information entropy)

$$H = - \sum_{k=1}^n P(x_k) \log(P(x_k)) \quad (8.13)$$

We define the **uncertain information entropy** Q for the unobserved wave function $f(x)$ as follows.
(Uncertain information entropy)

$$P(x) = |f(x)|^2 \quad (8.14)$$

$$Q = - \int P(x) \log(P(x)) dx \quad (8.15)$$

We define uncertain information entropy Q for the unobserved angular momentum of the k -th particle as follows.

(Uncertain information entropy)

$$P_1(k) = |\langle f|x+\rangle + \langle f|x-\rangle|^2 \quad (8.16)$$

$$P_2(k) = |\langle f|y+\rangle + \langle f|y-\rangle|^2 \quad (8.17)$$

$$P_3(k) = |\langle f|z+\rangle + \langle f|z-\rangle|^2 \quad (8.18)$$

$$Q = - \sum_{s=1}^3 \sum_{k=1}^n P_s(k) \log(P_s(k)) \quad (8.19)$$

We define the **general information entropy** G .

(General information entropy)

$$G = H + Q \quad (8.20)$$

This general information entropy conserves.

(Law of general information entropy conservation)

$$\delta G = \delta H + \delta Q = 0 \quad (8.21)$$

Certain information entropy always increases by a thermodynamics second law.

(Law of entropy increase)

$$\delta H > 0 \quad (8.22)$$

However, certain information entropy has the following upper limit because of uncertainty principle.

(Upper limit of certain information entropy)

$$H < \frac{1}{3} Q \quad (8.23)$$

All the particle's angular momentums of y -direction and z -direction become uncertain, when all the particle's angular momentums of x -direction are observed.

Certain information entropy increases when the wave function collapses.

Uncertain information entropy increases when the wave function diffuses (anti-collapses).

If we make a status uncertain status, uncertain information entropy increases.

8.3.4 Time in the hierarchical universe

To consider time in the hierarchy universe, we consider the one-dimensional universe that has only one photon. The circumference is 1 meter. We make a stationary state of a photon in the universe.

We suppose that the speed of light and the Planck's constant is one.

Table 8-2: Constant

Quantity	Before expansion	After expansion
Speed of light	1 [m/s]	1 [m/s]
Planck's constant	1 [g m ² /s]	1 [g m ² /s]

Now, we suppose that the universe expands slowly. The circumference is 2 meters. The each quantity changes as follows.

Table 8-3: Variation of the momentum of photon in the expansion of the universe

Quantity	Before expansion	After expansion
Circumference of the universe	1 [m]	2 [m]
Wave number of the photon	1	1
Wave length of the photon	1 [m]	2 [m]
Frequency of the photon	1[1/s]	0.5[1/s]
Momentum of the photon	1 [g m/s]	0.5 [g m/s]
Time to go around the universe	1[s]	2 [s]

I deduce the frequency of the photon is proportional to the radius of the universe from the above table.

Next, we consider the one-dimensional universe that has only one electron. The circumference is 1 meter. We make a stationary state of an electron in the universe. For easy calculation, we suppose that the electron's mass is 1 gram.

Now, we suppose that the universe expands slowly. The circumference is 2 meters. The each quantity changes as follows.

Table 8-4: Variation of the momentum of electron in the expansion of the universe

Quantity	Before expansion	After expansion
Circumference of the universe	1 [m]	2 [m]
Wave number of the electron	1	1
Wave length of the electron	1 [m]	2 [m]
Frequency of the electron	0.1[1/s]	0.05[1/s]
Momentum of the electron	0.1 [g m/s]	0.05 [g m/s]
Time to go around the universe	10[s]	20 [s]
Velocity of the electron	0.1 [m/s]	0.1 [m/s]
Mass of the electron	1[g]	0.5[g]

I deduce the mass of the electron is reverse proportional to the radius of the universe from the above table.

9 Appendix

9.1 Definition of Terms

We define terms in the following table.

Table 9-1: Normal space and so on

Term	Definition
Normal space	Three-dimensional normal space
Normal space-time	Fore-dimensional normal space-time
Closed path	Closed path of the particle generated by pair production and destroyed by pair annihilation

Table 9-2: Elementary domain and so on

Term	Definition
Elementary domain	The minimum domain of the wave space
Elementary position	Position of the wave space
Elementary path	An arrow from any elementary position to any elementary position
Normal domain	The minimum domain of the normal space
Normal position	Position of the normal space
Normal path	An arrow from any normal position to any normal position

Table 9-3: Elementary state and so on

Term	Definition
Elementary state	Point of the wave space
Position certain state	State having a certain position (position eigenstate)
Localized state	State that the distribution is a normal distribution
Elementary event	A transition from any elementary state to any elementary state
Path certain event	A transition from any position certain state to any position certain state
Localized event	A transition from any localized state to any localized state
Elementary world	Elementary state of the universe
Position certain world	Position certain state of the universe
Localized world	Localized states of the universe
Elementary history	Elementary event of the universe
Path certain history	Path certain event of the universe
Localized history	Localized events of the universe

Table 9-4: Localized displacement and so on

Term	Definition
Localized displacement	Localized transition of short-distance
Localized transition	Localized transition
Localized shift	Localized transition of long distance
Localized teleportation	Localized transition of ultra-long distance

9.2 Arrangement of Terms

We arrange terms in the following table.

Table 9-5: Elementary domain and so on

Category	Elementary	Normal
Domain	Elementary domain	Normal domain
Position	Elementary position	Normal position
Path	Elementary path	Normal path

Table 9-6: Elementary state and so on

Category	Elementary	Certain	Localized
State	Elementary state	Position certain state	Localized state
Event	Elementary event	Path certain event	Localized event
World	Elementary world	Position certain world	Localized world
History	Elementary history	Path certain history	Localized history

Table 9-7: First normal space-time and so on

Category	First universe	Second universe	Third universe
Normal space-time	First normal space-time	Second normal space-time	Third normal space-time
Particle	First particle	Second particle	Third particle
Position	First position	Second position	Third position
Path	First path	Second path	Third path
Wave function	First wave function	Second wave function	Third wave function

10 Acknowledgment

In writing this paper, I thank from my heart to TY and NS who gave valuable advice to me.

11 References

-
- 1 Mail: sugiyama_xs@yahoo.co.jp
- 2 Hugh Everett III, "Relative state" formulation of quantum mechanics, *Reviews of Modern Physics*, 29 (3):454-462, (1957).
- 3 James B. Hartle, 'Quantum Mechanics of Individual Systems', *American Journal of Physics*, 36, 704-712, (1968).
- 4 Bryce S. DeWitt (1970), 'Quantum mechanics and Reality', *Physics Today* 23, No. 9, pp. 30-35.
- 5 Neil Graham (1973), 'The Measurement of Relative Frequency', in De Witt and N. Graham (eds.) *The Many-Worlds Interpretation of Quantum Mechanics*, Princeton NJ: Princeton University Press.
- 6 Adrian Kent, "Against Many-Worlds Interpretations", *Int. J. Mod. Phys. A*5 (1990) 1745, (<http://arxiv.org/abs/grqc/9703089>).
- 7 David Deutsch (1999), "Quantum Theory of Probability and Decisions", *Proceedings of the Royal Society of London A* 455, 3129-3137, (<http://arxiv.org/abs/quant-ph/9906015>).
- 8 Sumio Wada, "Derivation of the Quantum Probability Rule without the Frequency Operator", *J. Phys. Soc. Jpn*, Vol.76, 094004 (2007), (<http://arxiv.org/abs/0707.1623>).
- 9 Max Born (1926), *Zur Quantenmechanik der Stoßvorgänge*, *Zeitschrift für Physik*, 37, #12, pp. 863-867 (German); English translation, *On the quantum mechanics of collisions*, in *Quantum theory and measurement*, section I.2, J. A. Wheeler and W. H. Zurek, eds., Princeton, NJ: Princeton University Press, 1983, ISBN 0-691-08316-9.
- 10 Andrew M. Gleason (1957) "Measures on the Closed Subspaces of Hilbert Space", *Journal of Mathematics and Mechanics* 6, 885-894.
- 11 Bryce S. DeWitt. (1967). "Quantum Theory of Gravity. I. The Canonical Theory". *Phys. Rev.* 160 (5): 1113-1148.
- 12 Julian Barbour (1999) "The End of Time ", W&N.
- 13 Pierre-Simon de Laplace (1812), "Théorie analytique des probabilités", Paris: Courcier Imprimeur.
- 14 Roger Penrose (1971), "Angular momentum: An approach to combinatorial space-time" in *Quantum Theory and Beyond* T. Bastin, ed. (Cambridge University Press, Cambridge).
- 15 Richard P. Feynman (1948). "The Space-Time Approach to Non-Relativistic Quantum Mechanics". *Reviews of Modern Physics* 20 (2): 367-387.
- 16 Paul A.M. Dirac (1927), "The Quantum Theory of the Emission and Absorption of Radiation". *Proceedings of the Royal Society of London A* 114: 243-265.
- 17 I. J. R. Aitchison (1989), A. J. G. Hey, *Gauge Theories in Particle Physics*, Second Edition, (Taylor & Francis; 2 edition).
- 18 Kaluza T., *On the Unification Problem of Physics (Zum Unitätsproblem in der Physik)*, *Sitzungsber. Preuss. Akad. Wiss. Berlin. (Math. Phys.)*, 96, 69-72, (1921).
- 19 Klein O., *Quantum theory and five dimensional theory of relativity (Quantentheorie und fünfdimensionale Relativitätstheorie)*, *Zeitschrift für Physik*, A37(12), 895-906, (1926).
- ²⁰ Cauchy, A.L. (1814), *Mémoire sur les intégrales définies*, *Oeuvres complètes Ser. 1* 1, Paris (published 1882), pp. 319-506.
- ²¹ Riemann, B. (1851), "Grundlagen für eine allgemeine Theorie der Funktionen einer veränderlichen komplexen Grösse", in H. Weber, *Riemann's gesammelte math. Werke*, Dover (published 1953), pp. 3-48.
- ²² Hamilton, William Rowan. [On quaternions, or on a new system of imaginaries in algebra](#). *Philosophical Magazine*. Vol. 25, n 3. p. 489-495. 1844.
- ²³ Fueter R.: *Die Funktionentheorie der Differentialgleichungen $\Delta u=0$ und $\Delta\Delta u=0$ mit vier reellen Variablen*. *Comment. Math. Helv.* 7 (1934), 307-330.
- ²⁴ A. Sudbery (1979) "[Quaternionic Analysis](#)", *Mathematical Proceedings of the Cambridge Philosophical Society* 85:199-225.
- 25 Elie Cartan (1899), "Sur certaines expressions différentielles et le problème de Pfaff", *Annales scientifiques de l'Ecole Normale Supérieure*: 239-332.
- 26 K. Sugiyama (1999), "Quantum probability from a geometrical interpretation of a wave function", (<http://arxiv.org/abs/quant-ph/9910108>).