

Numerical Solution of Radial Biquaternion Klein-Gordon Equation

Vic Christianto* and Florentin Smarandache†

*Sciprint.org — a Free Scientific Electronic Preprint Server, <http://www.sciprint.org>

E-mail: admin@sciprint.org

†Department of Mathematics, University of New Mexico, Gallup, NM 87301, USA

E-mail: smarand@unm.edu

In the preceding article we argue that biquaternionic extension of Klein-Gordon equation has solution containing imaginary part, which differs appreciably from known solution of KGE. In the present article we present numerical /computer solution of radial biquaternionic KGE (radialBQKGE); which differs appreciably from conventional Yukawa potential. Further observation is of course recommended in order to refute or verify this proposition.

1 Introduction

In the preceding article [1] we argue that biquaternionic extension of Klein-Gordon equation has solution containing imaginary part, which differs appreciably from known solution of KGE. In the present article we presented here for the first time a numerical/computer solution of radial biquaternionic KGE (radialBQKGE); which differs appreciably from conventional Yukawa potential.

This biquaternionic effect may be useful in particular to explore new effects in the context of low-energy reaction (LENR) [2]. Nonetheless, further observation is of course recommended in order to refute or verify this proposition.

2 Radial biquaternionic KGE (radial BQKGE)

In our preceding paper [1], we argue that it is possible to write biquaternionic extension of Klein-Gordon equation as follows:

$$\left[\left(\frac{\partial^2}{\partial t^2} - \nabla^2 \right) + i \left(\frac{\partial^2}{\partial t^2} - \nabla^2 \right) \right] \varphi(x, t) = -m^2 \varphi(x, t), \quad (1)$$

or this equation can be rewritten as:

$$(\diamond \diamond + m^2) \varphi(x, t) = 0, \quad (2)$$

provided we use this definition:

$$\diamond = \nabla^q + i \nabla^q = \left(-i \frac{\partial}{\partial t} + e_1 \frac{\partial}{\partial x} + e_2 \frac{\partial}{\partial y} + e_3 \frac{\partial}{\partial z} \right) + i \left(-i \frac{\partial}{\partial T} + e_1 \frac{\partial}{\partial X} + e_2 \frac{\partial}{\partial Y} + e_3 \frac{\partial}{\partial Z} \right), \quad (3)$$

where e_1, e_2, e_3 are *quaternion imaginary units* obeying (with ordinary quaternion symbols: $e_1 = i, e_2 = j, e_3 = k$):

$$\begin{aligned} i^2 = j^2 = k^2 = -1, \quad ij = -ji = k, \\ jk = -kj = i, \quad ki = -ik = j. \end{aligned} \quad (4)$$

and quaternion *Nabla operator* is defined as [1]:

$$\nabla^q = -i \frac{\partial}{\partial t} + e_1 \frac{\partial}{\partial x} + e_2 \frac{\partial}{\partial y} + e_3 \frac{\partial}{\partial z}. \quad (5)$$

(Note that (3) and (5) included partial time-differentiation.)

In the meantime, the standard Klein-Gordon equation usually reads [3, 4]:

$$\left(\frac{\partial^2}{\partial t^2} - \nabla^2 \right) \varphi(x, t) = -m^2 \varphi(x, t). \quad (6)$$

Now we can introduce polar coordinates by using the following transformation:

$$\nabla = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) - \frac{\ell^2}{r^2}. \quad (7)$$

Therefore, by substituting (7) into (6), the radial Klein-Gordon equation reads — by neglecting partial-time differentiation — as follows [3, 5]:

$$\left(\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) - \frac{\ell(\ell+1)}{r^2} + m^2 \right) \varphi(x, t) = 0, \quad (8)$$

and for $\ell = 0$, then we get [5]:

$$\left(\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + m^2 \right) \varphi(x, t) = 0. \quad (9)$$

The same method can be applied to equation (2) for radial biquaternionic KGE (BQKGE), which for the 1-dimensional situation, one gets instead of (8):

$$\left(\frac{\partial}{\partial r} \left(\frac{\partial}{\partial r} \right) - i \frac{\partial}{\partial r} \left(\frac{\partial}{\partial r} \right) + m^2 \right) \varphi(x, t) = 0. \quad (10)$$

In the next Section we will discuss numerical/computer solution of equation (10) and compare it with standard solution of equation (9) using Maxima software package [6]. It can be shown that equation (10) yields potential which differs appreciably from standard Yukawa potential. For clarity, all solutions were computed in 1-D only.

3 Numerical solution of radial biquaternionic Klein-Gordon equation

Numerical solution of the standard radial Klein-Gordon equation (9) is given by:

$$\begin{aligned} (\%i1) & \text{diff}(y,t,2) - \text{diff}(y,r,2) + m^2 y; \\ (\%o1) & m^2 \cdot y - \frac{d^2}{dx^2} y \\ (\%i2) & \text{ode2}(\%o1, y, r); \\ (\%o2) & y = \%k_1 \cdot \% \exp(mr) + \%k_2 \cdot \% \exp(-mr) \end{aligned} \quad (11)$$

In the meantime, numerical solution of equation (10) for radial biquaternionic KGE (BQKGE), is given by:

$$\begin{aligned} (\%i3) & \text{diff}(y,t,2) - (\%i+1) \cdot \text{diff}(y,r,2) + m^2 y; \\ (\%o3) & m^2 \cdot y - (i+1) \frac{d^2}{d^2 r} y \\ (\%i4) & \text{ode2}(\%o3, y, r); \\ (\%o4) & y = \%k_1 \cdot \sin\left(\frac{|m|r}{\sqrt{-\%i-1}}\right) + \%k_2 \cdot \cos\left(\frac{|m|r}{\sqrt{-\%i-1}}\right) \end{aligned} \quad (12)$$

Therefore, we conclude that numerical solution of radial biquaternionic extension of Klein-Gordon equation yields different result compared to the solution of standard Klein-Gordon equation; and it *differs appreciably* from the well-known Yukawa potential [3, 7]:

$$u(r) = -\frac{g^2}{r} e^{-mr}. \quad (13)$$

Meanwhile, Comay puts forth argument that the Yukawa lagrangian density has theoretical inconsistency within itself [3].

Interestingly one can find argument that biquaternion Klein-Gordon equation is nothing more than quadratic form of (modified) Dirac equation [8], therefore BQKGE described herein, i.e. equation (12), can be considered as a plausible solution to the problem described in [3]. For other numerical solutions to KGE, see for instance [4].

Nonetheless, we recommend further observation [9] in order to refute or verify this proposition of new type of potential derived from biquaternion Klein-Gordon equation.

Acknowledgement

VC would like to dedicate this article for RFF.

Submitted on November 12, 2007
Accepted on November 30, 2007

References

1. Yefremov A., Smarandache F. and Christianto V. Yang-Mills field from quaternion space geometry, and its Klein-Gordon representation. *Progress in Physics*, 2007, v. 3, 42–50.
2. Storms E. <http://www.lenr-canr.org>
3. Comay E. *Apeiron*, 2007, v. 14, no. 1; arXiv: quant-ph/0603325.
4. Li Yang. Numerical studies of the Klein-Gordon-Schrödinger equations. MSc thesis submitted to NUS, Singapore, 2006, p. 9 (<http://www.math.nus.edu.sg/~bao/thesis/Yang-li.pdf>).
5. Nishikawa M. A derivation of electroweak unified and quantum gravity theory without assuming Higgs particle. arXiv: hep-th/0407057, p. 15.
6. Maxima from <http://maxima.sourceforge.net> (using GNU Common Lisp).
7. http://en.wikipedia.org/wiki/Yukawa_potential
8. Christianto V. A new wave quantum relativistic equation from quaternionic representation of Maxwell-Dirac equation as an alternative to Barut-Dirac equation. *Electronic Journal of Theoretical Physics*, 2006, v. 3, no. 12.
9. Gyulassy M. Searching for the next Yukawa phase of QCD. arXiv: nucl-th/0004064.