

Numerical Solution of N^{th} -Order Fuzzy Differential Equation by Runge-Kutta Nystrom Method

T. Jayakumar, K. Kanagarajan, S. Indrakumar

Department of Mathematics
Sri Ramakrishna Mission Vidyalaya College of Arts and science
Coimbatore-641 020, Tamilnadu, India.
jayakumar.thippan68@gmail.com, kanagaraank@gmail.com
indrakumar1729@gmail.com

Abstract : In this paper we study a numerical solution for N^{th} -order fuzzy differential equations based on Seikkala derivative with initial value problem. The Runge-Kutta Nystrom method is used for the numerical solution of this problem and the convergence and stability of the method is proved. By this method, we can obtain strong fuzzy solution. This method is illustrated by solving examples.

Keywords : Fuzzy Numbers, N^{th} -order FDEs, RK-Nystrom method.

1 Introduction

The topic of fuzzy differential equations (FDEs) have been rapidly growing in recent years. The concept of fuzzy derivative was first introduced by Chang and Zadeh [10], it was followed up by Dubois and Prade [12] by using the extension principle in their approach. Other methods have been discussed by Puri and Ralescu [26] and Goetschel and Voxman [15]. Kandel and Byatt [25] applied the concept of fuzzy differential equation (FDE) to the analysis of fuzzy dynamical problems. The FDE and the initial value problem (Cauhy problem) were rigorously treated by Kaleva [21, 22], Seikkala [28], He and Yi [16], Kloeden [23] and by other researchers (see [6, 20]). The numerical methods for solving fuzzy differential equations are introduced in [1, 2, 3]. Buckley and Feuring [9] introduced two analytical methods for solving N^{th} -order linear differential equations with fuzzy initial value conditions. Their first method of solution was to fuzzy if the crisp solution and then check to see if it satisfies the differential equation with fuzzy initial value conditions; and the second method was the reverse of the first method, they first solved the fuzzy initial value condition and the checked to see if it defined a fuzzy function.

In this paper, a numerical method to solve N^{th} -order linear differential equations with fuzzy initial conditions is presented. The structure of the paper is organized as follows: In Section 2, we give some basic results on fuzzy numbers and define a fuzzy derivative and a fuzzy integral. then the fuzzy initial values is treated in Section 3 using the extension principle of Zadeh and the concept of fuzzy derivative. It is shown that the fuzzy initial value problem has a unique fuzzy solution when f satisfies Lipschitz condition which guarantees a unique solution to the deterministic initial value problem. In Section 4, the Runge-Kutta

Nystrom method for solving N^{th} -order fuzzy differential equations is introduced. In Section 5 Convergence and Stability are illustrated. In Section 6 the proposed method is illustrated by solving several examples, and the conclusion is drawn in section 7.

2 Preliminaries

An arbitrary fuzzy number is represented by an ordered pair of functions $(\underline{u}(r), \bar{u}(r))$ for all $r \in [0, 1]$, which satisfy the following requirements [12]:

- (i) $\underline{u}(r)$ is a bounded left continuous non-decreasing function over $[0, 1]$,
- (ii) $\bar{u}(r)$ is a bounded right continuous non-increasing function over $[0, 1]$,
- (iii) $\underline{u}(r) \leq \bar{u}(r) \forall r \in [0, 1]$,

let E be the set of all upper semi-continuous normal convex fuzzy numbers with bounded α -level intervals.

Lemma 2.1

Let $[\underline{v}(\alpha), \bar{v}(\alpha)]$, $\alpha \in (0, 1]$ be a given family of non-empty intervals. If

$$(i) [\underline{v}(\alpha), \bar{v}(\alpha)] \supset [\underline{v}(\beta), \bar{v}(\beta)] \text{ for } 0 < \alpha \leq \beta,$$

and

$$(ii) \left[\lim_{k \rightarrow \infty} \underline{v}(\alpha_k), \lim_{k \rightarrow \infty} \bar{v}(\alpha_k) \right] = [\underline{v}(\alpha), \bar{v}(\alpha)],$$

whenever (α_k) is a non-decreasing sequence converging to $\alpha \in (0, 1]$, then the family $[\underline{v}(\alpha), \bar{v}(\alpha)]$, $\alpha \in (0, 1]$, represent the α -level sets of a fuzzy number $v \in E$. Conversely if $[\underline{v}(\alpha), \bar{v}(\alpha)]$, $\alpha \in (0, 1]$, are α -level sets of a fuzzy number $v \in E$, then the conditions (i) and (ii) hold true.

Definition 2.1

Let I be a real interval. A mapping $v : I \rightarrow E$ is called a fuzzy process and we denoted the α -level set by $[v(t)]_\alpha = [\underline{v}(t, \alpha), \bar{v}(t, \alpha)]$. The Seikkala derivative $v'(t)$ of v is defined by

$$[v(t)]_\alpha = [\underline{v}'(t, \alpha), \bar{v}'(t, \alpha)],$$

provided that is a equation defines a fuzzy number $v'(t) \in E$.

Definition 2.2

Suppose u and v are fuzzy sets in E . Then their Hausdroff

$$D : E \times E \rightarrow R_+ \cup \{0\},$$

$$D(u, v) = \sup_{\alpha \in [0, 1]} \max \left\{ |\underline{u}(\alpha) - \underline{v}(\alpha)|, |\bar{u}(\alpha) - \bar{v}(\alpha)| \right\},$$

i.e., $D(u, v)$ is maximal distance between α -level sets of u and v .

3 Fuzzy Initial Value Problem

Now we consider the initial value problem

$$\begin{cases} x^{(n)}(t) = \psi(t, x, x', \dots, x^{(n-1)}), \\ x(0) = a_1, \dots, x^{(n-1)}(0) = a_n, \end{cases} \quad (1)$$

where ψ is a continuous mapping from $R_+ \times R^n$ into R and a_i ($0 \leq i \leq n$) are fuzzy numbers in E . The mentioned n^{th} order fuzzy differential equation by changing variables

$$y_1(t) = x(t), y_2(t) = x'(t), \dots, y_n(t) = x^{(n-1)}(t),$$

converts to the following fuzzy system

$$\begin{cases} y_1'(t) = f_1(t, y_1, \dots, y_n), \\ \vdots \\ y_n'(t) = f_n(t, y_1, \dots, y_n), \\ y_1(0) = y_1^{[0]} = a_1, \dots, y_n(0) = y_n^{[0]} = a_n, \end{cases} \quad (2)$$

where f_i ($1 \leq i \leq n$) are continuous mapping from $R_+ \times R^n$ into R and $y_i^{[0]}$ are fuzzy numbers in E with α -level intervals

$$[y_i^{[0]}]_\alpha = [\underline{y}_i^{[0]}(\alpha), \bar{y}_i^{[0]}(\alpha)] \quad \text{for } i = 1, \dots, n, \text{ and } 0 < \alpha \leq 1.$$

We call $y = (y_1, \dots, y_n)^t$ is a fuzzy solution of (2) on an interval I , if

$$\underline{y}'_i(t, \alpha) = \min\{f_i(t, u_1, \dots, u_n); u_j \in [\underline{y}_j(t, \alpha), \bar{y}_j(t, \alpha)]\} = \underline{f}_i(t, y(t, \alpha)), \quad (3)$$

$$\bar{y}'_i(t, \alpha) = \max\{f_i(t, u_1, \dots, u_n); u_j \in [\underline{y}_j(t, \alpha), \bar{y}_j(t, \alpha)]\} = \bar{f}_i(t, y(t, \alpha)), \quad (4)$$

and

$$\underline{y}_i(0, \alpha) = \underline{y}_i^{[0]}(\alpha), \quad \bar{y}_i(0, \alpha) = \bar{y}_i^{[0]}(\alpha).$$

Thus for fixed α we have a system of initial value problem in R^{2n} . If we can solve it (uniquely), we have only to verify that the intervals, $[\underline{y}_j(t, \alpha), \bar{y}_j(t, \alpha)]$ define a fuzzy number $y_i(t) \in E$. Now let $\underline{y}^{[0]}(\alpha) = (\underline{y}_1^{[0]}(\alpha), \dots, \underline{y}_n^{[0]}(\alpha))^t$ and $\bar{y}^{[0]}(\alpha) = (\bar{y}_1^{[0]}(\alpha), \dots, \bar{y}_n^{[0]}(\alpha))^t$, with respect to the above mentioned indicators, system (2) can be written as with assumption

$$\begin{cases} y'(t) = F(t, y(t)), \\ y(0) = y^{[0]} \in E^n. \end{cases} \quad (5)$$

With assumption $y(t, \alpha) = [\underline{y}(t, \alpha), \bar{y}(t, \alpha)]$ and $y'(t, \alpha) = [\underline{y}'(t, \alpha), \bar{y}'(t, \alpha)]$ where

$$\underline{y}(t, \alpha) = (\underline{y}(t, \alpha), \dots, \underline{y}(t, \alpha))^t, \quad (6)$$

$$\bar{y}(t, \alpha) = (\bar{y}(t, \alpha), \dots, \bar{y}(t, \alpha))^t, \quad (7)$$

$$\underline{y}'(t, \alpha) = (\underline{y}'(t, \alpha), \dots, \underline{y}'(t, \alpha))^t, \quad (8)$$

$$\bar{y}'(t, \alpha) = (\bar{y}'(t, \alpha), \dots, \bar{y}'(t, \alpha))^t, \quad (9)$$

and with assumption $F(t, y(t, \alpha)) = [\underline{F}(t, y(t, \alpha)), \overline{F}(t, y(t, \alpha))]$, where

$$\underline{F}(t, y(t, \alpha)) = (\underline{f}_1(t, y(t, \alpha)), \dots, \underline{f}_n(t, y(t, \alpha)))^t, \quad (10)$$

$$\overline{F}(t, y(t, \alpha)) = (\overline{f}_1(t, y(t, \alpha)), \dots, \overline{f}_n(t, y(t, \alpha)))^t, \quad (11)$$

$y(t)$ is a fuzzy solution of (5) on an interval I for all $\alpha \in (0, 1]$, if

$$\begin{cases} \underline{y}'(t, \alpha) = \underline{F}(t, y(t, \alpha)); \\ \overline{y}'(t, \alpha) = \overline{F}(t, y(t, \alpha)); \\ \underline{y}(0, \alpha) = \underline{y}^{[0]}(\alpha), \quad \overline{y}(0, \alpha) = \overline{y}^{[0]}(\alpha). \end{cases} \quad (12)$$

or

$$\begin{cases} y'(t, \alpha) = F(t, y(t, \alpha)), \\ y(0, \alpha) = y^{[0]}(\alpha). \end{cases} \quad (13)$$

Now we show that under the assumption for functions f_i , for $i = 1, \dots, n$ how we can obtain a unique fuzzy solution for system (2).

Theorem 3.1

If $f_i(t, u_1, \dots, u_n)$ for $i = 1, \dots, n$ are continuous function of t and satisfies the Lipschitz condition in $u = (u_1, \dots, u_n)^t$ in the region $D = \{(t, u) \mid t \in [0, 1], -\infty < u_i < \infty \text{ for } i = 1, \dots, n\}$ with constant L_i then the initial value problem (2) has a unique fuzzy solution in each case.

Proof. Denote $G = (\underline{F}, \overline{F})^t = (\underline{f}_1, \dots, \underline{f}_n, \overline{f}_1, \dots, \overline{f}_n)^t$ where

$$\underline{f}_i(t, u) = \min\{f_i(t, u_1, \dots, u_n); u_j \in [\underline{y}_j, \overline{y}_j], \text{ for } j = 1, \dots, n\}, \quad (14)$$

$$\overline{f}_i(t, u) = \max\{f_i(t, u_1, \dots, u_n); u_j \in [\underline{y}_j, \overline{y}_j], \text{ for } j = 1, \dots, n\}, \quad (15)$$

and $y = (\underline{y}, \overline{y})^t = (\underline{y}_1, \dots, \underline{y}_n, \overline{y}_1, \dots, \overline{y}_n)^t \in R^{2n}$. It can be shown that Lipschitz condition of functions f_i imply

$$\|F(t, z) - F(t, z^*)\| \leq L \|z - z^*\|.$$

This guarantees the existence and uniqueness solution of

$$\begin{cases} y'(t) = F(t, y(t)), \\ y(0) = y^{[0]} = (\underline{y}^{[0]}, \overline{y}^{[0]})^{2t} \in R^{2n}. \end{cases} \quad (16)$$

Also for any continuous function $y^{[1]} : R_+ \rightarrow R^{2n}$ the successive approximations

$$y^{[m+1]}(t) = y^{[0]} + \int_0^t F(s, y^{[m]}(s)) ds, \quad t \geq 0, \quad m = 1, 2, \dots, \quad (17)$$

converge uniformly on closed subintervals of R_+ to the solution of (16). In other word we have the following successive approximations

$$\underline{y}_i^{[m+1]}(t) = \underline{y}_i^{[0]} + \int_0^t \underline{f}_i(s, y^{[m]}(s))ds, \quad \text{for } i = 1, 2, \dots, n. \quad (18)$$

$$\bar{y}_i^{[m+1]}(t) = \bar{y}_i^{[0]} + \int_0^t \bar{f}_i(s, y^{[m]}(s))ds, \quad \text{for } i = 1, 2, \dots, n. \quad (19)$$

By choosing $y^{[0]} = (\underline{y}^{[0]}(\alpha), \bar{y}^{[0]}(\alpha))$ in (16) we get a unique solution $y^\alpha(t) = (\underline{y}(t, \alpha), \bar{y}(t, \alpha))$ to (3) and (4) for each $\alpha \in (0, 1]$.

Next we will show that the $y(t, \alpha) = [\underline{y}(t, \alpha), \bar{y}(t, \alpha)]$, defines a fuzzy number in E^n for each $0 \leq t \leq T$, i.e. that $y = (y_1, \dots, y_n)^t$ is a fuzzy solution to (14) and (15). Thus we will show that the intervals $[\underline{y}_i(t, \alpha), \bar{y}_i(t, \alpha)]$, for $i = 1, \dots, n$ satisfy the conditions of Lemma (2.1). The successive approximations $y^{[m]} = y^{[0]} \in E^n$,

$$y^{[m+1]}(t) = y^{[0]} + \int_0^t F(s, y^{[m]}(s))ds, \quad t \geq 0, \quad m = 1, 2, \dots, \quad (20)$$

where the integrals are the fuzzy integrals, define a sequence of fuzzy numbers $y^{[m]}(t) = (y_1^{[m]}(t), \dots, y_n^{[m]}(t))^t$ for each $0 \leq t \leq T$. Hence $[y_i^{[m]}(t)]_\beta \subset [y_i^{[m]}(t)]_\alpha$, if $0 < \alpha \leq \beta \leq 1$, which implies that

$$[\underline{y}_i^{[m]}(t, \beta), \bar{y}_i^{[m]}(t, \beta)] \subset [\underline{y}_i^{[m]}(t, \alpha), \bar{y}_i^{[m]}(t, \alpha)], \quad 0 < \alpha \leq \beta \leq 1,$$

since, by convergence of sequences (16) and (19), the end points of $[y_i^{[m]}(t)]_\alpha$ converge to $\underline{y}_i(t, \alpha)$ and $\bar{y}_i(t, \alpha)$ that means

$$\underline{y}_i^{[m]}(t, \alpha) \rightarrow \underline{y}_i(t, \alpha) \quad \text{and} \quad \bar{y}_i^{[m]}(t, \alpha) \rightarrow \bar{y}_i(t, \alpha). \quad (21)$$

Thus the inclusion property (i) of Lemma (2.1) holds for the intervals $[\underline{y}_i(t, \alpha), \bar{y}_i(t, \alpha)]$, for $0 < \alpha \leq 1$. For the proof of the property (ii) of Lemma (2.1), let (α_p) be a non-decreasing sequence in $(0, 1]$ converging to α . Then $y^{[0]}(\alpha_p) \rightarrow y^{[0]}(\alpha)$ and $\bar{y}^{[0]}(\alpha_p) \rightarrow \bar{y}^{[0]}(\alpha)$, because of $y^{[0]} \in E^n$. But by the continuous dependence on the initial value of the solution (16), $y(t, \alpha_p) \rightarrow y(t, \alpha)$ and $\bar{y}(t, \alpha_p) \rightarrow \bar{y}(t, \alpha)$, this means (ii) holds for the interval $[\underline{y}(t, \alpha), \bar{y}(t, \alpha)]$, for $0 < \alpha \leq 1$. Hence by Lemma (2.1), $y(t) \in E^n$ and so y is a fuzzy solution of (1). the uniqueness follows from the uniqueness of the solution of (16).

4 The Runge-Kutta Nystrom Method

With before assumptions, the initial values problem (2) has a unique solution, such as $y = (y_1, \dots, y_n)^t \in E^n$. for found an approximate solution for (2) with the Runge-Kutta Nystrom method, we first define

$$\begin{aligned} \underline{y}(t_{n+1}; \alpha) - \underline{y}(t_n; \alpha) &= \sum_{i=1}^3 w_i k_{i,1}(t_n, y(t_n; \alpha)), \\ \bar{y}(t_{n+1}; \alpha) - \bar{y}(t_n; \alpha) &= \sum_{i=1}^3 w_i k_{i,2}(t_n, y(t_n; \alpha)), \end{aligned} \quad (22)$$

$$\begin{aligned}
 [k_i(t, y(t; \alpha))]_{\alpha} &= [k_{i,1}(t, y(t; \alpha)), k_{i,2}(t, y(t; \alpha))], \quad i = 1, 2, 3 \\
 k_{i,1}(t_n, y(t_n; \alpha)) &= h.f(t_n + a_i h, y_1(t_n) + \sum_{j=1}^{i-1} b_{ij} k_{j,1}(t_n, y(t_n; \alpha))), \\
 k_{i,2}(t_n, y(t_n; \alpha)) &= h.f(t_n + a_i h, y_2(t_n) + \sum_{j=1}^{i-1} b_{ij} k_{j,2}(t_n, y(t_n; \alpha))),
 \end{aligned} \tag{23}$$

and

$$\begin{aligned}
 k_{1,1}(t, y(t; \alpha)) &= \min\{h.f(t, s_1, \dots, s_n) | s_i \in [y_1(t; \alpha), y_2(t; \alpha)]\}, \quad (i = 1, 2, \dots, n) \\
 k_{1,2}(t, y(t; \alpha)) &= \max\{h.f(t, s_1, \dots, s_n) | s_i \in [y_1(t; \alpha), y_2(t; \alpha)]\}, \\
 k_{2,1}(t, y(t; \alpha)) &= \min\{h.f(t + \frac{2}{3}h, s_1, \dots, s_n) | s_i \in [z_{1,1}(t, y(t; \alpha), h), z_{1,2}(t, y(t; \alpha), h)]\}, \\
 k_{2,2}(t, y(t; \alpha)) &= \max\{h.f(t + \frac{2}{3}h, s_1, \dots, s_n) | s_i \in [z_{1,1}(t, y(t; \alpha), h), z_{1,2}(t, y(t; \alpha), h)]\}, \\
 k_{3,1}(t, y(t; \alpha)) &= \min\{h.f(t + \frac{2}{3}h, s_1, \dots, s_n) | s_i \in [z_{2,1}(t, y(t; \alpha), h), z_{2,2}(t, y(t; \alpha), h)]\}, \\
 k_{3,2}(t, y(t; \alpha)) &= \max\{h.f(t + \frac{2}{3}h, s_1, \dots, s_n) | s_i \in [z_{2,1}(t, y(t; \alpha), h), z_{2,2}(t, y(t; \alpha), h)]\},
 \end{aligned} \tag{24}$$

where in the Runge-Kutta Nystrom method,

$$\begin{aligned}
 z_{1,1}(t, y(t; \alpha), h) &= y_1(t; \alpha) + \frac{2}{3}k_{1,1}(t, y(t; \alpha)), \\
 z_{1,2}(t, y(t; \alpha), h) &= y_2(t; \alpha) + \frac{2}{3}k_{1,2}(t, y(t; \alpha)), \\
 z_{2,1}(t, y(t; \alpha), h) &= y_1(t; \alpha) + \frac{2}{3}k_{2,1}(t, y(t; \alpha), h), \\
 z_{2,2}(t, y(t; \alpha), h) &= y_2(t; \alpha) + \frac{2}{3}k_{2,2}(t, y(t; \alpha), h).
 \end{aligned} \tag{25}$$

Define,

$$\begin{aligned}
 F(t, y(t; \alpha), h) &= 2k_{1,1}(t, y(t; \alpha), h) + 3k_{2,1}(t, y(t; \alpha), h) + 3k_{3,1}(t, y(t; \alpha), h), \\
 G(t, y(t; \alpha), h) &= 2k_{1,2}(t, y(t; \alpha), h) + 3k_{2,2}(t, y(t; \alpha), h) + 3k_{3,2}(t, y(t; \alpha), h).
 \end{aligned} \tag{26}$$

and suppose that the discrete equally spaced grid points $\{t_0 = 0, t_1, \dots, t_N = T\}$ is a partition for interval $[0, T]$. If the exact and approximate solution in the i -th α cut at $t_m, 0 \leq m \leq N$ are denoted by $[y_i^{[m]}(\alpha), \bar{y}_i^{[m]}(\alpha)]$ and $[\underline{Y}_i^{[m]}(\alpha), \bar{Y}_i^{[m]}(\alpha)]$ respectively, then the numerical method for solution approximation in the i -th coordinate α cut, with the Runge-Kutta Nystrom method is

$$\begin{aligned}
 \underline{Y}_i^{[m+1]}(\alpha) &= \underline{Y}_i^{[m]}(\alpha) + \frac{h}{8}F_i(t_m, Y^m(\alpha), h), & \underline{Y}_i^{[0]}(\alpha) &= \underline{y}_i^{[0]}(\alpha), \\
 \bar{Y}_i^{[m+1]}(\alpha) &= \bar{Y}_i^{[m]}(\alpha) + \frac{h}{8}G_i(t_m, Y^m(\alpha), h), & \bar{Y}_i^{[0]}(\alpha) &= \bar{y}_i^{[0]}(\alpha),
 \end{aligned} \tag{27}$$

where $[Y_i(t)]_\alpha = [\underline{Y}_i(t, \alpha), \bar{Y}_i(t, \alpha)]$, $Y^{[m]}(\alpha) = [\underline{Y}^{[m]}(\alpha), \bar{Y}^{[m]}(\alpha)]$

$$\underline{Y}^{[m]}(\alpha) = (\underline{Y}_1^{[m]}(\alpha), \dots, \underline{Y}_n^{[m]}(\alpha))^t, \text{ and } \bar{Y}^{[m]}(\alpha) = (\bar{Y}_1^{[m]}(\alpha), \dots, \bar{Y}_n^{[m]}(\alpha))^t. \quad (28)$$

Now we input

$$\begin{aligned} F^*(t, Y^{[m]}(\alpha), h) &= \frac{1}{8}(F_1(t, Y^{[m]}(\alpha), h), \dots, F_n(t, Y^{[m]}(\alpha), h))^t, \\ G^*(t, Y^{[m]}(\alpha), h) &= \frac{1}{8}(G_1(t, Y^{[m]}(\alpha), h), \dots, G_n(t, Y^{[m]}(\alpha), h))^t. \end{aligned} \quad (29)$$

The Runge-Kutta Nystrom method for solutions approximation α -cut of differential equation (13) is as follows

$$Y^{[m+1]}(\alpha) = Y^{[m]}(\alpha) + hH(t_m, Y^{[m]}(\alpha), h), \quad Y^{[0]}(\alpha) = y^{[0]}(\alpha) \quad (30)$$

where

$$H(t_m, Y^{[m]}(\alpha), h) = [F^*(t_m, Y^{[m]}(\alpha), h), G^*(t_m, Y^{[m]}(\alpha), h)], \quad (31)$$

and

$$\begin{aligned} F^*(t_m, Y^{[m]}(\alpha), h) &= \frac{1}{8}[2\underline{k}_1(t_m, Y^{[m]}(\alpha), h) + 3\underline{k}_2(t_m, Y^{[m]}(\alpha), h) \\ &\quad + 3\underline{k}_3(t_m, Y^{[m]}(\alpha), h)], \end{aligned} \quad (32)$$

$$\begin{aligned} G^*(t_m, Y^{[m]}(\alpha), h) &= \frac{1}{8}[2\bar{k}_1(t_m, Y^{[m]}(\alpha), h) + 3\bar{k}_2(t_m, Y^{[m]}(\alpha), h) \\ &\quad + 3\bar{k}_3(t_m, Y^{[m]}(\alpha), h)] \end{aligned}$$

and also

$$\begin{aligned} \underline{k}_j(t, Y^{[m]}(\alpha), h) &= (\underline{k}_{1j}(t, Y^{[m]}(\alpha), h), \dots, \underline{k}_{nj}(t, Y^{[m]}(\alpha), h))^t, \\ \bar{k}_j(t, Y^{[m]}(\alpha), h) &= (\bar{k}_{1j}(t, Y^{[m]}(\alpha), h), \dots, \bar{k}_{nj}(t, Y^{[m]}(\alpha), h))^t. \end{aligned} \quad (33)$$

5 Convergence and Stability

Definition 5.1

A one-step method for approximating the solution of a differential equation

$$\begin{cases} y'(t) = F(t, y(t)), \\ y(0) = y^{[0]} \in R^n, \end{cases} \quad (34)$$

which F is a n -th ordered as follows $f = (f_1, \dots, f_n)^t$ and $f_i : R_+ \times R^n \rightarrow R(1 \leq i \leq n)$, is a method which can be written in the form

$$Y^{[n+1]} = Y^{[n]} + h\psi(t_n, Y^{[n]}, h), \quad (35)$$

where the increment function ψ is determined by F and is a function $t_n, Y^{[n]}$ and h only.

Theorem 5.1

If $\psi(t, y, h)$ satisfies a Lipschitz condition in y , then the method given by (35) is stable.

Theorem 5.2

In relation (5), if $F(t, y)$ satisfies a Lipschitz condition in y , then the method given by (30) is stable.

Theorem 5.3

If

$$Y^{[m+1]}(\alpha) = Y^{[m]}(\alpha) + h\psi(t_m, Y^{[m]}(\alpha), h), \quad Y^{[0]} = y^{[m]}(\alpha) \quad (36)$$

where $\psi(t_m, Y^{[m]}(\alpha), h) = [\psi_1(t_m, Y^{[m]}(\alpha), h), \psi_2(t_m, Y^{[m]}(\alpha), h)]$ is a numerical method for approximation of differential equation (13), and ψ_1 and ψ_2 are continuous in t, y, h for $0 \leq t \leq T, 0 \leq h \leq h_0$ and all y , and if they satisfy a Lipschitz condition in the region $D = \{(t, u, v, h) | 0 \leq t \leq T, -\infty < u_i \leq v_i, -\infty < v_i \leq +\infty, 0 \leq h \leq h_0 \quad i = 1, \dots, n\}$, necessary and sufficient conditions for convergence above mentioned method is

$$\psi(t, y(t, \alpha), h) = F(t, y(t, \alpha)). \quad (37)$$

Proof. Suppose that $\psi(t, y(t, \alpha), 0) = F(t, y(t, \alpha))$, since, $F(t, y(t, \alpha))$ satisfying the conditions of theory (1), then the following equation

$$\begin{cases} y'(t) = F(t, y(t)), \\ y(0) = y^{[0]}(\alpha), \end{cases} \quad (38)$$

has a unique solution such as $y(t, \alpha) = [y(t, \alpha), \bar{y}(t, \alpha)]$, where $\underline{y}(t, \alpha) = (\underline{y}(t, \alpha), \dots, \underline{y}(t, \alpha))^t$ and $\bar{y}(t, \alpha) = (\bar{y}(t, \alpha), \dots, \bar{y}(t, \alpha))^t$. We will show that the numerical solutions given by (36) convergent to the $y(t)$. By the mean value theorem,

$$\underline{y}_i^{[m+1]} = \underline{y}_i^{[m]} + h\underline{f}_i(t_m + \underline{\theta}_i h, y(t_m + \underline{\theta}_i h)), \quad \text{for } 0 < \underline{\theta}_i < 1, \quad (39)$$

$$\bar{y}_i^{[m+1]} = \bar{y}_i^{[m]} + h\bar{f}_i(t_m + \bar{\theta}_i h, y(t_m + \bar{\theta}_i h)), \quad \text{for } 0 < \bar{\theta}_i < 1, \quad (40)$$

with assumption $\underline{\psi} = (\underline{\psi}_1, \dots, \underline{\psi}_n)^t$ and $\bar{\psi} = (\bar{\psi}_1, \dots, \bar{\psi}_n)^t$. From equation (36) obtain the following relations

$$\underline{Y}_i^{[m+1]}(\alpha) = \underline{Y}_i^{[m]}(\alpha) + h\underline{\psi}_i(t_m, Y^{[m]}(\alpha), h), \quad (41)$$

$$\bar{Y}_i^{[m+1]}(\alpha) = \bar{Y}_i^{[m]}(\alpha) + h\bar{\psi}_i(t_m, Y^{[m]}(\alpha), h), \quad (42)$$

and subtracting (39), (40) from (41), (42) respectively, and setting

$$e^{[m]}(\alpha) = [\underline{e}^{[m]}(\alpha), \bar{e}^{[m]}(\alpha)],$$

where

$$\underline{e}^{[m]}(\alpha) = \underline{e}(t_m, \alpha) = \underline{Y}^{[m]}(\alpha) - \underline{y}^{[m]}(\alpha) \quad \text{and} \quad \bar{e}^{[m]}(\alpha) = \bar{e}(t_m, \alpha) = \bar{Y}^{[m]}(\alpha) - \bar{y}^{[m]}(\alpha),$$

we get

$$\begin{aligned} \underline{e}_i^{[m+1]}(\alpha) &= \underline{e}_i^{[m]}(\alpha) + h\{\underline{\psi}_i(t_m, Y^{[m]}(\alpha), h) - \underline{\psi}_i(t_m, y^{[m]}(\alpha), h) + \underline{\psi}_i(t_m, y^{[m]}(\alpha), h) \\ &\quad - \underline{\psi}_i(t_m, y^{[m]}(\alpha), 0) + \underline{\psi}_i(t_m, y^{[m]}(\alpha), 0) - \underline{f}_i(t_m + \theta_i h, y(t_m + \theta_i h))\} \end{aligned}$$

on the other way, with respect to the relation of $\underline{\psi}_i(t_m, y^{[m]}(\alpha), 0) - \underline{f}_i(t_m, y^{[m]}(\alpha))$ we can write

$$\begin{aligned} &|\underline{\psi}_i(t_m, y^{[m]}(\alpha), 0) - \underline{f}_i(t_m + \theta_i h, y(t_m + \theta_i h))| \\ &\leq hL_1\theta_i + L_1 \sum_{i=1}^n |y_i(t_m + \theta_i h) - y_i(t_m)| + L_1 \sum_{i=1}^n |\bar{y}_i(t_m + \bar{\theta}_i h) - \bar{y}_i(t_m)| \\ &= hL_1\theta_i + L_1 \sum_{i=1}^n |y'_i(t_m + \xi_i \theta_i h)\theta_i h| + L_1 \sum_{i=1}^n |\bar{y}'_i(t_m + \bar{\xi}_i \theta_i h)\theta_i h| = hL_2, \end{aligned}$$

then

$$\begin{aligned} |\underline{e}_i^{[m+1]}(\alpha)| &\leq |\underline{e}_i^{[m]}(\alpha) + hL_1 \left\{ \sum_{j=1}^n |\underline{e}_j^{[m]}(\alpha)| + \bar{e}_j^{[m]}(\alpha) \right\}| + h^2 L_1 + h^2 L_2 \\ &\leq |\underline{e}_i^{[m]}(\alpha) + nhL_1 \max_{1 \leq j \leq n} \{|\underline{e}_j^{[m]}(\alpha)|\}| + nhL_1 \max_{1 \leq j \leq n} \{|\bar{e}_j^{[m]}(\alpha)|\}| + h^2(L_1 + L_2). \end{aligned}$$

On the other hand

$$\max_{1 \leq j \leq n} \{|\underline{e}_j^{[m]}(\alpha)|\} = k_i |\underline{e}_i^{[m]}(\alpha)|, \quad \max_{1 \leq j \leq n} \{|\bar{e}_j^{[m]}(\alpha)|\} = k'_i |\bar{e}_i^{[m]}(\alpha)|$$

with assumption $k_1 = \max_{1 \leq i \leq n} \{k_i, k'_i\}$ and $M = L_1 + L_2$, we can write

$$\begin{aligned} |\underline{e}_i^{[m+1]}(\alpha)| &\leq |\underline{e}_i^{[m]}(\alpha) + nhk_1 L_1 \{|\underline{e}_i^{[m]}(\alpha)| + |\bar{e}_i^{[m]}(\alpha)|\}| + M_1 h^2 \\ &\leq |\underline{e}_i^{[m]}(\alpha) + 2nhk_1 L_1 \max\{|\underline{e}_i^{[m]}(\alpha)|, |\bar{e}_i^{[m]}(\alpha)|\}| + M_1 h^2, \end{aligned} \quad (43)$$

similarly, we can obtain the following relation

$$|\bar{e}_i^{[m+1]}(\alpha)| \leq |\bar{e}_i^{[m]}(\alpha) + 2nhk_2 L'_1 \max\{|\underline{e}_i^{[m]}(\alpha)|, |\bar{e}_i^{[m]}(\alpha)|\}| + M_2 h^2. \quad (44)$$

Now, we input $L = \max\{L_1, L'_1\}$ and $M = \max\{M_1, M_2\}$ so the relations (43) and (44) can be written as follow

$$\begin{aligned} |\underline{e}_i^{[m+1]}(\alpha)| &\leq |\underline{e}_i^{[m]}(\alpha) + 2nhkL \max\{|\underline{e}_i^{[m]}(\alpha)|, |\bar{e}_i^{[m]}(\alpha)|\}| + Mh^2, \\ |\bar{e}_i^{[m+1]}(\alpha)| &\leq |\bar{e}_i^{[m]}(\alpha) + 2nhkL \max\{|\underline{e}_i^{[m]}(\alpha)|, |\bar{e}_i^{[m]}(\alpha)|\}| + Mh^2 \end{aligned}$$

Denote $e_i^{[m]} = |\underline{e}_i^{[m]}(\alpha) + |\bar{e}_i^{[m]}(\alpha)|$. Then by virtue of lemma (5.7)

$$e_i^{[m]}(\alpha) \leq (1 + 4nhkL)^m e_i^{[0]}(\alpha) + 2Mh^2 \frac{(1 + 4nhkL)^m - 1}{4nhkL},$$

where $e_i^{[0]} = |\underline{e}_i^{[0]}(\alpha)| + |\bar{e}_i^{[0]}(\alpha)|$. Then

$$\begin{aligned} |\underline{e}_i^{[m]}(\alpha)| &\leq e^{4mnkh} \times e_i^{[0]} + M \frac{e^{4mnklh}}{2nhkL} h \\ |\bar{e}_i^{[m]}(\alpha)| &\leq e^{4mnkh} \times e_i^{[0]} + M \frac{e^{4mnklh}}{2nhkL} h. \end{aligned}$$

In particular

$$|\underline{e}_i^{[N]}(\alpha)| \leq e^{4Nnkh} \times e_i^{[0]} + M \frac{e^{4Nnklh}}{2nhkL} h \quad \text{and} \quad |\bar{e}_i^{[N]}(\alpha)| \leq e^{4Nnkh} \times e_i^{[0]} + M \frac{e^{4Nnklh}}{2nhkL} h.$$

Since $\underline{e}_i^{[0]}(\alpha) = \bar{e}_i^{[0]}(\alpha) = 0$ and $h = \frac{T}{N}$ we obtain $\|e_1^{[N]}(\alpha)\| \leq M \frac{e^{4Nnklh}}{2nhkL} h$ and $\|e_2^{[N]}(\alpha)\| \leq M \frac{e^{4Nnklh}}{2nhkL} h$. Then $\|e^{[N]}(\alpha)\| \leq 2M \frac{e^{4Nnklh}}{2nhkL} h$ if $h \rightarrow 0$ we get $\|e^{[N]}(\alpha)\| \rightarrow 0$, so the numerical solution (36) converge to the solutions (38). Conversely, suppose that the numerical method (36) convergent to the solution of the system (38). With absurd hypothesis we suppose that (37) is not correct. Then $\psi(t, y(t, \alpha), 0) = g(t, y(t, \alpha)) \neq F(t, y(t, \alpha))$. Similarly, we can proof that the numerical method of (36) is convergent to the solution of following system

$$\begin{cases} u'(t) = g(t, y(t)), \\ u(0) = y^{[0]}(\alpha), \end{cases} \quad (45)$$

then $y(t, \alpha) = u(t)$. Since $g(t, y(t, \alpha)) \neq F(t, y(t, \alpha))$, suppose that F and g differ at some point $(t_\alpha, y(t_\alpha, \alpha))$. If we consider the initial value problem (38) and (45) starting from $(t_\alpha, y(t_\alpha, \alpha))$ we have

$$y'(t_\alpha, \alpha) = F(t_\alpha, y(t_\alpha, \alpha)) \neq g(t_\alpha, y(t_\alpha, \alpha)) = g(t_\alpha, u(t_\alpha)) = u'(t_\alpha)$$

which is a contradiction.

Corollary 5.1

The Runge-Kutta proposed method by (30) and is convergent to the solution of the system (13) respectively.

6 Numerical Example

Example 6.1 Consider the following fuzzy differential equation with fuzzy initial value

$$\begin{cases} y'''(t) = 2y''(t) + 3y'(t) & (0 \leq t \leq 1), \\ y(0) = (3 + \alpha, 5 - \alpha), \\ y'(0) = (-1 - \alpha, -3 + \alpha), \\ y''(0) = (8 + \alpha, 10 - \alpha). \end{cases}$$

the eigen value -eigen vector solution is as follows:

$$y(t; r) = \left(-\frac{1}{3} + \frac{7}{12}e^{3t} + \left(\frac{11}{4} + \alpha\right)e^{-t}, -\frac{1}{3} + \frac{7}{12}e^{3t} + \left(\frac{19}{4} - \alpha\right)e^{-t} \right)$$

The solution of Runge-Kutta Nystrom method is as follows and Figure 1 and Table 1 show the obtain the results:

$$\begin{aligned} \underline{Y}_1^{[m+1]} &= \underline{Y}_1^{[m]} + \left(h + \frac{h^3}{2}\right)\underline{Y}_2^{[m]} + \left(\frac{h^2}{2} + \frac{h^3}{3}\right)\underline{Y}_3^{[m]}, \\ \overline{Y}_1^{[m+1]} &= \overline{Y}_1^{[m]} + \left(h + \frac{h^3}{2}\right)\overline{Y}_2^{[m]} + \left(\frac{h^2}{2} + \frac{h^3}{3}\right)\overline{Y}_3^{[m]}, \\ \underline{Y}_2^{[m+1]} &= \underline{Y}_2^{[m]} + \left(\frac{3h^2}{2} + h^3\right)\underline{Y}_2^{[m]} + \left(h + h^2 + \frac{7h^3}{6}\right)\underline{Y}_3^{[m]}, \\ \overline{Y}_2^{[m+1]} &= \overline{Y}_2^{[m]} + \left(\frac{3h^2}{2} + h^3\right)\overline{Y}_2^{[m]} + \left(h + h^2 + \frac{7h^3}{6}\right)\overline{Y}_3^{[m]}, \\ \underline{Y}_3^{[m+1]} &= \underline{Y}_3^{[m]} + \left(3h + 3h^2 + \frac{7h^3}{2}\right)\underline{Y}_2^{[m]} + \left(2h + \frac{7h^2}{2} + \frac{10h^3}{3}\right)\underline{Y}_3^{[m]}, \\ \overline{Y}_3^{[m+1]} &= \overline{Y}_3^{[m]} + \left(3h + 3h^2 + \frac{7h^3}{2}\right)\overline{Y}_2^{[m]} + \left(2h + \frac{7h^2}{2} + \frac{10h^3}{3}\right)\overline{Y}_3^{[m]}. \end{aligned}$$

Table 1.

r	Approximation		Exact	
	$y_1(t_i; r)$	$y_2(t_i; r)$	$y_1(t_i; r)$	$y_2(t_i; r)$
0.1	12.3925	13.0544	12.4317	13.0939
0.2	12.4293	13.0175	12.4685	13.0571
0.3	12.4659	12.9807	12.5053	13.0203
0.4	12.5025	12.9439	12.5421	12.9835
0.5	12.5394	12.9073	12.5788	12.9467
0.6	12.5763	12.8692	12.6156	12.9099
0.7	12.6130	12.8338	12.6524	12.8731
0.8	12.6499	12.7971	12.6892	12.8363
0.9	12.6865	12.7600	12.7260	12.7996
1	12.7233	12.7233	12.7628	12.7628

7 Conclusion In this paper a numerical method for solving n^{th} -order fuzzy differential equation with fuzzy initial condition is presented. In this method n^{th} -order fuzzy differential equation is converted to a fuzzy system which will be solved with the *Runge-Kutta Nystrom* method.




Figure 1: h=0.1

References

- [1] S. Abbasbandy, T. Allah Viranloo, Numerical solution of Fuzzy Differential Equations by Taylor method, J. Comput. Methods Appl. Math. 2(2002)113-124.
- [2] S. Abbasbandy, T. Allah Viranloo, Oscar Lopez-Pouso, J.J. Neito, Numerical Methods for Fuzzy Differential Inclusions, Comput. Math. Appl. 48(2004) 1633-1641.
- [3] T. Allah Viranloo, N. Ahmady, E. Ahmady, Numerical solution of Fuzzy Differential Equations by predictor-corrector method, Infor. Sci. 177(2007) 1633-1647.
- [4] T. Allah Viranloo, N. Ahmady, E. Ahmady, A method for solving n^{th} -order Fuzzy Linear Differential Equations, Comput. Math. Appl 86(2009) 730-742.
- [5] S. Abbasbandy, T. Allah Viranloo, Numerical solution of N^{th} -Order Fuzzy Differential Equations by Runge-Kutta Method Comput. Methods Appl. Math. 4 (2011) 935-946.
- [6] B. Bede, I.J. Rudas, A. L. Bencsik, First order Linear Fuzzy Differential Equations generalized differentiability, Inform. Sci. 177 (2007) 1648-1662.
- [7] J. J. Buckley, Thomas Feuring, Fuzzy Differential Equations, Fuzzy Sets and Systems 105 (1999), 241-248.
- [8] J. J. Buckley, Thomas Feuring, Introduction to Fuzzy Partial Differential Equations, Fuzzy Sets and Systems 110 (2000), 43-54.
- [9] J. J. Buckley, T. Feuring, Fuzzy initial value problem for N^{th} -order fuzzy linear differential equations, Fuzzy Sets and Systems 121 (2001) 247-255.
- [10] S. L. Chang and A. Zadeh, On Fuzzy Mapping and Control, IEEE Trans. Systems Man Cybernet., 2 (1972) 30-34.

- [11] W. Congxin, S. Shiji, Existence theorem to the Cauchy problem of Fuzzy Differential Equations under compactness-type conditions, *Infor. Sci.* 108(1998) 123-134.
- [12] D. Dubois and H. Prade, Towards Fuzzy Differential Calculus: Part 3, Differentiation, *Fuzzy Sets and Systems*, 8 (1982) 225-233.
- [13] M. Friedman, M. Ming, A. Kandel, Fuzzy linear systems, *Fuzzy Set and Systems* 96 (1998) 201-209.
- [14] C.W. Gear, *Numerical Initial Value Problem In Ordinary Differential Equations*, Prentice Hall(1971).
- [15] R. Goetschel and W. Voxman, Elementary Fuzzy Calculus, *Fuzzy Sets and Systems*, 18 (1986) 31-43.
- [16] O. He, W. Yi, On fuzzy differential equations, *Fuzzy Sets Systems*, 24 (1989) 321-325.
- [17] T. Jayakumar, K. Kanagarajan, Numerical Solution for Hybrid Fuzzy System by Improved Euler Method, *International Journal of Applied Mathematical Science* 38 (2012) 1847-1862.
- [18] T. Jayakumar, D. Maheshkumar, K. Kanagarajan, Numerical Solution of Fuzzy Differential Equations by Runge-Kutta Method of Order Five, *International Journal of Applied Mathematical Science* 6 (2012) 2989-3002.
- [19] T. Jayakumar, K. Kanagarajan, Numerical Solution for Hybrid Fuzzy System by Runge-Kutta Method of Order Five, *International Journal of Applied Mathematical Science* 6 (2012) 3591-3606.
- [20] L.J. Jowers, J.J. Buckley, K.D. Reilly, Simulating continuous fuzzy systems, *Infor. Sci.* 177 (2007) 436-448.
- [21] O. Kaleva, Fuzzy Differential Equations, *Fuzzy Sets Systems*, 24 (1987) 301-317.
- [22] O. Kaleva, The Cauchy Problem for Fuzzy Differential Equations, *Fuzzy Sets and Systems*, 35 (1990) 389-396.
- [23] P. Kloeden, remarks on Peano-like theorems for fuzzy differential Equations, *Fuzzy Sets Systems*, 44 (1991) 161-164.
- [24] K. Kanagarajan, M.Sambath, Runge-Kutta Nystrom, method for solving fuzzy differential equations, *Journal of computational methods in Applied mathematics* 10 (2010) 195-203.
- [25] A. Kandel, W.J. Byatt, Fuzzy differential equations, in *Proceedings of the International Conference on Cybernetics and Society*, Tokyo, (1978) 1213-1216.
- [26] M. L. Puri, D. A. Ralescu, Differentials of fuzzy functions, *J. Math. Anal. Application* 91 (1983),321-325.
- [27] D. Ralescu, A survey of the representation of fuzzy concepts and its applications, in: M.M Gupta, R.K. Ragade, R.R. Yager, Eds., *Advances in fuzzy Set Theory and Applications*, North-Holland, Amsterdam(1979) 77-91.
- [28] S. Seikkala, On the Fuzzy Initial Value Problem, *Fuzzy Sets Systems*, 24 (1987) 319-330.