

Polynomial time integer factorization

Yuly Shipilevsky

Toronto, Ontario, Canada

E-mail address: yulysh2000@yahoo.ca

Abstract

A polynomial-time algorithm for integer factorization, wherein integer factorization reduced to a convex polynomial-time integer minimization problem.

Keywords: integer factorization, integer programming, polynomial-time, NP-hard.

1. Introduction

Cryptography, elliptic curves, algebraic number theory have been brought to bear on integer factorization problem.

Until now, no algorithm has been published that can factor in deterministic polynomial time. For an ordinary computer the best published asymptotic running time is for the general number field sieve (GNFS) algorithm [8,10].

The purpose of this paper is to develop a polynomial-time integer factorization algorithm, factoring in deterministic polynomial time.

The plan of this paper is as follows. In Section 2 we reduce integer factorization problem to some 2-dimensional integer minimization problem and show that if there exists a nontrivial divisor of N , those divisor is a minimizer of those 2-dimensional integer minimization problem, and any minimizer of those integer minimization problem is a nontrivial divisor of N .

In Section 3 we introduce and investigate a notion of U-equivalent conversion of minimization problems for changing properties of the objective functions and preserving the set of minimizers of the original problem.

In Section 4 we reduce integer factorization problem to the convex integer minimization 2-dimensional problem solvable in time polynomial in $\log(N)$.

Finally, we conclude that since we found a polynomial-time algorithm to

solve an NP-hard problem(an integer minimization problem, U-equivalent to the problem, mentioned in Section 2), it would mean that P is equal to NP.

2. Reduction to the Integer Programming problem

Let us reduce integer factorization problem to some integer minimization problem, so that any minimizer that is found solves integer factorization problem.

The key idea is to construct the objective function and constraints so that any minimizer satisfies the equation: $x_1x_2 = N$ and therefore is a solution of the integer factorization problem.

Let us consider the following integer minimization problem:

$$\begin{aligned}
 &\text{minimize} && x_1x_2 \\
 &\text{subject to} && x_1x_2 \geq N, \\
 &&& 2 \leq x_1 \leq N-1, \\
 &&& 2 \leq x_2 \leq N-1, \\
 &&& x_1 \in \mathbf{N}, x_2 \in \mathbf{N}, N \in \mathbf{N}.
 \end{aligned} \tag{1}$$

Let $\Omega := \{ x \in \mathbf{R}^2, x = (x_1, x_2) \mid x_1x_2 \geq N, 2 \leq x_1 \leq N-1, 2 \leq x_2 \leq N-1, x_1 \in \mathbf{R}, x_2 \in \mathbf{R} \}$ for a given $N \in \mathbf{N}$.

Hence, $\Omega^1 = \Omega \cap \mathbf{Z}^2$ is a feasible set of the problem (1).

It is clear that if there exists a nontrivial solution of integer factorization problem $x_1x_2 = N$, the objective function: $f(x_1, x_2) = x_1x_2$ reaches minimum at the integer point of the border $x_1x_2 = N$ of the region Ω and if there exists a nontrivial solution of integer factorization problem, any minimizer of the problem (1) provides a (nontrivial) solution of integer factorization problem.

Thus, in this case, any minimizer of the problem (1) guarantees solution of integer factorization problem and there exists at least one such minimizer.

Theorem 1. *If there exists a nontrivial solution of integer factorization problem, that solution is a minimizer of problem (1) and if there exists a nontrivial solution of integer factorization problem, any minimizer of the problem (1) is a nontrivial solution of integer factorization problem.*

As a result, we obtain the following Integer Factorization Algorithm.

Algorithm 1(Integer Factorization Algorithm).

Input: A positive integer number N.

Output: A nontrivial divisor of N(if it exists).

Solve the problem (1):

Based on the input data compute a minimizer $(x_{1 \min}, x_{2 \min})$ of the problem (1).

if $(x_{1 \min} x_{2 \min} = N)$

then

Return a nontrivial divisor $x_{1 \min}$ of N

else

Return “N is a prime”

Let us determine the complexity of the problem (1).

Despite in general integer programming is NP-hard or even incomputable, see, e.g., Hemmecke et al. [5], for some subclasses of the objective functions and constraints it can be computed in time polynomial.

Let us recall some well-known definitions.

A function $f: \mathbf{R}^n \rightarrow \mathbf{R}$, defined on the convex set S is called concave on the set S if for all $x, y \in S$, and all $\alpha \in (0,1)$ we have:

$$f((1 - \alpha)x + \alpha y) \geq (1 - \alpha)f(x) + \alpha f(y)$$

A function $f: \mathbf{R}^n \rightarrow \mathbf{R}$, defined on the convex set S is called convex on the set S if for all $x, y \in S$, and all $\alpha \in (0,1)$ we have:

$$f((1 - \alpha)x + \alpha y) \leq (1 - \alpha)f(x) + \alpha f(y)$$

A function $f: \mathbf{R}^n \rightarrow \mathbf{R}$ is called quasiconcave if all the upper level sets $\{x \in \mathbf{R}^n, f(x) \geq \alpha\}$, $\alpha \in \mathbf{R}$ are convex subsets of \mathbf{R}^n .

A function $f: \mathbf{R}^n \rightarrow \mathbf{R}$ is called quasiconvex if all the lower level sets $\{x \in \mathbf{R}^n, f(x) \leq \alpha\}$, $\alpha \in \mathbf{R}$ are convex subsets of \mathbf{R}^n .

Note that the dimension of the problem (1) is fixed and is equal to 2.

A fixed-dimensional polynomial minimization in integer variables, where

the objective function is a convex polynomial and the convex feasible set is described by arbitrary polynomials can be solved in time polynomial, - see, e.g., Khachiyan and Porkolab [6].

A fixed-dimensional polynomial minimization over the integer variables, where the objective function $f_0(x)$ is a quasiconvex polynomial with integer coefficients and where the constraints are inequalities $f_i(x) \leq 0$, $i = 1, \dots, k$ with quasiconvex polynomials $f_i(x)$ with integer coefficients, $f_i: \mathbf{R}^n \rightarrow \mathbf{R}$, $f_i(x)$, $i = 0, \dots, k$ are polynomials of degree at most $p \geq 2$, can be solved in time polynomial in the degrees and the binary encoding of the coefficients, - see, e.g., Heinz [4], Hemmecke et al. [5], Lee [7].

A mixed-integer minimization of a convex function in a convex, bounded feasible set can be done in time polynomial, according to Baes et al. [2], Oertel et al. [9].

Since the objective function $f(x_1, x_2) = x_1x_2$ of the problem (1) is a quasiconcave function in the feasible set Ω^I of the problem (1), we cannot use the results described in Baes et al. [2], Heinz [4], Hemmecke et al. [5], Khachiyan and Porkolab [6], Oertel [9] in order to solve the problem (1) in time polynomial in $\log(N)$. Note that Ω^I is described by quasiconvex polynomials, since $(-x_1x_2 + N)$ is a quasiconvex function for $x = (x_1, x_2) > 0$.

In general, since variables $x_1 \in \mathbf{N}$, $x_2 \in \mathbf{N}$ are bounded by the finite bounds: $2 \leq x_1 \leq N-1$, $2 \leq x_2 \leq N-1$, the problem (1) and the respective Algorithm 1 are computable, but at least not solvable in time polynomial in $\log(N)$.

3. U-equivalent minimization

The following results give us a possibility to change the properties of the objective function with preservation of the set of minimizers of the original problem.

Theorem 2. *Let O be the minimization problem:*

$$O = \{\text{minimize } g(x) \text{ subject to } x \in G\}, \quad g: X \rightarrow \mathbf{R}, \quad G \subseteq X.$$

Let E be the minimization problem:

$$E = \{\text{minimize } U(g(x)) \text{ subject to } x \in G\}, \quad G \subseteq X,$$

where $U: \mathbf{R} \rightarrow \mathbf{R}$, $U = U(u)$ is any increasing function.

Let M_O be a set of minimizers of problem O and

let M_E be a set of minimizers of problem E. Then:

$$M_O = M_E.$$

Proof. If $x_0 \in M_O$ then $g(x_0) \leq g(x)$ for any $x \in G$. Hence, $U(g(x_0)) \leq U(g(x))$ for any $x \in G$, since function U is the increasing function and therefore $x_0 \in M_E$ and $M_O \subseteq M_E$. If $x_0 \in M_E$ then we have: $U(g(x_0)) \leq U(g(x))$ for any $x \in G$ and therefore $g(x_0) \leq g(x)$ for any $x \in G$, as otherwise there exists $y_0 \in G$ such that $g(x_0) > g(y_0)$ and since function U is the increasing function it would mean that $U(g(x_0)) > U(g(y_0))$ in contradiction to the original supposition that $U(g(x_0)) \leq U(g(x))$ for any $x \in G$. So, since $g(x_0) \leq g(x)$ for any $x \in G$ then $x_0 \in M_O$ and $M_E \subseteq M_O$ and finally: $M_O = M_E$. \square

Definition 1. We say that the minimization problem:

$$E = \{\text{minimize } U(g(x)) \text{ subject to } x \in G\},$$

is U -equivalent to the minimization problem:

$$O = \{\text{minimize } g(x) \text{ subject to } x \in G\}, g: X \rightarrow \mathbf{R}, G \subseteq X,$$

where $U: \mathbf{R} \rightarrow \mathbf{R}$, $U = U(u)$ is some increasing function.

Corollary 1. If E is U -equivalent to O then E and O have the same set of minimizers.

Proof. It follows from Theorem 2 and Definition 1. \square

Thus, using U -equivalence we can convert original minimization problem into minimization problem that has objective function with desired properties, so that both problems, - the original one, and U -equivalent have the same set of minimizers and share the same feasible set.

Hence, as a result of the U -equivalent conversion the original feasible set and the original set of minimizers remain unchanged, whereas the objective function is being changed to obtain desired properties (e.g., faster minimiza-

tion), which can consider it(U-equivalence) as a flexible and effective tool.

U-equivalent conversion can be considered as unary operation defined on the set of minimization problems, having the same feasible set.

Example. Suppose, the problem (1) is the original minimization problem. Let q be e^u -equivalent to the problem (1). The objective function of the problem (1) is xy , whereas the objective function of q is $f(x, y) = e^{xy}$.

Both problems, due to the Theorem 2 have the same set of minimizers (and each such minimizer is a solution of the integer factorization problem, according to the Theorem 1). Note that if N is not a prime, minimum $q = e^N$.

However, no U-equivalent conversion applied to the original problem (1) in order to get a quasiconvex objective function exists, since if a function g is quasiconcave and a function U is increasing, then a function f , defined as $f(x) = U(g(x))$ is still quasiconcave.

We will use U-equivalence and results obtained in Section 2 at the end of Section 4.

4. Convexification. Polynomial-time integer factorization. Minimum Principle.

We are going now to reduce integer factorization to some convex integer minimization problem.

Baes et al. [2] have shown that minimizing an arbitrary nonnegative proper convex function in two integer variables provided that the feasible set is contained in a known finite box $[-B, B]^2$ can be done in time polynomial in $\log(B)$ as well as the corresponding oracle.

Theorem 3(Theorem 2 in Baes et al. [2]). *Let $f: \mathbf{R}^2 \rightarrow \mathbf{R}$ and $g_i: \mathbf{R}^2 \rightarrow \mathbf{R}$ with $i = 1, \dots, m$ be convex functions. Let $B \in \mathbf{N}$ and let $x \in [-B, B]^2$ such that $g_i(x) \leq 0$ for all $i = 1, \dots, m$. Then, in a number of evaluations of f and g_1, \dots, g_m that is polynomial in $\log(B)$, one can either*

- (a) *find an $x_0 \in [-B, B]^2 \cap \mathbf{Z}^2$ with $f(x_0) \leq f(x)$ and $g_i(x_0) \leq 0$ for all $i = 1, \dots, m$ or*
- (b) *show that there is no such point.*

Let us construct a convex target function, so that if N is not a prime, any minimizer of the target function in the feasible set solves integer factorization problem.

Let $g: \mathbf{R}^2 \rightarrow \mathbf{R}$ be analytic, convex and positive function ($g > 0$). Let us construct a function: $g(x)(x_1x_2 - N)/(x_1x_2 - N + \varepsilon)$, $x = (x_1, x_2) \in \Omega$, $\varepsilon \geq 0$ (see the definition of feasible region Ω in Section 1).

If $\varepsilon = 0$, this function is equal to $g(x)$, $x \in \Omega$ at points $x: x_1x_2 - N > 0$. Let us define this function equal to $g(x)$ at points $x: x_1x_2 - N = 0$ ($\varepsilon = 0$) as well. Since function g is convex and analytic, there exists a vicinity: $V(0, \varepsilon_0) = \{\varepsilon \in \mathbf{R}: 0 < \varepsilon < \varepsilon_0, \varepsilon_0 \in \mathbf{R}\}$ of $\varepsilon = 0$, where the above-constructed function preserves convexity on Ω due to the smoothness of the constructed function.

The corresponding vicinity $V(0, \varepsilon_0)$ can be determined using a necessary and sufficient condition of convexity by determining when the corresponding Hessian matrix of the target function is positive semidefinite.

As a result, the following convex integer minimization problem solves the problem:

$$\begin{aligned}
 &\text{minimize} && g(x)(x_1x_2 - N)/(x_1x_2 - N + \varepsilon), \quad \varepsilon \in V(0, \varepsilon_0), \quad \varepsilon \text{ is fixed,} \\
 &\text{subject to} && \log(N) - \log(x_1x_2) \leq 0, && (2) \\
 &&& 2 \leq x_1 \leq N - 1, \\
 &&& 2 \leq x_2 \leq N - 1,
 \end{aligned}$$

$$x_1 \in \mathbf{N}, \quad x_2 \in \mathbf{N}, \quad N \in \mathbf{N}, \quad \varepsilon \in \mathbf{R}, \quad \varepsilon_0 \in \mathbf{R}.$$

Note that the feasible sets of (1) and (2) are identical: it's Ω^I .

Theorem 4(Minimum Principle). *If N is not a prime, any minimizer of (2) is a solution of integer factorization problem for N and any solution of integer factorization problem for N is a minimizer of (2).*

Proof. It follows from (2) and definition of g . If N is not a prime, the target function has a minimum value of 0 at integral points that satisfy $x_1x_2 - N = 0$, since $g(x) > 0$, $\varepsilon > 0$ and $x_1x_2 - N \geq 0$ in the feasible set Ω^I . \square

Problem (2) completely satisfies Baes et al. [2] and therefore (2) and integer factorization problem can be solved in time polynomial in $\log(N)$.

Finally, we obtain the following algorithm.

Algorithm 2(Integer Factorization Algorithm).

Input: A positive integer number N ,
positive convex function g , $\varepsilon > 0$, $\varepsilon \in V(0, \varepsilon_0)$.

Output: A nontrivial divisor of N (if it exists).
Solve the problem (2) using algorithms [2]:
Based on the input data compute
a minimizer $(x_{1 \min}, x_{2 \min})$
of the problem (2).
if $(x_{1 \min} x_{2 \min} = N)$
then
Return a nontrivial divisor $x_{1 \min}$ of N
else
Return “N is a prime”

So, Algorithm 2 runs in time polynomial in $\log(N)$.

Thus, factoring is in FP(the class FP is the set of function problems which can be solved by a deterministic Turing machine in polynomial time(see e.g. Cormen et al. [3]).

Theorem 5. *Integer factorization is in FP.*

Algorithm 2 can be modified to serve the decision problem version as well - given an integer N and an integer q with $1 \leq q \leq N$, does N have a factor d with $1 < d < q$?

Let $\Omega_q := \{x \in \mathbf{R}^2, x = (x_1, x_2) \mid \log(x_1 x_2) \geq \log(N), 2 \leq x_1 \leq N - 1, 2 \leq x_2 \leq q - 1, x_1 \in \mathbf{R}, x_2 \in \mathbf{R}\}$ for a given $q, 1 \leq q \leq N, N \in \mathbf{N}$.

Let $\Omega_q^1 = \Omega_q \cap \mathbf{Z}^2$.

Let us replace (2) by the problem over the feasible set Ω_q^1 and denote the modified minimization problem (corresponding to the problem (2)) as (3).

Algorithm 3(Integer Factorization Algorithm).

Input: Positive integer numbers $N, q < N$,
positive convex function g , $\varepsilon > 0$, $\varepsilon \in V(0, \varepsilon_0)$.

Output: Existence of a factor d with $1 < d < q$.
Solve the problem (3) using algorithms [2]:
Based on the input data compute
a minimizer $(x_{1 \min}, x_{2 \min})$
of the problem (3)

```

if ( $x_1 \min x_2 \min = N$ )
then
    Return “The corresponding factor exists”
else
    Return “The corresponding factor does not exist”

```

Hence, Algorithm 3 runs in time polynomial in $\log(N)$ as well. Thus, factoring is in P. The class P is the class of sets accepted by a deterministic polynomial-time Turing machines (see, e.g., Cormen et al. [3]).

Theorem 6. *Integer factorization is in P.*

Note that algorithms 2-3 can be considered as polynomial-time primality tests and the only provably polynomial-time primality test was developed by Agrawal et al. [1].

The above-obtained results lead to solution of the famous P vs. NP problem. So we do not solve it directly, we solve it using reductions of factoring to the integer minimization and maximization problems and using U-equivalence.

Recall that (see, e.g., Hemmecke et al. [5], Section 1, Theorem 2) the minimization problem of a degree-4 polynomial over the integer points of a convex polygon is NP-hard.

The following U-equivalent problem(see Section 3, Corollary 1) to the original problem (1) is NP-hard for all $n \geq n_0 > 2$ (otherwise, for all $n > 2$, it is not NP-hard, which cannot be true for unbounded $n \in \mathbf{N}$):

$$\begin{aligned}
 &\text{minimize} && (x_1 x_2)^n \\
 &\text{subject to} && x_1 x_2 \geq N, \\
 &&& 2 \leq x_1 \leq N - 1, \\
 &&& 2 \leq x_2 \leq N - 1, \\
 &&& x_1 \in \mathbf{N}, x_2 \in \mathbf{N}, N \in \mathbf{N}, n \in \mathbf{N}.
 \end{aligned} \tag{4}$$

We developed polynomial-time Algorithms 2-3 in order to find minimizers of (1) (due to Theorems 1 and 4), associated with NP-hard problem (4),

which is u^n -equivalent (and, therefore, has the same set of minimizers) to the problem (1). It is well known that if there is a polynomial-time algorithm for any NP-hard problem, then, there are polynomial-time algorithms for all problems in NP, and hence, we would conclude that P is equal to NP.

References

- [1] M. Agrawal, N. Kayal, N. Saxena, PRIMES is in P, *Annals of Mathematics* 160(2) (2004) 781–793.
- [2] M. Baes, T. Oertel, C. Wagner, R. Weismantel, Mirror-Descent Methods in Mixed-Integer Convex Optimization, in: M. Jünger, G. Reinelt (Eds.), *Facets of combinatorial optimization*, Springer, Berlin, New York, 2013, pp. 101–131, , available electronically from <http://arxiv.org/pdf/1209.0686.pdf>
- [3] T. Cormen, C. Leiserson, R. Rivest, C. Stein, *Introduction To Algorithms*, third ed, The MIT Press, Cambridge, 2009.
- [4] S. Heinz, Complexity of integer quasiconvex polynomial optimization, *J. Complexity* 21(4) (2005) 543–556.
- [5] R. Hemmecke, M. Köppe, J. Lee, R. Weismantel, Nonlinear Integer Programming, in: M. Jünger, T. Liebling, D. Naddef, W. Pulleyblank, G. Reinelt, G. Rinaldi, L. Wolsey (Eds.), *50 Years of Integer Programming 1958–2008: The Early Years and State-of-the-Art Surveys*, Springer-Verlag, Berlin, 2010, pp. 561–618, available electronically from <http://arxiv.org/pdf/0906.5171.pdf>
- [6] L. G. Khachiyan, L. Porkolab, Integer optimization on convex semialgebraic sets, *Discrete and Computational Geometry* 23(2) (2000) 207–224.
- [7] J. Lee, On the boundary of tractability for nonlinear discrete optimization, in: *Cologne Twente Workshop 2009, 8th Cologne Twente Workshop on Graphs and Combinatorial Optimization*, Ecole Polytechnique, Paris, 2009, pp. 374–383, available electronically from <http://www.lix.polytechnique.fr/ctw09/ctw09-proceedings.pdf#page=385>
- [8] A. K. Lenstra, H. W. Jr. Lenstra, (Eds.), *The development of the number field sieve*, Springer-Verlag, Berlin, 1993.
- [9] T. Oertel, C. Wagner, R. Weismantel, Convex integer minimization in fixed dimension, *CoRR* 1203–4175 (2012).
- [10] P. Stevenhagen, *The number field sieve*, *Algorithmic Number Theory: Lattices, Number Fields, Curves, and Cryptography*, Mathematical Sciences Research Institute Publications, Cambridge University Press, Cambridge, 2008.

