The CMB and the Last-Scattering Cosmological Spheres

Sylwester Kornowski

Abstract: According to the Scale-Symmetric Everlasting Theory (S-SET), the very early Universe was the loop composed of protogalaxies built of the neutron black holes. Due to the succeeding inflows of the dark energy, there was the exit of the loop from its black-hole state and the loop transformed into the expanding sphere. The Milky-Way Galaxy is very close to the centre of the expanding Universe so we cannot see directly the first stage of evolution of the early Universe. But due to some phenomena, the early binary systems and clusters of protogalaxies are imprinted on the CMB. Due to the mergers of the protogalaxies during the unseen period of evolution, the number of observed today clusters of galaxies should be smaller than it follows from the CMB and such conclusion is consistent with the observational facts. Here, I answered following question: Why we cannot see directly the evolution of the early Universe whereas we can see the CMB with coded initial number of clusters of protogalaxies?

Introduction and motivation

Here, I answered following question: Why we cannot see directly the evolution of the early Universe whereas we can see the CMB with coded initial number of clusters of protogalaxies?

According to the Scale-Symmetric Everlasting Theory (S-SET), [1], the very early Universe was the loop composed of protogalaxies built of the neutron black holes. Due to the succeeding inflows of the dark energy, there was the exit of the loop from its black-hole state and the loop transformed into the expanding sphere. To conserve stability of the protogalaxies, there was the radial polarization of their rotational and magnetic axes, especially of protogalaxies with higher radial speeds i.e. of protogalaxies placed closer to the surface of the expanding Universe. The loop was left-handed so angular velocity and velocity of protogalaxies were antiparallel.

Due to the very high initial temperature, the electrons were separated from protons whereas due to the rotations of the protogalaxies, there appeared vortices of electrons. There as well was the radial polarization of rotational and magnetic axes of the electron vortices. Generally, the maximum radial speed of the baryonic matter was and is 0.6415c [1]. But inertia of the electrons is much lower than protons so maximum radial speed of the electron vortices was equal to the speed of light c. Of course, it was possible only at the beginning of expansion of the Universe when temperature was very high and distances between the proton and electron vortices were not big (in the cosmic scale). We can see that above the expanding baryonic front there were concentric spheres composed of the electron vortices with radial speeds of the expanding spheres from 0.6415c to c. Of course, there were such spheres as well inside

the baryonic sphere and their radial speeds changed from zero to 0.6415c. The concentric spheres were the last-scattering spheres for photons emitted by the protogalaxies. In such a way there appeared the different CMBs.

The duality of relativity follows from the quantum entanglement [2]. The big mistake in the mainstream cosmology follows from the wrong interpretation of the Michelson-Morley experiment. In reality, due to the quantum entanglement, the speed of light c is the speed of photons in relation to their source or in relation to a last-interaction object. Detectors are always the last-interaction objects so measured speed of photons is the c always. But it is untrue that a photon has simultaneously the speed c in relation to all reference systems.

The last-scattering spheres were the last-interaction objects for the photons so speed of photons was the c in relation to the spheres. The observed motions of the large-scale Universe suggest that the Milky-Way Galaxy is close to the centre of the expanding Universe. This leads to conclusion that the radial speeds of photons scattered on the spheres composed of the electron vortices (the radial speeds of the different CMBs), that are moving towards the Milky Way, change from c to zero. The CMBs were produced when the last-scattering spheres were very close to the early Milky Way. Today we observe one of the different CMBs moving with different radial speeds towards Earth. The initial radius of the loop/very-early-Universe was about 0.2 Gyr [1] whereas the correct age of the Universe is about 21 Gyr (we can see the last 13.4 Gyr only [3]), [1], [2], so we can see today the CMB composed of photons with radial speeds towards Earth close to zero – of course, speed of such photons measured by our detectors should be the c because the detectors are the last-interaction objects.

Notice that on the observed CMB should be 'imprinted' almost all protogalaxies. During the unseen initial 7.5 Gyr, due to the mergers of protogalaxies, number of clusters of galaxies decreased about 2.4 times [4] and this theoretical result is consistent with observational facts [5].

The electron vortices and proton vortices produced the E-modes in the CMB whereas the expanding weak condensates composed of the Einstein-spacetime components, produced the B-modes [6].

References

 Sylwester Kornowski (14 March 2014). "The Everlasting Theory and Special Number Theory".

http://vixra.org/abs/1203.0021 [v3].

Sylwester Kornowski (2012 - 2014); http://vixra.org/author/sylwester_kornowski .

- [2] Sylwester Kornowski (16 December 2014). "The Correct Age of the Universe". http://vixra.org/abs/1412.0184 .
- [3] Sylwester Kornowski (21 June 2014). "The Evolution of Massive Galaxies". http://vixra.org/abs/1406.0014 [v4].
- [4] Sylwester Kornowski (16 August 2014). "The Origin of the Missing Galaxy Clusters". http://vixra.org/abs/1408.0100.
- [5] Planck Collaboration. "Planck 2013 results. XX. Cosmology from Sunyaev-Zeldovich cluster counts".

arXiv:1303.5080v2 [astro-ph.CO].

[6] Sylwester Kornowski (21 March 2014). "The E- and B-Modes in the CMB". http://vixra.org/abs/1403.0501.