

The Abstract

An Approximation to the Mass Ratio of the Proton to the Electron.

The mass ratio of the proton to the electron is a dimensionless physical constant. The following expression provides a good estimate.

$$(4\pi) \left(4\pi - \frac{1}{\pi}\right) \left(4\pi - \frac{2}{\pi}\right) = 1836.15 \quad (1)$$

The above expression is also the greatest lower bound (GLB) for a more general expression. A geometric persuasion is given and the more general expression is determined via the inversion of the spheres.

The Background

My name is Harry Watson. Today I am presenting a model for an approximation. I discovered the approximation on 28 June 1989. The idea begins with a ball of radius one and a line segment of length 4π . The line segment is attached to the ball at one end point and is tangent to the ball. This is referred to as *the system*. Assume that the system is isolated in 3-space.



Figure 1: The “Stick and Ball” System

Figure 2 shows the region of space traced out by the line segment. Define the point of attachment of the line segment to the ball as the origin $(0, 0, 0)$ and the center of the ball to be $(0, 0, 1)$. We rotate the system about the three axes. Rotation about the z -axis defines a disk of radius 4π in the xy -plane.




Figure 2: The Disk in the yz -plane

In Figure 3 we see the trace of the ball. The radius of 2 is used to do the Inversion of the Spheres, or more correctly: The Inversion of the Circles in two dimensions. This inversion creates a disk of radius 4π and an inner deleted disk of radius $1/\pi$.




Figure 3: Rotation about the y -axis (zx -plane)

Rotating the system about the y -axis followed by rotation about the x -axis doubly defines the interior disk of radius 2. To accommodate the area, doubling the area requires the interior disk in the yz -plane to have the effective radius of $2\sqrt{2}$.




Figure 4: Rotation about the x -axis (yz -plane)

Multiply the three radii measures: 4π , $4\pi - 1/\pi$, and $4\pi - 2/\pi$; this product nearly equals the mass ratio of the proton to the electron, a dimensionless physical constant. These operations underwrite an interesting result:

$$4\pi(4\pi - 1/\pi)(4\pi - 2/\pi) = 1836.15 \quad (2)$$

The Model

Preface: This short article defines a function which models some characteristics of the proton. A general function is created which gives a close approximation to the proton-electron mass ratio. The *greatest lower bound* (GLB) is derived as a limiting case for the ratio.

It has been said that all models are wrong, but some are useful.¹ The proton is no longer considered a fundamental particle; it admits the possibility of being modeled by fundamental particles. In particular, we consider the “ball and stick” model. Let M_e denote the mass of the electron and M_p denote the mass of the proton. In a gravitational field, $M_p/M_e \approx 1836.15$. Let $\chi(x)$

¹Box, George E. P., (1919-2013) British mathematician

and $\xi(y)$ be defined as follows.

$$\chi(x) = \begin{cases} 1 & \forall 2/\pi \leq x \leq 4\pi \\ 2\varepsilon(x/2) & \forall 0 \leq x < 2/\pi \end{cases}$$

$$\xi(y) = \begin{cases} 1 & \forall 1/\pi \leq y \leq 4\pi \\ \varepsilon(y) & \forall 0 \leq y < 1/\pi \end{cases}$$

The function ε maps the interval $[0, 1/\pi]$ into the interval $[0, 1]$ such that $\varepsilon(0) = 0$, $\varepsilon(1/\pi) = 1$, and $\forall y \in (0, 1/\pi)$, $\varepsilon(y)$ is the fraction of surface area that particles of positive mass intersect with a sphere of radius $4/y$ and center $(0, 0, 0)$. Recall that a sphere of radius r is the point set $\{(x, y, z) | \sqrt{x^2 + y^2 + z^2} = r\}$. The Inversion of the Spheres sends y to $4/y$. Thus we have an elementary equation.

$$M_p/M_e = \int_0^{4\pi} \int_0^{4\pi} \int_0^{4\pi} \chi(x) \xi(y) dx dy dz \quad (3)$$

The Environment The environment external to the closed ball of radius 4π affects the determination of the mass ratio. This is modeled in Equation (3) by the functions χ and ξ , which are, in turn, determined by the function ε , described above.

An Integral The function ε is dependent on the test environment. A related variable τ is defined as follows.

$$\int_0^{4\pi} \int_0^{4\pi} \int_0^{4\pi} \chi(x) \xi(y) dx dy dz = 4\pi \left(\int_0^{4\pi} \int_0^{4\pi} \chi(x) \xi(y) dx dy \right) \quad (4)$$

$$= 4\pi \left(\left(4\pi - \frac{1}{\pi} \right) + \int_0^{1/\pi} \xi(y) dy \right) \left(\left(4\pi - \frac{2}{\pi} \right) + \int_0^{2/\pi} \chi(x) dx \right) \quad (5)$$

$$= 4\pi ((4\pi - 1/\pi) + \tau) ((4\pi - 2/\pi) + 4\tau) \quad (6)$$

Let $M_p/M_e = 1836.15267245$. (This the current recommended value.) Plug in the value $\tau = 0.0000038295$ to give 1836.1526724595 . Therefore, a small change in τ results in a significant change in the result. This compares with $(1836.15267245 - 4\pi(4\pi - 1/\pi)(4\pi - 2/\pi))/1836.15267245 = 0.000000508$.

Setting χ and ξ equal to zero in the regions $[0, 1/\pi]$, and $[0, 2/\pi]$, respectively, an interesting integral can be obtained. This is the *Greatest Lower Bound* (GLB) that can be determined by experimentation.

$$M_p/M_e \approx \int_0^{4\pi} \int_{\frac{1}{\pi}}^{4\pi} \int_{\frac{2}{\pi}}^{4\pi} dx dy dz = 1836.15 \dots \quad (7)$$

$$\int_0^{4\pi} \int_{\frac{1}{\pi}}^{4\pi} \int_{\frac{2}{\pi}}^{4\pi} dx dy dz = (4\pi) \left(4\pi - \frac{1}{\pi}\right) \left(4\pi - \frac{2}{\pi}\right) \quad (8)$$

Setting χ and ξ equal to one in the regions $[0, 1/\pi]$, and $[0, 2/\pi]$, respectively, an another interesting integral can be obtained:

$$\int_0^{4\pi} \int_0^{4\pi} \int_0^{4\pi} dx dy dz = (4\pi)^3 = 1984.40 \dots \quad (9)$$

This is what would model the event where there was no inversion of the spheres. This model gives the appearance of inherent instability.

13 November 2014.
 Harry Watson
 Eastvale, CA 92880