Electric Dark Matter Black Holes Outside and Inside Galaxies possibly related to the creation and contraction of a Cyclic Universe.

Author: Leo Vuyk.
Architect,
Rotterdam, The Netherlands.

Abstract,

For years, Stephen Hawking (and Hartle) has been applying the No Boundary Proposal, or the idea that the universe is "closed" and can have a beginning, and an end, however still exist forever.

I will call this a closed cyclic universe and for symmetry reasons a cyclic entangled raspberry shaped multiverse based on my proposal for the Quantum Function Follows Form (FFF) model. Accordingly, the Form and Microstructure of elementary particles, is supposed to be the origin of Functional differences between Higgs- Graviton- Photon- and Fermion particles.

As a consequence, a new paradigm splitting, accelerating and pairing massless Dark Matter Black Hole (without Graviton production) is able to convert vacuum energy (ZPE) into real energy by entropy decrease and quark electron positron production at the black hole horizon. Secondly, a Chiral dark energy Higgs vacuum lattice, is able to explain quick Gas and Dust production without large annihilation at the black hole horizon and Galaxy- and Star formation.

Even microscopic black holes like Sunspots, (Micro) Comets, Lightning bolts, Sprite Fireballs and Ball Lightning seem to produce gas.

However only the largest Primordial Galaxy Anchor Black Holes (GABHs) located far outside galaxies, are supposed to stop gas and dust production and by merging with other black holes they should be one of the two origins of universal contraction into a Big Crunch and a cyclic multiverse.

A second origin of universal contraction seems to be the growing central black holes in the centres of massive galaxies even in the Milky Way.

The recent observation of a central cluster black hole of 660 BILLION times the mass of the sun is reason to suggest that these black holes are the real start of the big crunch.

Such huge black holes are supposed to be the result of merger supernova black holes and growing during the galaxy lifetime, but also by the merging of large GABHs (Galaxy Anchor Black Holes) just in the centres of galaxy clusters.

Black holes are supposed to be massless, because only fermions are able to emit gravitons. The recent LIGO merger of 29x36 solar mass BH result is therefore interesting and should be based on BH size or better: Higgs content, instead of real mass.

As a consequence, the size of the first growing central galaxy black hole has about the same size as two merger primordial supernova black holes of the first stars of the galaxy

Thus: Black Holes Outside Galaxies are primarily responsible for the creation of galaxies and growing central galaxy black holes are assumed to be responsible for the contraction start of our Cyclic Universe.

The Hubble redshift, in contrast with the mainstream, is supposed to originate local vacuum lattice dilution created by local dark matter black holes.

All black holes are supposed to eat the local Higgs vacuum lattice with a local Planck length extension- as the result and the base for frequency decrease of a passing photon.

GABHs are mostly only observable by their background lensing effect and as quasi-soft x-ray sources, to be found in particular around merger galaxies, due to swirling gas and dust expelled from the galaxies.

.

Introduction.

The NEW Plasma Creating Electric Dark Matter Black Hole versus the Hawking Black Hole, according to Quantum FFF Theory, the origin of Comet - Sunspot and Ball Lightning mysteries.

I am convinced that Stephen Hawking did not calculate with the possibility of a chiral oscillating Higgs field vacuum lattice combined with propeller shaped Fermions.

Then, due to Vacuum chirality, Electron- and Positron propellers could both pushed away from the BH horizon after spin flip polarization at different distances, forming two charged separated spheres. With quark (plasma) formation in between.

Based on such a simple object (propeller and process) oriented ontology, Black Holes could be imagined as charge splitters violating the 2e law af thermodynamics, combined with a continuous microscopic big bang plasma creation process!

According to the new BH paradigm, electrons and positrons are both pushed away from the Black Hole horizon at different distances, forming TWO different charged separated spheres, with Quark (plasma) formation in between.

How is this possible?

Only if the oscillating Higgs vacuum Lattice (tetrahedral shaped) has a preferred chirality. (figure: 7,8,).

So in our Material universe, all the Higgs oscillations follow spiral vacuum lattices. The vacuum lattices form together the edges of a 3 dimensional tetrahedral system. Each lattice is supposed to be shaped as a left handed spiral and divided into at least three connected parts.

As a result all neutrinos are left handed! because the neutrino information travels in the form of an Higgs oscillation along these edges.

Each lattice is equipped with two opposite oscillating Higgs field particles, able to transfer photonic and neutrino information with the local speed of light.

Neutrinos and photons are suppose to travel not as particles but as specific Higgs oscillations along different lattice paths of the chiral vacuum lattice, with the local speed of light..

At the same time, positrons must have a preferred spin (OLO left=L) they catch a Gluon to form a u-quark (OLO+ROR) which is better protected for the aggressive oscillations of the left handed Higgs field than an electron ORO combined with the mirror Gluon LOL forming a right handed anti- d-Quark! (figure: 3 and ref: 4)

As a consequence, New paradigm Black Holes are CHARGE SPLITTERS violating the 2e law, combined with a micro big bang plasma creation process!

Explanation: it is assumed that Quantum fluctuations around strongly curved (small) BH horizons produce pairs of leptons. e-,e+ and even compound quarks (d, u, etc, see Fermion 3D string propeller theory: Fermion repelling by Lorentz polarized spin flip). (**figure: 9**)

As a consequence, the Big Bang did not produce instantly all the Fermions in the universes. Even now lots of Fermions are produced- in the form of negative charged plasma- by most BHs (see Comets, ball lightning.)

Small Black holes (in Starspots, Ball Lightning and Comets like 67P- CG) are hot and produce gas as observed: $. [N II], [OIII], and H\alpha$.

The result can be described as follows:

- 1: Black holes are the same as Dark Matter, they all consume photons, even gravitons and the Higgs field, but REPEL Fermions due to their propeller shape. They produce electric charged plasma.
- 2: Dark Energy is the oscillating (Casimir) energy of the Higgs Field equipped with a tetrahedron lattice structure with variable Planck length..
- 3: Quantum Gravity = Dual Push gravity = Attraction (Higgs-Casimir opposing Graviton push).
- 4: The Big Bang is a Splitting dark matter Big Bang Black Hole (BBBH), splitting into smaller Primordial Big Bang Splinters (PBBS) forming the Fractal Lyman Alpha forest and evaporating partly into a zero mass energetic oscillating Higgs particle based Higgs field.
- 5: Dual PBBSs hotspots, produce central plasma concentration in electric Herbig Haro systems as a base for star formation in open star clusters as a start for Spiral Galaxies.
- 6: Spiral Galaxies will keep both Primordial Dark Matter Black Holes as Galaxy Anchor Black Holes (GABHs) at long distance.
- 7: After Galaxy Merging, these GABHs are the origin of Galaxy- and Magnetic field complexity and distant dwarf galaxies .
- 8: Black Holes in general produce Plasma direct out of the Higgs field because two Higgs particles are convertible into symmetric electron and positron (or even dual quark-) propellers (by BH horizon fluctuations).
- 9: The chirality of the (spiralling) vacuum lattice is the origin our material universe. (propeller shaped positrons merge preferentially first with gluons to form (u) Quarks to form Hydrogen. 10: The first Supernovas produce medium sized Black Holes as the base for secondary Herbig Haro systems and open star clusters.
- 11: ALL Dark Matter Black Holes are supposed to be CHARGE SEPARATORS with internal positive charge and an external globular shell of negative charged Quark electron plasma.
- 12: The lightspeed is related to gravity fields like the earth with long extinction distances to adapt with the solar gravity field.
- 13. Quantum FFF Theory states that the raspberry shaped multiverse is symmetric and instant entangled down to the smallest quantum level. Also down to living and dying CATS in BOXES.
- 14 Large Primordial Big Bang Splinters (PBBS) are responsible for the creation of the Lyman Alpha forest structure and first spiral galaxy forming of the universe, but they seem to be also responsible for the implosion of the universe at the end in the form of Galaxy Anchor Black Holes (GABHs) located mainly outside galaxies. see: (Quasisoft Chandra sources)

15, As a consequence of 12-14, Time is always local time in combination with cyclic period time of the multiverse.

16, If our material universes has a chiral left handed oscillating Higgs field, then our material Right Handed DNA helix molecule could be explained.

However it also suggests that in our opposing symmetric ANTI-MATERIAL entangled neighbor universe the DNA helix should have a Left handed spiral.

According to Max Tegmark: in an entangled multiverse we may ask: is there COPY PERSON over there, who is reading the same lines as I do?

If this COPY person is indeed living over there, then even our consciousness should be shared in a sort of entangled DEMOCRATIC form.

Then we are not alone with our thoughts and doubts, see:

Democratic Free Will in the instant Entangled Multiverse.

17, All particles and waves should be entangled with anti-copy particles and Higgs field based waves in our opposing anti universes.

Then wave particle duality and the so called one particle double slit interference could be explained by accepting that real shaped 3D photon particles only exist in their function of Gluons. However Gluons should transfer form information to the oscillating luminiferous Higgs field ether, within about 1 cm distance from a fermion.(see: Babock and Bergman anomaly,1964)

- 18, Different elementary particles have different qualities by their different complex stringy shape. Leptons and Quarks have a propeller shape with left or right handed pitch creating charge difference. Gluons, Photons and Neutrino particles have no pitch.
- 20 All particles can be converted to one or more Higgs particles, Lepto Quarks can be compound particles.
- 21 All particles are assumed to be present as Higgs particles already in the cyclic Big Bang Black Hole (BBBH) after the former multiverse has collapsed into a Big Crunch Black Hole.
- 22, We seem to live inside one material universal bubble of an instant entangled (Charge Parity Time) CPT symmetric raspberry bubble multiverse. Each quantum jump or wave function collapse or human choice is guided by this instant entanglement at long distant.

If our material universes has a chiral oscillating Higgs field, then our material Right Handed DNA helix molecule could be explained.

However it also suggests that in our opposing ANTI-MATERIAL multiverse neighbour universe the DNA helix should have a LEFT HANDED spiral.

According to Max Tegmark: in an entangled multiverse we may ask: is there COPY PERSON over there, who is reading the same lines as I do?

If this COPY person is indeed living over there, then even our consciousness should be shared in a sort of DEMOCRATIC form,


Then we are not alone with our thoughts and doubts, see:

Democratic Free Will in the instant Entangled Multiverse.

http://vixra.org/pdf/1401.0071v2.pdf

Dual entropy trajectories of each cyclic universe


Big Bang and Big Crunch originated by the splitting pairing and recycling via supernova black holes, merging into each growing Galaxy central black hole leading to the cyclic Big Crunch.



Figuur 1. Imaging the cyclic multiverse, according to Quantum FFF Theory.


Super Massive Black Holes, Young and Old

Old SMBHs are supposed to be created instantly by the BH splitting Big Bang inflation epoch. Young SMBHs are the result of merger S-Nova black holes located in the center of galaxies, starting the Big Crunch. (according to Quantum FFF Theory)



Figuur 2. Imaging the cyclic multiverse. according to Quantum FFF Theory


Galaxy Anchor Black Hole (GABH)-Effect on High speed gas ejection.



Massive molecular outflows and negative feedback in ULIRGs observed by Herschel-PACS

Figure 3 Massive molecular outflow, observed by Hershell, assumed to be related with GABHs (primordialGalaxy Anchor BlackHoles.)

Galaxy Anchor Black Holes (GABHs) are supposed to be the centers of Dwarf spheroid star clusters located in-and around the Milky Way according to Quantum FFF-theory. A recent detected 27.000 light-year long remnant x-ray jet is the tell tale of a primordial HH system.



Below: Merging spiral galaxies with dual Herbig Haro object (Black Hole) systems, original connected by x-ray jets between multiple Electric Dark Matter Black Holes.





Figure 4. Milky way X-ray jet 27.000 lightyear long assumed to connect two giant primordial GABHs..

Elliptical NGC 1275 with red colored linear xray jet structure between dual Galaxy Anchor Black Holes (GABHs: Former Herbig Haro Objects) according to Quantum FFF Theory.



Below: Merging spiral galaxies with dual Black Hole systems, (former HH objects) become ellipticals like NGC 1275, with multiple Electric Dark Matter Black Holes




Figure 5. NGC 1275 with clear ion jets into different directions.

A new class of Quasisoft X-ray sources found around elliptical and spiral galaxies. According to Quantum FFF theory, these sources should be the tell tales of the creator black holes of the universe and the Lyman Alpha forest distribution of galaxies. However they also could be the future executioner of the universe. (photos: Chandra/Harvard)




Figure 6. A new class of Quasisoft X-ray sources found around elliptical and spiral galaxies. According to Quantum FFF theory, these sources should be the tell tales of the creator black holes of the universe and the Lyman Alpha forest distribution of galaxies.

Possible candidates for GABHs are a new class of Dark Matter Black Holes found by Chandra observations. (figure 6)

http://chandra.harvard.edu/photo/2004/m101/

Ouotation;

"Chandra observations of the spiral galaxy M101 and several other nearby galaxies have revealed a possible new class of X-ray sources. These mysterious X-ray sources, marked with green diamonds in the image, are called "quasisoft" sources because they have a temperature in the range of one to four million degrees Celsius.

The power output of quasisoft sources is comparable to or greater than that of neutron stars or stellar-mass black holes fuelled by the infall of matter from companion stars. This implies that the region that produces the X-rays in a quasisoft source is dozens of times larger.


One explanation is that these sources are produced by intermediate-mass black holes that have masses a hundred or more times greater than the mass of the Sun. Such objects would have much larger event horizons, which would account for the larger sizes and lower temperatures associated with the quasisoft sources. Alternatively, they could be standard neutron stars or stellar black holes where the associated region of hot gas is for some as yet unknown reason much larger than usual.

"Hard" X-ray sources, such as neutron stars or stellar-mass black holes have temperatures greater than ten million degrees. The "supersoft" X-ray sources due to white dwarfs have temperatures of several hundred-thousand-degrees. A total of 72 quasisoft sources were discovered in the four galaxies: M101, M83, M51, and NGC 4697.


Until a few years ago, astronomers only knew of two sizes of black holes: stellar black holes with masses about ten times that of the Sun, and supermassive black holes located at the centers of galaxies, with masses ranging from millions to billions times that of the Sun. Recent evidence, however, suggests that a class of "intermediate-mass" black holes may also exist. Theories for the formation of intermediate-mass black holes involve the collapse of extremely massive stars, or the merger of many stellar-mass black holes.

As more quasisoft sources are discovered, the types of galaxies in which they reside and where they are located in a galaxy should give astronomers additional clues as to their nature and how they are formed. The present study indicates that quasisoft occur in various locations in elliptical as well as spiral galaxies." (End of the quote) (**figure 6**)

New Dark Matter Black Hole, "Eat" the chiral oscillating Higgs vacuum.(Quantum-FFF Theory)



Symplified tetrahedral vacuum lattice structure (left), constructed by complex chiral tetrahedral building blocks (right). Each universe has to choose between a left- or righthanded rotation of its tetrahedral building blocks. Each chiral lattice has the Planck length and is supposed to be equipped with at least two oscillating massless new Higgs particles.



Black holes EAT the vacuum lattice with the oscillating Higgs particles, So, the vacuum lattice structure around a black hole could be represented like

this.

As a result, the near vacuum is thinner or diluted, it has holes, which could be the origin of the Universal HUBBLE redshift. By contrast, my new black hole paradigm says: the origin of universal redshift is Dark matter. Dark matter is the proliferation of vast numbers of Black holes around Galaxies. Conclusion, based on Hubble redshift, we don't know anything about expansion or contraction of the universe. (Author Leo Vuyk)

Figure 7, The new pardigm black hole with chiral Higgs field vacuum lattice.




Figure 8, Fermion propellers (green) Hggs (red) Graviton (blue) and 5x Gluons/ Photons (Yellow).

In contrast with mainstream physics, Fermion / GAS production around Cometary BH nuclei is observed at comet 67P CG.

This is also observed around small (Herbig Haro: HH systems) black hole horizons and is assumed to happen during and after fractal inflation of the multiverse.

After Supernova explosion, at least one BH is created, able to form a new HH systems after meeting with a second BH of the same size. Meeting with a larger BH will split the larger BH into two equal parts, as observed in the Eagle nebula.

At the end of the multiverse, it is supposed that most smaller dark matter black holes (hot dark matter, like Herbig Haro objects) have merged into large Monster BHs. (cold dark matter). Giant BHs (Galaxy Anchor Black Holes: GABHs) do not produce gas or dust due to their large angle of horizon curvature.

(Ref. 15) viXra:1111.0061

"Black Hole Horizon Curvature Dependent Balance Between Plasma Creation and e-e+ Annihilation in Quantum FFF Theory"

It is said: "Our work strongly reinforces the theory that supernovae are producing the dust seen in galaxies of the early universe,"

The leading theory is that supernovae - stars that explode at the end of their lives - contain large amounts of metal-enriched material that, in turn, harbours key ingredients of dust, like silicon, iron and carbon.

The NEW Plasma Creating Electric Black Hole versus the Hawking Black Hole, according to Quantum FFF Theory, the origin of Comet - Sunspot –BL mystery.





Figure 9, The New paradigm: the Charge splitting and plasma producing dark matter black holes.

The Beginning and End of Time. Multiverse cycle time versus quantities of real (SM) particles, Dark Matter Black Hole (BH) particles and Dark Energy Higgs /Casimir field particles.

The cyclic Multiverse depicted as continuous flow of increasing and decreasing numbers of Higgs particles located inside Dark Matter Black Hole Nuclei, or free oscillating in the vacuum lattice or derived as real SM particles.



A= The End of Time of the former multiverse.

All SM (StandardModel) particles equipped with internal pressurized tension are located inside the Big Crunch Black Hole Nucleus, compressed by the decreasing- pressure and number of the oscillating Higgs /Casimir vacuum particles, with the Big Bang explosion as result.

B= The beginning of Time and expansion by Big Crunch/ Big Bang Nucleus splitting into primordial BHs into a fractal shaped Lyman Alpha forest, the start of pairing and plasma forming dual Herbig Haro BH hotspots with bullet stars in between, forming galaxies by a fast Super Nova proces. Primordial BHs partly evaporate into the Dark Energy Higgs field.

C= The start of the Multiverse vacuum contraction due to vacuum absorbtion by the growing numbers of dark matter (Supernova) black holes.

The HUBBLE REDSHIFT is supposed to be A SIDE EFFECT of proliferated dark matter black holes of various sizes ABSORBING and DILUTING the Higgs vacuum lattice, the origin of local variable PLANCK lengths and LIGHTSPEED.

Examples are (star-)sunspot BHs- Cometary nuclear BHs, DUAL Herbig Haro BHs as presursors of SABHs and GABHs (Stellar- or Galaxy Anchor BHs).

D= The total number of particles during the cyclic process is constant!

According to the Quantum Function Follows Form model.

See: https://tudelft.academia.edu/LeoVuyk/Papers

Figure 10., the beginning and end of time in a cyclic multiverse.

The Beginning and End of Time. (figure 10)

Multiverse cycle time versus quantities of real (SM) particles, Dark Matter Black Hole (BH) nuclear particles and dual oscillating Dark Energy Higgs /Casimir field particles.

The entangled cyclic Multiverse depicted as continuous flow of increasing and decreasing numbers of Higgs particles located inside Dark Matter Black Hole Nuclei, or free oscillating in the vacuum lattice or derived as real SM particles.

A= The End of Time of the former multiverse.

All SM (Standard Model) particles equipped with internal pressurized tension are located inside the Big Crunch Black Hole Nucleus, compressed by the decreasing- pressure and number of the oscillating Higgs /Casimir vacuum particles, with the Big Bang explosion as result.

B= The beginning of Time and expansion by Big Crunch/ Big Bang Nucleus splitting into primordial BHs into a fractal shaped Lyman Alpha forest, the start of pairing and plasma forming dual Herbig Haro BH hotspots with bullet stars in between, forming galaxies by a fast Super Nova proces. Primordial BHs partly evaporate into the Dark Energy Higgs field. C= The start of the Multiverse vacuum contraction due to vacuum absorption by the growing numbers of dark matter Supernova black holes.

The Hubble Redshift is supposed to be a side effect of proliferated dark matter black holes of various sizes absorbing and diluting the Higgs vacuum lattice, the origin of local variable Planck lengths and local variable gravity direction related lightspeed.

See also: Cause of galactic death: Strangulation:

http://www.sciencedaily.com/releases/2015/05/150513132650.htm

Black hole Examples according to Quantum FFF Theory are: (star-)sunspot BHs- Cometary nuclear BHs, DUAL Herbig Haro BHs as precursors of SABHs and GABHs (Stellar- or Galaxy Anchor BHs).

D= The total number of particles during the cyclic process is constant! According to the Quantum Function Follows Form model.

The Raspberry geometry of the 12 fold Dodecahedron Multiverse according to Quantum FFF Theory.





Figure 11. The reaspberry multiverse.


Plasma production by trailing dual related starspot dark matter black holes, expelled inside GK Persei Nebula. According to Quantum FFF Theory.



Rd is [N II], green is [OIII], and blue is Hα. See: http://iopscience.iop.org/1538-3881/143/6/143/article#aj429218f1e

Figure 12., Plasma/gas production by former Starspots.

Bi-polar (A) and multipolar (B) Planetary Nebula related to Dark Matter Black Hole based (DMBH) Starspots ejection (C). DMBHs aim by ZPE to pair (D) and form Herbig Haro systems creating binary stars in between these DMBHs called SABHs. (E)



According to Quantum FFF Theory, dual Herbig Haro hotspots are former DMBHs called SABHs Stellar Anchor Black Holes with binary star creation in between, most at the top of dust pillars of creation inside older nebula.(E) This process is also the base of galaxy formation in the early universe.

Figure 13., Bi and multipolar nebula the result of former Herbig Haro objects depicted as Stellar Anchor Black Holes. (SABHs).

Charge splitting black holes and the mysterious dual pairing to form Herbig Haro objects, galaxies and starclusters. Black holes create negative charged plasma and




Figure 14 Plasma production around Herbig Haro objects and early galaxy Anchor Black Holes (GABHs).



Figure 15, Examples of micro black holes on earth.

The origin of Gamma Ray Bursts, by splitting black holes of different size, leaving three BHs behind. Two of them could form a Herbig Haro Object with a dumbbell structure. (Quantum FFF theory@Leo Vuyk).

TWO REPULSIVE ION TAILS around electron jets (bar shaped) between two DIFFERENT sized black holes. THE SMALLER BLACK HOLE WILL SPLIT THE LARGER ONE BY ITS CONCENTRATED ELECTRON JET AND VACUUM INFLUENCE. Author: lee Vuyk.

SPLITTED

CASIMIR FORCE

SPLITTED

GAMMA RAY BURST

CASIMIR FORCE


SPLITTED

Only the smallest BH is able to split the larger black hole, leading to a proliferation of smaller equally sized BHs and Herbig haro Objects.

Figure 16, Splitting BH process observed in nebula like the Eagle nebula.

Typical form example of pairing black holes of the same size, leaving ion tails and a bar behind before separating to a stable dumbbell position@Leo Vuyk





The casimir pressure structure of the Higgs dark energy vacuum around Pairing black holes

Figure 17, Even in galaxy structures we may find remnants of the large scale BH splitting proces

Splitting and merging boundaries.

Dual Black Hole merger and splitter Higgs content "size" ratios. See Quantum FFF Theory based new black hole paradigm.

LIGO ratio: 0.8x (29:36)

$$\bigcirc$$
---- \bigcirc

Herbig Haro creation ratio: 1x

BH Splitter ratio <0.8x >0.01?

Galaxy central merger ratio: <0.01?

Figure 18, According to LIGO, the "virtual mass" or Higgs content ratio of the merger BHs was 0.8x. If Herbig Haro systems are only created by a 1.0 ratio BH system, then we are interested in what would be the ratio boundary for Central merger BHs (perhaps 0.01x?)

As a consequence, the size of the first central galaxy black hole has about the same size as two merger primordial supernova black holes of the first stars of the galaxy.




Figure 19,, the 3D geometrical proposal for the Standard Model.




Figure 20,, Birkeland and Alfven current circuits around galaxies and Herbig Haro systems, for quick star formation in the centre.

Black holes create negative charged plasma and Herbig Haro Objects to form stars in between.

TWO pairing and receding micro black holes, propelled by unstable ionised gas tails, as observed in many Nebula: Carina, Eagle, Crab etc. creating HH objects or Gas Pillars. Black hole leaving the Tail B back place for stable Herbig Haro tail location (A) or unstable gas PILLAR creation (B). The unstable tail swing **√** Jet is guided by external Tail B new micro blac gravity sources like the Tail A swing into a RECEDING opposing micro black BAR between both Bhs Author: Leo Vuyk hole gravity and other more distant (Nebula) **√** Jet sources. 2 pairing Black hole leaving the meeting В place for stable Herbig Haro micro black location (A) or unstable gas holes PILLAR creation (B)

Quantum FFF Theory says: Infalling observers (in BHs) and self created plasma will be pushed out (by fermion spin flip) and negative charged. The result is that the plasma will act as pushing accelerator to accelerate the BH away from nearby larger (BH) gravity sources.

Consequiently, equal sized black holes will find each other and form a so called plasma BAR, at a critical distance, but split in succession to form an electric Herbig Haro system as the base for an open star cluster.

In the early universe however, these star cluster will grow into primordial galaxies. Those systems are observed as radio galaxies (below)




Figure 21, (Below) Galaxy formation in the early universe by dual Big Bang Black Hole Splinters. Able to form dual systems. (Above) The Eagle nebula seems to be an example of pairing and splitting black holes to form Herbig Haro systems with star formation in between.

FLAT Pillars of Creation in 3D, support the Electric Dark Matter Black Hole pairing and splitting process into Herbig Haro systems, according to Quantum FFF Theory.




Figure 22. Flat Pillars of Creation in 3D. compared to New paradigm Dark Matter Black Hole pairing and splitting into a Herbig Haro System, according to Q-FFF Theory.

The Contracting Universe.

Diluted Higgs field Lattice structure (pink) with current circuits between two Dark Matter Black Hole based Galaxy Clusters (blue), merging in the future! According to Quantum FFF Theory.




Figure 23, The Contracting Universe.

Diluted Higgs field Lattice structure (pink) with Birkeland Alfven current circuits between two Dark Matter Black Hole (DMBH) based Galaxy Clusters (blue), expected to merge in the future! According to Quantum FFF Theory. Compared with simple dual DMBH systems of stars.

References:

[59] The New God Particle and Free Will. Author: Leo Vuyk, LuLu publishers, 2008. ISBN number 978-1-4092-1031-3 http://www.lulu.com/spotlight/LeoVuyk

[58] Quantum FFF Theory is also published in the form of POSTERS at the Flickr site:

 $\underline{http://www.flickr.com/photos/93308747@N05/}$

[57] Numbered listing of Vixra essays by Leo Vuyk (see below).

http://vixra.org/author/leo_vuyk

[56] viXra:1502.0086

Parity and Chirality as a Base for Quark-Gluon Plasma Creating and Charge Splitting Black Holes.

[55] viXra:1411.0039

Qauntum FFF Theory, Poster News 2

[54] viXra:1410.0039

Evidence for Dark Matter Black Hole Based Plasma, Dust and Ice Production Inside Comets Like 67P,C-G. and the Growing Earth.

[53] viXra:1409.0164

The O'Connell Effect in Eclipsing Binaries Explained by Mass Related Light Speed Extinction Distances (Lasof) of Stars and Even Planets.

[52] viXra:1407.0001

Evidence for Evaporating Dark Matter Particles in Silicon Fireballs.

[51] viXra:1406.0076

Galaxy Formation and Evolution According to Quantum FFF Theory.

[50] viXra:1406.0019

Quantum FFF Theory Proposals for Some Unsolved Physics Problems.

[49] viXra:1405.0224

Dual Star Ejection in Open Star Clusters by the Central Star.

[48] viXra:1404.0151

Future Free Energy of New Physics Ball Lightning Black Holes, the Origin of a Super Social Society or the Opposite?

[47] viXra:1404.0002

Proposal for the Origin of Unexpected Large B-Modes Found in the Bicep2 Measurements.

[46] viXra:1402.0132 Reconciliation of GR and QM by a New Gravity-Black Hole and Lightspeed Model Called Quantum- FFF Theory.

[45] viXra:1402.0044

The Impossible Zero Point Electric Black Hole as the Origin of New Physics even for Comets like 67P,C-G.

[44] viXra:1401.0115

Calabi Yau Shaped Double Fermion Spin States.

[43] viXra:1401.0071

Democratic Free Will in the Instant Entangled Multiverse.

[42] viXra:1312.0143

The Navel Cord Multiverse with Raspberry Shape.

[41] viXra:1312.0076

Are Sunspots Made of Gravitating Dark Matter Black Holes?

[40] viXra:1310.0101

The New Nuclear Magic Number (34) Explained by the Polar Coaxial Ring System of

Category: Nuclear and Atomic Physics

[39] viXra:1310.0059

The Semi Relativistic Higgs Field Aether with Mass Related Lightspeed Adaptation.

[38] viXra:1309.0081

New Physics by Table Top Experiments.

[37] viXra:1308.0083

The Raspberry Shaped Super Symmetric Multiverse

[36] viXra:1307.0068

Quantum Function Follows Form Theory, the Small Scale, Posters part 1.

[35] viXra:1307.0067

Quantum Function Follows Form Theory, the Large Scale, Posters part 2.

[**34**] viXra:1306.0218

Dark Matter UFO Dust Particles or Quantum Knots in the LHC at CERN?

33, viXra:1306.0065

The Splitting Dark Matter- Black Hole- Big Bang and the Cyclic Multiverse.

32, viXra:1305.0140

Quantum FFF Theory Experiments, a Summary.

31, viXra:1305.0041

The Shnoll Effect Explained by Quantum FFF Theory.

30, viXra:1304.0022

The Self Organizing Universe in the Carina Nebula.

29, viXra:1303.0211

Hubble Redshift Combined with Universal Contraction is Possible.

28, viXra:1303.0053

Micro Birkeland Currents Between Two RF Generated Plasma Balls in the Laboratory.

27, viXra:1302.0172

Silicon Based Ball Lightning Globule Structures and Signs for Accumulation and Retarded Decrease of Tunnelling Energy Bullets.

26. viXra:1301.0088

Two or Three Large Quasar Groups (LQGs) Located at the Start of Two or Three Lyman Alpha Systems and a Part of the Raspberry Multiverse?

25, viXra:1301.0050

Dwarf Galaxies and Their Relation with Dark Matter Based Galaxy Anchor Black Holes (Gabhs).

24, viXra:1210.0177

Instant Broglie Bohm Pilot Waves, the Origin of All Entanglement Effects in the Lab and Wavefunction Collapses in Our Universe as Related to Our Opposing Anti-Copy Universe(s) According to Quantum FFF Theory.

23, viXra:1209.0092

New Dark Matter Black Holes and a New Dark Energy Higgs Field, Lead to a Bouncing CP Symmetrical Multiverse, and New Experiments.

22, viXra:1209.0061

Birkeland Currents, Sunspots, Comets and Ball Lightning Originated by New Paradigm Dark Matter Black Holes.

21, viXra:1209.0030

Majorana and Sterile Neutrino Solutions in the Quantum-FFF Model.

20, viXra:1208.0031

Clumpy Dark Matter Around Dwarf Galaxies a Support for an Alternative Black Hole Theory According to the Quantum Function Follows Form Model.

19, viXra:1202.0091

Earth Magnetic Monopole Array Field Interaction with Cyclotron –Synchrotron Electrons and Muon Conversion Used for Levitation Systems.

18, viXra:1201.0092

Earth Magnetic Monopole Array Field Interaction with Cyclotron Electrons used for Levitation Systems.

17, viXra:1112.0065

LHC Signals Between 121-130 Gev Interpreted with Quantum-FFF Theory

16, viXra:1111.0096

Reconciliation of QM and GR and the Need for a Pulsating Entangled CPT Symmetric Raspberry Shaped Multiverse.

15, viXra:1111.0061

Black Hole Horizon Curvature Dependent Balance Between Plasma Creation and e-e+ Annihilation in Quantum FFF Theory.

14, viXra:1108.0036

Artificial Ball Lightning Production and Exploitation Device for Zero Point Electric Energy Usage.

13, viXra:1108.0006

Mass in Motion in Quantum FFF Theory

12, viXra:1104.0083

Quantum FFF Theory in Posters.

11. viXra:1104.0044

Ball Lightning, Micro Comets, Sprite-Fireballs and X-Ray/gamma Flashes According to Quantum FFF Theory.

10, viXra:1104.0002

Stellar Anchor Black Holes as the Remnants of Former Herbig Haro Objects

9, viXra:1103.0097

ZPE Zero Point Energy Examples Around Black Holes.

8, viXra:1103.0068

Funktion Follows Form, at the Quantum Scale and Beyond.

7, viXra:1103.0024

Ouantum Gravity and Electro Magnetic Forces in FFF Theory

6, viXra:1103.0015

Wavefunction Collapse and Human Choice-Making Inside an Entangled Mirror Symmetrical Multiverse.

5, viXra:1103.0011

An Alternative Black Hole, Provided with Entropy Decrease and Plasma Creation

- **4,** viXra:1103.0002
- 3- Dimensional String Based Alternative Particles Model
- **3,** viXra:1102.0056

Experiments to Determine the Mass Related Lightspeed Extinction Volume

2, viXra:1102.0054

Atomic Nuclear Geometry Based on Magic Number Logic.

1, viXra:1102.0052

Construction Principles for Chiral "atoms of Spacetime Geometry
