UNIFIED MATTER THEORY

(Cosmology, astrophysics, physics) Original text and figures (except references) © 2015 by Prof. Solomon Budnik

President, UTG-PRI LTD., Israel budnik1@013.net s.b0246@gmail.com

Significance

This article portrays space and time as conformal projections of motion, and reveals the co-relation between the universality of spacetime (relativistic model) and energy-frequency (quantum model). This is a new theory of cosmology with a concept of **unified matter**, which created supermatter previously defined as black hole, thus negating the assumption of a gravitational collapse of a star to form a black hole. In general, there is no cosmic hole in the fabric of space where the density of space and not that of matter should be comprehended. Accordingly, the assumption that light cannot escape from the "black hole" is wrong, for to emit light, the dark matter must evolve into a bright matter when it reaches its bifurcation point.

Moreover, this article contributes to the understanding of the formation of twin accretion discs and the origin of twin relativistic jets in **gauge/gravity duality**. This concept is in conformity with the basic principle of duality in nature, where no singular event of the alleged "Big Bang" could have taken place, for there must always be <u>the cause</u> and <u>the effect</u> in a dual interaction, as in the wave-particle duality.

The Theory

1. Two parallel lines never meet in geometry, but superstrings of matter in cosmos might converge in a bow-like superstructure and produce a shock wave, as in fig. 4 and ref. 2 far below. Compare: on earth a pair of lines of longitude is parallel at the equator but converge toward the poles. So, let's consider two parallel superstrings of matter with integrated accretion discs in the middle of each one. (Fig.4 far below). Those discs are formed by two attracted linear superstrings of matter, whose central core inflates to create an accretion disc (fig. 1) in a quasi super-Planckian effect (fig 2.)



Fig. 1




Fig. 2

The accretion disc of matter in fig. 1 consists of **nuclei** (see ref. 3 below) and membrane that oscillate in unison in **unified resonant frequency** in a given frame of spacetime, which is nit trapped in geometrical dimensions, but oscillates/modulates in different atomic frequencies we cannot perceive or detect, for we're not yet "tuned" to them, as a common radio cannot pick up radio waves it is not tuned to. Therefore, our superstrings model and concept differ from the string theory model, which envisages a structure of different integrated, not unified and inharmonic shapes, which cannot co-vibrate in a general pitch. (For **pitch** see Ref. 3 below)

In our opinion, a cosmic superstring resembles a hexagonal musical string which has "core" of one material, with encircled by other materials. A musical string is linearly stretched along the fingerboard of a musical instrument, where the fingers of the musician press/short them in different intervals, while the bow pulls the string sideways to make it vibrate in various sounds. Similarly in cosmos, a superstring of matter stretches itself linearly in space, where applied gravity presses/shorts it in spacetime intervals/knots of mater and then pulls it sideways to make it resonate/oscillate in various frequencies and gravitational waves which create a supercavitation (not a black hole) in space caused by the density of coupled superstrings, as in fig. 1.

Accordingly, our superstring is not a macrojet of matter but a chain of intertwined gauge/gravity microstrings- cosmic tubes that funnel the invisible superfluid dark matter (as **continuum fluid in the phase space**) and highly compressed energy in thereby twisted spacetime (Fig. 3 a, b), which **expands and contracts** in pulsation due to inherent gravity and fluorescent mirror reflections of subatomic particles.



(**Fig. 3**, a.b. Time and space, according to Einstein's theories of relativity, are woven together). Our concept image of intertwined/woven superstring fabric as in fig. 3 has its further confirmation in **ref. 4** below.

So in our unified model, coupled superstrings unify in space via supercore (fig. 1) of the overlapping, oscillating condensed matter (fig. 2), which creates in supercavitation a superdensity vortex, previously termed the black hole that had puzzled scientists who couldn't comprehend how matter spiraling around a black hole in an **accretion disk** suddenly plunges into the void without gravitational collapse.

Nota bene: the intertwined superstring in fig. 3 has open opposite ends through which matter traverses spacetime in superconductivity contrary to the speed of light constant, for said superstring acts as a supercollider and superaccelerator in space. The inherent shape of our superstring and its internal succession of multiple micro-barriers with supercold vacuum chambers explain the phenomena of neutrinos, which propagate unobstructed through space and earth, originate in one part of the universe and immediately appear in a remote part by piercing the spacetime. For **neutrinos** the **spin** is always **opposite** the linear momentum, so in our superstring accelerator we might deal with the Casimir effect and particles that emit the virtual photons which lose momentum **p** in the recoil, and the other particles get the momentum.

The idea is that albeit a neutrino has no single mass, by passing through each termonuclear combustion chamber in our superstring, a swarm/block of neutrinos accelerates **while in the jets** (see ref. 3 below) with each fusion cycle and accumulates enormous cumulative energy to warp the spacetime where the entry point is the instant exit point for neutrinos at any part of the universe. Hence, the universe might have been built by such blocks of primordial matter to then evolve into current universe per my theory of twin (non-parallel) universes.


2. With regard to the **unified matter**, as depicted in fig.4, the ends of the upper string of matter are gravitationally pulled/shifted (for **shift** see ref. 3 below) sideways downward the corresponding ends of the base string of matter, creating thereby a **bow of matter** (fig. 4 and ref. 2 below) with the jet arrow of matter emitted along the axis of rotation from the base disc via the upper disc over its event horizon, as in fig. 4. So in our bow of matter we deal with the induced curvature of space per theory of general relativity, which predicted that a sufficiently compact mass can deform spacetime.

Large vertical "**arrow**" in our model represents a relativistic jet of matter which is forced from the twin superstrings structure due to accumulated tension applied by the bow of matter and rotational acceleration around the axis of the disc core. A similar phenomenon

occurs on earth, where the jets of magma erupt from volcanoes due to accumulated internal pressure and earth' occasional acceleration around its axis which might had caused in the past the death of the dinosaurs, whose skeleton couldn't sustain their enormous weight in increased gravity. The 2009 experiments conducted by Professor Sergey Lebedev's team in the Department of Physics at Imperial College London (www.imperial.ac.uk) confirm our theory by suggesting that the jets of matter are fired out more like bullets or buckshot,"they don't break into pieces—they are formed in pieces."

Note that our jet-arrow of matter points in one forward direction and isn't emitted in the opposite direction as suggested by the astronomers in ref. 3 far below. In fact the info in ref. 3 confirms our twin accretion disks concept as shown in **fig. 4** below, where the jet of matter from invisible lower disc is emitted via upper visible twin, in accordance with the **constant of duality** in the nature and cosmos. (See twin galaxies or single ones after separation).

The discussed bow of matter has its reflection in a bow-like shock wave in cosmos in NASA image (see ref. 3 below).





One might ask how the gravitational pull was created in the parallel superstrings of matter, with no external force. The answer is that we deal here with the warps of matter as the set of lengthwise streams of particles held in opposing tension in said strings and **twisting** around those strings of matter, as a certain metal encapsulates a violin string in a spiral. Our <u>warp concept</u> is supported by a practical experiment conducted by Professor Sergey Lebedev's team in the Department of Physics at Imperial College London that sent a high-powered pulse of energy into an aluminum disk. In less than a few billions of a second, the aluminum began to evaporate, creating a cloud of plasma very similar to the plasma cloud surrounding a young star. Where the energy flowed into the center of the disk, the aluminum eroded completely, creating a hole through which a magnetic field from beneath the disk could penetrate. The field initially pushes aside the plasma, forming a bubble within it. As the field penetrates further and the bubble grows, however, the magnetic fields begin to **warp and twist, creating a knot in the jet**.

To discuss further our superstrings of matter, we come to the notion of **repellent gravity**, as observed in galaxies. Due to anti-gravity between two superstrings of matter in our

model, matter **shifts** toward the density points/loops at the end of unified strings. (For **shifting** se ref. 3 below). This corresponds with Max Planck's idea that a harmonic oscillator (classically, anything that wiggles like a mass bobbing on the end of an ideal spring). Hence, our superstrings are "**plugged**" at their low density vertex representing a **plasma conductor in a vacuum loop**, which can withstand large gravitational pull. Dynamics of plasmas are often the sources of electromagnetic fields. ((Loop quantum gravity and **Aharonov–Bohm effect** might be applicable). This corresponds also with the **superfluid vacuum theory** (SVT).

<u>Ref. 1</u>

Twin Black Holes Discovered

Posted October 3rd, 2012


Astronomers recently made a surprising find while searching for a unique <u>black hole</u> in a tight cluster of stars 10,000 <u>light years</u> away from Earth. Instead of finding one black hole, scientists with the <u>National Radio Astronomy Observatory (NRAO)</u> found two – twins – something that surprised them because, according to modern theory, there should only be one black hole in a cluster.

Ref: 2 below

Supernova-powered bow shock creates cosmic spectacle

12:31 21 January 2013

Picture of the DaySpace Victoria Jaggard, space and physical sciences news editor



(Image: NASA/JPL-Caltech)

Ref. 3 below

2013: Using the European Space Agency's (ESA) <u>XMM-Newton</u> observatory, lead author <u>Dr. María Díaz Trigo</u> and colleagues studied a black hole binary system located in our own galaxy. This system, called 4U1630–47, has been known to show X-ray outbursts over the period of several months or even years, the agency explained in a statement. "In our observations, we found signs of <u>highly ionized **nuclei**</u> of two heavy elements, iron and nickel," said Trigo, who works at the European Southern Observatory

in Munich, Germany. "The discovery came as a surprise – and a good one, since it shows beyond doubt that the composition of black hole jets is much richer than just electrons."

"Intriguingly, we found the lines were not where they should be, but rather <u>were shifted</u> <u>significantly</u>," said <u>Dr. James Miller-Jones</u>, who led the radio observations and is a member of the International Centre for Radio Astronomy Research (ICRAR). That shift was similar to that of the **pitch change** of a vehicle's siren as it moves towards or away from the hearer, and signified that the length of the sound wave is becoming <u>shorter or</u> <u>longer due to the movement</u>. The phenomenon led the study authors to believe that the **particles were being accelerated to high speeds while in the jets** – one towards the Earth, and the other in the **opposite** direction. According to Dr. Miller-Jones, this is the first strong evidence of the presence of such particles in typical black hole jets. "We've known for a long time that jets contain electrons, but haven't got an overall negative charge, so there must be something positively charged in them too," he explained.


"Until now it wasn't clear whether the positive charge came from positrons, the antimatter 'opposite' of electrons, or positively charged atoms," the Dr. Miller-Jones added. "Since our results found nickel and iron in these jets, we now know ordinary matter must be providing the positive charge."

Read more at <u>http://www.redorbit.com/news/space/1113003040/black-hole-jet-contents-</u> <u>111413/#i7XX4sDVdvHdPqrQ.99</u>

Ref. 4 below

Monster black hole from early cosmos challenges physics

Thursday, 26 February 2015 Genelle Weule and Stuart Gary ABC



New physics are needed to explain an ancient supermassive black hole quasar 12 billion times the mass of the Sun (NASA/Caltech). Super big discovery **The discovery of a supermassive black hole from the early cosmos is set to rewrite physics, say scientists.**

An international team of astronomers detected a black hole 12 billion times the mass of our Sun, they report today in the journal <u>Nature</u>. The black hole, which formed just 900 million years after the Big Bang, is the source of a powerful beam of bright material known as a quasar.

"When we found this supermassive black hole we got very excited because we had found something that we never thought we could find," says Dr Fuyan Bian of the <u>Australian National University</u>. The team, led by Xue-Bing Wu at Peking University, discovered the black hole and quasar -- known as SDSS JO100+2802 -- using the Sloan Digital Sky Survey, then followed up with three other telescopes. With a luminosity of 420 trillion that of our Sun's, the new quasar is seven times brighter than the most distant quasar known.

"This quasar is very unique. Just like the brightest lighthouse in the distant universe, its glowing light will help us probe more about the early Universe," says Wu.