

PCT, Spin, Lagrangians, Part II

Hans Detlef Hüttenbach

*They're building Church and University,
Deceiving the people continually.
Tell the children the truth!*

Babylon System, Bob Marley

Abstract. I promised a second part. Here it is. I hesitated, because I don't like being destructive, and unfortunately, to deal with the current state of affairs in quantum theory means just doing this: I show you that the current gauge theory is fundamentally ill-designed, along with it fall Weinberg model and Wightman theory.

1. Continuing with the Lagrangian Formalism

Let me continue from where I left off in the former part: As we saw, the Lagrangian formalism for what is conceived to be the free quantum field amounts to evaluate the extremal for $\delta \langle \phi, A\phi \rangle$ with $A := (1/2)(\square\phi - m^2)$. Now, A is a self-adjoint operator, and for that operator it is plain vanilla that the kernel of A is the space of the extremals of $\langle \phi, A\phi \rangle$, and in this case happens to be the space of solutions of the Klein-Gordon equation $\square\phi - m^2\phi = 0$.

The problem in here is that from $\square\phi - m^2\phi = 0$ it cannot be concluded that

$$\mathcal{L} := (-1/2)(\langle \partial_\mu\phi, \partial^\mu\phi \rangle - \langle \phi, m^2\phi \rangle)$$

or any of its equivalent reformulations is the correct Lagrangian: A can be anything, for example the square or n -th power of A have the same kernel, but they would define different Lagrangians. (In [6] I showed that there are two distinct Lagrangians solving the very same equation of motion. Now we see that there are arbitrarily many.

Besides this: Does m^2x_μ for $0 \leq \mu \leq 3$ possess the dimension of an action? No! But mx_μ does! - The Lagrangian Formalism is simply meaningless!

2. Gauge Fields

It is commonly held that the electromagnetic field was a $U(1)$ gauge field. Let me disprove it:

I refer to [8, Ch.7, p.125ff.] to discuss the things in a non-abstract mathematical way: Neuenchwander introduces a $U(1)$ gauge transformation as mapping $\phi(x) \mapsto e^{i\chi(x)}\phi(x)$ where χ is a real-valued function in [8, eq. 7.2][eq. 7.2] and defines gauge invariance as invariance of the equations of motion under that gauge transformation. He then shows that this gauge transformation transforms the partial derivatives ∂_μ to $D_\mu := \partial_\mu + iA_\mu$, where $A_\mu := \partial_\mu\chi$ (see: [8, eq. 7.1.7] and proves the gauge invariance of the Maxwell equations. So far, everything is fine. The detrimental error comes in [8, Sec. 7.4], where Neuenchwander argues that the equations of motions for a gauged transformed free field theory would be the same as the free Lagrange equation with minimal coupling $\phi \mapsto \phi - A$, where A is the 4-potential of an external electromagnetic field; so, both would be equivalent, i.e.: the theory of electromagnetism was a $U(1)$ gauge theory.

What is wrong is that the 4-potential $A_\mu = \partial_\mu\chi$ of a $U(1)$ transformation $\phi \mapsto e^{i\chi}\phi$ is integrable (within \mathbb{R}^4) to a scalar function χ , which is χ . However, due to its non-zero rotation, a non-trivial electromagnetic 4-potential never is integrable within \mathbb{R}^4 . So, whatever $U(1)$ -gauge is chosen, no gauge field matches a non-zero electromagnetic 4-potential!

Therefore, whatever is described in a $SU(2) \times U(1)$ gauge theory, known as Salam-Weinberg model (see:[10]), it cannot be the unification of weak and electromagnetic theory!

Let's dig deeper and work out, what could be missing to fix this:

The space X of all complex linear combinations of the four Dirac matrices $\gamma_0, \dots, \gamma_3$ is a four-dimensional vector space with a (continuous) inner product, defined by:

$$\langle \sum_{\mu} \lambda_{\mu} \gamma_{\mu}, \sum_{\nu} \kappa_{\nu} \gamma_{\nu} \rangle := \sum_{\mu, \nu} \gamma_{\mu}^* \bar{\lambda}_{\mu} \kappa_{\nu} \gamma_{\nu} = \sum_{\mu} \bar{\lambda}_{\mu} \kappa_{\mu}.$$

This space X then is isometric with \mathbb{C}^4 , although the notion of orthonormality is lost within X , because $\gamma_{\mu}^* \gamma_{\nu} \neq 0$ for $\mu \neq \nu$. So, a unitary mapping on \mathbb{C}^4 maps into a unitary mapping on X . Similarly, the restriction of $\iota : \mathbb{C}^4 \ni (\lambda_0, \dots, \lambda_3) \mapsto \sum_{\mu} \lambda_{\mu} \gamma_{\mu} \in X$ to real values of λ_{μ} is an isometry of \mathbb{R}^4 to some (yet still complex) subspace $Y \subset X$, say. Both spaces \mathbb{R}^4 and Y as well as \mathbb{C}^4 and X induce a different concept of differentiation: The differentials dx_0, \dots, dx_3 on \mathbb{R}^4 or \mathbb{C}^4 become $dx_0\gamma_0, \dots, dx_3\gamma_3$ on X and Y . Now, considering an electromagnetic for potential $A = (A_0, \dots, A_3)$, then by gauge invariance, as a function on \mathbb{R}^4 , we can ensure that $A_{\mu} dx_{\nu} = -A_{\nu} dx_{\mu}$ for $\mu \neq \nu$ (which is not integrable). However, if I map this over to X (or Y), A becomes $\iota A = \gamma_0 A_0 + \dots + \gamma_3 A_3$, and path integration along the μ -th coordinate comes with an extra factor γ_{μ} . That means, we can now integrate ιA along paths $\omega : [0, 1] \ni \xi \mapsto X$ to some scalar function χ , which is real-valued if all the A_{μ} are, so $e^{i\chi}$ will be the desired gauge transformation.

So, the trick played is to integrate $\iota A = \sum_{\mu} \gamma_{\mu} A_{\mu}$ within X , but not A within \mathbb{R}^4 . But, where can we get the extra factors γ_{μ} from? - Let's have a look:

A current is described in electrodynamics as $j = \rho(u_0, \dots, u_3)$, where u is the 4-velocity. If that was to read instead $j = \sum_{\mu} j_{\mu} \gamma_{\mu}$, not only we would get the invariant $j^2 = j_0^2 - \dots - j_3^2$ for its square, also, by leaving out the Dirac matrices, we would erroneously turn the spinor into a scalar. Now j comes as a factor in the resulting force on a test charge (density) in motion, j' , say. Since we dropped the Dirac matrices in j , integrating the force in \mathbb{R}^4 will run us in problems: What would come out as a scalar energy function after path integration the spinor force will become a non-integrable, complicated vector-potential of scalar components. And to enforce the integrability of that vector-potential, it would need the artificial insertion of the Dirac matrices again, which will give us a scalar action function.

But watch out: if we had begun with a spinor j , we'd get a spinor force F , would path integrate that to a scalar potential, and a second path integration will give the action as a spinor function, and not as a scalar function! Now, what is correct? Is the potential a scalar and the action a spinor, or is it the other way round?

As will be shown in the last section, we can get rid of integration within X just by letting the charges become Dirac spinors, and path integrate the charge flux in \mathbb{R}^4 to its action, which then will be a spinor. So, the first alternative will win over the second.

And that implicitly means that as a gauge theory, electromagnetism cannot be a $U(1)$ -gauge theory: instead, the gauge transformation comes out as $\phi \mapsto e^i \sum_{\mu} \gamma_{\mu} X_{\mu}$, which, according to the metrics chosen in isometry with \mathbb{C}^4 is a unitary group, albeit one of dimension four, and not one! And there is only one unitary group (up to isomorphism) of this dimension, which is $U(2)$.

Again, this is satisfactory: In a charge symmetric world there is no absolute means to tell what charge is positive and what negative: all we know is that one observer may look at an electron as being negatively charged, while another observer may determine its positive charge. A $U(2)$ model fits perfectly (and again advocates that it might be a good idea to accept the co-existence of positrons with electrons).

I have one further point to make: The gauge invariance allows us to get rid of the symmetric part of A , i.e. that part, that can be integrated into a scalar function within the Euclidean metrics. In the first part I showed that there is this antisymmetric part A , and that there is also this symmetric part, which I denoted as the "neutral mass". Because the group of unitary mappings on \mathbb{C}^4 is the group $U(4)$, that would mean that, given all fields were gauge theories, all neutral fields would go into a $U(4)/U(2)$ gauge field, where the divisor denotes the quotient class. It would also then imply that we would only have two long-ranged forces, one for charges, and the rest for

neutral matter. I am rather sceptical as this being *ultima ratio*, and I would expect either progress and change as to this.

3. Wightman Axioms

In the mid 60's Arthur Wightman condensed what he thought were the basic assumptions and ingredients of a Quantum Field Theory in a set of axioms (see e.g.: [9] for a good online source other than the original book [11]).

In it, he states under the (W0) axiom, that the energy-momentum operator is to have its spectrum contained in the forward light cone.

That seems fair - at first sight, because of causality reasons. But, it is detrimental:

Not only does it disallow time (or energy) inversion, which is so important in classical physics, it disallows parity inversion either: Parity inversion is the transformation of the location coordinates $x \mapsto -x$. It is an error to think that parity inversion maps the forward light cone into itself: it does not: when you turn the coordinate system around a location coordinate for 180° , such that the time axis flips to its opposite direction, then the parity gets also inverted! That means, you can't have parity symmetry without the symmetry of time inversion. So, not only would you lose the principle of time invariance, you also would lose momentum conservation!

Sorry: That's not what it's supposed to be! It is just the old Platonic error, be it Dirac, be it Wightman: A symmetry does not ask for exclusion, but demands coexistence!

(When you theoretically cut off three of a baby's limbs, would you expect this theoretical child to behave like a real child?)

Neither is it comprehensible to exclude the interchange of time and location coordinates for the sake of a dogma of positivity of the Minkowski metrics, whereas in classical mechanics the replacement of any generalized location coordinate with the time coordinate is allowed, nor is it comprehensible to better speak of virtual, positive energetic particles moving backwards in time, rather than frankly of real particles with negative energy, moving forward in time!

4. Mass Field

By the many years of its existence, classical mechanics is commonly held to be a homogenous, self-consistent theory. Inside, however, it separates into different concepts: one is the motional, another one a dynamical theory. As to the motional theory, the system under consideration is a time curve $t \mapsto (q_1, \dots, q_N)$ of the N generalized location coordinates in space and time. In the dynamical part, then, that's not enough: we also need their canonically conjugated N momentum coordinates plus the total energy (which is the canonical conjugate of time). This then leads straight into the Hamiltonian mechanics, in which $2N$ independent momentum and location parameters are

coupled by N Hamiltonian equations (of first order).

Interestingly, a different concept emerged even before that: The dynamical system could be replaced by a mass-valued function $m(\vec{x}(t)) = m(t, \vec{x})$ in space and time. Since the masses of all bodies are positive, they all add up, so all positons, their speed, their momenta, and their energy can be identified, just through a mass-valued time curve in a three dimensional Euclidean space. That means that a dynamical mechanical system of n objects would be completely determined through the mass density curve $t \mapsto \rho(\vec{x}(t))$, and the famous Poisson equation, $\nabla^2 \Phi = \rho$, would then state the equivalence of the gravitational field with the mechanical system. (And it was then really understood that way by former physics.) Now I replace the mass m by a spinor of Pauli matrices, $m \mapsto m(\sigma_1 + \sigma_2 + \sigma_3)$, and ρ becomes a function of Pauli spinors $\rho_1 \sigma_1 + \rho_2 \sigma_2 + \rho_3 \sigma_3$. Then, given that this function is harmonic, the path integration of this spinor-function along any closed loop in the three dimensional space of location coordinates is zero. Therefore, I can path integrate that spinor function into yet another spinor function $S(\vec{x}(t))$, say, for which $\nabla S(\vec{x}(t)) = \rho_1(\vec{x}(t))\sigma_1 + \dots + \rho_3(\vec{x}(t))\sigma_3$ holds. Multiplying Poisson's equation from the left by another spinor test mass, integrating over space, and demanding that the masses are all contained in a bounded region, which ensures that the integral exists and boundary terms from partial integration vanish, Poisson's equation becomes a bilinear form, and it rewrites into:

$$-\nabla S(\vec{x}(t)) = \sum_{1 \leq k \leq 3} \rho_k \sigma_k.$$

Now, mass is energy (divided by the square of the speed of light), so S is the action. Once again: The knowledge of the action of a mechanical system in space and time completely determines the mechanical system itself (and vice versa).

Remark 4.1. If the non-relativistic gravitational field is a gauge theory at all, then it must be an $SU(2)$ -field, since the Pauli spinors are 2×2 -matrices. And it looks like to become one, if only one accepts the formal relation $1/(\sum_k \lambda_k \sigma_k) = d(\ln(\sum_k \lambda_k \sigma_k))/d(\sum_k \lambda_k \sigma_k)$ for $\lambda_k \neq 0$, ($1 \leq k \leq 3$).

Then, when it comes to relativistics, we can do the analogous thing as above, we just need to replace the Pauli spinors by the Dirac spinors, and the closed loops in space become closed loops in space-time, albeit outside the light cone. Maxwell's equations become the natural extension of Poisson's equation to the relativistic theory, in which spinors represent a unified concept of matter, including charges.

References

- [1] Clay Institute (A. Jaffe, E. Witten), *First Millenium Problem: Yang-Mills Fields And Mass Gap*, <http://www.claymath.org/sites/default/files/yanigmills.pdf>, 2000

- [2] A. Einstein, *Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?*, <http://www.physik.uni-augsburg.de/annalen/history/Einstein-in-AdP.htm>, 1905.
- [3] R. P. Feynman, *Lectures on Physics*, Vol. I-III, Addison Wesley, 1977.
- [4] H. D. Hüttenbach, *Analytic Functions For Clifford Algebras*, <http://vixra.org/pdf/1408.0084v3.pdf>, 2014.
- [5] H. D. Hüttenbach, *The Action Function of Adiabatic Systems*, <http://vixra.org/pdf/1404.0441v2.pdf>, 2014.
- [6] H. D. Hüttenbach, *In/Equivalence of Klein-Gordon and Dirac Equation*, <http://vixra.org/pdf/1410.0154v2.pdf>, 2014.
- [7] H. Kleinert, *Particles and Quantum Fields*, <http://users.physik.fu-berlin.de/~kleinert/b6/>, book publishing in preparation, 2014.
- [8] D. E. Neuenschwander, *Emmy Noether's Wonderful Theorem*, John Hopkins University Press, 2011.
- [9] NLab, n.n. *Wightman Axioms*, <http://ncatlab.org/nlab/show/Wightman+axioms>, 2015
- [10] V. Kaplunowski, *Glashow-Weinberg-Salam-Theory*, <http://bolvan.ph.utexas.edu/~vadam/classes/2012f/GWS.pdf>, 2012.
- [11] R.F. Streater, A. Wightman, *PCT, Spin and Statistics and All that*, Princeton University Press, 1978.
- [12] S. Weinberg, *Cosmology*, Oxford University Press, 2008.
- [13] Wikipedia, *Wheeler-Feynman Absorber Theory*, http://en.m.wikipedia.org/wiki/Wheeler-Feynman_absorber_theory, 2014.
- [14] Wikipedia, *Creation and annihilation operators*, http://en.m.wikipedia.org/wiki/Creation_and_annihilation_operators, 2014.

Hans Detlef Hüttenbach

e-mail: detlef.huettenbach@computacenter.com