

# Discovery of Quantum Gravity

Gene H Barbee, July 2015

## Table of Contents

Discovery of Quantum Gravity.....	1
FOREWORD .....	3
INTRODUCTION.....	3
GRAVITATIONAL FORCE ON THE EARTH.....	5
CIRCLES.....	6
Gamma.....	7
LARGE ORBITS and QUANTUM CIRCLES .....	8
CELLULAR COSMOLOGY .....	9
THE SOURCE OF GRAVITY .....	10
Calculation of gravitational force with accepted coupling constant .....	11
Calculation of Gravitational Constant from the Proton Mass Model .....	12
Two candidates for the correct gravitational energy scale.....	13
Candidate #1 the conventional Planck scale .....	13
Candidate #2 quantum gravity (the “dark horse” candidate) .....	14
Advantages of this proposal.....	15
Unification of the Special Theory of Relativity with the General Theory of Relativity.....	15
Quantum gravity reconciles gravity with the remaining forces .....	18
EXPANSION EQUATIONS .....	22
<b>Cell Diagram</b> .....	23
Transition from quantum behavior .....	25
What actually expands the universe?.....	26
Proof of non-quantum behavior .....	27
TIME AND COSMOLOGY.....	28
Time associated with kinetic energy.....	30
THERMODYNAMICS AND COSMOLOGY .....	30
Conservation of Kinetic and Potential Energy.....	30
Gravitational Accumulation.....	31

Formation of Black Holes .....	31
<b>Gravitational accumulation forms orbits</b> .....	32
Is Velocity Relative?.....	35
Potential and kinetic energy distribution.....	36
<b>Non-ideality involved in reconversion of kinetic energy to potential energy</b> .....	37
<b>Accounting for friction heat</b> .....	38
Velocity is not relative .....	39
Origin of the Second Law of Thermodynamics .....	41
DARK MATTER.....	42
<b>Galaxy formation</b> .....	42
BLACK HOLES AND QUANTUM GRAVITY .....	45
<b>The Schwarzschild radius</b> .....	46
<b>Approximate Solar Model</b> .....	46
<b>Analysis of black hole with 4.7 solar mass (9.9e30 Kg)</b> .....	49
<b>Compression thermodynamics</b> .....	50
<b>Pressure inside cells prevents black hole collapse</b> .....	51
<b>More massive black holes</b> .....	54
<b>Supermassive Black Holes</b> .....	55
<b>The BH surface</b> .....	58
Summary Black Holes and Quantum Gravity.....	58
Explanation of dark field compact galaxies .....	58
CONCLUSIONS.....	59
SELECTED REFERENCES .....	60
Appendix 1 Natural Logarithms .....	62
Appendix 2 Neutron Model .....	64
Abbreviated Neutron Model.....	65
Abbreviated Proton Model .....	67
Appendix 2 the Schwarzschild equation .....	67
Appendix 3 Black hole model results.....	69
Appendix 4 Information based reality .....	72

## FOREWORD

Have you wondered if there was “DNA” for the universe, where it is stored and how it influences everything?

Do you want to know where the space around you originates?

Are you concerned that science lacks answers to fundamental questions like “Why can’t scientists find quantum gravity? What exactly is time? What is dark matter and dark energy?”

Do you want to know how the universe can be created *ex nihilo* (out of nothing) and what is inside black holes?

Do you want science to contribute to our understanding of whom we are and why we are here?

This work builds on the best science available but correlates data with tools from the field of information theory. This approach is new and leads to a fresh view of physics data. Can the universe be viewed as a large information source that interacts with our mind to produce reality?

## INTRODUCTION

Gravity is still an active research area for scientists even though English mathematician Isaac Newton published his work entitled *Principia* in 1687. For the last twenty years or so some of the best theorists have been working on quantum gravity. There is a basic disconnect at the heart of gravity. The problem is that although Isaac’s equations are correct they describe large scale behavior of objects. Einstein’s general theory of relativity is the modern theory of gravity but again, it describes large scale behavior of objects following paths curved by mass. This leaves small scale (quantum scale) gravity a subject of research. In the author’s opinion, a solution to the problem was delayed by general acceptance of an old relationship between fundamental constants. Literature states that the gravitational constant (G) originates at the Planck scale. The Planck length (Compton wavelength)  $L = (\hbar^2 G / C^3)^{0.5} = 1.61 \times 10^{-35}$  meters where  $\hbar$  is Heisenberg’s reduced constant ( $\hbar = h / 2\pi = 6.58 \times 10^{-22}$  Mev-Sec) and C is of course the speed of light. The energy associated with the Planck length is  $1.2 \times 10^{22}$  MeV. This energy is far greater than the energy of a proton and the Planck length is incredibly small. Many physicists are reluctant to give up the equation that contains G,  $\hbar$  and C in what appears to be a defining relationship. Theorists are exploring alternatives like string theory in search of unification but have failed to gain acceptance of a new theory describing the origin of the gravitational constant.

The proverbial apple that Isaac was watching when he conceived of gravity was a bit of Hollywood. He actually rolled objects down an incline and measured time and distance. He was an observational based scientist. But we have high standards for present day physics and should expect someone to find the source of the gravitational constant and its relationship to the other forces. During Newton’s lifetime the concept of a small scale was not taken seriously although

the idea of an atom came much earlier. We will explore the possibility that the Planck length is too low and the energy  $1.2e22$  MeV is too high. A new theory of quantum gravity will be proposed at the scale of the proton.

What is space and time? It is interesting to the author that there isn't agreement on these basic concepts. The difficulties with gravity extend to the origin of space. Furthermore, we all experience elapsed time but some physicists don't believe in time that moves forward. They do believe that time cycles at the small scale and of course they know that energy is related to these cycles.

A proposal will be made regarding the origin of space and time and its relationship to expansion of the universe. Most cosmologists believe that expansion exceeds the speed of light. They say that space is being created and carries light with it. This appears to violate one of our most sacred concepts; that particles cannot travel at the speed of light? There is no need to violate  $C$  in the author's proposal.

Thermodynamics is one of the most advanced sciences but again there are disconnects at the core. Most physicists agree that velocity is relative but does this mean that kinetic energy is also relative? Kinetic energy is just a form of energy and kinetic energy ( $1/2*m*V^2$ ) is conserved. Energy according to the first law of thermodynamics can't be created or destroyed. The second law of thermodynamics says that systems of particles containing heat energy always "run down". Entropy is a one way downward street but where is the origin? Unfortunately there appears to be no answer that everyone agrees on. We must tie cosmology and thermodynamics together.

During Sir Isaac's lifetime, the concept of a universe was pretty much limited to a solar system. He wouldn't have believed how large the universe is but he would have been fascinated as we all are about findings in the last 100 years. He probably couldn't even conceptualize of gravity being so strong that even light can't escape an object but this is exactly what happens in a black hole. Many have been discovered and they seem to come in mainly two types: Burnt out stars and supermassive black holes almost as massive as an entire galaxy. Scientists need a quantum theory of gravity because it might help them understand black holes. One question that keeps surfacing is "do black holes collapse?" If they do it suggests a singularity where physics breaks down.


There is another "physics breaks down" possibility at the heart of physics. What was the beginning condition? Some physicists believe that the big bang was a singularity, so hot that relationships we know and use might have been different and there is even speculation that the four forces of nature might have been unified at that point. The author believes that the laws of physics existed before the big bang and are based on information theory [1]. There are interesting parallels between information theory and thermodynamics suggesting that physics is information based like our minds [30].

Note: The document is written in language I call "engineering 101", not very scholarly, but practical. I would recommend following the calculations with a Microsoft excel® spreadsheet. All the equations will use meters for distance (m), kilogram for mass (kg) and seconds (sec) for time. Newton was rewarded by using his name for the force unit. It is abbreviated N. Recall that numbers can be written in scientific notation, i.e.  $5.98e24$  kg is the earth's mass in kilograms

and e24 means move the decimal place 24 places to the right. Conversely, the electron orbits a proton at 5.29e-11 meters meaning that you move the decimal place 11 places to the left.

## GRAVITATIONAL FORCE ON THE EARTH

The earth pushes up on our feet but astronauts don't need this help. Here on earth, the reason we feel force upward is that our velocity is too low to be on an orbit (geodesic) defined by the radius of the earth. A geodesic is a combination of velocity  $V$ , radius  $R$  and mass  $M$  that give  $G$ , the gravitational constant;  $G=R \cdot V^2/M=6.6742e-11 \text{ N}\cdot\text{m}^2/\text{kg}^2$ . Astronauts are in orbit and on a geodesic. We can calculate the velocity  $V$  required to be on a geodesic. First we calculate our acceleration at the surface of the earth. We need to know that the earth's mass is  $M= 5.98e24 \text{ kg}$  and that the radius of the earth  $R = 6.39e6 \text{ meters}$ . This gives us the gravitational acceleration on the surface of the earth  $a=G \cdot M_{\text{earth}}/R_{\text{earth}}^2=9.8 \text{ meters/second}^2$  (abbreviated  $\text{m/sec}^2$ ). Next we calculate velocity  $V= (a \cdot R)^{0.5}= 7909 \text{ m/sec}$ . The force upward on our feet is  $F=\text{mass} \cdot a=\text{mass} \cdot V^2/R$ . Your mass in kg is your weight in pounds divided by 2.2. If your weight is 198 lb =90 kg, the earth is pushing up on you with the force  $F =90 \cdot 7909^2/6.38e6=883 \text{ Newton (N)}$ . Force upward from the earth is making up for the outward inertial force you are missing because your velocity is too low. The equation could also be written  $F=\text{mass} \cdot (7909^2 - V_{\text{low}}^2)/R$ . In this equation  $V_{\text{low}}$  is fixed by us being on earth. Inertial force is outward force in an orbit and gravitational forces are inertial forces. The diagram below describes the situation.



Kids feel inertial force when you they are on a merry-go-round and you calculate it by  $F=m \cdot a$  where acceleration  $a= V^2/r$ . Gravitational force is also inertial force and Newton recognized this because *Principia*, his book on physics also stated that  $F=m \cdot a$ . Force is mass times acceleration. But if an astronaut has this velocity, why is she “weightless”? The essence of Einstein’s general theory of relativity (GR) is that mass follows curved space-time and “doesn’t know” about the forces involves. It simply says that the earth curves space-time and the astronaut follows the curve. When the astronaut has the right velocity,  $V=7909 \text{ m/sec}$  in this case, she feels no force. She is falling but also circling the earth fast enough that she never becomes closer to the earth. There is a statement regarding this concept called equivalence of

acceleration and gravity. The story goes “if you are in a free falling elevator, how would you know about the force on you?” Since you don’t feel it, you don’t measure it.

Kinetic energy is converted to potential energy and visa-versa. The equation that applies is kinetic energy (ke) plus potential energy (pe) is a constant, i.e.  $ke+pe = \text{constant}$ . Potential energy is force times the distance the force pushes through, i.e.  $PE=F*R$ . To find the potential energy from the orbital kinetic energy we must get the origin (initial condition) correct. The origin is the big bang when particles with kinetic energy separate. Gravitation resists expansion and kinetic energy is converted to potential energy. Later when the mass starts its fall, potential energy is reconverted to kinetic energy. It either accumulates in bodies from a position established by expansion of the universe or it establishes an orbit. In both cases it has potential and kinetic energy. At the orbital position (the geodesic) the outward inertial force is balanced. We calculated the inertial force 883 N in the orbit assuming that the astronaut had gained 7909 m/sec from the rocket. Here on earth, we are “off the geodesic”. The earth must push up on our feet to make up for the inertial force that we are lacking. Yes, there is energy of position (potential energy) but this does not produce the gravitational force. Think about climbing the stairs. Where does your energy go as you climb? It goes into overcoming force F and the potential energy is  $PE=F*R$ . It is the result of a force moving through a distance (force units are Newtons N and distance units are meters and energy is N-m). To calculate potential energy you need a conversion factor ( $PE=F*R$  (N-m)\*conversion factor) to know the potential energy in Million electron Volts (MeV). MeV is a convenient energy unit and represents the energy required to move an electron through a one volt potential (eV) but since it is a small energy, it is multiplied by one million. The conversion  $6.24e12$  MeV/(N-m) is the main conversion used in the document.

Although this explains our gravitational situation here on earth there are still several questions: 1) how did we get on this curve? 2) Why is our kinetic energy low? 3) Deeper yet, where does gravity originate? 4) What is the resisting force that allows the earth to support us? Overall, we haven’t gone beyond what Newton wrote and Einstein taught us, but you will if you keep reading.

## **CIRCLES**

Quantum mechanics (QM) deals with small circles. The circle is a model and the radius is probabilistic. The basic concept is that energy is related to a circle by the equation  $E=H*v$ . Frequency  $v$  is the number of times per second energy travels around a quantum circle at the speed of light and H is Heisenberg’s constant  $4.136e-21$  MeV-sec. The speed of light is  $3e8$  meters/second. Frequency can be a large number but cycle time  $t$  will be  $1/\text{frequency}=1/v$ . How much time does it take to move around a circle (R) at velocity C? The time  $t= 2*\pi*R/C$  equals time  $t= H/E$ . The important constant H (Heisenberg’s constant) relates time and energy. Knowing the constant relationship, radius is defined. If we are to understand quantum circles, we need to know their radii. Below, we will find a radius from accepted facts about an electron circling a proton (the element hydrogen) and then generalize the equation for other quantum circles.

**$t=H/E$  and  $t=2\pi R/V$  are equal for a little quantum circle.**

**$2\pi R/C=1/\text{frequency}$**

**$2\pi R/C=H/E$**

**where  $H$ =Heisenberg's Constant  $4.136\text{e-}21$  mev-sec.**

<b>Electromagnetic field</b>		<b>2.72E-05 MeV</b>	
<b>t=H/E</b>	<b>t=4.14e-21/27.2e-6</b>	<b>1.52E-16</b>	<b>seconds</b>
<b>2*pi*R/V</b>	<b>equal but V?</b>	<b>1.52E-16</b>	<b>seconds</b>

**If we know V above, we can calculate R**

<b>Known</b>	<b>1.36E-05 MeV</b>	<b>kinetic energy</b>
<b>Known</b>	<b>0.511 Mev</b>	<b>electron mass</b>
<b>g</b>	<b>0.999973 g=0.511/(0.511+13.6e-6)</b>	
<b>V/C</b>	<b>0.007296 V/C=(1-g^2)^0.5</b>	

**R calculated from  $H/E=2\pi R/V$**

**$R=H/E\cdot V/(2\pi)$**

**$R=4.136\text{E-}21/27.2\text{e-}6\cdot 0.00729\cdot 3\text{e}8/(2\cdot \text{PI}())=5.29\text{e-}11$  meters**

The equation for radius R is central to fundamental forces with different inputs.

Where:  $H$ =Heisenberg's constant

$M$ =mass of the particle. If the particle is moving fast relativistic mass is  $m/\gamma$

$E$ =field energy that helps define the radius of the circle.

$R$ =maximum probabilistic position of the particle.

$R=HC/(2\pi)/(E\cdot m/g)^{0.5}$

$R=1.973\text{e-}13/(E\cdot m/g)^{0.5}$  and sometimes  $=1.973\text{e-}13/E$

where  $HC/(2\pi)=1.973\text{e-}13$  MeV-m

## Gamma

Velocity is distance/time with a small shift in time. Gamma is a ratio indicating the shift and the reason physicists refer to space-time instead of space alone. Gamma can be calculated from kinetic energy or velocity. It is simply  $\gamma(g) = \text{mass}/(\text{mass}+\text{kinetic energy})$ . We already know that the astronaut's velocity was 7909 m/sec so it is easier to find gamma from the special relativity equation  $(g)=(1-(V/C)^2)^{0.5}=(1-(7909/3\text{e}8)^2)^{0.5} = 0.9999999997$ .

Newton couldn't have discovered that he was dealing with space-time as the very source of gravity based on gamma  $(g) = 0.9999999997$ . For all practical purposes, gamma was 1 but you can see that velocity  $V$  is only a function of gamma. Is this enough to curve space-time? The curvature we are talking about for an astronaut in earth orbit is the radius of the earth  $6.38\text{e}6$  meters (actually a small amount larger so the astronaut orbits above the earth). What does the earth radius have to do with gamma? To understand this we have to turn to one of Einstein's

equations and another mathematician by the name Schwarzschild. This story will continue under the heading “Unification of special relativity and general relativity”.

## LARGE ORBITS and QUANTUM CIRCLES

We are face to face with where gravity originates. If it originates in quantum circles like the other forces, there is a huge gap. Consider what the orbital radius of the earth might be on a proton by proton basis (I am aware that the earth is not made of protons with a circling electron but we can use protons as an example). Speculate for a moment that we can consider the volume of the earth broken into smaller volumes around each proton. The mass of the earth is  $4e24$  kg and the mass of a proton is  $1.67e-27$  kg. There are  $4e24/1.67e-27=3.58e51$  protons in the earth and the volume of the earth is  $4/3\pi*6.3e6^3=1.1e21$  m<sup>3</sup>. Dividing these we find the volume surrounding each proton ( $3e-31$  m<sup>3</sup>) and then determine the radius for this volume. The answer is  $r = (\text{vol}/(4/3*\pi))^{1/3} = 4.17e-11$  meters. Actually, this makes sense. We know that the electron orbits a proton at  $5.29e-11$  meters. The electrons repel each other and resist being compressed. Even with considerable mass building up pressure down through the layers of the earth, the electrons are only on average compressed to  $4.17e-11$  meters. Remember the concept because later in the document we will try to determine what resists the enormous pressures down through a black hole.

This helps us understand why the earth pushes up on our feet, but it doesn’t help us reconcile large orbits like earth with  $6.4e6$  meters and little orbits like those encountered with quantum mechanics.

Newton could not estimate the number of particles in the universe. Further the whole concept of expansion of the universe was centuries in the future. In Newton’s wildest dreams, he would not have anticipated that the sky temperature contains clues regarding the beginning we call the big bang. After cosmic microwave background (CMB) radiation was discovered around 1950, cosmologists started to analyze what the signature of the cosmic background radiation might reveal. They proposed and later received funding for a balloon project called COBE and satellite projects called WMAP [7] and PLANCK. We can analyze how many protons there are in the universe from the data. The data gave us increasingly accurate estimates of rho, the density of the present universe and its radius [4][6][17].

Question about number of particles in universe			
Critical Density Predictions (kg/M <sup>3</sup> )			
Density	8.93E-27	pg 337 isHughes	
Density	3.73E-26	rho zero pg 103 Peebles at H=.71	
Density rf	9.5E-27	WMAP basic results Table 3	
R	6.30E+25	meters	
N protons	1.61E+78	$N=\rho*0.27*(4/3)*PI()*R^3/1.67e-27$	
ln (N)	180.0759		



What could be the possible meaning of  $\exp(180) = 1.61e78$  protons? (The nomenclature  $\exp(180)$  stands for the natural number e to the power 180.

## CELLULAR COSMOLOGY

Consider large mass  $M$  (for our purposes the mass of the universe although the term universe seems a little presumptive) broken into  $\exp(180)$  small cells, each with the mass of a proton labelled lower case  $m$  below. The mass ( $m$ ) of a proton is  $1.67e-27$  kg. Fill a large spherical volume with  $\exp(180)$  small spheres we will call cells. Consider the surface area of many small cells as a model of the surface of one large sphere with the same surface area. For laws of nature to be uniform throughout the universe there can be no preferred position. A surface offers this property but the equivalent surfaces of many small spheres also offer this property as long as we do not distinguish an edge. As such a surface model equivalent to the surface of many small cells is useful if the fundamentals of each cell are known.

In general relativity [6] the metric tensor (scholarly matrix equations from general relativity) is based on  $(ds^2 = \text{three distances}^2 + (C \cdot \text{time})^2)$ . Note that  $ds^2$  is a surface area and it is this surface that we will break into  $\exp(180)$  small spheres. Let small  $r$  represent the radius of each small cell and big  $R$  represent the radius of one large sphere containing  $\exp(180)$  cells with the same surface area. Position a proton like mass on the surface of each cell. The total energy will be that of one protons/cell plus a small amount of kinetic energy. We will evaluate the gravitational constant  $G$  of a large sphere and compare it with  $G$  of small cells.

$$\begin{aligned} \text{Area} &= 4 \cdot \pi \cdot R^2 \\ \text{Area} &= 4 \cdot \pi \cdot r^2 \cdot \exp(180) \\ A/A &= 1 = R^2 / (r^2 \cdot \exp(180)) \\ R^2 &= r^2 \cdot \exp(180) \\ r &= R / \exp(90) \quad \text{surface area substitution} \\ M &= m \cdot \exp(180) \quad \text{mass substitution} \end{aligned}$$

For gravitation and large space, we consider velocity  $V$ , radius  $R$  and mass  $M$  as the variables (capital letters for large space) that determine the geodesic. With  $G$  constant,  $M = m \cdot \exp(180)$  and the surface area substitution  $R = r \cdot \exp(90)$ , the gravitational constant would be calculated for large space and cellular space as follows (lower case  $r, v$  and  $m$  below are for cellular space):

<b>At any time during expansion</b>		
<b><u>Large space</u></b>		<b><u>Cellular Space</u></b>
		<b>With substitutions:</b>
		<b><math>R = r \cdot \exp(90)</math> and <math>M = m \cdot \exp(180)</math></b>
<b><math>R \cdot V^2 / M =</math></b>	<b><math>G = G</math></b>	<b><math>r \cdot \exp(90) \cdot V^2 / (m \cdot \exp(180))</math></b>
<b><math>R \cdot V^2 / M =</math></b>	<b><math>G = G</math></b>	<b><math>(r \cdot v^2 / m) / \exp(90)</math></b>

The extremely small value  $1/\exp(90)$  is the coupling constant for gravity. When measurements are made at the large scale as must done to measure G, the above derivation indicates that we should multiply cell scale values ( $r*v^2/m$ ) by  $1/\exp(90)$  if we expect the same G. Geometric and mass relationships give the cell “cosmological properties”. I call this cellular cosmology.

It must be recognized that for equal gravitational constant the radius of curvature and mass are vastly different between the large and small scale. It was unfortunate that the great physicists of the 1900’s did not have the advantage of WMAP [7] and Cmagic [8] expansion models, nor did they have the advantage of knowing the approximate number of protons in the universe. Perhaps they couldn’t compare cellular scale space to large space because they lacked information.

## THE SOURCE OF GRAVITY

A simplified Neutron mass model [10][13][24] is shown below. It is derived from a full Neutron mass model reviewed in Appendix 2.  $Mass+ke=959.92$  MeV is overall equal and opposite field energy 959.92 MeV with the net zero (further comments later). The value of interest for gravity is the field energy 2.723 MeV. (The neutron mass is 939.57 MeV made up of quarks and kinetic energy). Add down through the values in each column of the table below to see the components that total 959.92 MeV. Note that nature uses the value 10.15 MeV many times.

<b>r20 uc2</b>				
<b>Mass and Kinetic Energy</b>			<b>Field energy</b>	
<b>Mass</b>	<b>KE</b>	<b>Strong</b>	<b>Strong</b>	<b>Gravitational</b>
<b>Quarks</b>		<b>Residual</b>	<b>field energy</b>	<b>Energy</b>
<b>MeV</b>	<b>MeV</b>	<b>Field</b>	<b>MeV</b>	<b>MeV</b>
<b>Strong</b>	<b>130.16</b>	<b>799.25</b>	<b>-957.18</b>	<b>-2.73</b>
<b>Strong Residual KE</b>		<b>10.15</b>		
<b>Neutron</b>		<b>939.57</b>	<b>-20.35</b>	<b>-959.92</b>
<b>neutrinos</b>		0.05		
<b>Gravitational ke</b>		<b>10.15</b>		
<b>Gravitational pe</b>		<b>10.15</b>		
<b>Total</b>		<b>959.92</b>		

$$R(\text{meters}) = \frac{HC}{(2*\pi)*(E*E)^{.5}} = \frac{HC}{(2*\pi)*2.723} = 7.224e-14$$

$$HC/(2*\pi) = 1.973e-13 \text{ MeV-m}$$

**The radius of a quantum circle for gravity is 7.224e-14 meters with E=2.723 MeV.**

Large scale gravity is the result of a body with mass  $m$  falling into curved space time defined by a central mass  $M$ . As a body falls it loses potential energy, gains kinetic energy and finds a geodesic where it feels no force. Again, the geodesic variables  $R, V$  and  $M$  combine to give  $G$ , the gravitational constant;  $G=R \cdot V^2/M$ . Quantum gravity follows the same physics except the curvature is the radius  $7.224 \times 10^{-14}$  meters above and the geodesic is  $G=r \cdot V^2/m/\exp(90)$ . The body that falls to the geodesic curvature  $r=7.224 \times 10^{-14}$  meters is the neutron. It has mass  $1.675 \times 10^{-27}$  kg and according to the model above, it initially has potential energy 20.3 MeV but when it achieves an orbit, its kinetic energy has increased to 10.14 MeV and its potential energy has decreased to 10.16 MeV.

### Calculation of gravitational force with accepted coupling constant

Literature [2][3][28] regarding a coupling constant for gravity is reviewed below. The gravitational coupling constant  $\alpha_G$  is the coupling constant characterizing the gravitational attraction between two elementary particles having nonzero mass.  $\alpha_G$  is a fundamental physical constant and a dimensionless quantity, so that its numerical value does not vary with the choice of units of measurement:

$$\alpha_G = Gm_e^2 / (\hbar c) = (m_e^2 / m_p^2) = 1.752 \times 10^{-45}$$

where  $G$  is the Newtonian constant of gravitation;  $m_e$  is the mass of the electron;  $C$  is the speed of light in a vacuum;  $\hbar$  is the reduced Planck constant;  $m_p$  is the Planck mass.

This coupling constant can be understood as follows:

<a href="http://en.wikipedia.org/wiki/alphaG=(mp/me)^2=1.752e-45">http://en.wikipedia.org/wiki/alphaG=(mp/me)^2=1.752e-45</a>
$\alpha_G = (m_p/m_e)^2 = 1.752 \times 10^{-45}$
$m_p/m_e = 1836$ , where $m_p/m_e = \text{proton/electron}$
$\alpha_G = 1836.15^2 \cdot 1.752 \times 10^{-45} = 5.907 \times 10^{-39}$
$F = (5.9068 \times 10^{-39}) \cdot \hbar c / R^2$

If  $R$  for the force calculation is  $7.22 \times 10^{-14}$  meters, as proposed above, the force is:

$F = (5.9068 \times 10^{-39}) \cdot \hbar c / R^2$		
$\hbar c$	$6.5821 \times 10^{-22}$	mev-sec
$\hbar c$ in NT-m-sec	$1.05 \times 10^{-34}$	NT m sec
$\hbar c$ in NT-m <sup>2</sup> =K	$3.16 \times 10^{-26}$	NT m <sup>2</sup>
$F = (5.9068 \times 10^{-39}) \cdot K / R^2$		
$F = (5.9068 \times 10^{-39}) \cdot 3.16 \times 10^{-26} / (7.22 \times 10^{-14})^2 = 3.58 \times 10^{-38}$		
<b><math>3.579 \times 10^{-38}</math></b>	<b>NT</b>	

This result agrees with the simple Newtonian force within adjustments for gamma:

$$F=Gmm/R^2 \text{ (nt)}=6.67428\text{e-}11*1.6726\text{e-}27^2/7.224\text{e-}14^2=3.666\text{e-}38 \text{ nt}$$

## Calculation of Gravitational Constant from the Proton Mass Model

Using values for the proton mass model that the author believes unify nature's forces (6), the gravitational constant is calculated below and agrees with the published constant,  $G=6.674\text{e-}11$  N meters<sup>2</sup>/kg<sup>2</sup>.


The following table follows a format that will be used several times. The goal is to use the fundamental radius  $7.224\text{e-}14$  meters to calculate the gravitational inertial force. The inputs listed at the top of the table originate in the neutron model above. Firstly, the mass of a proton in MeV and its mass in kg are specified in the table. The gravitational field energy  $2.723$  MeV gives  $R=7.224\text{e-}14$  but there is kinetic energy ( $10.14$  MeV) in the orbit that the neutron falls into. With mass and kinetic energy,  $\gamma$  and  $V/C$  can be calculated. Next the inertial force is determined for the mass orbiting at radius  $R$ .

GRAVITY	proton	neutron
<b>Neutron Mass (mev)</b>	<b>938.2720</b>	<b>939.565</b>
<b>Neutron Mass M (kg)</b>	<b>1.673E-27</b>	<b>1.675E-27</b>
<b>Field Energy E (mev)</b>	<b>2.732</b>	<b>2.732</b>
<b>Kinetic Energy ke (mev)</b>	<b>10.111</b>	<b>10.140</b>
<b>Gamma (g)=M/(M+ke)</b>	<b>0.9893</b>	<b>0.9893</b>
<b>Velocity Ratio v/C=(1-g^2)^0.5</b>	<b>0.1456</b>	<b>0.1457</b>
<b>R (meters) =(HC/(2pi)/(E*E)^0.5</b>	<b>7.224E-14</b>	<b>7.224E-14</b>
<b>Inertial Force (F)=(M/g^2V^2/R)*1/EXP(90) N</b>	<b>3.656E-38</b>	<b>3.666E-38</b>
<b>HC/(2pi)=1.97e-13 mev-m</b>		
<b>Calculation of gravitational constant G</b>		
<b>G=F*R^2/(M/g^2)=NT m^2/kg^2</b>	<b>6.6739E-11</b>	<b>6.6743E-11</b>
<b>Published by Partical Data Group (PDG)</b>	<b>6.67E-11</b>	<b>6.6743E-11</b>

The measured gravitation constant  $G$  [16] is calculated above from fundamentals. The constant  $1/\exp(90)$  scales the quantum level to the large scale we observe around us. It has the effect of dramatically reducing the force between neutrons and makes gravity very long range compared to the other forces. The inertial force  $3.66\text{e-}38$  N is the same force as the literature above and confirms the radius  $7.22\text{e-}14$  as the radius for quantum gravity.

Note: There is a small difference in kinetic energy between the proton and neutron. The literature above is based on a proton. The author believes that gravity is based on a neutron. Later in the document I will refer to the kinetic energy  $10.11$  MeV. The value  $10.14$  MeV is the neutron kinetic energy and  $10.11$  MeV ke is for the proton.

The calculation for G above indicates that nature uses the general theory of relativity at the quantum level. The author calls this cellular cosmology. A cell is diagrammed below. The inertial force  $F=m*V^2/R=3.66e-38$  N. The proton “feels” no force just like our astronaut travelling on the geodesic. This geodesic is a “quantum circle” but it is related through cellular cosmology to a large orbit.



### Two candidates for the correct gravitational energy scale

We will further explore below whether we can call this circle of radius  $7.224e-14$  meters the scale for quantum gravity.

### Candidate #1 the conventional Planck scale

There are tests for quantum gravity: We will compare the Planck scale relationships [2] with the relationships above.

Nomenclature and review of the Planck scale

	<b>Constants</b>				
<b>h</b>	<b>6.58E-22</b>	<b>MeV-sec</b>		<b>reduced</b>	
<b>E</b>	<b>1.22E+22</b>	<b>MeV</b>		<b>Planck Energy</b>	
<b>M</b>	<b>2.18E-08</b>	<b>kg</b>		<b>Compton mass</b>	
<b>G</b>	<b>6.67E-11</b>	<b>Nt m<sup>2</sup>/kg<sup>2</sup></b>		<b>gravitational</b>	
<b>C</b>	<b>3.00E+08</b>	<b>m/sec</b>			
	<b>Relationships</b>				
	<b>L=G*M/C<sup>2</sup></b>	<b>Compton wavelength</b>			
	<b>L=G*M/C<sup>2</sup></b>	<b>6.67e-11*2.18e-8/3e8<sup>2</sup></b>	<b>1.62E-35</b>		<b>meters</b>
	<b>L=C*h/E</b>	<b>3e8*6.58E-22/1.22E+22</b>	<b>1.62E-35</b>		<b>meters</b>
	<b>L=h/(M*C)</b>		<b>1.61E-35</b>		<b>meters</b>
	<b>G=h*C/M<sup>2</sup></b>	<b>6.58E-22*3e8/2.18e-8<sup>2</sup>*1.6</b>		<b>6.67E-11</b>	<b>Nt m<sup>2</sup>/kg<sup>2</sup></b>

The criteria for quantum level is quantum mechanical “action” [2]. Action must be 1.0 to be at the quantum level (it just tests whether the variables make a quantum circle.) Action is the value

$P \cdot L / h$  where  $P$  is momentum,  $L$  is the wavelength and  $h$  is Heisenberg's reduced constant ( $H/(2 \cdot \pi)$  labelled  $\hbar$ ,  $hbar$  or just lower case  $h$ ). Compare action for two energy levels, the Planck scale (1.22e22 MeV and the much lower level 938.27 MeV proposed above.

	<b>action= <math>p \cdot L / h</math></b>			
	<b>Planck energy</b>	<b>(MeV)</b>	<b>1.22E+22</b>	
	<b>Planck L</b>	<b>(meters)</b>	<b>1.62E-35</b>	
	<b>Momentum</b>	<b><math>p = E / C</math></b>	<b>4.07E+13</b>	
	<b><math>p \cdot L</math></b>	<b>Mev-sec</b>	<b>6.58E-22</b>	
	<b>action= <math>p \cdot L / h</math></b>		<b>1.00E+00</b>	

Yes, the Planck scale meets the criteria for being at the quantum level because  $\text{action} = p \cdot L / h$  is 1.

### Candidate #2 quantum gravity (the "dark horse" candidate)

<b>Proposal ( cell d305 "unified")</b>			
Field Energy		2.732 mev	
constant	$HC / (2\pi)$	1.97E-13	mev-m
	$R = \text{constant} / E$	7.22E-14	m
	Field side	R side	
	$H/E$	$2 \cdot \pi \cdot r / C$	
time (t)	1.51E-21	1.51E-21	sec
Proposal p ( $p = E / C$ )		9.11E-09	mev-sec/m
$p \cdot R / h$		1.00	
qm test	$M / C^2 \cdot R^2 / t$	6.58E-22	mev-sec
qm test/h	$M / C^2 \cdot R^2 / t / h$	1.00	

The proposal also meets the  $\text{action} = 1$  requirement for a quantum level relationship since  $\text{action} = P \cdot R / h = 1$ .

Further comparison:

The proton mass is 938.27 MeV, not 1.22e22 MeV (1.67e-27 kg, not 2.17e-8 kg). Compare the calculation for gravitational constant for the Planck scale and the quantum gravity mass level and note that they differ by a large factor.

$$G = h \cdot C / M^2$$

$$G = (6.58e-22 \cdot 3e8 / (2.18e-8)^2 \cdot 1.603e-13)$$

$$6.66E-11 \quad N \cdot m^2 / kg^2$$

$$G = h \cdot C / M^2$$

$$\text{Proposed mass } 1.67e-27 \text{ kg}$$

$$G = (6.58e-22 * 3e8 / (1.67e-27)^2 * 1.603e-13) / \exp(88.03)$$

$$6.66E-11 \quad \text{N m}^2/\text{kg}^2$$

Gravity, defined the Planck way requires a large divisor  $\exp(88.03)$ . Both candidates use a large divisor but there is a huge difference between  $\exp(88.03)$  and  $\exp(90)$ . A divisor is required because gravity is shared among  $\exp(180)$  protons and the surface area of each cell is  $1/\exp(90)$  of the surface area of the universe but this makes  $1/\exp(90)$  the correct coupling constant. The correct scale for quantum gravity is the radius  $7.22e-14$  meters not the Planck scale  $7.62e-35$  meters.

### **Advantages of this proposal**

- 1) Cellular cosmology unifies the special theory of relativity with the general theory of relativity.
- 2) It reconciles gravity with the remaining forces.
- 3) Quantum gravity at the scale  $7.22e-13$  meters is fundamental to expansion equations that give the size of the universe. It gives us a clear description of the space around us. We are walking around in expanded quantum gravity.
- 4) Combined with cosmology, quantum gravity shows that time advances and explains the twin paradox.
- 5) Quantum gravity produces realistic and important results for the study of black holes.
- 6) Cosmology and quantum gravity explain where the first and second law of thermodynamics originates.
- 7) Quantum gravity allows us to track potential and kinetic energy changes.

### **Unification of the Special Theory of Relativity with the General Theory of Relativity**

Before Einstein's special theory of relativity (SR), space and time were separate. Experiments had shown (Aether experiments) that the speed of light was a constant but Einstein understood the implications. Basically, SR is the origin of gamma and its relation to the velocity ratio  $V/C$  where  $C$  is the speed of light. Gamma is defined as  $\gamma = \text{mass}/(\text{mass} + \text{kinetic energy})$ . For example if a proton with mass  $938.272 \text{ Mev}$  has kinetic energy  $10.11 \text{ Mev}$ ,  $\gamma = 938.27 / (938.27 + 10.11) = 0.989$ . Gamma is always less than 1.0 and is considered the shift into space-time for a particle with kinetic energy. Velocity is directly related to gamma in the SR equation  $V/C = (1 - \gamma^2)^{.5}$ . For  $10.11 \text{ Mev}$  of kinetic energy,  $V/C$  would be  $0.145$ . The new variable gamma is related to a quantity called time dilation. Time dilation  $dt = 1/\gamma - 1$ . In the example above  $dt = 1/0.989 - 1 = 0.0103$ . The saying "moving clocks run slow" describes the situation. When a particle has kinetic energy and if it is moving relative to the observer, time

runs 1.03% slower for the particle. This is the basis of the Lorentz transformation [2]. After the development of SR, everyone started combining space and time together as space-time.

The theory of General relativity (GR) came later. Einstein modernized our understanding of gravity and stated that gravity could be considered the geometry of space-time. Thereafter, matrix equations called metric tensors defined relationships between four dimensions. The saying “mass curves space-time and objects follow the curvature” describes how objects move in space-time. Basically, acceleration and gravitational acceleration are equivalent (known as the equivalence principle). Objects do not respond to gravitational fields, they just move along a path defined for them by the space-time they are embedded in. Schwarzschild solved the metric equations for a simple case. His solution was very similar to the equation we use for a geodesic. His solution was  $G=S \cdot V^2/M$  but his S is twice the geodesic radius R. A geodesic is the relationship between velocity, mass and radius that give the gravitational constant G, i.e.  $G=R \cdot V^2/M$ . Schwarzschild also developed equation for time dilation for the central mass. The equations are in derived in Appendix 2 with examples below:

Schwarzschild’s time dilation (dt) is calculated below for earth Mass M and earth Radius R:

<b>Astronaut (kg)</b>	<b>90</b>
<b>Mass M kg (earth)</b>	<b>5.98E+24</b>
<b>earth R (m)</b>	<b>6378100</b>
<b>dt=1/((1-G*m/(R*C^2)))^0.5-1</b>	
<b>3.47E-10</b>	<b>mass curves space</b>

The Schwarzschild equation can’t be used for the Astronaut’s time dilation. It has to be calculated from special relativity using her velocity 7908 m/sec and mass 90 kg.

$$\text{Gamma (g)}=0.9999999997$$

$$V/C=2.64e-5 \text{ and } V=7909 \text{ m/sec}$$

$$dt=1/g-1=3.47e-10$$

In cellular cosmology variables of interest are related to cell radius. In this case cell radius  $r=R_{\text{earth}} \cdot 1.67e-27/M_{\text{earth}}=6.38e6 \cdot 1.67e-27/5.98e24=2.17e-6$  meters. This characterizes the quantum level with a central mass of  $1.67e-27$  kg and a velocity of 7909 m/sec. Below the Schwarzschild equation is used to calculate dt:



<b>Astronaut (kg)</b>	<b>90</b>
<b>Mass M kg (earth)</b>	<b>5.98E+24</b>
<b>earth R (m)</b>	<b>6378100</b>
<b><math>r=R*1.67e-27/M=2.17e-6</math> meters</b>	
<b><math>dt=1/((1-EXP(90)*G*1.67e-27/(r*C^2)))^0.5-1</math></b>	
<b>3.47E-10 mass curves space</b>	
<b>and particle follows curve</b>	

The SR equation  $(g)=(1-(V/C)^2)^{.5}=(1-(7909/3e8)^2)^{.5} = 0.9999999997$  also gives  $dt=1/g-1=3.47e-10$ . See the difference? Schwarzschild's equation for general relativity doesn't work for the astronaut's mass because it is not the central mass. With cellular cosmology, special relativity and general relativity always give the same correct result because the mass of a proton is always the central mass ( $1.67e-27$  Kg).

Above in the section entitled "Gamma", we asked the question: "How could Newton know that gravity is the curvature of space time?" In fact, we calculated the small gamma  $(g)=0.9999999997$  and wondered how such a small number could curve space time. Continue the story but use cellular cosmology. In cellular cosmology, the geodesic is  $G=(r*V^2/m)/exp(90)$  but  $m$  is the mass of a proton. Solved for radius, the equation is  $radius\ r=(G*m/V^2)/exp(90)$ . Velocity  $V^2$  is the only variable that indicates that space-time is curved because the other values in the geodesic equation are constants (mass  $m$  is the mass of a proton  $1.67e-27$  Kg). When  $V$  is small,  $r$  is large. When the radius of curvature is large, the geodesic would be almost flat (meaning space time is not curved). Please don't misunderstand the cause of the curvature. It is not mass and it is not velocity, it is a cellular radius established by a field. We can now answer the question: "How does gamma as low as  $g=0.9999999997$  curve space?" It doesn't, but it is an indication that velocity is involved. **General relativity continues right down to the quantum level.** A 2.723 MeV field shapes space and protons fall into the curvature and orbit with velocity  $V$ . Velocity is only an indication that there is curvature. Could Isaac have known this? Well maybe if he had read about Albert Einstein's work, known Schwarzschild's solution to the metric equations, been dis-satisfied with the Planck scale, read papers regarding the results of satellite missions that allowed him to know the approximate number of protons in the universe and developed cellular cosmology. Newton, like all of us "stands on the shoulders of giants" and others were asking question about gravitation. He knew that experiments could answer the questions and he deserves to be a hero. Most breakthroughs [5] in science take a long time to become accepted because there is a network that protects the status quo. New work can become recognized only if key people recognize the value of your work and it is correct (Many don't like this system but basic physics doesn't change every day).

Unified relativity will be useful when we use cells to model expansion of the universe. Einstein and Schwarzschild extended Newton's understanding of gravity with their equations for curved space-time. Schwarzschild's equation for the bending of light around massive objects is a tool used to measure distant masses such as black holes.

## Quantum gravity reconciles gravity with the remaining forces

Quantum gravity is only slightly different than the other forces of nature [3][16][22]. Before considering gravitation more thoroughly, it is instructive to review other interactions (forces) supported by information extracted from the proton mass model (Appendix 2). The fundamental forces of nature are simply Force= Field energy divided by Radius ( $F=E/R$ ) and this is equal to inertial force  $F=m*V^2/R$  but only gravity contains the coupling constant  $1/\exp(90)$ . This makes the equation for radius R basic to force unification. The equation was derived in the section entitled “Quantum Circles”.


$$R=1.973e-13/(E*m/g)^{0.5} \text{ (called the R equation below)}$$

where  $HC/(2*\pi)=1.973e-13 \text{ MeV-m}$

The R equation is arranged in a column of calculations below for the Proton. The inputs are: M (energy of the mass), E (energy of the capturing field) and kinetic energy.

Force Table			<b>Strong MeV</b>	<b>Strong Residual</b>	<b>Gravity proton</b>	<b>Electro- Magnetic</b>
		(Higgs ref)	128992.05			
Field Energy E (mev)			957.18	20.303	2.732	2.72E-05
Particle Mass (mev)			129.54	928.120743	938.272	0.511
Mass M (kg)			2.31E-28	1.6545E-27	1.673E-27	9.11E-31
Kinetic Energy (mev)			798.58	10.1513	10.110	1.361E-05
		Rydberg energy from PDG				1.361E-05
Gamma (g)=m/(m+ke)			0.1395	0.989	0.989	1.000
Velocity Ratio		$v/C=(1-(g)^2)^{.5}$	0.9902	0.1467	0.1456	0.0073
R (meters) =((HC/(2pi))/(E*M/g)^0.5)			<b>2.0928E-16</b>	<b>1.430E-15</b>	<b>7.224E-14</b>	<b>5.291E-11</b>
Force	$F=E/R*1.6022e-13$		7.1E+05	2275.20	**	8.242E-08
Inertial	$F=M/g*V^2/R \text{ nt}$		7.1E+05	2262.9	<b>3.656E-38</b>	8.24E-08
					1/exp(90)	

These four forces are aspects of the energy interaction involving the exchange of the logarithm 2.0 [10][24] and are referred to as the strong force, residual strong (weak force), electromagnetic force, and the gravitational force. The R equation gives the size of the quantum circle but there are different particles and different kinetic energies for each of the forces. The radial position of the particle can be anywhere near the surface but is probably at R.



The four forces are diagrammed above. First note the similarity. They all have quantum circles of radius R (in meters) associated with them. The proton mass table is summarized below. Note that mass+kinetic energy is opposite and equal to the combined field energies. The total is 959.92 MeV. In the beginning, a separation from zero occurred:

$$0 = 959.92 \text{ MeV} - 959.92 \text{ MeV}$$

For the strong forces, mass+kinetic energy is equal and opposite the field energy. But the quarks+ke total (928.21 MeV) falls into a 20.3 MeV strong residual field energy and with 10.15 MeV of kinetic energy. Nature is built on nested orbits. The resulting neutron has 20.3 MeV of potential energy and falls to the gravitational radius  $7.224 \times 10^{-14}$  meters where it has 10.11 MeV of kinetic energy. After the neutron decays, the proton and electron have opposite  $27.2 \times 10^{-6}$  MeV fields and the electron falls into a field with  $13.6 \times 10^{-6}$  MeV of kinetic energy. The fundamental point is that field energy characterizes the *space* a particle exists in and *mass+kinetic* energy characterizes the particle that orbits on the surface of the space. The exact mass of the proton (938.272 MeV) results from neutron decay.

Mass and Kinetic Energy			Field energy	
Mass	KE	Strong Residual	Strong field energy	Gravitational Energy
MeV	MeV		MeV	MeV
<b>Strong</b>	<b>130.16</b>	<b>799.25</b>	<b>-957.18</b>	-2.73
<b>Strong Residual</b>		<b>10.15</b>		
<b>Neutron</b>		<b>939.57 (-20.3)</b>		-959.92
<b>Below the Neutron decays ejecting an anti-neutrino and an electron with ke</b>				
<b>ejected neutrino</b>		<b>-0.67</b>		
<b>Electron</b>	<b>-0.51</b>	<b>-0.11</b>	<b>2.72E-05</b>	<b>E/M charge splits</b>
<b>Proton</b>		<b>938.27</b>	<b>-2.72E-05</b>	
<b>neutrinos</b>		0.05		
<b>Gravitational ke</b>		<b>10.15</b>	<b>10.11</b>	
<b>Gravitational pe</b>		<b>10.15</b>	<b>10.19</b>	
<b>Total</b>		959.92		
(959.92=938.27+.67+.511+.67+.05+20.3)				

Each of the forces is a little different.

**Strong Force:** There are actually three quarks and confining forces added together in the strong force information above. It confines the quarks inside the atom. High energy colliders can't knock the three individual quarks of the proton out which means they aren't observed independently. Their masses, kinetic energies and associated fields are listed below for a proton.

	MeV	MeV	MeV	MeV
<b>Quark S</b>	<b>101.947</b>	<b>641.880</b>	<b>-753.291</b>	<b>-0.687</b>
<b>Quark U</b>	<b>13.797</b>	<b>78.685</b>	<b>-101.947</b>	<b>-0.687</b>
<b>Quark D</b>	<b>13.797</b>	<b>78.685</b>	<b>-101.947</b>	<b>-0.687</b>
	<b>129.541</b>	<b>799.251</b>	<b>-957.185</b>	<b>-2.061</b>

**Strong Residual:** This circle is associated with fusion. A mass of  $129.5+799.25-.67=928.12$  (for the proton) orbits a 20.3 MeV residual energy field with kinetic energy 10.15 MeV. When protons and neutrons are forced into the nucleus, they give up a portion of this energy depending on the atoms fusing and this binds the protons and neutrons into a nucleus [14]. This appears to be the main function of the strong residual force but it is involved in particle decay. (Note: The strong residual 20.3 MeV is 959.92 minus the neutron mass 939.56).


**Gravity:** The diagram above shows that there was about 20.3 MeV of kinetic energy available from the proton model. A quantum orbit is formed with 10.11 MeV of kinetic energy. Since the proton is attracted to and separated from the center of the field, there was also 10.11 MeV of internal potential energy when the orbit is established at  $7.22e-14$  meters for 2.723 MeV of gravitational field energy. It is fundamental to gravity, space-time and the starting point for

expansion. As the kinetic energy of the proton on the surface of this radius decreases, the radius increases and the orbit becomes non-quantum. For the proton, it started at 10.11 MeV and is now 1.29e-12 MeV. This increases the radius to 0.54 meters for the cosmological cells. Together with all the other  $\exp(180)$  cells, they create the space we call the universe. The gravitational coupling constant  $1/\exp(90)$  makes gravity weak and long range. The fundamental force associated with gravity is extremely weak ( $3.66e-38$  N). Gravity acts through long distances and the gravity of all the protons in a mass like the earth act together to affect distant objects. Einstein taught us that gravity is the geometry of space-time but actually all the forces are similar. Field energy for each of the four forces shapes space-time into quantum circles and mass falls to the surface of the circle (they all are so similar we can even use the same unit for their radius (meters)).

**Electromagnetic (E/M):** Electrons circle a proton at the radius  $5.29e-11$  meters. This is the only circle that shifts in quantum steps and releases light. The E/M force is the force that underlies the thermodynamics of gases. It is also the reason we can't walk through a wall. Our electrons repel electrons in the wall. All the other forces attract but the E/M force repels and attracts. The reason for this is that the electromagnetic energy is separation energy. When a neutron decays, the proton takes a positive E/M field energy and the electron takes a negative E/M field energy.


The following diagram illustrates how the circles are arranged by nature.

The quarks are "bundled" by the strong field and orbit the strong residual field energy with kinetic energy 10.15 MeV. Together this makes the proton mass 938.27 MeV after it ejects a neutrino with mass 0.671 MeV.



And of course the 0.511 MeV electron circles the proton with kinetic energy  $13.6e-6$  MeV. The proton orbits the gravitational field with kinetic energy 10.11 MeV. Originally this was 20.3 MeV of potential energy but the proton fell into the field and gained kinetic energy.

### Four Nested Orbits for the Proton



## EXPANSION EQUATIONS

Quantum gravity is fundamental to expansion equations that give the size of the universe. Simple kinetic energy and potential energy equations are applied to expanding cells. Each particle of mass  $m$  has kinetic energy and an associated velocity  $V$  tangential to the cell surface. The model shows protons with about 20.3 MeV that fall into “orbits” with 10.11 MeV of kinetic energy and 10.11 MeV of potential energy. Initially the proton on the cell surface has high velocity (0.14C) and inertial force, the basis of quantum gravity. Tangential kinetic energy ratio decreases directly with expansion ratio and can be modeled as orbit that maintains the gravitational constant at  $G$ .

$G$  remains constant  $G=rv^2/(M)$

$RV^2/(M/g)=rv^2/(M/g0)$	$RV^2/M=rv^2$	10.11 ke	$ke=.5 (m/g)v^2$
$RV^2*g=rv^2*g0$	$RV^2=rv^2$	↓	$ke0=.5 (m/g0)V^2$
$(v/V)^2=(r/R)*g0/g$	$(v/V)^2=(r/R)$		$ke/ke0=(m/g)v^2/((m/g0)V^2)=r/R$
$(v/V)=(r/R)^.5*(g0/g)^.5$			$ke/ke0=(g0/g)(v/V)^2$
<b><math>ke=ke0*(r/R)</math></b>	Ke decreases with r		$ke=ke0*(g0/g)(r/R)$

One set of information defines each proton but duplication produces  $\exp(180)$  produces. Initially the universe has probability 1.0. With each proton’s probability  $1/\exp(180)$  the duplication process re-establishes probability  $1.0=1/\exp(180)*\exp(180)$ . There is one proton per cell. After duplication and expansion the cells are no longer quantum. (Magnetic moments or other forces may disrupt the proton forcing it to move throughout the cell). This is explored below in the section entitled “Proof of non-quantum behavior”. The kinetic energy of the proton can be considered pressure and pressure expands the cell. The integral  $PdV$  does work against gravity converting kinetic energy to potential energy ( $dV$  is a small volume change and  $PdV$  is a small energy change that must be added (integrated) since  $P$  and  $V$  are both changing). During expansion, gravity causes some protons to fall (accelerate) toward each other and accumulate, exchanging potential energy for kinetic energy. It is this energy that we see when orbits are


established around galaxies and planetary systems. It is also this energy that provides pressures and temperatures high enough to initiate fusion.

### Cell Diagram

Initial cell radius is  $7.22e-14$  meters. Initial forces in the cell are balanced and initially are  $3.66e-38$  N. With an initial kinetic energy of 10.11 MeV, the initial expansion velocity can be calculated.

$$\text{Gamma (g)} = 938.27 / (938.27 + 10.11) = 0.9897$$

$$V/C = (1 - 0.9897^2)^{.5} = 0.146$$



$$\text{PE expansion} = \int F \, dR$$

The goal below is to model expansion of a small cell that provides values scalable to the universe.

### Nomenclature

(all calculations are MKS)

t-time

g=dimensionless time=time/alpha time

Lower case r is a cell radius

Upper case R=r\*exp(60)

R1 radius is first expansion component

R3 radius is second expansion component

H3 is Hubble's constant for R3


First expansion component; R1  
 $(r/r_0)^3$  increases as  $(t/\alpha)^2$  (kinetic energy requirement)  
 $r=r_0 \cdot g^{(2/3)}$   
 $R=r_0 \cdot \exp(60) \cdot g^{(2/3)}$   
 $r_0=1.93e-13/(2.723 \cdot 2.723)^{.5}=7.22e-14$  m  
 $R1=(7.22e-14 \cdot \exp(60)) \cdot g^{(2/3)}$

Second expansion component: R3  
 $dr/(r \cdot dt)=H3$   
 $dr=H3 \cdot r \cdot dt$   
 $dr=H3 \cdot \alpha \cdot r \cdot dg$  (dt=alpha dg)  
 $dr=H3 \cdot \alpha \cdot r_0 \cdot g^{(2/3)} \cdot dg$   
 $r=H3 \cdot \alpha \cdot r_0 \cdot g^{(5/3)}/1.6666$   
 $R3=H3 \cdot \alpha \cdot (7.35e-14 \cdot \exp(60)) \cdot g^{(5/3)}/1.666$

$r1+r3=(7.22e-14) \cdot g^{(2/3)}+(7.22e-14) \cdot g^{(5/3)} \cdot H1 \cdot \alpha/1.666$   
 $R1+R3=r1 \cdot \exp(60)+r3 \cdot \exp(60)$   
 $t=\alpha \cdot (R1/7.22e-14)^{(3/2)}$   
 $(R1/7.22e-14)^{(3/2)}=g=t/\alpha$

Alpha and H1 are evaluated to fit WMAP data. Alpha=0.0529 seconds and H1=3.1e-18/sec.

The R1+R3 model can be compared with the concordance [6][7] and Cmagic [8] models.




The above chart is established physics and uses the cosmology presented in Peebles [6]. It is called the Lambda Cold Dark Matter (CDM) model with data gathered from missions COBE,



WMAP [7] and PLANCK. It compares favorably with expansion data from supernovae (Cmagic [8]). However, there are differences that should be pointed out. The CDM model assumes that velocity is larger than the speed of light. Early in expansion, the accepted model involves several thousand times the speed of light. The author's cellular model's maximum expansion velocity is  $V/C=0.146$ . Cells move apart at greater than  $C$  but expansion of the cell is below the speed of light (sub-luminal).

In the author's work, the speed of light cannot be exceeded because it is just the ratio between space dimension and time dimension. Here is the explanation:

The velocity of light is the ratio of dimensions, the distance dimension and time dimension. Both dimensions are on a quantum circle. A quantum circle is defined by field energy juxtaposed by light energy. (It also repeats at  $\psi^*\psi=1$  where  $\psi$  is the wave function).



**Energy is defined by a quantum circle**

**$E=H/t$  where  $C=3e8$  m/sec and  $t$  is time around the circle**

**$t=H/E$**

**Time is measured around a circle of radius  $r$**

**$t=2*\pi*r/C$**

**$d=2*\pi*r$**

**The speed of light is the ratio distance/time**

**$C=d/t$**

### **Transition from quantum behavior**

The author believes that the space we walk around in is defined by gravity at the quantum level ( $r=7.22e-14$  meters) by cells that expanded to a present radius of about 0.55 meters/cell. In three dimensions  $\exp(180)$  cells give large  $R=0.54*\exp(60)=6.2e25$  meters. You might be wondering, as I did, whether a cell that expands from  $7.22e-14$  meters to 0.54 meters is still quantum. In quantum mechanics, quantum particles move in circles and are statistically "everywhere" at once near a surface and movement into the interior of the sphere that defines them is very limited. For example, the electron does not normally move inside the sphere  $5.29e-11$  meters and if it is forced to, it is relativistic or de-generate. For expanded cells, all mass is associated a cell but mass with kinetic energy travels slower than the velocity of light and is no longer quantum in nature.


<b>cell uc2 c198</b>			
<b>Field Energy</b>		<b>r cell (m)</b>	<b>r' cell (m)</b>
	<b>2.732</b>	<b>7.22E-14</b>	<b>3.14E-12</b>
<b>KE (MeV)</b>		<b>10.11</b>	<b>0.233</b>
<b>Velocity (m/s)</b>		<b>3.00E+08</b>	<b>6.66E+06</b>
<b>V/C</b>		<b>1</b>	<b>0.022</b>
<b>gamma</b>		<b>1</b>	<b>0.9998</b>
<b>Field side</b>	<b>R side</b>		<b>R side</b>
<b>H/E</b>	<b>2*pi*r/C</b>		<b>2*pi*r/V</b>
<b>1.514E-21</b>	<b>1.51E-21</b>		<b>2.96E-18</b>
<b>p=E/C (mev-s)</b>	<b>9.11E-09</b>		<b>4.10E-07</b>
<b>p*R/h</b>	<b>1.000</b>		<b>1.95E+03</b>

The quantum test  $pR/h=1$ , where  $p$  is momentum is maintained by the left side of the diagram above.  $H/E=4.13e-21/2.73$  seconds equals  $2\pi r/C=1.51e-21$  seconds (fundamental time) but  $pR/h$  is greater than 1 for the right side, indicating that as expansion occurs we are no longer dealing with quantum size circles. The right side of the diagram shows that as the cell expands from  $r$  to  $r'$  velocity decreases.

### What actually expands the universe?

Pressure is the collective action of particles with kinetic energy (temperature) that collide with each other in all directions. The fact that protons are colliding and able to move throughout the space created by expanding the fundamental radius  $7.22e-14$  meters indicate that a critical transition has occurred. Protons enter the radius that defines gravity and pressure expands space itself. Particles exhibit non-quantum behavior (perhaps because the force is very low and it is easy to force particles into the interior of the volume). The gravitational kinetic energy 10.11 MeV can now be considered the source of velocity (pressure and temperature) inside cells. The protons and their kinetic energy exist in the space around us. The energy can't escape because there is nowhere for it to go but cells have a specific kinetic energy and potential energy depending on their history.

The Boltzmann relationship  $T(K)=ke/(1.5*B)$  with  $B=8.62e-11$  MeV/K assigns a temperature to kinetic energy. Cosmologists use the expansion ratio  $z$  to scale temperatures and the horizontal axis in the plot below is the natural logarithm 45 progressing to about 90. Large scale time progresses from  $\exp(45)*1.514e-21=0.0529$  seconds to approximately  $\exp(88.5)*1.51e-21$ seconds= approximately 14 billion years presently.



The discontinuity in temperature is explained in reference 19, but the temperature is 2.73K at the current stage of expansion.

### Proof of non-quantum behavior


There is a critical concept at stake that needs our understanding. If the particles are non-quantum and the expansion kinetic energy is temperature, it is no longer limited to a surface. Particles with kinetic energy bounce off of one another and create pressure. Is it this pressure that expands the universe? Can the particles fill all of space or are they quantum like and limited in their travel? If we calculate what a gas would do perhaps we can answer the above two questions.

The gas constant  $R$ , is 8.317 Joule/K/Mole. (Joule=Nt-m and 1000 Mole/Kg for H). If we assume an ideal gas for hydrogen the gas constant  $R=8317$  Nt-m/K/Kg and the pressure  $P$  would be:

Pressure  $P=8317 \cdot \text{density} \cdot \text{temperature}$  (Nt-m/K/kg\*kg/m<sup>3</sup>\*K=Nt/m<sup>2</sup>) where density is kg/m<sup>3</sup> and temperature is degrees Kelvin (K).

With density based on one proton for half the cells (the other half is probably cold dark matter [13]) and an initial radius of 7.22e-14 meters, the above initial pressure is 2.97e26 Nt/m<sup>2</sup> where initial temperature=7.58e10 K. The following table models integral  $P \cdot dV$  (an integral means we add increments together since  $P$  and volume change as  $R$  increases) for the first few steps in expansion.

Volume/cell (m <sup>3</sup> )	1.67E-39	2.72E-39	4.43E-39	7.24E-39	1.18E-38
Density (kg/m <sup>3</sup> )	5.02E+11	3.08E+11	1.89E+11	1.16E+11	7.09E+10
Temperature (K)	7.56E+10	1.01E+10	1.08E+10	1.05E+10	9.84E+09
Pressure (NT/m <sup>2</sup> )	2.97E+26	2.44E+25	1.60E+25	9.54E+24	5.46E+24
Pressure (lb/in <sup>2</sup> )		3.54E+21	2.32E+21	1.38E+21	7.92E+20
Pdv (MeV)		2.11E+00	4.33E-01	4.47E-01	4.28E-01
Integral Pdv (MeV)	6.6	2.11E+00	2.55E+00	2.99E+00	3.42E+00



The integral of  $P \cdot dv$  quickly saturates at a level consistent with the initial kinetic energy of 10.11 MeV (the gas is not ideal and the constant is somewhat uncertain). Overall, *pressure* can be considered the driver for expansion. The net effect is the proton receives gravitational potential energy against a resisting gravitational force. For expansion, the kinetic energy term is initially 10.11 MeV of kinetic energy but decreases as integral Pdv increases. Defined this way, we expect the equation  $10.11 = ke + Pdv$  to be satisfied. Although the proton model and cellular cosmology define kinetic energy for the cell, it is pressure and temperature that expand the universe. Rather than being limited to a quantum mechanical orbit, particles are free to move throughout space. After two additional early transitions (equality of photon and mass density and decoupling of electrons [7]), gravitation is locally able to dominate gas pressure. This gas does not act like the one that thermodynamics normally describes. The particles are gravitationally “sticky” and small accumulations of matter grow and eventually form clusters, galaxies, stars and planets [6][12].

### Current kinetic energy per proton

Each cell (small  $r$ ) that is still expanding is now about 0.54 meters and large  $R = 0.54 \cdot \exp(60) = 6.2e25$  meters. The initial  $KE = 10.11$  MeV has diminished by the ratio of current radius to initial radius ( $0.54 / 7.22e-14 = 7.48e12$ ) and is now  $10.11 / 7.48e12 = 1.35e-12$  MeV/proton. Of course, we do not see the cells and they blend together to form the space around us.

## TIME AND COSMOLOGY

There is a strange situation in fundamental physics regarding time. Well respected physicists [Julian Barbour for example] point out that all quantum mechanical equations are cyclical with time. Common sense tells us that time advances and tension exists between fundamentals and

what we observe. This situation extends to fundamentals of space as well as fundamentals of time. Special relativity and curvature of space time is known to be the source of gravity at the large scale but the author's approach to quantum gravity is not generally known. Further, the concept that velocity is relative seems to be accepted but velocity is related to kinetic energy that is conserved.

The author believes that the cycle time  $1.51e-21$  seconds has repeated many times since the beginning. In other words, a quantum mechanical fundamental time is defined that cycles *and* counts forward (fundamental time\*exp(N)).

<b>Identify the radius and time for the gravitational orbit described above</b>	
<b>Fundamental radius</b>	<b>=<math>1.93e-13/(2.732*2.732)^{.5}=7.224e-14</math> meters</b>
<b>Fundamental time</b>	<b>=<math>7.224e-14*2*PI()/((3e8)^2)=h/E=4.13e-21/2.732</math></b>
<b>Fundamental time</b>	<b><math>1.514E-21</math> seconds</b>

Consider why the universe expands. Kinetic energy (ke) must be turned into gravitational potential energy (pe=Fr) over *time*. Time enters physics through cosmology! The derivation below indicates that the increasing radius of the universe and increasing time are related through expansion.

ke	pe
ke	F r
$1/2M(v)^2$	GMM/r
$1/2M(r/t)^2$	GMM/r
$1/2Mr^3/t^2$	GMM
$1/(2GM)*r^3$	$t^2$
$(r/r0)^3$ increases as	$(t/t0)^2$

The above derivation contains only radius and time. If we believe that expansion occurred we must believe that time advances (perhaps in snapshots of fundamental time  $1.5e-21$  sec).

$(r/r0)^3$  increases as  $(t/\alpha)^2$  (kinetic energy requirement)

With the understanding that the large scale we observe is made of cells defined by gravity and the further understanding that fundamental time cycles, counts and moves everything forward we can simplify our understanding of nature. This cycle is established by the quantum mechanics of the gravitational field inside each proton (the proton model in Appendix 2) and each proton is identical and none occupy a preferred position. All protons advance in elapsed time simultaneously ready for the next count. Elapsed time is the primary variable for the expansion equations and they determine the expanded radius we will label radius prime (r'). It determines how much kinetic energy has been converted to potential energy. In fact, expanded kinetic energy is simply  $ke=10.11*7.22e-14/r'$  MeV.

## **Time associated with kinetic energy**

According to special relativity time runs slow for particles with kinetic energy ( $\gamma = m/(m+ke)$  and  $V/C = (1-\gamma^{-2})^{0.5}$ ). The slowness of time is expressed by  $dt = 1/\gamma - 1$  and is about 1% slow for an astronaut orbiting the earth. But  $\gamma$  is established by kinetic energy and kinetic energy is maintained by a particle unless something changes its velocity. Time runs slow for the moving mass even though a new fundamental time cycle has started. This means there are two measures for time. A time phenomenon called the twin paradox has been carefully explored by physicists.

The twin paradox has been proven with synchronized atomic clocks in satellites. A clock in motion runs slower and when they are brought together they read differently. .


Explanation of twin paradox: Fundamental time is controlled at the quantum level. When the journey is complete both twins are together in fundamental time. However, the traveling twin is a little younger because his  $\gamma$  was established and maintained by his kinetic energy.

## **THERMODYNAMICS AND COSMOLOGY**

### **Conservation of Kinetic and Potential Energy**

Particles initially have kinetic energy related to the proton mass model. Kinetic energy is converted to potential energy as expansion occurs and is reversed when mass accumulation starts. The total energy is  $PE+ke=10.11$  MeV/particle. There is a new understanding of space and time in cellular cosmology. General relativity extends to the quantum level and we are walking around in space that has expanded from that level.

Gravity is defined at the quantum scale by the energy 10.11 MeV/particle and gravitational field energy 2.723 MeV at Radius =  $7.22e-14$  meters with  $\exp(180)$  cells. Since expansion is well characterized (and agrees with the author's calculated expansion), one can simply calculate expansion kinetic energy and expansion potential energy as a function of time and determine if initial KE is in fact converted to potential energy. The initial resisting force is the inertial force  $F=M*V^2/R=3.66e-38$  NT where  $V$  is the tangential velocity  $V/C=0.15$  that decreases with increasing  $R$ . Final potential energy (integral of  $F*dR$ ) is 10.11 MeV but the initial kinetic energy is reduced to the current value of  $1.29e-12$  MeV. To obtain this result the author reduced the number of protons to one half  $\exp(180)$ . Why one half? Details presented in reference 3 suggest that protons (baryons) make up one half of the total mass. The value 10.11 MeV/particle is only enough energy to expand half the cells. Since cold dark matter (CDM) is found surrounding galaxies the other half of the mass apparently has a similar energy source (a mirror of the proton perhaps). Changes in energy are plotted below (the horizontal axis units are increments of time and they quickly saturate).



The graph above indicates that kinetic energy and potential energy are indeed conserved.


### Gravitational Accumulation

Expanded particles do not simply reverse expansion as they fall due to gravitation. At about 200K years after the beginning a condition known as equality [6][7] of photon density and matter density occurred and gravity became an important force determining their behavior. Initially, gravitational accumulation was aided by acoustic waves but as particles collided, their gravitational attractive forces started to dominate and particles no longer behaved like gases we are familiar with. The pressure at equality was about  $5e-8$  psi (pounds per square inch) and the temperature was 9100 K. The gas was low pressure plasma. A later critical juncture in thermodynamics occurred as the plasma cleared (this condition is called decoupling and electrons assume orbits around protons). The temperature at this point was about 3300 degrees K and the pressure was  $6e-14$  psi. At the present time it is  $3.7e-27$  psi and 2.7 K.

### Formation of Black Holes

The author uses a spreadsheet (fallmodel.xls) that combines Newtonian kinetics with the expansion equations. Starting at equality of matter and radiation acoustic waves develop and concentrate mass. WMAP [7] measured the red shift of (spots) that we can now associate with clusters. When the universe was about  $1e22$  meters in size waves were no longer dampened (equality) and started travelled until they were spots of about  $3e21$  meters. This dividing matter into approximately  $2.6e4$  clusters each containing  $1e46$  kg mass. The Jeans length is a natural

wavelength associated with temperature and the state of matter. At decoupling the plasma cleared and the Jeans length transitioned to a much lower value. It went from Jeans high  $5e22$  meters to Jeans low  $3e19$  meters. This low Jeans number is “empirical” [12] but divides the WMAP spot size of  $3e21$  meters into about  $1e6$  smaller spots that we associate with galaxies. The smaller spots contained about  $1e42$  kg mass. According to the fall model, some mass fell quickly inward. One possible reason the first masses form black holes is that their fall velocity becomes very high and the absence of other bodies do not deflect the particles into orbits. Apparently some black holes formed soon after decoupling of mass and radiation and aided galaxy formation. Below the fall velocity as a function of time is shown from fallmodel.xls:




A black hole can be an attractor that brings mass into a galaxy. As new mass falls, it falls into orbits around the black hole and the galaxy builds mass from the inside out since later mass gathers from further away. As it falls to approximately 42% of its origination height, it adds to the outer portion of the galaxy.

### **Gravitational accumulation forms orbits**


First review how orbits are formed.




### Expansion determined height



Particles expand to a height determined by the expansion equations. As accumulation starts, particles fall and in general are deflected into an orbit. The orbit (geodesic) is by definition the combination of velocity, mass and radius that gives the gravitational constant  $G$ . Orbits, also by definition, are force balanced. In other words inertial forces are balance on a particle in a way that it feels no force. The radius is, in general, less than half the original height. The graph below is a simulation of a particle falling into a galaxy of  $1e41$  Kg to a position similar to our sun in the Milky Way. The force of gravity on the particle starting for a high position increases as it falls toward the center ( $F=GMm/R^2$ ) since  $R$  is decreasing. But as the particle falls it is deflected into a circle ideally and the inertial force increases. The forces are balanced when the particle reaches 42% its original height. The velocity is calculated for a particle given enough time to fall through the distances and the graph below shows the acceleration driving the fall. The gravitational acceleration inward minus inertial forward outward is shown below. The particle achieves orbit when the forces are balanced (zero acceleration).




The chart below shows the kinetic and potential energy changes. Potential energy changes from 10.11 MeV is shown for graphing convenience and kinetic energy is calculated from the velocity increases due to excess gravitational acceleration (gravitation minus inertial). Of course potential energy is converted into kinetic energy.



The proposed model for expansion is based on an orbiting proton with 10.11 MeV of kinetic energy (b). As expansion occurred (process (b)→(c) below), 10.11 MeV of kinetic energy was converted to 10.11 MeV of external potential energy. At a much later point in expansion (c), although there is motion (temperature) of the proton on the surface of the expanding cell, there is no motion between cells (protons) except for expansion. With the proton velocity nil between

cells geodesics will be extremely flat (on the order of  $5e38m$ ) compared to  $6.2e25m$ . This causes acceleration of particles toward one another (process 2 below) and external kinetic energy (between protons) increases as protons fall back toward the geodesic (d)→(e). Theoretically, 10.11 MeV of external potential energy could be reconverted to 10.11 MeV of kinetic energy as particles fall toward one another but as we will show below, only a fraction of 10.11 is found in galactic and planetary orbits. (The example below is for 0.01 MeV of kinetic energy re-conversion but later we will show that this it is only  $1.6e-3$  MeV). Overall, process (b)→(c)→(d)→(e) converts cellular kinetic energy to external potential energy between cells and a small amount of kinetic energy.



What actually happened during expansion was a transition occurred and acoustic waves broke the total mass into about 27000 clusters. After equality of photon density and mass density, process (d)→(e) occurred; protons accumulated and eventually fell into orbits that we observe as clusters of galaxies, galaxies, etc. As an engineer one cannot help but be impressed with the approximate energy conservation of combined processes 1 and 2. These processes represent the largest construction project in nature (the universe) and almost no energy is consumed. The “neat trick” seems to be cells that expand and are able to move and fall relative to each other after they are far apart. Contraction is limited to a fraction of 10.11 MeV since orbits or solid objects are formed. Although gravitation is based on the mass of  $\exp(180)$  protons, this may be a combination of protons and cold dark matter.

### Is Velocity Relative?

To properly address this subject we must show a complete energy balance/neutron as the universe expands. The balance of mass with kinetic (959.91 MeV) exactly balances field energy (959.91 MeV). Zero energy is separated into these components at the beginning and as expansion occurs the energy components change slightly but still balance to zero. About the only thing that changes is the conversion of potential energy to kinetic energy. As expansion starts, using this condition as a reference, there is 10.15 MeV of kinetic energy.

ENERGY BALANCE PER PARTICLE AS EXPANSION OCCURS									
simple cell bk9	Fusion		Fusion		Gravitational	Gravitational	Electron		Total
begin	Atoms/Strong	ke	Heat	e/m	Kinetic Energy	Potential E	neutrinos		mev
	131.4566								131.4566 mass
0.000	797.9575	10.15127013		0	0.00	20.30254027	4.85E-02		828.4598 ke
-2.732	-957.1848								-959.9164 pot
									0 total
<b>after decay to P &amp; fusion to helium</b>				e/m					
(released as K	129.5409						1.2789		130.8197 mass
	798.5799	8.521270135	1.63	2.720E-05	10.15	10.15	0.1114		829.1451 ke
-2.7316	-957.1848								-959.9164 pot
									0 total
					Grav Kinetic E	Grav Pot E	CMB		
<b>Now</b>	6.20E+25								0 total
	129.5409						1.2789		130.8197 mass
	798.5799	8.351270135	1.8	2.720E-05	1.40E-12	20.30	0.1114		829.1451 ke
-2.7316	-957.1848								-959.9164 pot
									0 total
<b>Near end</b>	2.36E+26								
	129.5409						1.2789		130.8197 mass
	798.5799	8.351270135	1.8	2.720E-05	0.00	20.30	0.1114		829.1451 ke
-2.7316	-957.1848								-959.9164 pot
									0 total

The first law of thermodynamics is straightforward. It states that energy can be converted from one form into another but not created or destroyed. The concept that velocity is relative seems to be accepted but velocity is related to kinetic energy ( $ke=0.5*m*V^2$ ) that is conserved. Some would point out that special relativity indicates that simultaneity is dependent on motion and therefore, since motion is relative, time is relative. They use the Lorentz transformation to calculate how time is changed by velocity relative to some other particle. If velocity is relative there is tension between this statement and kinetic energy. How can the first law of thermodynamics be satisfied (no destruction of energy) if velocity is relative but kinetic energy is not relative? Furthermore, the above diagram is violated if energy is not conserved. Using tools associated with cellular cosmology, we will account for kinetic energy/per proton for orbits associated with earth. This will help understand how potential energy and kinetic are now distributed and provide information regarding the “velocity is relative” concept.

### Potential and kinetic energy distribution

On earth we feel acceleration. In the section entitled “Gravitational Force on the Earth” an astronaut’s velocity was determined as follows:

<b>R (m)</b>	<b>orbit R =earth R</b>	<b>6.38E+06</b>
<b>M (kg)</b>	<b>Earth Mass</b>	<b>5.98E+24</b>
<b>a (m/sec<sup>2</sup>)</b>	<b>a=GM/R<sup>2</sup></b>	<b>9.8</b>
<b>V</b>	<b>V=(aR)<sup>0.5</sup></b>	<b>7909</b>

Her kinetic energy/proton is  $5.32e-15*7909^2=3.3e-7$  MeV. Velocity, kinetic energy, gamma and cell radius associated with orbits that involve the earth are listed below. From cell r above, the large scale geodesic is scaled up by the ratio of the central mass/proton mass. For example, for the earth, the cellular r geodesic scales to the earth geodesic as follows: The earth geodesic based on one cell radius is  $R=r*M/m=2.14e-6*5.98e25/1.67e-27=6.4e6$  meters.

Note: All calculations below are per proton. The reason to do this is we can compare the energy to kinetic and potential energy changes/proton for expansion and compare them to the total 10.11 MeV.

Orbit		Vel m/sec	ke (MeV)		gamma	cell r (m)	Mass M	Orbital R
P/earth	6378100	7.91E+03	3.28E-07	ke from 7898	0.9999999997	2.163E-06	5.98E+24	6.35E+06
earth surface		4.64E+02	1.13E-09	ke earth spin				
earth/sun	1.55E+11	2.92E+04	4.48E-06	ke in earth orbit	0.9999999952	1.583E-07	1.99E+30	1.54E+11
sun/galaxy	2.5098E+20	5.58E+05	1.63E-03	ke from 8.29 kps gala	0.9999982590	4.345E-10	1.17E+42	2.50E+20
	galaxy	5.60E+05	1.65E-03	ke galaxy	0.9999982464	4.314E-10		
			3.28E-03	total ke	0.9999965003	2.161E-10	4.34E-10	

Above we add earth orbit that should have  $3.3e-7$  MeV/proton adding to that the kinetic energy/proton of the earth orbit around the sun and adding to that the kinetic energy of the sun's orbit around the galactic center. The sun's kinetic energy in a flat galaxy velocity profile is included from the graph below in the topic entitled "Flat galactic velocity profiles". The next orbit should be a galaxy orbiting the center of a cluster but in general this is not observed (the possible reason is below but you can skip if you desire).

A possible reason that galaxies do not move in orbits around a cluster may be that galaxies have mass all around them and the forces are balanced. This condition may date back to the high to low Jeans length transition that forms the galactic centers on almost equal spacing. Only if a galaxy moves toward another galaxy does gravitation force start the galaxy moving (this appears to be the case for Andromeda and the Milky Way that are moving toward one another). Another possible reason is that galaxies form relatively fast because black holes accelerate their formation and clusters do not have this advantage. Galaxies probably have velocity but a value similar to the stars themselves seems reasonable for the table above.

Adding together the kinetic energies contributions from earth orbit, solar orbit and galactic kinetic energy we can identify only about  $3.3e-3$  MeV/proton. Compare this to the expanded cell potential of 10.11 MeV. Only a small amount of potential energy has been converted back to kinetic energy by falling. (Black holes are an exception since they develop about 10.11 MeV of internal kinetic energy by compression).

### Non-ideality involved in reconversion of kinetic energy to potential energy

Not all cells fall from the geodesic established by the expansion equation and are still expanding. For protons that fall, potential energy is reconverted to kinetic energy (ke) and heat dQ. We will try to account for the energy and determine where the original 10.11 MeV resides. The non-ideal case for normal matter is that collisions occur that cause particles to lose some of their kinetic energy to "friction" between particles. The equation that applies is  $10.11=(ke+dQ)+Pdv$ . The term ke and Pdv are converted back and forth but the term dQ contains the friction energy (heat) and we expect to find this energy in the temperature of the planets and the stars (before fusion adds to the energy). All particles that form a central mass like a star or planet fall are limited in their fall. The electron orbit  $5.29e-11$  meter limit recompression and establishes the following

theoretical limit. The temperature associated with recompression to  $5.29 \times 10^{-11}$  meter cell size is about  $10^8$  degrees K and 0.013 MeV. We do not find this large amount of energy in solid bodies indicating that the recompression process is highly non-ideal. The process of gravitational accumulation (asteroids and comets) converts kinetic energy to heat but particles in the vicinity of the sun have no more than 0.0013 MeV and when they hit a solid body like the earth the earth heats slightly but subsequently radiates its energy to space. Even at the core temperature of the earth (6000K) the kinetic energy is only  $10^{-6}$  MeV.


### Accounting for friction heat

We found 0.0016 MeV in the orbits we know about but there is also heat energy that needs to be accounted for. Heat energy is related to temperature in the earth and sun. Recall that the particles have  $5 \times 10^5$  m/sec and can fall quite hard into solid bodies, i.e. comets and asteroids. The estimates below account for the fact that some protons are more abundant than others.

	Temp	ke/particle	mass each	number	num*mass	num*N*ke	ke/particle (mev)
sun with	1E+06	0.0001293	2.0000E+30	1.0000E+11	2.00E+41	1.55E+64	1.2929E-04
planets	6000	7.758E-07	5.98E+24	1E+12	5.98E+36	2.78E+57	2.3195E-11
dust	2.73	3.5299E-10	5.98E+24	1E+12	5.98E+36	1.26E+54	1.0554E-14
					2.00E+41	1.55E+64	
					1.20E+68		
					average ke/particle		1.2929E-04

Our orbital energy/proton is about 0.003. The total heat energy “found” for the sun, planets and dust was 0.00013 MeV/proton (excluding fusion). Small black holes are compressed and hold more energy but there are relatively few of these. Overall most of the energy we were searching only amounts to about 0.003 MeV/proton. This energy is only a fraction of the 10.11 MeV available.


Here is a plot indicating why we only find a small amount of the original energy. Starting from the right side of the potential energy axis, the average cellular radius would have to become quite low to convert a large amount of potential energy back into kinetic energy.




Most of the protons in the universe, on average, are surrounded by a large amount of space. In this condition, they have about 10.11 MeV of potential energy. Most of the potential energy was gained by overcoming strong gravitational attraction when particles were compressed. Falling into orbits and colliding with other particles does not produce much kinetic energy per particle. Even fusion produced energy is relatively low compared to 10.11. For example, the sun's interior temperature of  $2 \times 10^7$  K has only 0.0026 MeV/proton.

#### Velocity is not relative

This concept should be re-labelled "The speed of light  $C$  is always measured  $3 \times 10^8$  m/sec". The idea that velocity is relative came from experiments that showed object velocities do not add to the speed of light. In addition, velocity is an incomplete specification for kinetic energy. Most of the energy around us comes from gravitational potential energy that has been reconverted to kinetic energy and then to compression energy and heat when bodies like the earth are formed. The total of potential energy and kinetic energy is constant (10.11 MeV). The ke portion is really only 0.003 MeV but it wouldn't show in the graph.



Clearly the energy is accounted for since it totals 10.11 MeV. However, some particles can transfer energy to other particles. This is a local effect because the particles must interact (bump) to transfer energy.



If particles interact, some particles will have lower energy and some higher. However, the average of all particles will be 10.11 MeV. The statement that velocity is relative becomes “on average the potential and kinetic energy of all particles is 10.11 MeV”. The statement regarding relativity is only for kinetic energy that we lose track of because we don’t know the history of every particle. If you calculate time dilation and the Lorentz transformation on the basis that velocity is relative, remember that your calculation is relative locally and you do not have enough information to apply it globally.



## Origin of the Second Law of Thermodynamics

The second law is not as straightforward as energy conservation. A quantity called entropy describes the probability of energy states for systems with many particles. The second law states that more probable energy states become filled over time and energy differences that can be used to carry out work become less available. The source of a high original state that can continually “run down” has been difficult to identify.

### Friction, heat and entropy

Thermodynamics is the science of groups of particles. Entropy,  $S$  is defined as follows and helps characterize the second law of thermodynamics.

The cyclic integral of change in heat energy/divided by temperature is equal or less than  $S$  where  $S$  is defined as entropy, i.e. cyclic integral of  $dQ/T < \text{ or } = dS$ .

The change in the entropy of a system as it undergoes a change of state may be found by integrating:  $S_2 - S_1 = \text{integral } (dQ/T) \text{ from state 1 to state 2}$  [29]. The overall change in  $dQ/T$  will always be less than entropy  $dS$ . In other words entropy, defined this way, always increases. There is a limiting (ideal or reversible) condition where entropy might be equal.

The thermodynamics of a gravitational potential has not been developed to the author's knowledge. After expansion, a very improbable (high information) state has been established. Expanded particles separated from one another are free to accumulate due to gravity. As they do, they fall to lower energy (more probable) states as they collide and accumulate. Mass will accumulate and bodies will fall into orbits around other bodies. As they fall, collisions occur. The collisions/friction causes heat and the temperature rises. As particles form large bodies the temperature and pressure can become so high that they fuse, subsequently explode spewing out elements [15][20] that can combine into molecules and life [24]. Conventional thermodynamics describes the behavior of gases that gathers around planets and stars. There are a lot of potential states awaiting particles that fall and collide due to gravitation potential. In general when particles collide during accumulation, heat will be generated and entropy  $dS$  will exceed  $S_2 - S_1$ . This is the origin of the universe's initial low entropy state. The “zeroth” law of thermodynamics states that entropy is zero at absolute zero. It might be better to state the “zeroth” law by referring to cosmology. Entropy is zero in the expanded condition before non-ideal collisions occur. The one way downward street begins as the expanded state and the heat we deal with originates from gravitational potential energy.


The topic below entitled “Compression thermodynamics” indicates that there are two levels of thermodynamics at play. The particles around are in the electromagnetic controlled region of the chart. The electron is circling the atom and controls the thermodynamics. The proton itself had external energy but most of the 10.11 MeV is now potential energy and a small amount of is the kinetic energy of large body orbits. Under extreme compression, like that in the core of small black holes, the proton's gravitational kinetic energy comes back into play.

# DARK MATTER


There is matter in the universe that can't be easily observed. It is called cold dark matter (CDM) because it appears to interact only gravitationally. We know about CDM matter in several ways. The velocity in the outer part of galaxies can be explained only if there is dark matter. It doesn't light up with fusion so the light coming from the outer part of galaxies doesn't reveal its presence. The other way we know about dark matter is Schwarzschild's equations for the bending of light due to gravity. Further, expansion data compared to expansion expectations indicate that there is extra "density" in the universe. In the author's work [13], cold dark matter is proton like (one hot matter proton for every cold dark matter proton) except it is transparent to normal particles. CDM is gravitationally active and this aids accumulation. Apparently it never acts like a gas and is free to accumulate earlier than normal matter.

## Galaxy formation


For our Sun in the Milky Way and ideal conversion of potential energy to kinetic energy the fall model gives the velocity profile for orbits from the center of the galaxy out to the sun. The chart below shows what happens to "ideal mass" as it falls. Ideal mass is mass that does not lose kinetic energy from collisions. It could be "cold dark matter" or just particles that don't happen to collide.



This is a simulation of a velocity in a galaxy like the Milky Way that has 5.58e5 m/sec orbital velocity of a 2e30 kg sun at 8.29 kiloparsec (2.5e20 meters) from the center. The velocity falls off with radius due to Newtonian mechanics.



The velocity profile below represents galaxies that are actually observed (the simulation is for a galaxy similar to our Milky Way with a sun in about the same position).




One difference is that the flat profile galaxy starts with  $1e40$  Kg and the ideal galaxy starts with  $5.5e41$  Kg but the mass must also be distributed differently. Both end up with  $1.19e42$  Kg inside the position of the sun.

### Galaxies with dark matter

Galaxies contain cold dark matter and this “hidden” mass exists in a halo and causes the velocity to be approximately  $5.6e5$  m/sec from near the center to the edge. This galaxy will be a combination of cold dark matter and hot matter. The normal matter lost considerable kinetic energy by friction as it fell into the galaxy. The only way that the mass distribution for the flat


profile could have developed was interaction with another galaxy (see studies of barred spiral galaxies).



There are differences in the way cold dark matter behaves gravitationally. The CDM particles fall with gravity but they accumulate around normal matter. Their conservation of PE and ke should be ideal. The expansion equations apply to them as well and there should be many that expanded to 0.54 meters/mass in size and have about 16 m/min velocity. The theory that they do not interact means that there can never be any compression, fusion, friction related heat (dQ) or transfer of energy between particles.

### Dark energy

Before we finish energy accounting, we must address the subject of dark energy. Some say that this is a huge unknown energy source so large that accelerating expansion eventually reduces density to near zero. The second expansion component (R3 mentioned in the section entitled "EXPANSION EQUATIONS") is late stage expansion and causes the acceleration observed [8]. Using cellular cosmology the potential energy can be calculated for each expansion component (R1) and (R3). The calculation shows that the potential energy required for expansion component R3 is 1.5e-11 MeV. It is very low because late stage expansion is resisted by small forces. (Stated the other way, when the radius is low forces are high but delta radius is low). The value 1.5e-11 MeV is a small portion of 10.11 MeV and is negligible. One can make a strong case against large amounts of dark energy based on these results.



The traditional [6] derivation  $(V/R)^2 = 8/3 \cdot \pi \cdot G \cdot \rho C$  where  $\rho C$  is critical density is based on initial kinetic energy being converted to potential energy as expansion occurs and the assumption that density characterizes the energy. The equation  $H = V/R = (4/3 \cdot \pi \cdot G \cdot \rho C)^{0.5}$  incorrectly assumes that the second component of expansion consumes a large amount of kinetic energy. Critical density  $9.5 \cdot 10^{-27} \text{ kg/m}^3$  should not be used to calculate cosmological parameters for the second component. If we remove dark energy from critical density [12] the lowered density is  $9.5 \cdot 10^{-27} \text{ kg/m}^3 \cdot (1 - 0.718) = 2.7 \cdot 10^{-27} \text{ kg/m}^3$ . Total mass/m<sup>3</sup> at the end of the author's expansion curves  $= 1.67 \cdot 10^{-27} \cdot \exp(180) / (\text{Volume}) = 2.47 \cdot 10^{-27} \text{ kg/m}^3$  with  $\text{Volume} = 4/3 \cdot \pi \cdot 6.2 \cdot 10^{25}^3$ . Based on only 10.11 MeV of kinetic energy being available for expansion, the proton density is one half the lowered critical density value and cold dark matter is the other one half. All energy and mass appear to be accounted for. Implications are further discussed in reference 13.

## BLACK HOLES AND QUANTUM GRAVITY

Black holes are interesting to the scientific community. Observations indicate that remnants of stars that become black holes range from about 4 solar masses through about 20 solar masses (Wiki). A supermassive black hole with mass  $4.2 \cdot 10^{40} \text{ Kg}$  has also been observed. Astronomers use x-ray emission and the bending of light around massive objects to study them and observations to date challenge physicists to extend their knowledge.

A layer by layer model of pressure, temperature, density and fusion power (MeV/sec) of the sun was developed and compared to a supermassive black hole with similar density and fusion kinetics. To simulate smaller black holes, the author developed a procedure to estimate the pressure and density as a function of radius. Conditions at the core of small black holes are

extreme enough to cause degenerate (relativistic) behavior of know forces. It was unknown however, whether collapse actually occurs. The author uses a cellular model of gravity, space, time, expansion, kinetic and potential energy at the quantum level. Quantum gravity indicates that black holes do not involve a singularity. Kinetic energy inside cells resists the pressure at the core of black holes similar to the way electromagnetic kinetic energy controls the thermodynamics of gases.

### The Schwarzschild radius

Schwarzschild solved the metric equations for general relativity. His solution defining the radius S of a black hole was  $S=2 \cdot G \cdot M / C^2$  where:  $G=6.67e-11 \text{ N m}^2/\text{kg}^2$ , M is the mass of the central body and C is the speed of light. S' is half the S value defined and agrees with the geodesic radius. The geodesic is defined by the gravitational constant  $G=S' \cdot C^2 / M$ . In cellular cosmology a cell is the volume surrounding a single proton. The table below compares three black holes and calculates critical parameters for comparison.


<b><math>S=2GM/c^2=1.48e-27 \cdot m</math></b>					
<b>My Geodesic at C and High M</b>					
<b><math>S=2 \cdot \text{Geodesic}</math></b>					
			<b>Minimum</b>	<b>20x solar</b>	<b>smassive</b>
<b>G</b>	<b>nt m<sup>2</sup>/kg'</b>	<b>6.674E-11</b>			
<b>C</b>	<b>m/sec</b>	<b>299792458</b>	<b>Kg</b>	<b>9.94E+30</b>	<b>4.00E+31</b>
<b>S'</b>	<b>=MC<sup>2</sup>/2</b>		<b>S' meters</b>	<b>7.38E+03</b>	<b>2.97E+04</b>
<b>Density</b>	<b>M/vol</b>		<b>kg/m<sup>3</sup></b>	<b>5.90E+18</b>	<b>3.64E+17</b>
			<b>N cells</b>	<b>5.95E+57</b>	<b>2.40E+58</b>
			<b>Vol cell (m<sup>3</sup>)</b>	<b>2.83E-46</b>	<b>4.58E-45</b>
<b>Average Cell Radius</b>			<b>Cell radius (n</b>	<b>4.087E-16</b>	<b>1.034E-15</b>
<b>S'/m=G/C<sup>2</sup></b>	<b>7.426E-28</b>				<b>1.067E-09</b>
<b>proton m (Kg)</b>	<b>1.67E-27</b>				

Using S' to determine the radius of a black hole (BH), the average density, cell radius, cell kinetic energy and temperature can be determined for a series of BH's with increasing mass. Volume BH= $4/3 \cdot \pi \cdot S'^3$ . Volume cell= (Volume BH/N cells) with N cells=M black hole/1.67e-27 kg. This gives the critical parameter discussed below: cell radius = (Volume cell/(4/3\*pi))^(1/3). Radius can also be calculated from density rho assuming hydrogen. Cell radius= (1/rho\*1.67e-27/(4/3\*pi))^(1/3). Cell radius is very low for black holes, but larger for supermassive black holes.


### Approximate Solar Model


An approximate model of the pressure and temperature of the sun as a function of radius is shown below. The reason for this exercise is to develop a procedure for analyzing black holes.

Since data exists for the sun, we can “tune-up” the model before it is used for black holes. It is based on breaking the sun into 10 radial shells each 1/10 of the full radius.



The mass above a shell surface is gravitationally attracted to the mass below the shell yielding acceleration  $a=G*M_{\text{central}}/R^2$ . At the bottom of each shell, the pressure is calculated as  $P_{\text{shell}}=\text{acceleration}*\text{density}*dR$ . Pressure adds down through the 10 shells until a central pressure is calculated. Because the sun is about 4B years old, most of its early history has dissipated and temperature increases from the outside to the core due to nuclear power in the core. A linear relationship was assumed even though the radiation and convection zones conduct heat differently. The surface temperature is known ( $6e3$  K) and the interior temperature is approximately  $1.7e7$  K. Density of each shell is determined by the equation  $\rho=P/(R*T)$ , where  $R$  is the gas constant for hydrogen  $7980$  N-m/(Kg-K). Our sun has an interior density of about  $1.2e5$  N/m<sup>2</sup> and the temperature and pressure profiles below produce this density in the interior. Here are plots for the model of the  $2e30$  Kg sun.





The core of the 10 shell model is in the chart below along with the equations. All 10 layers are in Appendix 3. The top of the core ( $h_9$ ) is at radius  $6.92e7$  meters and  $R$  is the sun's overall radius  $5.9e8$  meters.

<b>P9 (nt/m<sup>2</sup>)</b>		<b><math>P_9 = P_8 + \rho_8 \cdot a_8 \cdot R/10</math></b>	<b>1.41E+15</b>
<b>rho9 kg/m<sup>3</sup></b>		<b><math>\rho_9 = P_9 / (R \cdot T_8)</math></b>	<b>3.946E+04</b>
<b>h9 (m)</b>		<b><math>h_9 = h_8 - R/10</math></b>	<b>6.92E+07</b>
<b>T (K)</b>		<b><math>T_9 = T_8 + dT</math></b>	<b>2.160E+07</b>
<b>a10 (m/sec<sup>2</sup>)</b>		<b><math>a_9 = 6.673e-11 \cdot (M - \text{cum}m_{10}) / (h_9)^2</math></b>	<b>7.90E+02</b>
<b>m10 shell Kg</b>		<b><math>m_{10} = 4/3 \cdot \pi \cdot (h_9^3 - (h_9 - R/10)^3) \cdot \rho_9</math></b>	<b>1.94E+27</b>
<b>cum mass10</b>		<b><math>\text{cum}m_{10} = \text{cum}m_9 + m_{10}</math></b>	<b>1.943E+30</b>
<b>P10 (nt/m<sup>2</sup>)</b>		<b><math>P_{10} = P_9 + \rho_9 \cdot a_9 \cdot R/10</math></b>	<b>3.57E+15</b>
<b>rho10 kg/m<sup>3</sup></b>		<b><math>\rho_{10} = P_{10} / (R \cdot T_9)</math></b>	<b>8.869E+04</b>
<b>h10 core</b>		<b><math>h_{10} = h_9 - R/10</math></b>	<b>0</b>
<b>r cell (m)</b>		<b><math>r_{\text{cell}} = (1 / \rho_{10} \cdot 1.67E-27 / (4/3 \cdot \pi))^{1/3}</math></b>	<b>1.65E-11</b>

Again, the cell radius is the radius around each proton and is a function of density. The sun is dense enough at the core ( $8.9e6$  kg/m<sup>3</sup> above) to make the electron degenerate because the cell radius is lower than the electron radius  $5.29e-11$  meters. The author uses a probability based model [14] [15] for fusion diagramed in Appendix 3. The following table gives the fusion kinetics based on the pressure and temperature model above:



<b>2e30 Kg Solar Model</b>		
	<b>Max Dens</b>	<b>5.0E+11</b>
<b>Temp</b>	<b>deg K</b>	<b>2.0E+07</b>
<b>Density</b>	<b>kg/m<sup>3</sup></b>	<b>1.2E+05</b>
<b>KE</b>	<b>1.5*B*T</b>	<b>2.6E-03</b>
<b>degeneracy</b>		<b>3.6E+00</b>
<b>Degenerate radius (DR)</b>		<b>1.5E-11</b>
<b>v/c</b>		<b>1.0E-01</b>
<b>Barrier</b>		<b>-1.4E-02</b>
<b>P barrier</b>		<b>4.5E-03</b>
<b>Pdensity</b>		<b>2.4E-07</b>
<b>Rate/sec</b>		<b>2.8E-09</b>
<b>Probability/sec</b>		<b>3.1E-18</b>
<b>burn time (Byrs)</b>		<b>1.0E+01</b>
<b>sun N</b>		<b>1.2E+57</b>
<b>fract burning</b>		<b>1.0E-01</b>
<b>burn rate N/sec*mev/N</b>		<b>3.7E+38</b>

It has a burn time of about 10 Billion years and generates about 4e38 MeV/sec.

#### **Analysis of black hole with 4.7 solar mass (9.9e30 Kg)**

A black hole (BH) with 4.7 solar masses is quite different than the sun. To achieve BH status, the Schwarzschild radius  $S' = G \cdot M / C^2$  for this 9.9e30 Kg black hole was 7800 m, not 6.9e8 meters like the sun. This BH has a density of approximately 1.8e19 kg/m<sup>3</sup> and much higher temperature and pressure. Power generation in the sun increases the core temperature and lowers its density. But BH's of this size are burned out. The lack of nuclear power allowed the density to increase and it became a black hole.


If we insist that  $C$  is not exceeded, the acceleration in each shell is  $a = C^2/R$ , where  $R$  is radius to the shell. This means  $R$  is the Schwarzschild radius  $S'$  for each layer. In other words, each layer of the model is like the surface of the black hole and the velocity is  $C$ , i.e.  $(a \cdot S')^{0.5} = C$ . Consider the alternatives. If we let  $a = C^2/R$  be less than  $C$  for each layer, the mass will not fit inside  $S'$  and it won't be a black hole. The other alternative is to let velocity exceed  $C$ . The speed of light,  $C$  is just the ratio between space and time and can't be violated. In special relativity gravity is the geometry of space-time and gravitational force is inertial force. The net velocity of particles inside the BH is too low (near zero since they are moving in all directions) to be on a geodesic defined by the radius of the layer. We can calculate  $V$  to be on the geodesic. Inside a black hole,  $V = C = (a \cdot S')^{0.5}$  and  $V_{low} = \text{zero}$ ; fixed by the fact that particles are inside the black hole. The equation becomes  $F = \text{mass} \cdot (C^2 - V_{low}^2) / S'$ . The force is large and associated with velocity  $C$  because that would put the particle on a geodesic. The particles in the BH have no net velocity that would give them inertial force. But they do have temperature because the gravitational kinetic energy is trapped inside the cell.

## Compression thermodynamics

For gases at low pressure, the electron is circling the nucleus at a radius of approximately  $5.29 \times 10^{-11}$  meters. Electromagnetic kinetic energy  $13.6 \times 10^{-6}$  MeV is associated with temperature and pressure that resist compression. The origin of gas thermodynamics for hydrogen is shown below. The form is once again a column of calculations with inputs listed toward the top, calculation of relativistic effects further down and finally a calculation for radius  $r$  with the R equation.

<b>0.511 Electron</b>	<b>mass MeV</b>
<b>9.11E-31</b> $.511 * 1.78 \times 10^{-30}$ kg/MeV	<b>mass kg</b>
<b>2.72E-05</b> E/M field	<b>Field MeV</b>
<b>1.36E-05</b> E/M field/2	<b>Kinetic Energy MeV</b>
<b>1.00E+00</b> $0.511 / (0.511 + ke)$	<b>gamma</b>
<b>7.30E-03</b> $V/C = (1 - \gamma^{-2})^{0.5}$	<b>V/C</b>
<b>5.291E-11</b> $r$	<b>Radius (meters)</b>
<b>8.24E-08</b> $F = (9.11 \times 10^{-31} / g) * (v/C)^2 / R$	<b>Force Nt</b>
<b>3.52E-20</b> $area = 4 * \pi * r^2$	<b>Surface area <math>m^2</math></b>
<b>2.34E+12</b> $P = F / area$	<b>Pressure nt/<math>m^2</math></b>
<b>1.05E+05</b> $T = ke / 1.5 / 8.6 \times 10^{-11}$	<b>Temperature based on ke (K)</b>
<b>2.69E+03</b> $\rho = (1 / r^3 * 1.67 \times 10^{-27} / (4/3 * F$	<b>Density = <math>P / (RT)</math></b>
<b>8269.77</b> $R = P / (\rho * T)$ <b>Nt-K/(m-kg)</b>	<b>Gas constant</b>

In the model above  $\rho$  is calculated from radius  $r$  and the gas constant is derived from  $R = P / (\rho * T)$ . Pressure  $P$  is calculated from force/area of the electron orbit. The low pressure gas constant derived agrees with literature. As radius  $r$  decreases with compression of the gas,  $ke = 13.6 \times 10^{-6} * 5.29 \times 10^{-11} / \text{radius}$  MeV and this increases the pressure and temperature. When the radius decreases to the point where compression kinetic energy = 0.11 MeV the electron is no longer sustained in the shell and transition of the proton to a neutron occurs (the Proton model in Appendix 2 justifies the use of 0.11 MeV. (Also read Wiki concerning White Dwarfs and Neutron Stars). For neutrons at high pressures gravitational kinetic energy (10.11 MeV) controls pressure and a chart similar to the one above is required. It is developed below under the heading "Pressure inside cells prevents black hole collapse" but results are shown below. The red line is for neutrons and the blue line to the left of 0.11 MeV is for a gas. The resistance pressure of a gas and neutrons differ by at least three orders of magnitude for the same radius.




We concluded above that pressure expands the universe and it was shown that 10.11 MeV of kinetic energy had been converted to potential energy integral  $Pdv = 10.11 \text{ MeV}$  for the cells that expanded fully (small  $v$  stands for volume). The kinetic energy after expansion is very low and  $r$  for the cell diagram above is 0.54 meters. This gives the current radius of the universe  $= 0.54 * \exp(60) = 6.2e25$  meters. Further, reference 13 shows that kinetic KE in the cell diagram can be considered temperature. Using 10.11 MeV as the beginning energy and temperature  $T = ke/1.5/B$  (and adjusting it upward for primordial He4 fusion), the kinetic energy is now associated with the cosmic microwave background temperature (CMB)  $= 2.73 \text{ K}$ . Using  $KE = 1.5 * B * T$  ( $B$  is Boltzmann's constant  $8.6e-11 \text{ MeV/K}$ ); the kinetic energy is now  $3.5e-10 \text{ MeV}$ . Knowing that KE in the cell diagram is temperature and that there is pressure inside each cell are key understandings used below. We will now explore what happens when expansion is thrown into reverse gear and the radius is forced to decrease. The value 10.11 MeV was associated with radius  $7e-14$  meters but the black hole comparison table above shows that their cellular radius can be two orders of magnitude lower than this.

### Pressure inside cells prevents black hole collapse

At extremely high pressure, the proton plus the electron (and a neutrino) become a neutron. Neutrons have no charge and degenerate electrons no longer exist. It is believed by some that strong forces and the Pauli exclusion principle prevents collapse. Others believe that collapse occurs and a singularity forms. The neutron has gravitational kinetic energy and velocity but it is disorganized into temperature and pressure. It will be shown below that the cell internal pressure and temperature become high enough to resist collapse of a black hole. The black hole consists of many cells and there is no place for the 10.11 MeV/neutron of energy to go when compression of the cell squeezes it below  $7.22e-14$  meters. (Recall that the gravitational cellular space is the *only* space outside the neutron). The circle diagram is repeated below that was labeled "you are here". Thankfully your cells weren't among the cells that fell into the black hole.

## Three Nested Orbits for the Neutron



As the radius becomes small, the kinetic energy of the neutrons increases the pressure in the cell. The neutrons bounce off of each other because of Pauli exclusion. The neutron has only random velocity it has net velocity zero, i.e.  $V_n=0$  and is not on the geodesic. In fact the required geodesic velocity approaches  $C$  if the neutrons fall into a black hole. Force  $F=m*C^2/r-m*V_n^2/r$  is very large because  $C$  is large,  $r$  is small and  $V$  is zero. The pressure in the cells must exert outward force to make up for the inertial force that the neutrons need to be on the geodesic.

When cell radius  $r$  becomes less than  $6.5e-15$  meters the electromagnetic force is overwhelmed (see graph above in the section entitled "Compression thermodynamics") and the resisting pressure is based on cell pressure. Neutrons collide and the change in momentum is force. Pressure is determined by dividing the force by the cell surface area. Originally the star that collapsed into a black hole was hot and when the nuclear fuel was expended, it contracted creating compression energy. The gravitational model for the core pressure below is based on setting  $(a*S')^{0.5}=C$ . The equations shown are the same for each layer.  $P$  stands for pressure,  $\rho$  stands for density,  $h$  stands for the height from the center of the BH,  $T$  stands for temperature,  $a$  stands for acceleration,  $m$  stands for mass, cum mass is the mass above the layer and  $S$  stands for the radius of the BH. Layers are labelled 1 through 10, with 10 being the core. Part of layer 9 and all of layer 10 (the core) are shown below. The full model results are in the appendix 3 but the radius of a cell at the core of a  $9.94e30$  Kg black hole is  $2.93e-16$  meters.

<b>P9 (nt/m<sup>2</sup>)</b>		<b>P9=P8+rho8*a8*h0/10</b>	<b>8.82E+35</b>
<b>rho9 kg/m<sup>3</sup></b>		<b>rho9=1.67E-27/(4/3*PI()*(((4.9e-25)/(P9)))</b>	<b>9.42E+18</b>
<b>h9 (m)</b>		<b>h9=h8-h0/10</b>	<b>7.38E+02</b>
<b>T (K)</b>		<b>T9=(9.7*7.34e-14/rcell)/1.5/8.6e-11</b>	<b>1.58E+13</b>
<b>a10 (m/sec<sup>2</sup>)</b>		<b>a9=C<sup>2</sup>/h9</b>	<b>1.22E+14</b>
<b>m10 shell Kg</b>		<b>ms10=4/3*PI()*((h9<sup>3</sup>-(h9-S/10)<sup>3</sup>)*rho9</b>	<b>1.59E+28</b>
<b>cum mass10</b>		<b>cumm10=a9*h9<sup>2</sup>/G</b>	<b>9.96E+29</b>
<b>P10 (nt/m<sup>2</sup>)</b>		<b>P10=P9+rho9*a9*h0/10</b>	<b>1.73E+36</b>
<b>rho10 kg/m<sup>3</sup></b>		<b>rho10=1.67E-27/(4/3*PI()*(((4.9e-25)/(P10)))</b>	<b>1.58E+19</b>
<b>h10 core</b>			<b>0</b>
<b>r cell (m)</b>	<b>rcell=(1/rho10*1.67E-27/(4/3*PI()))^(1/3)</b>		<b>2.93E-16</b>
		<b>C=(a*R9)^.5</b>	<b>3.00E+08</b>
			<b>smallest BH</b>
			<b>proton</b>
	<b>mass MeV</b>	<b>Proton</b>	<b>938.2720</b>
	<b>mass kg</b>	<b>938.27*1.78e-30 kg</b>	<b>1.673E-27</b>
	<b>Field MeV</b>	<b>(See Proton Model)</b>	<b>2.73E+00</b>
	<b>ke</b>	<b>KE=10.11*7.22e-14/</b>	<b>2489.197</b>
	<b>g=m/(m+ke)</b>	<b>g=938.27/(938.27+k</b>	<b>0.2738</b>
	<b>V/C</b>	<b>V/C=(1-(g)^2)^0.5</b>	<b>0.9618</b>
	<b>rcell (meters)</b>	<b>r</b>	<b>2.932E-16</b>
	<b>F=m*V<sup>2</sup>/R/exp(90)</b>	<b>F=(1.67e-27/g)*(v/C)</b>	<b>1.420E-33</b>
			<b>1</b>
	<b>area=4*pi*R<sup>2</sup></b>	<b>area=4*pi*r<sup>2</sup></b>	<b>1.081E-30</b>
	<b>Pressure (nt/m<sup>2</sup>)</b>	<b>P=F/area</b>	<b>1.603E+36</b>
	<b>Temperature (K)</b>	<b>T=ke/1.5/8.6e-11</b>	<b>2.4404E+13</b>
	<b>rho</b>	<b>rho=(1/r<sup>3</sup>*1.67E-27</b>	<b>1.58E+19</b>
	<b>"gas constant" R</b>	<b>R=P/(rhoT) Nt-K/(m-</b>	<b>4.16E+03</b>
		<b>r=(1/rho*1.67E-27/(4</b>	<b>2.93E-16</b>

Note: the equation for density is developed in reference 27. The important calculation for pressure is the force/area, where area is the surface area of the layer.

Recognize the bottom part of the above diagram as the quantum gravity calculation except radius has been reduced by compression in the core of the black hole. It follows the same form as the basis of low pressure gas thermodynamics presented above in the section entitled "Compression thermodynamics". Since cellular quantum gravity kinetic energy varies with radius, the kinetic

energy has increased to  $10.11 \cdot 7.22e-14 / 2.93e-16 = 2489$  MeV and this is associated with compression temperature of  $2492 / 1.5 / B = 2.5e13$ . Pressure in the cell is calculated by the change in momentum of neutrons colliding. This is equivalent to inertial force/cell surface area. The outward pressure  $1.6e36$  N/m<sup>2</sup> exerted by the cell resists the pressure near the center of the black hole ( $1.7e36$  N/m<sup>2</sup>) and prevents it from collapsing. The energy is a natural outcome of the pressure and temperature resulting from compression and does not need to be associated with degeneracy. Pressure and density are intimately linked. The density increases until the pressure is resisted similar to the way gases resist compression. This allowed the author to develop the equation below for rho. It is based on solving the gravity equation for cell radius and determining rho from cell radius. It is based on known equations but fits a computationally friendly relationship within 8% over small black hole pressures investigated.

$$\rho = 1.67E-27 / (4/3 * \pi * ((4.8E-25/P)^{0.2565})^3)$$

### More massive black holes



Stars that complete their aging cycle can be about 20 solar masses but not much greater according to Wiki. The analysis above was repeated for this size black hole.

<b>P9 (nt/m<sup>2</sup>)</b>		<b>P9=P8+rho8*a8*h0/10</b>	<b>7.47E+33</b>
<b>rho9 kg/m<sup>3</sup></b>		<b>rho9=1.67E-27/(4/3*PI()*(((4.9e-25)/(P9)))</b>	<b>2.39E+17</b>
<b>h9 (m)</b>		<b>h9=h8-h0/10</b>	<b>4.22E+04</b>
<b>T (K)</b>		<b>T9=(9.7*7.34e-14/rcell)/1.5/8.6e-11</b>	<b>4.66E+12</b>
<b>a10 (m/sec<sup>2</sup>)</b>		<b>a9=C<sup>2</sup>/h9</b>	<b>2.13E+12</b>
<b>m10 shell Kg</b>		<b>ms10=4/3*PI()* (h9<sup>3</sup>-(h9-S/10)<sup>3</sup>)*rho9</b>	<b>1.48E+31</b>
<b>cum mass10</b>		<b>cumm10=a9*h9<sup>2</sup>/G</b>	<b>5.70E+31</b>
<b>P10 (nt/m<sup>2</sup>)</b>		<b>P10=P9+rho9*a9*h0/10</b>	<b>8.98E+33</b>
<b>rho10 kg/m<sup>3</sup></b>		<b>rho10=1.67E-27/(4/3*PI()*(((4.9e-25)/(P1</b>	<b>2.76E+17</b>
<b>h10 core</b>			<b>0</b>
<b>r cell (m)</b>	<b>rcell=(1/rho10*1.67E-27/(4/3*PI()))^(1/3)</b>		<b>1.13E-15</b>
		<b>C=(a*R9)<sup>.5</sup></b>	<b>3.00E+08</b>
			<b>20x sol BH</b>
			<b>proton</b>
	<b>mass MeV</b>	<b>Proton</b>	<b>9.38E+02</b>
	<b>mass kg</b>	<b>938.27*1.78e-30 kg</b>	<b>1.67E-27</b>
	<b>Field MeV</b>	<b>(See Proton Model)</b>	<b>2.73E+00</b>
	<b>ke</b>	<b>KE=10.11*7.22e-14/</b>	<b>645.775</b>
	<b>g=m/(m+ke)</b>	<b>g=938.27/(938.27+k</b>	<b>5.92E-01</b>
	<b>V/C</b>	<b>V/C=(1-(g)<sup>2</sup>)<sup>0.5</sup></b>	<b>8.06E-01</b>
	<b>rcell (meters)</b>	<b>r</b>	<b>1.130E-15</b>
	<b>F=m*V<sup>2</sup>/R/exp(90)</b>	<b>F=(1.67e-27/g)*(v/C)</b>	<b>1.19E-34</b>
			<b>1.00E+00</b>
	<b>area=4*pi*R<sup>2</sup></b>	<b>area=4*pi*r<sup>2</sup></b>	<b>1.61E-29</b>
	<b>Pressure (nt/m<sup>2</sup>)</b>	<b>P=F/area</b>	<b>9.08E+33</b>
	<b>Temperature (K)</b>	<b>T=ke/1.5/8.6e-11</b>	<b>6.3311E+12</b>
	<b>rho</b>	<b>rho=(1/r<sup>3</sup>*1.67E-27</b>	<b>2.76E+17</b>
	<b>"gas constant" R</b>	<b>R=P/(rhoT) Nt-K/(m-</b>	<b>5.19E+03</b>
		<b>r=(1/rho*1.67E-27/(4</b>	<b>1.13E-15</b>

## Supermassive Black Holes

Supermassive black holes are more similar to our sun than small mass black holes because their density is low. In fact, analysis shows that supermassive black holes have not burned out and their fusion kinetics are similar to the sun. The author is aware that there are short comings to applying the solar model to a black hole. Nevertheless, it is worth showing results even if the model is inaccurate because it reveals large differences involved. First, we will apply the model

to a BH of  $4.2 \times 10^{40}$  Kg. Here is a plot of the pressure and temperature for a BH of  $4.2 \times 10^{40}$  Kg with a horizon  $3.1 \times 10^{13}$  meters. The average density of the BH is quite low, only  $6 \times 10^{-2}$  Kg/m<sup>3</sup>. The pressure at the core is about  $6 \times 10^{17}$  N/m<sup>2</sup>. The thermodynamics are controlled by electromagnetic kinetic energy since the pressure is well below the electromagnetic to gravity transition  $1 \times 10^{28}$  N/m<sup>2</sup>.



The hydrogen fusion kinetics is estimated below:



		<b>largest BH</b>
<b>4.2e40 Kg Solar Model</b>		
	<b>Max Dens</b>	<b>5.020E+11</b>
<b>Temp</b>	<b>deg K</b>	<b>6.00E+10</b>
<b>Density</b>	<b>kg/m^3</b>	<b>3.31E+01</b>
<b>KE</b>	<b>1.5*B*T</b>	<b>7.76E+00</b>
<b>degeneracy</b>		<b>2.31E-01</b>
<b>Degenerate radius (DR)</b>		<b>5.290E-11</b>
<b>v/c</b>		<b>0.998</b>
<b>Barrier</b>		<b>-0.0139</b>
<b>P barrier</b>		<b>9.98E-01</b>
<b>Pdensity</b>		<b>6.60E-11</b>
<b>Rate/sec</b>		<b>7.888E-09</b>
<b>Probability/sec</b>		<b>5.19E-19</b>
<b>burn time (Byrs)</b>		<b>61</b>
<b>N=m/1.67e-27</b>		<b>5.988E+67</b>
<b>fract burning</b>		<b>0.15</b>
<b>burn rate N/sec</b>		<b>4.67E+48</b>

The 4.2e40 Kg BH is burning so slowly that it could last 60 billion years. Its heat generation can be compared to our sun (1e39 MeV/sec) but is 2.1e10 more massive. One difference from a star is that heat may not escape as readily. This would increase its overall temperature and lower its density. Density can't be lowered much or the mass will not fit within the Schwarzschild radius. The pressure profile depends on the mass and again can't change much. One might ask if the contained mass becomes a "pressure cooker" since nuclear energy cannot easily escape. According to the proton mass model (appendix 3), there is only 10.15 MeV/particle of nuclear energy available per particle. The effect of burning this fuel and containing it in the BH is shown below.

<b>Nuclear energy release 10 MeV</b>	
<b>T core=6e10K</b>	
<b>dQ=10 MeV</b>	
<b>dt=10/(1.5 B)</b>	
<b>7.73E+10 K</b>	
<b>T core with dQ=6e10+7.8e10</b>	
<b>1.37E+11 K</b>	
<b>P=7829 T rho</b>	
<b>Pressure may double</b>	

The pressure may double if all the nuclear energy per particle is released. However pressure is reasonably low and easily contained.

## **The BH surface**

What are the conditions at the surface of the black hole? As new mass enters the black hole and collides with the spherical surface it carries about 10.11 MeV of kinetic energy ( $7.6 \times 10^8$  K) with it as a result of falling toward the black hole at high velocity. There is a lot of photon energy produced by a hot surface. One question might be: Where does the photon energy go? The Schwarzschild equation tells us that photons are attracted to the large gravitational pull of the BH and photons travel in curved paths around the BH horizon. They travel at the speed of light but cannot readily escape (read Hawking regarding BH evaporation). This means that at the horizon there is a large flux of photons. This flux probably photo-disintegrates bonds (electromagnetic and nuclear) of atoms that cross the horizon.

## **Summary Black Holes and Quantum Gravity**

Black holes that range from 4.7 to 20 solar masses involve high core pressures. The author addressed the question “What keeps these black holes from collapsing?” Based on the author’s theory of quantum gravity the answer to this question is “pressure and temperature inside the cell”. Thermodynamic pressure in low pressure gases is based on compression of the electron radius. The density in the gas adjusts in such a way to resist compression. When a radius is compressed beyond  $4.6 \times 10^{-15}$  meters, the electron is overwhelmed since the kinetic energy involved converts a proton back into a neutron. Black holes involve radii as low as  $2.93 \times 10^{-16}$  meters. The model for gravitation kinetic energy resisted pressure is similar to standard gas thermodynamic pressure except the gravitational kinetic energy is 10.11 MeV, not the electromagnetic kinetic energy  $13.6 \times 10^{-6}$  MeV. The transition between these two zones is about  $1 \times 10^{28}$  N/m<sup>2</sup> for a gas but a small BH is a hot soup of neutrons. The equation given in Wiki for density for dense objects like White Dwarfs can’t be used for black holes.


Supermassive black holes as large as  $4.2 \times 10^6$  solar masses have been observed. These black holes involve temperature and pressures in their cores that strongly suggest nuclear energy generation. However, analysis shows that they burn very slowly and can contain the power generated. Their pressure is low and well within the electromagnetic controlled thermodynamics zone.

## **Explanation of dark field compact galaxies**

Dark field Hubble photographs show that highly redshifted galaxies appear more compact. They are far away and could have formed earlier than other galaxies. At a particular time in expansion, expanded cells have a certain amount of potential and kinetic energy. The section entitled “Gravitational accumulation”, above, indicated that particles achieve an orbit at 42% their original height. The larger the universe is, the higher the particles will be when they start their fall. This means that galaxies formed late in expansion will appear larger than ones formed early.

We can estimate gravitational accumulation using the spreadsheet fallmodel.xls. Study of a model galaxy ( $1.2 \times 10^{42}$  kg) allows us to determine where energy resides as it is re-converted to kinetic energy by falling. Some particles leave the radius defined by the expansion equation and fall into orbits around a gravitational center that will become a galaxy. In the graph below, the later galaxies form the lower their star velocities will be. Since they have lower velocities they

will orbit at a larger distance from the center. These galaxies will appear less compact. The sun is about 5 billion years old and is half way through its hydrogen fuel.



## CONCLUSIONS

The gravitational constant  $G$  can be calculated from a new concept called cellular cosmology and information from an energy model of the neutron. The quantum gravitational scale is the radius  $7.22 \times 10^{-14}$  meters and the neutron mass,  $1.67 \times 10^{-27}$  Kg with a coupling constant  $1/\exp(90)$ . The neutron model contains four field energies and associated masses consistent with data regarding nature's four fundamental interactions. In each case a field curves space and a mass falls into an orbit with kinetic energy, defining the quantum radius. This indicates that general relativity continues down to the quantum level but in cellular cosmology, a field curves space and there is initially one neutron in each cell. The large scale space-time we walk around in is defined by gravity at the quantum scale, the gravitational coupling constant and expansion. During expansion, rather than being limited to a quantum mechanical orbit, protons are free to move throughout space. This means that although the proton model defines kinetic energy in a gravitational orbit, it is pressure and temperature that expands the universe. After two early transitions (equality of photon and mass density and decoupling of electrons), gravitation is locally able to dominate gas pressure. This gas acts according to thermodynamics but the particles are gravitationally "sticky" and small accumulations of matter grow and eventually form clusters, galaxies, stars and planets.

The first law of thermodynamics deals with groups of particles interacting through electromagnetic energy and in extreme cases with gravitational kinetic energy. The top of the "one-way downward street" we call entropy is the expanded universe. As accumulation begins there are many potential states for particles to fall into. As they falls collide and produce heat, the second law of thermodynamics describes their behavior.

Elapsed time enters physics through cosmology. If we believe  $(r/r_0)^3$  increases we must believe that time advances  $(t/\alpha)^2$  ( $\alpha$  is a constant). It appears that time cycles at the quantum level and counts forward for all particles. The cycle time for one count is the fundamental time defined by quantum gravity. Elapsed time, expansion equations and

conversion of potential energy to kinetic energy define the space around us. Free particles have velocity and an associated gamma.

Black holes that range from 4.7 to 20 solar masses involve high core pressures. The author addressed the question “What keeps black holes from collapsing?” Based on quantum gravity the answer to this question is “pressure and temperature originating from 10.11 MeV of gravitational kinetic energy resists compression”. Thermodynamic pressure in low pressure gases is based on compression of the electron radius. The density in the gas adjusts in such a way to resist compression. When a radius is compressed beyond  $4.6e-15$  meters, the electron is overwhelmed since the kinetic energy involved converts a proton back into a neutron. Black holes involve radii as low as  $2.93e-16$  meters. Cellular pressure is similar to standard gas thermodynamic pressure except the gravitational kinetic energy is 10.11 MeV, not the electromagnetic kinetic energy  $13.6e-6$  MeV. The transition pressure between these two zones is about  $1e28$  N/m<sup>2</sup>.

Supermassive black holes as large as  $4.2e40$  Kg have been observed. These black holes involve temperature and pressures in their cores that strongly suggest nuclear energy generation. However, analysis shows that they burn very slowly and can contain the power generated. Their pressure is low and well within the electromagnetic controlled thermodynamics zone.

What about the argument that velocity is relative? Each proton in nature has a specific energy and can be associated with a cell. Expansion potential energy can be re-converted to compression energy, orbital kinetic energy and heat. We can account for energy and find out interesting things about our history using cellular cosmology. The author believes that nature maintains zero energy, made of two opposite energy values. Since particles interact some protons contain more energy than others but they average to zero. It is well known that the velocity light is constant and that time varies slightly. Equations related to the Lorentz transformation should be considered “constant C equations” rather than relativity equations. It can be misleading to compare one velocity with another because we do not have enough information to know where all the energy resides. It is the author’s view that velocity (and associated kinetic energy) is not relative.

Cold dark matter exists and makes up about half the mass of the universe. Most cosmologists use critical density as a predictor of energy components but it does not match calculations of potential and energy conservation. Dark energy exists but is almost negligible according to these calculations. The author believes that all energy is accounted for.

## SELECTED REFERENCES

1. Brillouin, L., *Science and Information Theory*, New York: Academic Press, Inc., 1956.
2. Feynman, R., Leighton, R., Sands, M., *the Feynman Lectures on Physics*, California Institute of Technology, Addison-Wesley Publishing Company, 1965.
3. Glashow, S., *Interactions*, Warner Book, Inc., 1988.
4. Hughes, I. S., *Elementary particles Third Edition*, Cambridge University Press, 1991

5. Weinberg, S, *A Unified Physics by 2050?*, Scientific American End of the Millennium Special Issue, December 1999.
6. P.J.E. Peebles, *Principles of Physical Cosmology*, Princeton University Press, 1993.
7. Bennett, C.L. et al. *First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Preliminary Maps and Basic Data*, Astrophysical Journal, 2001 L. Verde, H. V. Perris, D. N. Spergel, M. R.olta, C. L. Bennett, M. Halpern, G. Hinshaw, N. Jarosik, A. Kogut, M. Limon, S. S. Meyer, L. Page6, G. S. Tucker, E. Wollack, E. L. Wright
8. Conley, et al. (*THE SUPERNOVA COSMOLOGY PROJECT*), *Measurement of Omega mass and Omega lambda from a blind analysis of TypeIa supernovae with CMAGIC*.
9. <http://arxiv.org/pdf/1212.5226v3.pdf> Table 2.
10. Barbee, Gene H., *A Top-Down Approach to Fundamental Interactions*, FQXi essay, June 2012 and vixra:1307.0082 revised Mar 2014. Reference Microsoft ® spreadsheet Unifying concepts of nature.xls.
11. Barbee, Gene H., *On the Source of the Gravitational Constant at the Low Energy Scale*, vixra:1307.0085, revised Feb 2014. Prespacetime Journal Vol. 5 No. 3 March 2014.
12. Barbee, Gene H., *Application Proton Mass Model to Cosmology*, vixra:1307.0090, revised Mar 2014. Reference spreadsheet simple1c.xls.
13. Barbee, Gene H., *On Expansion Energy, Dark Energy and Missing Mass*, Prespacetime Journal Vol. 5 No. 5, May 2014. Previously vixra:1307.0089, revised Mar 2014.
14. Barbee, Gene H., *A Simple Model of Atomic Binding Energy*, vixra:1307.0102, revised Feb 2014. Reference spreadsheet atom.xls.
15. Barbee, Gene H., *Semi-Fundamental Abundance of the Elements*, vixra:1308.0009, revised Nov 2013.
16. D. E. Groom et al. (Particle Data Group). Eur. Phys. Jour. C15, (2000) (URL: <http://pdg.lbl.gov>)
17. Bergstom, L. and Goobar, A., *Cosmology and Particle Astrophysics*, 2<sup>nd</sup> Edition, Springer-Praxis Books in Astrophysics and Astronomy, 2004.
18. Barbee, Gene H., *Baryon and Meson Masses Based on Natural Frequency Components*, vixra:1307.0133.
19. Barbee, Gene H., *The Effect of He4 Fusion on Primordial Deuterium*, vixra:1404.0465, May 2014.
20. Atomic masses (for binding energy) [http://physics.nist.gov/cgi-bin/Compositions/stand\\_alone.pl?ele=&ascii=html&isotype=some](http://physics.nist.gov/cgi-bin/Compositions/stand_alone.pl?ele=&ascii=html&isotype=some) (National Institute of Standards and Technology, <http://www.nist.org>)
21. Shannon, Claude. *A mathematical Theory of Communication*, 1948.
22. <http://www.lbl.gov/abc/wallchart/chapters/04/1.html>
23. [www.FQXI.org](http://www.FQXI.org)
24. Barbee, Gene H., *Unification*, vixra:1410.0028.
25. Barbee, Gene H., *Cosmology, Thermodynamics and Time*, vixra:1407.0187, September 2014.
26. [http://en.wikipedia.org/wiki/Black\\_hole](http://en.wikipedia.org/wiki/Black_hole)
27. Barbee, Gene H., *Black Holes and Quantum Gravity*, vixra:1409.0242, September 2014.
28. [http://en.wikipedia.org/wiki/Gravitational\\_coupling\\_constant](http://en.wikipedia.org/wiki/Gravitational_coupling_constant).
29. Gordon J. Van Wylen, *Thermodynamics*, New York: John Wiley & Sons, Inc., 1959.
30. Gene H Barbee, EPR, dual slit and “back from the future” observations, Academia.edu, April 2015.

- 31. Barbee, Gene H., *Life from Information*, vixra:1311.0124v1, FQXi 2013 essay contest.
- 32. Barbee, Gene H., *Camp Four*, FQXi 2014 essay contest.
- 33. Barbee, Gene H., *Energy 0, Probability 1, Exchange 2*, FQXi essay, January 2015.

## Appendix 1 Natural Logarithms

The following “information code” was a result of correlating fundamental energy data [4][14]. I do not know why this code is used by nature but it anchors energy values. The numbers are natural logarithms. There are four sets and total 90.

<b>Mass &amp;</b>			
<b>Ke</b>		<b>Field</b>	
<b>N values</b>		<b>N values</b>	
↓		↓	
15.432	↘ set1	17.432	
12.432		10.432	
13.432	↘ set2	15.432	
12.432		10.432	
13.432	↘ set3	15.432	
12.432		12.432	
0.075	↘ set4	0.075	
10.333		10.333	
90		90	
<b>P=1/exp(90)</b>		<b>P=1/exp(90)</b>	
	<b>P=1/exp(180)</b>		


Set 2 is used in the example below. The code doesn't represent energy until after the following exchange on each set. Set 2 starts with the natural logarithms 13.43 and 12.43. The number 2 is added to 13.43 to become 15.43 and 2 is subtracted from 12.43 to become 10.43. This will be called an energy interaction and the four values of N involved in the exchange will be called a quad.

Before adding and subtracting 2				After adding and subtracting 2			
		MeV			MeV		
		E=e0*exp(N)			E=e0*exp(N)		
<b>N1</b>	<b>13.43</b>	<b>13.8</b>	<b>E1 mass</b>	<b>N3</b>	<b>15.43</b>	<b>101.95</b>	<b>E3 field</b>
<b>N2</b>	<b>12.43</b>	<b>5.1</b>	<b>E2 ke</b>	<b>N4</b>	<b>10.43</b>	<b>0.69</b>	<b>E4 field</b>

Information N is conserved (13.43+12.43=15.43+10.43). Each of the four positions has a specific meaning. N1 is always a mass, N2 is kinetic energy, N3 is a strong field and N4 is a component of the gravitational field. Energy is evaluated by the equation  $E=e_0 \cdot \exp(N)$  where  $e_0=2.025e-5$  MeV. E0 was determined from N for the electron ( $10.136=10.333-2 \cdot \ln(3/e)$ ) and its known energy 0.511 MeV.

Results of the above energy interaction				
	(difference ke)			
	E3+E4-E1-E2		E3 field1	
E1 mass	ke	E2 ke		E4 field2
mev	mev	mev	mev	mev
13.80	83.76	5.08	-101.95	
				-0.69
Sum of energy from above table				
E1+difference ke	102.63	E3+E4	-102.63	
Energy is conserved since 102.634=102.634				

Total energy is conserved to zero (102.634 MeV-102.634 MeV) using the convention that fields are negative. The numbers represent two orbits. The 13.8 MeV mass orbits with 83.76 MeV of kinetic energy in a 101.95 MeV strong field energy and a 0.69 MeV gravitational field energy component. Here is the strong orbit:



The particle mass 13.8 MeV is one of the quarks in a neutron. The neutron model below adds three quark energies together from quads 1 through 3. When these quads are treated the same way and

added together they make the neutron of mass 939.57 MeV within measurement error [10]. Their masses total 130.163 MeV and their kinetic energies total 799.25 MeV.

The proton is thought to be a primary manifestation of the underlying laws and as such contains information that determines many aspects of nature. The Proton Mass model is the source of constants for unification of forces in the table above.

## Appendix 2 Neutron Model

The Neutron table starts with a left side showing the quads and associated energies after the exchange 2 operation.

Unified.xls cell g191					
	mass	Energy-mev	S field	Energy	
Charge	ke		G field	mev	
0.667	15.432	101.95	17.432	753.29	
	12.432	5.08	10.432	0.69	
-0.333	13.432	13.80	15.432	101.95	
	12.432	5.08	10.432	0.69	
-0.333	13.432	13.80	15.432	101.95	
	12.432	5.08	10.432	0.69	
	-10.333	-0.62	-10.333	-0.62	
	10.408	0.67	10.408	0.67	
	10.33	0.62	10.333	0.6224	
	0.000	0.0000	0	2.02E-05	
	90.000	sum	90.000		

The right hand side of the Neutron model below lists the total mass, kinetic energy and fields associated with the neutron. Quad 4 of the code also gives us the 4<sup>th</sup> component of the gravitational field energy (-0.671 MeV) which totals -2.73 MeV.



					Gravitational	
		Residual ke				Field
Mass	Difference KE		Expansion		Strong field	
mev	mev	mev	mev	KE	MeV	MeV
101.95	641.88				-753.29	
						-0.69
13.80	78.69				-101.95	
						-0.69
13.80	78.69				-101.95	
						-0.69
		10.15		20.30	expansion pe	
				0.00	expansion ke	
	-0.67		0.67	v neutrino		
			0.05	neutrinos		-0.67
129.54	798.58	938.27	<b>PROTON MASS</b>			
0.51	0.11	e ke				
<b>ELECTRON</b>						
			Total m+ke		Total Negative	
	0.62	938.27	0.72	959.92	-959.92	-2.73
				MeV	MeV	

The neutron decays into a proton, an electron and a neutrino. This gives the measured proton mass 938.27 MeV below. As the proton and electron split they develop opposite fields of  $27.2 \times 10^{-6}$  MeV. When the electron falls into the proton field it develops  $13.6 \times 10^{-6}$  MeV.

### Abbreviated Neutron Model

Next an abbreviated model is shown for the neutron. The abbreviated model adds the three quark masses together. They total 130.163 MeV. Then, we add together the three quark kinetic energies. They total 799.25 MeV. The three quarks are imbedded in a strong field labelled -957.185 and a gravitational field energy labelled -2.723 MeV. The total field energy for the entire table is -959.92 MeV. This is lower than the known Neutron mass 939.565 MeV and the “missing energy” is a potential energy field -20.3 MeV. The quarks with their kinetic energy fall into the field and gain 10.15 MeV from their fall. The neutron subsequently falls to a radius  $7.22 \times 10^{-14}$  meters. It originally had kinetic energy 20.3 MeV but when the gravitational orbit was established, the ke was about 10.15 and the potential energy was about 10.15. The neutron plus its associated gravitational energy balances the total table energy 959.92 MeV.

<b>r20 uc2</b>			<b>Field energy</b>	
<b>Mass and Kinetic Energy</b>			<b>Strong</b>	<b>Gravitational</b>
<b>Mass</b>	<b>KE</b>	<b>Strong</b>	<b>field energy</b>	<b>Energy</b>
<b>Quarks</b>		<b>Residual</b>		
<b>MeV</b>	<b>MeV</b>	<b>Field</b>	<b>MeV</b>	<b>MeV</b>
<b>Strong</b>	<b>130.16</b>	<b>799.25</b>	<b>-957.18</b>	<b>-2.73</b>
<b>Strong Residual KE</b>		<b>10.15</b>		
<b>Neutron</b>		<b>939.57</b>	<b>-20.35</b>	<b>-959.92</b>
<b>neutrinos</b>		0.05		
<b>Gravitational ke</b>		<b>10.15</b>		
<b>Gravitational pe</b>		<b>10.15</b>		
<b>Total</b>		<b>959.92</b>		

More detail is shown below for the Quarks. To understand the origin of values in each column the reader should access reference 10. The steps leading up to this are briefly described below:

	<b>Mass and Kinetic Energy</b>		<b>Field energy</b>	
	<b>Mass</b>	<b>ke</b>	<b>Strong</b>	<b>Gravitational</b>
			<b>field energy</b>	<b>Energy</b>
	<b>MeV</b>	<b>MeV</b>	<b>MeV</b>	<b>MeV</b>
<b>Quark S</b>	<b>101.947</b>	<b>631.729</b>	<b>-753.291</b>	<b>-0.687</b>
<b>Quark D</b>	<b>13.797</b>	<b>83.761</b>	<b>-101.947</b>	<b>-0.687</b>
<b>Quark D</b>	<b>13.797</b>	<b>83.761</b>	<b>-101.947</b>	<b>-0.687</b>
	<b>129.541</b>	<b>799.251</b>	<b>-957.185</b>	
		<b>10.151</b>		
	<b>0.622</b>			-0.671
		<b>939.565</b>	<b>Mev</b>	<b>-2.732</b> MeV

The two energies 0.671 MeV and 0.622 MeV come from quad 4 of the information code. The logarithm 12.432 gives  $E=e0*\exp(N)=2.025e-5*\exp(12.432)=5.076$  MeV. Multiples of this energy appears several times in the tables above and below (10.15 MeV and 20.30 MeV).

## Abbreviated Proton Model

Mass and Kinetic Energy			Field energy	
Mass	KE	Strong Residual	Strong field energy	Gravitational Energy
MeV	MeV		MeV	MeV
<b>Strong</b>	<b>130.16</b>	<b>799.25</b>	<b>-957.18</b>	-2.73
<b>Strong Residual</b>		<b>10.15</b>		
<b>Neutron</b>		<b>939.57 (-20.3)</b>		-959.92
<b>Below the Neutron decays ejecting an anti-neutrino and an electron with ke</b>				
<b>ejected neutrino</b>		<b>-0.67</b>		
<b>Electron</b>	<b>-0.51</b>	<b>-0.11</b>	<b>2.72E-05</b>	<b>E/M charge splits</b>
<b>Proton</b>		<b>938.27</b>	<b>-2.72E-05</b>	
<b>neutrinos</b>		0.05		
<b>Gravitational ke</b>		<b>10.15</b>	<b>10.11</b>	
<b>Gravitational pe</b>		<b>10.15</b>	<b>10.19</b>	
<b>Total</b>		959.92		
(959.92=938.27+.67+.511+.67+.05+20.3)				

Information from the proton mass model underlies the four interactions of nature.

	Mass (m) (MeV)	Ke (MeV)	gamma (g)	R meters	Field (E) (MeV)
Gravity	938.272	10.11	0.990	7.224E-14	-2.732
Electromag	0.511	1.36087E-05	1.000	5.291E-11	-2.722E-05
Strong	129.541	798.580	0.140	2.093E-16	-957.185
Strong	928.121	10.151	0.989	1.430E-15	-20.303

## Appendix 2 the Schwarzschild equation

Derivation of the Schwarzschild equation is given below:

<b>Show that gamma for General Relativity is derived</b>	
<b>from gamma from Special Relativity</b>	
<b>(Derivation for Schwarzschild equation)</b>	
<b>constants from gravity above:</b>	
<b>ke</b>	<b>10.11 MeV</b>
<b>m</b>	<b>1.67E-27 kg</b>
<b>m</b>	<b>938.27 MeV</b>
<b>r</b>	<b>7.244E-14 meters</b>
<b>G</b>	<b>6.674E-11 nt m<sup>2</sup>/kg<sup>2</sup></b>
<b>C</b>	<b>299792458 m/sec</b>
<b>gravitational coupling constant 1/exp(90)</b>	
<b>gamma=m/(m+ke)</b>	
<b>gamma=938.272/(938.22+10.11)</b>	
<b>0.9893</b>	<b>gamma</b>
<b>dt=1/gamma-1</b>	
<b>0.0108</b>	<b>dt</b>
<b>dt=1/((1-(v/C)<sup>2</sup>)<sup>.5</sup>-1</b>	
<b>G=Rv<sup>2</sup>/m</b>	
<b>v=(Gm/R)<sup>.5</sup></b>	
<b>dt=1/((1-G*m/(R*C<sup>2</sup>)))<sup>0.5</sup>-1</b>	

Apply the above equation to a 90kg astronaut orbiting the earth:

<b>Astronaut (kg)</b>	<b>90</b>
<b>Mass M kg (earth)</b>	<b>5.98E+24</b>
<b>earth R (m)</b>	<b>6378100</b>
<b>dt=1/((1-G*m/(R*C<sup>2</sup>)))<sup>0.5</sup>-1</b>	
<b>3.47E-10</b>	<b>mass curves space</b>

Mass in the equation above is earth Mass, not the Astronaut. But we can find time dilation (dt) for the Astronaut in a separate (SR) calculation for dt:

<b>ke (MeV)</b>	<b>1.76E+22</b>
<b>N</b>	<b>5.38922E+28</b>
<b>ke/N</b>	<b>3.26E-07 object moving</b>
<b>g</b>	<b>9.999999997E-01</b>
<b>V/C</b>	<b>2.64E-05</b>
	<b>7908 m/sec</b>
<b>dt=1/g-1</b>	<b>3.47E-10</b>
	<b>1.00E-16</b>
	<b>3.34E-10 object has same dt</b>

With cellular cosmology, we can find the equivalent quantum radius:

<b>Astronaut (kg)</b>	<b>90</b>
<b>Mass M kg (earth)</b>	<b>5.98E+24</b>
<b>earth R (m)</b>	<b>6378100</b>
	<b><math>r=R*1.67e-27/M=2.17e-6</math> meters</b>
	<b><math>dt=1/((1-EXP(90)*G*1.67e-27/(r*C^2)))^0.5-1</math></b>
	<b>3.47E-10 mass curves space</b>
	<b>and particle follows curve</b>

And using the following Schwarzschild equation for cellular cosmology, we can find time dilation (dt) for both the curved space and the orbiting proton associated with the cell. This means that General Relativity has been combined with Special Relativity for cellular cosmology.

It may be surprising that the cell radius simulates the earth orbit, but here is proof that the cell radius is a proper geodesic for velocity 7909 m/sec.

$$G=R*V^2/m*\exp(90)=2.17e-6*7909^2/1.67e-27*\exp(90)=6.67e-11 \text{ N-m}^2/\text{kg}^2.$$

### Appendix 3 Black hole model results

Central mass (Kg)	2.00E+30	9.94E+30	4.20E+40	4.00E+31
	sun model	smallest BH	Largest BH	20x sol cmax
Avg Density	1.44E+03	5.90E+18	3.31E-01	3.64E+17
	6.000E+03	8.90E+12	5.000E+06	3.80E+12
	2.399E+06	7.90E+11	3.000E+12	7.2E+11
	2.400E+07	1.68E+13	3.000E+13	1.10E+13
		1.40E+19		4.70E+15
h0 (m)	6.92E+08	7.38E+03	3.12E+13	2.97E+04
T0 (K)	6.000E+03	9.000E+12	5.307E+06	1.04E+12
a (m/sec^2)	2.05E+02	1.22E+13	2.10E+03	3.03E+12
m1 (Kg)	5.27E+29	6.39E+30	1.14E+40	1.40E+29
cum mass1	5.266E+29	9.96E+30	1.136E+40	4.01E+31
P1 (Nt/m^2)	1.99E+13	1.26E+35	2.16E+15	4.23E+31
rho1 (kg/m^3)	1.400E+03	2.11E+18	3.300E-01	4.47E+15
h1	6.23E+08	6.64E+03	2.81E+13	2.67E+04
T	2.405E+06	9.61E+12	3.000E+12	1.24E+12
a	1.81E+02	1.35E+13	1.82E+03	3.37E+12
mass shell	4.22E+29	7.70E+29	9.10E+39	1.06E+29
cum mass2	9.483E+29	8.96E+30	2.046E+40	3.61E+31
P2	3.74E+13	1.47E+35	4.04E+15	8.70E+31
rhonew	8.345E+03	2.37E+18	1.720E-01	7.79E+15
h2	5.54E+08	5.91E+03	2.50E+13	2.38E+04
T	4.805E+06	1.00E+13	6.000E+12	1.49E+12
a	1.57E+02	1.52E+13	1.55E+03	3.79E+12
mass shell	3.28E+29	6.76E+29	7.09E+39	1.44E+29
cum mass	1.277E+30	7.97E+30	2.755E+40	3.21E+31
P	1.28E+14	1.74E+35	4.87E+15	1.75E+32
rhonew	1.433E+04	2.70E+18	1.037E-01	1.33E+16
h3	4.84E+08	5.17E+03	2.18E+13	2.08E+04
T	7.204E+06	1.04E+13	9.000E+12	1.78E+12
a	1.36E+02	1.74E+13	1.28E+03	4.33E+12
mass shell	2.47E+29	5.77E+29	5.33E+39	1.86E+29
cum mass	1.524E+30	6.97E+30	3.288E+40	2.81E+31
P	2.63E+14	2.08E+35	5.28E+15	3.46E+32
rhonew	1.957E+04	3.10E+18	7.499E-02	2.25E+16
h4	4.15E+08	4.43E+03	1.87E+13	1.78E+04
T	9.604E+06	1.09E+13	1.200E+13	2.12E+12
a	1.16E+02	2.03E+13	1.01E+03	5.05E+12
mass shell	1.77E+29	4.76E+29	3.82E+39	2.25E+29
cum mass	1.700E+30	5.98E+30	3.669E+40	2.40E+31
P	4.20E+14	2.55E+35	5.52E+15	6.83E+32
rhonew	2.345E+04	3.62E+18	5.876E-02	3.80E+16
h5	3.46E+08	3.69E+03	1.56E+13	1.49E+04

Central mass (Kg)	2.00E+30	9.94E+30	4.20E+40	4.00E+31	
	sun model	smallest BH	Largest BH	20x sol cmax	
<b>h5 (continued from above)</b>	3.46E+08	3.69E+03	1.56E+13	1.49E+04	
<b>T</b>	1.200E+07	1.15E+13	1.500E+13	2.52E+12	
<b>a</b>	1.01E+02	2.44E+13	7.54E+02	6.06E+12	
<b>mass shell</b>	1.19E+29	3.72E+29	2.56E+39	2.55E+29	
<b>cum mass</b>	1.819E+30	4.98E+30	3.925E+40	2.00E+31	
<b>P</b>	5.84E+14	3.20E+35	5.66E+15	1.37E+33	
<b>rhonew</b>	2.608E+04	4.32E+18	4.819E-02	6.49E+16	
<b>h6</b>	2.77E+08	2.95E+03	1.25E+13	1.19E+04	
<b>T</b>	1.440E+07	1.22E+13	1.800E+13	3.01E+12	
<b>a</b>	9.51E+01	3.05E+13	5.13E+02	7.57E+12	
<b>mass shell</b>	7.19E+28	2.69E+29	1.55E+39	2.64E+29	
<b>cum mass</b>	1.891E+30	3.98E+30	4.080E+40	1.60E+31	
<b>P</b>	7.55E+14	4.17E+35	5.74E+15	2.83E+33	
<b>rhonew</b>	2.813E+04	5.30E+18	4.070E-02	1.13E+17	
<b>h7</b>	2.08E+08	2.21E+03	9.36E+12	8.91E+03	
<b>T</b>	1.680E+07	1.31E+13	2.100E+13	3.63E+12	
<b>a</b>	1.12E+02	4.06E+13	3.05E+02	1.01E+13	
<b>mass shell</b>	3.69E+28	1.70E+29	7.97E+38	2.37E+29	
<b>cum mass</b>	1.928E+30	2.99E+30	4.160E+40	1.20E+31	
<b>P</b>	9.73E+14	5.76E+35	5.77E+15	6.23E+33	
<b>rhonew</b>	3.107E+04	6.79E+18	3.512E-02	2.08E+17	
<b>h8</b>	1.38E+08	1.48E+03	6.24E+12	4.52E+04	
<b>T</b>	1.920E+07	1.42E+13	2.400E+13	4.45E+12	
<b>a</b>	2.04E+02	6.10E+13	1.82E+02	1.99E+12	
<b>m8 shell Kg</b>	1.36E+28	8.00E+28	2.94E+38	1.49E+31	
<b>cum mass9 Kg</b>	1.941E+30	1.99E+30	4.189E+40	6.10E+31	
<b>P9 (nt/m^2)</b>	1.41E+15	8.82E+35	5.79E+15	7.47E+33	
<b>rho9 kg/m^3</b>	3.946E+04	9.42E+18	3.084E-02	2.39E+17	
<b>h9 (m)</b>	6.92E+07	7.38E+02	3.12E+12	4.22E+04	
<b>T (K)</b>	2.160E+07	1.58E+13	2.700E+13	4.66E+12	
<b>a10 (m/sec^2)</b>	7.90E+02	1.22E+14	4.41E+02	2.13E+12	
<b>m10 shell Kg</b>	1.94E+27	1.59E+28	4.19E+37	1.48E+31	
<b>cum mass10</b>	1.943E+30	9.96E+29	4.194E+40	5.70E+31	
<b>P10 (nt/m^2)</b>	3.57E+15	1.73E+36	5.84E+15	8.98E+33	
<b>rho10 kg/m^3</b>	8.869E+04	1.58E+19	2.761E-02	2.76E+17	
<b>h10 core</b>	0	0	0	0	
<b>r cell (m)</b>	r <sub>cell</sub> =(1/rho	1.65E-11	2.93E-16	2.44E-09	1.13E-15
			3.00E+08	3.71E+07	3.00E+08

<b>Solar example</b>		<b>B=8.62e-11</b>	
<b>Temp deg K</b>	<b>2.39E+07</b>	<b>Dmax kg/m<sup>3</sup></b>	<b>5.020E+11</b>
<b>Density kg/m<sup>3</sup></b>	<b>1.22E+05</b>		
<b>KE temp 1.5*B*T</b>	<b>3.084E-03</b>		
<b>degeneracy</b>	<b>3.56E+00</b>		
<b>Degenerate radius (D)</b>	<b>1.485E-11</b>		
<b>v/c</b>	<b>0.109</b>		
<b>Barrier</b>	<b>-0.0139</b>		
<b>Example calculation for above conditions</b>			
<b>rate</b>	<b>Pbarrier</b>	<b>Pd=(dens/m</b>	<b>Preaction rate R/sec</b>
<b>Probability/sec</b>	<b>exp(-.0139/.</b>	<b>(1.2e5/5e11</b>	<b>v/r/exp(62.87)</b>
<b>8.19E-18</b>	<b>0.0109429</b>	<b>2.43E-07</b>	<b>1.09667E-09</b>
<b>burn time (Byrs)</b>	<b>3.9</b>		<b>3.078E-09</b>
<b>sun N</b>	<b>1.198E+57</b>		
<b>fract burning</b>	<b>0.15</b>		
<b>burn rate N/sec*mev/</b>	<b>1.47E+39</b>		

## Appendix 4 Information based reality

Observations regarding unexpected connections in quantum mechanically entangled systems are revealing a new understanding of our position in the universe. Experiments known by the initials EPR (Einstein, Podolsky and Rosen) show a statistical correlation between separated particle properties. If two particles are produced with opposite spins and move in different directions, it is observed that changes induced in one particle cause immediate changes in its partner. The classic “dual slit experiment” demonstrates that quantum photons can either produce a spot pattern or an interference pattern depending on whether an observer can “measure” which of two slits the photon travels through. More recently, a Discovery Magazine article by Zeeva Merali (Aug 26, 2010) indicates that an entangled particle responds to future changes in its partner (called “back from the future” observations by Jack Sarfatti of Cornell University). The author discusses the implications of taking this data at face value. Three FQXi contest essays [31][32][33] lead up to what I believe to be the explanation for EPR results and back from the future observations.

A recent paper [30] stating that in view of EPR, double slit and back from the future observations we should suspect that time and distance are largely products of our perception. It appears that reality can be viewed as an interaction between an information source and the information we are comprised of. I call this information based reality. The neutron mass model is a result of using information theory to correlate fundamental energy values for the particles of nature. The particles can be viewed as information. Further, it was shown that the light our eye receives can be correlated with an equation attributed to Feynman. The response of our eye can be viewed as information and the same unit of information that correlates fundamental energy values gives the



information value of the four wavelengths of light our eye is tuned to. Quantum mechanics utilizes complex numbers and a “particle” is the point where the math allows the complex numbers to be a real number. This is its source of quantum mechanics, particle/wave duality and the Heisenberg uncertainty principle. The Copenhagen interpretation makes quantum mechanics an information science “...the only information we can have about a particle position is probabilistic”. If particles are information, light is information and the only thing we can know about a particles positions is information, one has to ask the question what isn't information?

One may ask how information based reality relates to the present paper. There is no doubt that the information source has a beginning and history. The big bang describes this well but the language we use needs to change. The history of the universe may be a series of “snapshots” that we interpret as time. It seems logical that information related to the history is refreshed once per cycle around the quantum circle that defines time and distance (See section above entitled “TIME AND COSMOLOGY” that states a radius of  $7.2e-14$  meters). However, these circles may contain only information (zero distance and time) because we can identify the information that defines the circle. Each snapshot in our history can be characterized by energy because energy is information. The exact relationship is probability  $P$  equals an energy ratio  $e_0/e$ , i.e.  $P=e_0/e$  and information  $I$  equals natural logarithm of probability, i.e.  $I= \ln P$  [10]. Information can be spread out but the separation does not need to be distance, it could be just information. For example, our mind might “unpack” the information related to the quantum circle and create a perception of distance and time. Our body is information “adopted” from the information source and our perceptions center our perspective locally. There are two types of time. One is related to the sequence defined by fundamental time  $1.5e-21$  seconds (See section above “Time and Cosmology” and the other is related to gamma. Gamma is maintained as the snapshots sequence and this explains the twin paradox. In information based reality, the explanations of EPR, the double slit experiment and back from the future observations are as follows:

The essence of EPR is that separated up spin and down spin particle are correlated. You flip the spin of one particle and the other will flip appearing to communicate in a way that violates our concept of distance.

Explanation: The information source is built from “zero energy” and “zero properties” [1][7] and two entangled particles preserve zero. They are separated by information only. You create a space like reality by accessing the information source. This gives you the impression they are physically separated and can't communicate.

The double slit experiment has a similar explanation. This phenomenon occurs because reality is an interaction between preexisting information and the observer's thoughts. With the information the observer has been given two perceptions are possible and an interference pattern is observed. If the individual photon's path can be observed (measured), it will act as an individual photon, not an interference pattern. The perception is completed by acting on unambiguous information from the source.

The essence of back to the future observation is similar to EPR, except the particles are separated in time and violate our concept of order.

Explanation: The information source separates zero into two related energy ratios (probabilities) and the entangled particles preserve zero. To contain energy the particles must have a time base but it does not

exist in the information source. The impression that they contain energy and that time has passed is your constructed reality based on multiple accesses of the information source.