

Each unidimensional system is Hamiltonian, so that each unidimensional system is quantizable

Abstract

I prove that the field of classical trajectories can be a field Hamiltonian projection of higher dimension.

I hypothesize that the same is valid for any dimension: each system is Hamiltonian, and each system is quantizable using the correspondence principle

Unidimensional Hamiltonian System

Each unidimensional trajectory can be described by a differential equation of high order, and high degree (each derivable function can be approximate by a sum of Taylor, Fourier and Laplace series, that is a solution of a linear differential equation, but an improved approximation is a nonlinear differential equation):

$$0 = \mathcal{F}(y, \dot{y}, \ddot{y}, \dots) = a_{10\dots} + a_{010\dots}y + a_{0010\dots}\dot{y} + \dots + a_{0101\dots}y\dot{y} + \dots \quad (1)$$

$$0 = \mathcal{F}(y, \dot{y}, \ddot{y}, \dots) = \sum_{i_0, \dots, i_n} a_{i_0, \dots, i_n} \frac{d^{i_0}y}{dt^{i_0}} \dots \frac{d^{i_n}y}{dt^{i_n}} \quad (2)$$

the derive of the differential equation is linear in the higher derivative:

$$0 = \frac{d\mathcal{F}(y, \dot{y}, \ddot{y}, \dots)}{dt} = \sum_{i_0, \dots, i_n} a_{i_0, \dots, i_n} \frac{d^{i_0}y}{dt^{i_0}} \dots \frac{d^{i_n+1}y}{dt^{i_n+1}} + \dots \quad (3)$$

$$\frac{d^N y}{dt^N} = \mathcal{G}\left(y, \frac{dy}{dt}, \dots, \frac{d^{N-1}y}{dt^{N-1}}\right) \quad (4)$$

so that each polynomial differential equation can be write linearly in the maximum derivative; so that:

$$\left\{ \begin{array}{l} y = y_0 \\ \dot{y}_0 = y_1 \\ \dot{y}_1 = y_2 \\ \vdots \\ \dot{y}_{N-2} = y_{N-1} \\ \dot{y}_{N-1} = \mathcal{G}(y_0, \dots, y_{N-1}) \end{array} \right. \quad (5)$$

this system can be the half of an Hamiltonian system $H = \sum_i p_i f_i$, that have N new momenta:

$$H = \sum_{i=0}^N p_i f_i = \sum_{j=0}^N p_i \{y_{i+1} + \delta_{iN} [-y_{i+1} + \mathcal{G}]\} = \sum_{i=0}^{N-1} p_i y_{i+1} + p_N \mathcal{G}$$

$$\begin{cases} \dot{y}_{j \neq N} = \frac{\partial H}{\partial p_j} = f_j = y_{j+1} \\ \dot{y}_N = \frac{\partial H}{\partial p_N} = f_N = \mathcal{G} \\ \dot{p}_{j \neq N} = -\frac{\partial H}{\partial y_j} = -p_N \frac{\partial \mathcal{G}}{\partial y_j} - p_{j-1} \\ \dot{p}_N = -\frac{\partial H}{\partial y_N} = -p_{N-1} \end{cases} \quad (6)$$

the trajectories in the coordinates are ever the same, for each momentum initial condition; the volume of the phase space is an invariant in the space (coordinates, momenta) and the sum of the areas is invariant, because of there is a momenta compensation.

In this case, each quantum system is equal to the classical system:

$$\begin{aligned} H &= p_i f_i \\ i\hbar \partial_t \psi &= -i\hbar \sum_i f_i \partial_i \psi \\ 0 &= \partial_t \psi + \sum_i f_i \partial_i \psi \end{aligned} \quad (7)$$

there are two classical solutions:

$$\begin{cases} \dot{y}_1 = f_1 \\ \vdots \\ \dot{y}_N = f_N \end{cases} \quad (8)$$

$$\frac{dy_1}{f_1} = \dots = \frac{dy_N}{f_N} = dt$$

$\partial_t \psi + f_i \partial_i \psi = 0$

that is a surface solution, in an N+1 dimensional space (coordinates and times), and ψ is the solution of the differential equation.

Another solution is the Hamilton-Jacobi equation, that give the classical solution of the Hamiltonian:

$$\begin{aligned} H &= p_i f_i \\ \partial_t \psi + H(p_i = \partial_i \psi, y_i) &= 0 \end{aligned} \quad (9)$$

$\partial_t \psi + f_i \partial_i \psi = 0$

in this case the function ψ permit to calculate the momenta values like a gradient of the ψ function. Also in this case the classical solution, and the quantum solution, coincide; and the equation for the amplitude, or the probability, are equal because of the linearity of the equation.