

# **Examination of a multiple universe version of the Partition function of the universe, based upon Penrose's cyclic conformal cosmology. Leading to uniform values of $\hbar$ (Planck's constant) and invariant physical laws in each Universe of the 'multiverse'**

**Andrew Walcott Beckwith**

Physics Department, Chongqing University, College of Physics, Chongqing University Huxi Campus, No. 44 Daxuechen Nanlu, Shapinba District, Chongqing 401331, People's Republic of China

[Rwill9955b@gmail.com](mailto:Rwill9955b@gmail.com); [abeckwith@uh.edu](mailto:abeckwith@uh.edu)

## **Abstract**

We look at, starting with Shankar's treatment of the partition function, inserting in the data of the modified Heisenberg uncertainty principle as to give a role of the inflaton in the formation of a partition of the universe. The end result will be, even with the existence of a multiverse, i.e. simultaneous universes, uniform physical laws throughout the Multiple universes.

**Key words, Cyclic Conformal cosmology (Penrose), Modified HUP**

## 1. What is Unique about our application of modification of the Penrose Cyclic cosmology theory?

We review the modification of the Penrose cyclic conformal cosmology paradigm given in [1] and include in a partition function given in [2], as a way to include in the modified Heisenberg Uncertainty principle, [3] as a way to ascertain the role of the inflaton, as we write it up in using Padmanabhan's reference [4].

Modification of the HUP, and including in the inflaton, in the Partition function will then lead to, after we are including in the results from [2] a way to discuss how to get a uniform value for Planck's constant,  $\hbar$ , in, per cycle creation of new universes.

Contrary to the supposition given by Susskind and others, [5] as to up to  $10^{100}$  universes, with only say  $10^6$  of them surviving due to sufficiently 'robust' cosmological values, for stable physical law. The end result is that we would have instead is a 'multiverse' which is dynamic and stable over time. And so we review our present modification of the Penrose cyclic conformal cosmology model to take into account multiple universes

## 2. Extending Penrose's suggestion of cyclic universes, black hole evaporation, and the embedding structure our universe is contained within. This multiverse embeds BHs and may resolve what appears to be an impossible dichotomy.

That there are no fewer than  $N$  universes undergoing Penrose 'infinite expansion' [1] contained in a mega universe structure. Furthermore, each of the  $N$  universes has black hole evaporation, with the Hawking radiation from decaying black holes. If each of the  $N$  universes is defined by a partition function, called  $\{\Xi_i\}_{i=1}^{i=N}$ , then

there exist an information ensemble of mixed minimum information correlated as about  $10^7 - 10^8$  bits of information per partition function in the set  $\{\Xi_i\}_{i=1}^{i=N}$  before, so minimum information is conserved between a set of partition functions per universe. We are when following this using the notation of [1] while noting that there is a subsequent alteration of the notation used for partition functions.

$$\{\Xi_i\}_{i=1}^{i=N} \Big|_{\text{before}} \equiv \{\Xi_i\}_{i=1}^{i=N} \Big|_{\text{after}} \quad (1)(30)$$

However, there is non-uniqueness of information put into each partition function  $\{\Xi_i\}_{i=1}^{i=N}$ . Furthermore

Hawking radiation from the black holes is collated via a strange attractor collection in the mega universe structure to form a new big bang for each of the  $N$  universes represented by  $\{\Xi_i\}_{i=1}^{i=N}$ . Verification of this mega

structure compression and expansion of information with a non-uniqueness of information placed in each of the  $N$  universes favors ergodic mixing treatments of initial values for each of  $N$  universes expanding from a singularity beginning. The  $n_f$  value, will be using (Ng, 2008)  $S_{\text{entropy}} \sim n_f$ . [14]. How to tie in this energy expression, as in Eq. (1) (30) will be to look at the formation of a nontrivial gravitational measure as a new big bang for each of the  $N$  universes as by  $n(E_i)$ . the density of states at a given energy  $E_i$  for a partition function. (Poplawski, 2011) [6] [15]

$$\{\Xi_i\}_{i=1}^{i=N} \propto \left\{ \int_0^{\infty} dE_i \cdot n(E_i) \cdot e^{-E_i} \right\}_{i=1}^{i=N} \quad (2)$$

(31)

Each of  $E_i$  identified with Eq.(2) (31) above, are with the iteration for N universes (Penrose, 2006)[14] Then the following holds, namely, from [1]

**Claim 1,**

$$\frac{1}{N} \cdot \sum_{j=1}^N \Xi_j \Big|_{j\text{-before-nucleation-regime}} \xrightarrow{\text{vacuum-nucleation-transfer}} \Xi_i \Big|_{i\text{-fixed-after-nucleation-regime}} \quad (3)$$

(32)

For N number of universes, with each  $\Xi_j \Big|_{j\text{-before-nucleation-regime}}$  for  $j = 1$  to N being the partition function of each universe just before the blend into the RHS of Eq. (3) (32) above for our present universe. Also, each of the independent universes given by  $\Xi_j \Big|_{j\text{-before-nucleation-regime}}$  are constructed by the absorption of one to ten million black holes taking in energy. **I.e.** [1] [14]. Furthermore, the main point is similar to what was done in [18] in terms of general ergodic mixing

**Claim 2**

$$\Xi_j \Big|_{j\text{-before-nucleation-regime}} \approx \sum_{k=1}^{\text{Max}} \tilde{\Xi}_k \Big|_{\text{black-holes-jth-universe}} \quad (4)$$

(33)

What is done in **Claim 1 and Claim 2** is to come up with a protocol as to how a multi dimensional representation of black hole physics enables continual mixing of spacetime [16] largely as a way to avoid the Anthropic principle, as to a preferred set of initial conditions. How can a graviton with a wavelength  $10^4$  the size of the universe interact with a Kere black hole, spatially. Embedding the BH in a multiverse setting may be the only way out.

**Claim 1** is particularly important. The idea here is to use what is known as CCC cosmology, which can be thought of as the following.

First. Have a big bang ( initial expansion) for the universe. After redshift  $z = 10$ , a billion years ago, SMBH formation starts. Matter- energy is vacuumed up by the SMBHs, which at a much later date than today ( present era) gather up all the matter-energy of the universe and recycles it in a cyclic conformal translation, as follows, namely

$$\begin{aligned} E &= 8\pi \cdot T + \Lambda \cdot g \\ E &= \text{source for gravitational field} \\ T &= \text{mass energy density} \\ g &= \text{gravitational metric} \\ \Lambda &= \text{vacuum energy, rescaled as follows} \end{aligned} \quad (5)$$

$$\Lambda = c_1 \cdot [Temp]^\beta \quad (6)$$

$c_1$  is , here a constant. Then

The main methodology in the Penrose proposal has been in Eq.(6) evaluating a change in the metric  $g_{ab}$  by a conformal mapping  $\hat{\Omega}$  to

$$\hat{g}_{ab} = \hat{\Omega}^2 g_{ab} \quad (7)$$

Penrose's suggestion has been to utilize the following[1]

$$\hat{\Omega} \xrightarrow{ccc} \hat{\Omega}^{-1} \quad (8)$$

The infall into cosmic black holes has been the main mechanism which the author asserts would be useful for the recycling apparent in Eq.(8) above with the caveat that  $\hbar$  is kept constant from cycle to cycle as represented by

$$\hbar_{old-cosmology-cycle} = \hbar_{present-cosmology-cycle} \quad (9)$$

Eq. (9) is to be generalized, as given by a weighing averaging as given by Eq (3). where the averaging is collated over perhaps thousands of universes, call that number  $N$ , with an ergotic mixing of all these universes, with the ergodic mixing represented by Eq.(3) to generalize Eq.(9) from cycle to cycle.

### 3. Now for the mixing being put in, and Birkhoff's Ergodic mixing theorem

We will, afterwards, do the particulars of the partition function. But before that, we will do the 'mixing' of inputs into the Partition function of the Universe, i.e. an elaboration on Eq. (3) above. To do this, first, look at the following, from [ 7 ]

#### Birkhoff's Ergodic mixing theorem

**Let**  $T : (X, \mu) \rightarrow (X, \mu)$  be a measure preserving transformation of a probability space  $\varphi \in L^1(X, \mu)$

Then for  $\mu$  almost every  $x \in X$  the following time average exists

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \varphi(T^j(x)) =: \varphi_T(x) \quad (10)$$

In the end, we need to have a way to present how the bona fides of Eq. (9) can be established, and the averaging of both Eq. (10) and Eq. (4) above need to be put to a consistent general treatment for an invariant  $\hbar$  for cycle to cycle, of cosmological creation.

To do this, we also refer to the generalized treatment of , from [ 8 ]

$$\begin{aligned}
\text{Energy(average)} &= kT^2 \left( \frac{\partial \ln Z(T)}{\partial T} \right) \\
&\sim \hbar \omega \\
\Leftrightarrow \hbar(\text{averaged - value}) &= \frac{k_B T^2}{\omega} \left( \frac{\partial \ln Z(T)}{\partial T} \right) \\
&= \frac{k_B T^2}{\omega} \left( \frac{\partial \ln Z(T)}{\partial T} \right) = \frac{k_B \lambda T^2}{2\pi c \omega} \left( \frac{\partial \ln Z(T)}{\partial T} \right) \\
\Leftrightarrow \hbar(\text{averaged - value}) &= \frac{k_B \lambda T^2}{2\pi c} \left( \frac{\partial \ln Z(T)}{\partial T} \right)
\end{aligned} \tag{11}$$

Having said, that, the remaining constraint is to come up with a suitably averaged value of the Partition function in the above work. Our averaging eventually will have to be reconciled with the Birkhoff Ergodic Mixing theorem.

### 3. How to average out the Planck's constant, of Eq. partition function given in Eq. (11)

We begin with what is given in Shankar's treatment of the partition function of [9] as given by

$$Z(T) = Z(\beta) \simeq \int dr \cdot \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r)]}{\Delta r} \tag{12}$$

Using, for Pre Planckian space-time the approximation of

$$\begin{aligned}
Z(\beta) &\simeq \int dr \cdot \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r)]}{\Delta r} \sim \Delta r \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r_{midpoint})]}{\Delta r} \\
\Leftrightarrow Z(\beta) &\simeq \sqrt{\frac{1}{2\pi}} \cdot \exp[-\beta V(r_{midpoint})]
\end{aligned} \tag{13}$$

Approximate using Beckwith's treatment of the HUP, in Pre Planckian space-time [10]

$$\begin{aligned}
Z(\beta) &\simeq \int dr \cdot \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r)]}{\Delta r} \sim \Delta r \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r_{midpoint})]}{\Delta r} \\
\Leftrightarrow Z(\beta) &\simeq \sqrt{\frac{1}{2\pi}} \cdot \exp[-\beta V(r_{midpoint})] \\
V(r_{midpoint}) &\sim \Delta E \sim \frac{\hbar}{\delta t \cdot \delta g_{tt}} \sim \frac{\hbar}{\delta t \cdot (a_{\min})^2 \phi} \\
\Leftrightarrow Z(\beta) &\simeq \sqrt{\frac{1}{2\pi}} \cdot \exp \left[ -\beta \cdot \frac{\hbar}{\delta t \cdot (a_{\min})^2 \phi} \right]
\end{aligned} \tag{14}$$

Put in now the value of the inflaton given by Padmanbhan,[4] as for

$$\begin{aligned}
a &\approx a_{\min} t^\gamma \\
\Leftrightarrow \phi &\approx \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right\} \\
\Leftrightarrow V &\approx V_0 \cdot \exp \left\{ -\sqrt{\frac{16\pi G}{\gamma}} \cdot \phi(t) \right\}
\end{aligned} \tag{15}$$

Put in the value for the inflaton of Eq. (15) into the partition function of Eq. (14)

$$\begin{aligned}
Z(\beta) &= \sqrt{\frac{1}{2\pi}} \cdot \exp \left[ -\beta \cdot \frac{\hbar}{\delta t \cdot (a_{\min})^2 \phi} \right] = \sqrt{\frac{1}{2\pi}} \cdot \exp \left[ -\beta \cdot \frac{\hbar}{\delta t \cdot (a_{\min})^2 \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right\}} \right] \\
&= \sqrt{\frac{1}{2\pi}} \cdot \exp \left[ \beta \cdot \frac{\hbar}{\delta t \cdot (a_{\min})^2 \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ 1 / \left( \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}} \right] \\
&\& \ln \left\{ 1 / \left( \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\} = \left[ \frac{\left\{ 1 / \left( \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}^{-1}}{\left\{ 1 / \left( \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}} \right] + \frac{1}{2} \cdot \left[ \frac{\left\{ 1 / \left( \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}^{-1}}{\left\{ 1 / \left( \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}} \right]^2 + \dots
\end{aligned} \tag{16}$$

Then

$$\begin{aligned}
\frac{\partial \ln Z(T)}{\partial T} &= \frac{1}{Z(T)} \cdot \frac{\partial Z(T)}{\partial T} \\
&= \frac{-\hbar}{\delta t \cdot (a_{\min})^2 \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ 1 / \left( \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}} \cdot \frac{1}{k_B T^2} \\
&= \frac{\hbar}{\delta t \cdot (a_{\min})^2 \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ \left( \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}} \cdot \frac{1}{k_B T^2} \\
&\sim \frac{\hbar}{(\delta t)^2 \cdot (a_{\min})^2 \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \sqrt{\frac{\gamma}{4\pi G}}} \cdot \frac{1}{k_B T^2} \tag{17}
\end{aligned}$$

Then by use of Eq. (11) we obtain

$$\begin{aligned}
\hbar(\text{averaged - value}) &= \frac{k_B \lambda T^2}{2\pi c} \left( \frac{\partial \ln Z(T)}{\partial T} \right) \\
&\sim \frac{\lambda}{2\pi c} \frac{\hbar}{(\delta t)^2 \cdot (a_{\min})^2 \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \sqrt{\frac{\gamma}{4\pi G}}} \tag{18} \\
&\Leftrightarrow \frac{\lambda}{2\pi c} \frac{1}{(\delta t)^2 \cdot (a_{\min})^2 \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \sqrt{\frac{\gamma}{4\pi G}}} \sim 1
\end{aligned}$$

This is the baseline of the constraint which will make Planck's constant, a constant per universe creation cycle. As given by Eq. (9). I.e. Eq. (9) is confirmed by Eq. (18). We will next then go to how this ties into Eq. (10) above, via use of averaging is affecting the choice of the inputs into Eq. (18) above. Doing this will allow investigation as to how to falsify the Birkhoff Ergodic mixing theorem as mentioned next.

#### 4. Applying the Birkoff Ergodic averaging Eq.(10) to the inputs into Eq. (18)

To do this, we specifically look at the wavelength, namely , applying [7] to a wavelength  $\lambda$  . One over the wavelength is proportional to frequency, so if we have the wave length, as represented by the following situation, with invariance set in stone. Here we are assuming that the formulation is, as follows: With N the number of recycled “universes”

$$\begin{aligned}\varphi(T^j(x)) &= \lambda \\ \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=0}^{N-1} \varphi(T^j(x)) &= \lambda\end{aligned}\tag{19}$$

I.e. the averaging by the Burkhoﬀ theorem implies that there is a critical invariance. And this invariance should be linked, then, to the diameter of a non singular bounce point. A non singular bounce, i.e., beginning of an expansion of a new universe is the main point of [ 11] . Furthermore, we have that if we look at applying the insights of [12, 13] we obtain

$$\begin{aligned}V_0 &\sim m = \text{mass}(\text{tranferred} - \text{space} - \text{time}) \\ &= m_{\text{graviton}} \cdot \text{Entropy} = m_{\text{graviton}} n(\text{entropy}) \\ &\& \\ n(\text{entropy}) &\sim 10^{10} - 10^{20} \\ &\& \\ m_{\text{graviton}} &\sim 10^{-62} \text{ grams} \\ &\& \\ (a_{\text{min}})^2 &\sim 10^{-110} \\ &\& \\ \delta t &\sim 10^{-\beta} t_{\text{Planck}}; 0 < \beta < 10\end{aligned}\tag{20}$$

We will try to show, in a later date that these are invariant per cycle, but the upshot is that if there is a natural fit, as to Eq. (19) and if  $\lambda$  is fixed as an invariant per cycle, given by 19, then the invariance of  $\hbar$  per cycle is then maintained.

## 5. Conclusion: Implications of the invariance of $\hbar$ : Uniform physics laws per universe, and not the $10^{1000}$ created universes with only say $10^{10}$ surviving through a cosmic cycle.

In a word this demolishes the program of the cosmic land scape of string theory [5] , and gives credence to the possibility of an invariant multiverse, which would not be collapsing.

If this is confirmed, experimentally, it will do much to reduce what has been at times a post modern fragmentation of basic physics inquiry and to have physics, with a uniform set of laws, regardless of if there were many worlds, or just one, in terms of one universe, or many universes. As well as allow investigation of the information theory approach of [14] to event horizons and early universe cosmology. How [14] could influence a choice of partition functions is given in reference A below

## 6.. Acknowledgements

This work is supported in part by National Nature Science Foundation of China grant No. 11375279



## Appendix A: Highlights of J.-W. Lee's paper [14]

The following formulation is to highlight how entropy generation blends in with quantum mechanics, and how the breakdown of some of the assumptions used in Lee's paper coincide with the growth of degrees of freedom. What is crucial to Lee's formulation, is Rindler geometry, not the curved space formulation of initial universe conditions. First of all. [19] (Lee, 2010),

*“Considering all these recent developments, it is plausible that quantum mechanics and gravity has information as a common ingredient, and information is the key to explain the strange connection between two. If gravity and Newton mechanics can be derived by considering information at Rindler horizons, it is natural to think quantum mechanics might have a similar origin. In this paper, along this line, it is suggested that quantum field theory (QFT) and quantum mechanics can be obtained from information theory applied to causal (Rindler) horizons, and that quantum randomness arises from information blocking by the horizons*

To start this we look at the Rindler partition function, as by [14] (Lee, 2010)

$$Z_R = \sum_{i=1}^n \exp[-\beta H(x_i)] = \text{Trace} \cdot (\exp[-\beta H]) \quad (\text{A.1})$$

As stated by Lee [48] .we expect  $Z_R$  to be equal to the quantum mechanical partition function of a particle with mass  $m$  in Minkowski space time. Furthermore, there exists the datum that: Lee made an equivalence between Eq. (A1) and [19] (Lee, 2010)

$$Z_Q = N_1 \int \wp x \cdot \exp\left[\frac{-i}{\hbar} \cdot I(x_i)\right] \quad (\text{A2})$$

Where  $I(x_i)$  is the action ‘integral’ for each path  $x_i$ , leading to a wave function for each path?  $x_i$

$$\psi \sim \exp\left[\frac{-i}{\hbar} \cdot I(x_i)\right] \quad (\text{A3})$$

If we do a rescale  $\hbar = 1$ , then the above wave equation can lead to a Schrodinger equation,

**The example given by (Lee, 2010)** is that there is a Hamiltonian for which

$$H(\phi) = \int d^3x \cdot \left\{ \frac{1}{2} \cdot \left( \frac{\partial \phi}{\partial t} \right)^2 + \frac{1}{2} \cdot (\nabla \phi)^2 + V(\phi) \right\} \quad (\text{A4})$$

Here,  $V$  is a potential, and  $\phi$  can have arbitrary values before measurement, and to a degree,  $Z$  represent uncertainty in measurement. In Rindler co-ordinates,  $H \rightarrow H_R$ , in co-ordinates  $(\eta, r, x_2, x_3)$  with proper time variance  $ard\eta$  then

$$H_R(\phi) = \int dr dx_{\perp} ar \cdot \left\{ \frac{1}{2} \cdot \left( \frac{\partial \phi}{\partial r} \right)^2 + \frac{1}{2} \cdot \left( \frac{\partial \phi}{ar \partial \eta} \right)^2 + \frac{1}{2} \cdot (\nabla_{\perp} \phi)^2 + V(\phi) \right\} \quad (\text{A5})$$

Here, the  $\perp$  is a plane orthogonal to the  $(\eta, r)$  plane. If so then

$$Z = \text{tr} \exp[-\beta H] \mapsto Z_R = \text{tr} \exp[-\beta H_R] \quad (\text{A6})$$

**Now, for the above situation, the following are equivalent**

1.  $Z_R$  Thermal partition function is from information loss about field beyond the Rindler Horizon
2. QFT formation is equivalent to purely information based statistical treatment suggested in this paper
3. QM emerges from information theory emerging from Rindler co-ordinate

Lee also forms a Euclidian version for the following partition function, if  $I_E(x_i)$  is the Euclidian action for the scalar field in the initial frame. I.e.

$$Z_Q^E = N_1 \int \phi x \cdot \exp\left[\frac{-i}{\hbar} \cdot I_E(x_i)\right] \quad (\text{A7})$$

There exist analytic continuation of  $\tilde{t} \mapsto it$  leading to  $Z_Q^E \mapsto Z_Q =$  Usual zero temperature QM partition function of  $Z_Q$  for  $\phi$  fields.

**Important Claim:** The following are equivalent

1.  $Z_R$  And  $Z_Q$  are obtained by analytic continuation from  $Z_Q^E$
2.  $Z_R$  And  $Z_Q$  are equivalent.

## Reference

- [1] Penrose, Roger, (2010) Cycles of Time: An Extrardinary New View of the Universe. The Bodley Head, the Random House Group, Ltd., London
- [2] Beckwith, Andrew, "Analyzing Black Hole Super-Radiance Emission of Particles/Energy from a Black Hole as a Gedankenexperiment to Get Bounds on the Mass of a Graviton"; Advances in High Energy Physics Volume 2014 (2014), Article ID 230713, 7 pages <http://dx.doi.org/10.1155/2014/230713>; <http://www.hindawi.com/journals/ahp/2014/230713/>
- [3] Beckwith, A. (2016) Gedanken Experiment for Refining the Unruh Metric Tensor Uncertainty Principle via Schwarzschild Geometry and Planckian Space-Time with Initial Nonzero Entropy and Applying the Riemannian-Penrose Inequality and Initial Kinetic Energy for a Lower Bound to Graviton Mass (Massive Gravity). *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 106-124. doi: [10.4236/jhepgc.2016.21012](https://doi.org/10.4236/jhepgc.2016.21012).
- [4] Padmanabhan, T., "understanding our universe, current status and open issues", pp 175-204, of 100 Years of Relativity, Space-Time Structure: Einstein and Beyond, Editor, A. Ashatekar, World Scientific Publishing Co. Pte.Ltd, Singapore, Republic of Singapore, 2005; <http://arxiv.org/abs/gr-qc/0503107>
- [5] Susskind, L. "[The Cosmic Landscape: String Theory and the Illusion of Intelligent Design](#)", Little, Brown, and company, Time Warner Book Group, New York, New York, USA, 2006
- [6] Poplawski, N. , "Cosmological constant from QCD vacuum and torsion," *Annalen der Physik*, vol. 523, pp. 291–295, 2011. [View at Publisher](#) · [View at Google Scholar](#)
- [7] Katok, A., Hasselblatt, "Introduction to the Modern theory of Dynamical Systems," Cambridge University Press, 1999, Cambridge, UK.
- [8] Shankar, R. "Fundamentals of Physics, mechanics, relativity and thermodynamics", Yale University Press, Yale, USA, 2014
- [9] Shankar, R. "Principles of Quantum Mechanics, 2<sup>nd</sup> edition", Springer Verlag, Heidelberg, Federal Republic of Germany, 1994
- [10] Beckwith, A. (2016) Gedanken Experiment for Refining the Unruh Metric Tensor Uncertainty Principle via Schwartz Shield Geometry and Planckian Space-Time with Initial Nonzero Entropy and Applying the Riemannian-Penrose Inequality and Initial Kinetic Energy for a Lower Bound to Graviton Mass (Massive Gravity). *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 106-124. doi: [10.4236/jhepgc.2016.21012](https://doi.org/10.4236/jhepgc.2016.21012)

- [11] Beckwith, A. (2016) Gedanken Experiment for Delineating the Regime for the Start of Quantum Effects, and Their End, Using Turok's Perfect Bounce Criteria and Radii of a Bounce Maintaining Quantum Effects, as Delineated by Haggard and Rovelli. *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 287-292. doi:[10.4236/jhepgc.2016.23024](https://doi.org/10.4236/jhepgc.2016.23024).
- [12] . Corda, C. " Gravity's Primordial Breath, " Electronic Journal of Theoretical physics, EJTP 9, No. 26 (2012) 1–10; <http://www.ejtp.com/articles/ejtpv9i26p1.pdf>
- [13 ] Gao, C. (2012) A Model of Nonsingular Universe. *Entropy*, 14, 1296-1305.  
<http://dx.doi.org/10.3390/e14071296>
- [14] [ Lee, J.W. "Quantum Mechanics Emerges from Information Theory Applied to Causal Horizons", *Found Phys*, DOI: 10.1007/s10701-010-9514-3, pp 1-10 ( on line November 3, 2010)