

# Vir Theory of Elementary Particles

M. J. Germuska

Retired, 47 Clifden Road, Worminghall, HP18 9JR, UK.  
mgermuska@aol.com, phone: +(44) 1844 339754

## Abstract

A new theory of elementary particles is presented based on solid mathematical foundations of Variational Calculus, Euler's Equations of Motion and Special Relativity. The Vir Theory of Elementary Particles explains that a particle is a stationary circular wave created by the motion of twin vortices in the relativistic ether. Mathematical equations show that they must have integer or half-integer spin, explain why the electric charge must be plus, minus or zero, why neutral particles come in right-handed and left-handed pairs, why charge-parity-time (CPT) transformation are invariant and why there are anti-particles but no stable anti-matter. A simple formula for the relationship between particle spin and mass is also derived, that can be used to verify the theory using the existing PDG data.

**Keywords:** elementary particles, theory, spin, mass, electric charge, CPT, anti-matter

## 1. Introduction

All known matter has a hierarchical structure with smaller and smaller entities. The smaller entities orbit the centre of a larger entity, or each other. It is believed that in the centre of each galaxy is a black hole orbited by stars. Stars are orbited by planets. Planets are made of atoms that have a nucleus made of protons and neutrons, while the nucleus is orbited by electrons. It is also believed that protons, neutrons and other hadrons are made up of quarks that orbit inside them giving particles the spin and almost all of their mass.

It is conceivable that this structure goes on for ever. However, this paper explores the possibility that the process is finite, ending with some fundamentally different ultimate entity that must be unique, with the least resistance to some kind of motion. The theory must have solid mathematical foundations based on the principle of least effort, i.e. least action [1], [2], which is shared by all successful theories including Optics [3], Newtonian Mechanics [4], Special Relativity [5], General Relativity [6] and Quantum Mechanics [7].

For the ultimate entity to be fundamentally different it must have no orbiting parts, although it must have integer or half-integer spin like all elementary particles. Spinning motion is governed by Euler's equations [8] which tell us that a spinning hole in relativistic ether [9] creates a circular wave. The wave destroys itself, unless it has integer or half-integer spin. Hence the ultimate entity should be a hole having the shape with the least resistance to spinning. This shape is a spheroid [10], a convex object with a circular footprint and elliptical profile. This may explain why stars, planets and moons are all spheroids.

However, the integrals for mass and moments of inertia of spheroids do not lead to a formula connecting spin and mass, since there is no algebraic solution to polynomials with powers greater than three. On the other hand, if instead of convex shapes we consider concave we find that the shape with the minimum moment of inertia is that of a vortex [10] and we obtain an algebraic spin-mass formula that gives the mass for over 200 particles with the accuracy of the measurement errors, as will be shown in the next paper.

A particle may be imagined as twin vortices with the common spin axis connected at the large ends, a shape called Vir [11]. The eye of a hurricane is miniscule in comparison to the area over which a hurricane affects the weather. This may well explain why particles have the dual nature, bullet-like and wave-like. Vir Theory explains mathematically why spin is quantised, why electric charge is positive, negative or neutral; why neutral particles come in right-handed and left-handed pairs; why particles are invariant under the charge-parity-time (CPT) transformation; why there are anti-particles, but no stable anti-mass. Quarks are simply assigned spin and charge and even then do not provide an explanation for any of the above.

We will show by solving the Euler equations of motion for rotating bodies that a Vir with its spinning, precession and nutation can result in a circular wave, where the head chases its tail. Such a wave destroys itself after one precession if the head and tail are out of phase, but results in a stationary circular wave if they are in phase.

If a spinning top has a perfectly circular footprint then its spin axis precesses in a perfect circle, with no nutation and the spinning does not create any waves if its environment is friction-less. If the top has an elliptical footprint then its spin axis follows a wavy circular motion, but this motion is such that the top never returns to its initial state and thus it cannot produce a stationary wave. However, if the top is only very slightly elliptical and its axis almost coincides with the total angular momentum vector then its motion can produce an almost standing wave, if some additional conditions are met.

A spinning top that is symmetrical about axis  $z$  and about  $xy$  plane has three moments of inertia  $I_x$ ,  $I_y$  and  $I_z$ , where  $I_x \approx I_y$  if its footprint is almost circular. The additional conditions are that the ratio of its moments of inertia  $I_x/I_z$  is an integer or half-integer. If the ratio is an integer then the top (almost) returns to its initial state after one precession and if it is a half-integer then after two precessions. This restricts particles to the spin that we observe.

Electro-magnetic fields for two inertial observers moving with relative velocity  $v$  are related by Lorentz transformations used in Special Relativity. If the relative velocity  $v$  between two observers is along their  $z$  axes then the  $z$  components of the electric and magnetic fields are the same for both observers, but the  $x$  and  $y$  components are different [12]. If in a “stationary frame” the electric field is absent then the “moving” frame acquires electric field with non-zero  $x$  and  $y$  components. In Vir theory this leads to the explanation why particles have electric charge.

Lorentz transformations can be expressed by hyperbolic functions of a real argument that is zero for  $v = 0$  and infinity for  $v = c$  [13]. Alternatively we can use an imaginary argument with trigonometric functions. Thus we can consider these transformations as rotations by an imaginary angle in  $zt$  plane, where axis  $z$  is aligned with the velocity of the moving observer. If the rotation in  $zt$  plane is positive the ether flows out from the ends of the Vir and we have a positive particle. If the ether flows in through the ends of the Vir we have a negative particle. If it flows in from one end and out from the other we have a neutral particle.

This completes the description of all possible Vir rotations in space-time. These rotations reveal the origin and nature of spin and electric charge in particles and lead to explanations of other particle properties.

CPT is the most fundamental conservation law observed in particle physics, but there is no explanation for this in terms of quarks. The time transformation  $T$  reverses all motion and transforms a particle to an anti-particle. Using this concept of anti-particles Vir Theory shows that CPT transformation must be invariant.

Although there are anti-protons, anti-neutrons and anti-electrons (positrons) there are no stable anti-atoms, i.e. anti-matter. The conventional thinking is that this may be due to some asymmetry that occurred very soon after the big-bang. Vir Theory explains how this comes about without resorting to the big-bang.

The integrals for the mass and the moments of inertia for the Vir can be evaluated algebraically and using these expressions one can obtain an algebraic formula for the relationship between the spin and mass of Vir. It is a formula that allows several masses for a given spin and vice-versa. This formula can be tested using the existing particle data and thus verify the correctness of Vir Theory.

## 2. Vir moments of inertia and mass

It has been proven mathematically using the Euler-Lagrange equations that the symmetric concave spinning top with the minimum moment of inertia, i.e. with the least resistance to spinning, has the shape of a solid of rotation generated by the profile function  $r(z)$  called Vir [10].

$$r(z) = a \left( \frac{a}{|z|} \right)^\alpha \quad \text{where} \quad \alpha > 0 \quad a > 0 \quad r(0) = \infty \quad (2.1)$$

The parameter  $a$  gives the size, if  $z = a$  then  $r = a$  and vice versa, while the parameter  $\alpha$  gives the shape. It has also been shown experimentally [11] that water and air vortices have the Vir shape with  $\alpha = [0.6, 2.5]$ .

Vir Theory of elementary particles assumes that particles are symmetric twin vortices in the relativistic ether, i.e. consisting of two vortices spinning about a common axis that are joined at the large ends. For the purpose of calculating the volume, mass and the moments of inertia we can treat a particle as a symmetric solid of revolution generated by the profile function  $r(z)$  in (2.1).

Since for symmetric twin vortices  $r(z) = r(-z)$  it is sufficient to consider  $r(z)$  only on the interval  $[0, Z]$  where  $Z$  is half height of the Vir. For a profile curve  $r(z)$  the expressions for mass  $m$  and the moments of inertia around axes  $z$  and  $x$ , i.e.  $I_z$  and  $I_x$  are

$$m = 2\rho\pi \int_0^Z r(z)^2 dz \quad (2.2)$$

$$I_z = \rho\pi \int_0^Z r(z)^4 dz \quad (2.3)$$

$$I_x = \rho\pi \int_0^Z \frac{1}{2} r(z)^4 + 2r(z)^2 z^2 dz \quad (2.4)$$

Since the shape of the vortices is the solid of rotation around axis  $z$  generated by the profile function  $r(z)$  the moment of inertia around axis  $y$  is the same as around axis  $x$ . The evaluation of the above integrals gives the following formulas with finite values, provided the parameter  $\alpha$  is restricted as shown below.

$$m = 2\rho\pi \frac{a^{2+2\alpha}}{1-2\alpha} Z^{1-2\alpha} \quad \alpha < 1/2 \quad (2.5)$$

$$I_z = \rho\pi \frac{a^{4+4\alpha}}{1-4\alpha} Z^{1-4\alpha} \quad \alpha < 1/4 \quad (2.6)$$

$$I_x = \rho\pi \frac{a^{2+2\alpha}}{3-2\alpha} Z^{3-2\alpha} \quad \alpha < 3/2 \quad (2.7)$$

Using the formulas (2.5), (2.6) and (2.7) above we find the following formula for mass  $m$

$$m = b(2\sigma - 1)^\beta \quad (2.8)$$

where 
$$\sigma = \frac{I_x}{I_z} \quad (2.9)$$

and 
$$\beta = \frac{1 - 2\alpha}{2 + 2\alpha} \quad 1/2 > \beta > 1/5 \quad (2.10)$$

thus 
$$\alpha = \frac{1 - 2\beta}{2 + 2\beta} \quad 0 < \alpha < 1/4 \quad (2.11)$$

while 
$$b = 2\rho\pi a^3 \frac{1}{1 - 2\alpha} \left( \frac{1 - 3 - 2\alpha}{4 - 1 - 4\alpha} \right)^\beta \quad (2.12)$$

When we know the parameters  $b$  and  $\beta$  we can find the corresponding parameters  $a$  and  $\alpha$

$$a = \left( \frac{3b}{2\rho\pi} \frac{\beta}{1 + \beta} \right)^{\frac{1}{3}} \left( \frac{4 - 5\beta - 1}{5\beta + 2} \right)^{\frac{\beta}{3}} \quad (2.13)$$

Using the formulas (2.6), (2.7) for  $I_x$ ,  $I_z$  we can obtain the expression for  $\sigma$  in (2.9) in terms of  $a$ ,  $\alpha$  and  $Z$ . Then using the formula (2.5) for mass  $m$  we obtain the formula for  $Z$

$$Z = a \left( \frac{3 - 2\alpha}{1 - 4\alpha} \frac{2\sigma - 1}{4} \right)^{\frac{1}{2 + 2\alpha}} \quad (2.14)$$

Since  $\alpha$  and  $\sigma$  are dimensionless quantities we have the height  $Z$  in the units of the parameter  $a$ . The radius  $r$  in the formula (2.1) is also in units of  $a$ . In all the formulas shown in this section it is assumed that the radius  $r$  is infinite at  $z = 0$ . This is of course unrealistic, for a particle cannot span the whole of the universe. The infinite radius is assumed here because it leads to much simpler formulas and because it may give a sufficiently good approximation to our needs, but this will have to be found out experimentally.

### 3. Euler's equations of motion

To find the relationship between the moments of inertia and spin we start by investigating Euler's equations of motion for a solid rotating body [8]. We consider an inertia frame  $XYZ$  and a spinning top with the angular momentum vector  $\mathbf{M}$ . Since  $\mathbf{M}$  is a constant of motion we can align axis  $Z$  with  $\mathbf{M}$ . We also have a frame  $xyz$  with the origin at the centre of mass of the spinning top and axis  $z$  coincides with the spin axis.

From the Euler's equations of motion we find that a symmetrical spinning top, i.e. one with the moments of inertia  $I_x = I_y$ , rotates (spins) about its axis  $z$  with a constant angular velocity  $\Omega_z$ . Spin vector  $s$  precesses at a constant angular velocity  $\Omega_{Pr}$  and at a constant angle  $\theta$  relative to the total angular momentum  $\mathbf{M}$ .

The movement of axis  $z$  creates the surface of a cone and the end of the spin vector  $s$  describes a circle in a plane perpendicular to vector  $\mathbf{M}$ . The assumptions (3.1) and the solution (3.2) are below [14]:

$$\vec{M} = \text{constant} \quad I_x = I_y \quad \vec{I} = \text{constant} \quad (3.1)$$

$$\Omega_z = \frac{M}{I_z} \cos \theta \quad \theta = \text{constant} \quad \Omega_{Pr} = \frac{M}{I_x} \quad (3.2)$$

From (3.2) we find the following relationship between moments of inertia and angular velocities:

$$\frac{\Omega_z}{\Omega_{Pr}} = \frac{I_x}{I_z} \cos \theta \quad (3.3)$$

From the definition of the moments of inertia we have for any body the following inequality:

$$I_x + I_y \geq I_z \quad \text{equality for a planar body} \quad (3.4)$$

For a symmetrical top we have  $I_x = I_y$  and therefore from (3.4) we get the following restriction:

$$\frac{I_x}{I_z} \geq \frac{1}{2} \quad \text{for any symmetrical top} \quad (3.5)$$

There is a bottom limit of  $1/2$  on the ratio  $I_x/I_z$  for a symmetrical top. The limit is reached only by an idealized planar body, for a true three dimensional body the ratio is always greater than  $1/2$ . Now, from equation (3.3) we find that for very small angle  $\theta$  we have the following relationship:

$$\frac{\Omega_z}{\Omega_{Pr}} = \frac{I_x}{I_z} \quad \text{as } \theta \rightarrow 0 \quad (3.6)$$

Therefore, there is a bottom limit on the rotation velocity  $\Omega_z$  relative to the precession velocity  $\Omega_{Pr}$  of  $1/2$  as  $\theta$  tends to zero. There isn't any body that can rotate slower relative to precession.

#### 4. Return of the symmetric top to the initial state

Euler's equations give us also an expression for the angular velocity -  $\Omega_\zeta$  (note the minus sign) with which vector  $\mathbf{M}$  orbits an observer that is attached to the rotating body [15].

$$\Omega_\zeta = \Omega_z - \Omega_{Pr} \cos \theta \quad (4.1)$$

An observer attached to the top in such a way that in the initial state is facing vector  $\mathbf{M}$  will face vector  $\mathbf{M}$  again on completion of one precession if  $\Omega_\zeta/\Omega_{Pr} = n$ , where  $n$  is 0, 1, 2, 3, etc. The number  $n$  tells us how many times the observer will face vector  $\mathbf{M}$  during one precession, including the final state, but excluding the initial state. When  $n = 0$  then  $\Omega_\zeta = 0$  and the observer is facing vector  $\mathbf{M}$  all the time, never see any change. From (4.1) we get two equivalent return conditions:

$$\frac{\Omega_\zeta}{\Omega_{Pr}} = n \quad \frac{\Omega_z}{\Omega_{Pr}} - \cos \theta = n \quad n = 0, 1, 2, \text{ etc.} \quad (4.2)$$

From (3.3) and (4.2) we find another condition for the return to the initial state after one precession:

$$\left( \frac{I_x}{I_z} - 1 \right) \cos \theta = n \quad (4.3)$$

For very small precession angle  $\theta$  we get  $I_x/I_z = n + 1$  from (4.3), which means that the ratio  $I_x/I_z$  takes values of 1, 2, 3, 4 etc. Thus, when  $\theta$  is very small the top returns to its initial state on completion of one precession under the following conditions:

$$\frac{I_x}{I_z} = 1, 2, 3, \text{ etc. as } \theta \rightarrow 0 \quad (4.4)$$

The conditions for returning to its initial state only on completion of the second precession are:

$$\frac{\Omega_\zeta}{\Omega_{Pr}} = \frac{2n-1}{2} - 1 \quad \frac{\Omega_z}{\Omega_{Pr}} - \cos \theta = \frac{2n-1}{2} - 1 \quad n = 1, 2, 3, \text{ etc.} \quad (4.5)$$

$$\left( \frac{I_x}{I_z} - 1 \right) \cos \theta = \frac{2n-1}{2} - 1 \quad (4.6)$$

Number  $|2n-3|$  tells us how many times the observer will face vector  $\mathbf{M}$  during the first two precessions, including the final state, but excluding the initial state. This time there is no solution for  $n = 0$  since that would require negative  $I_x/I_z$ . For  $n = 1$  we have  $\Omega_\zeta/\Omega_{Pr} = -1/2$  which means that the observer orbits vector  $\mathbf{M}$  in the opposite direction to all other values of  $n$ . Thus, when  $\theta$  is very small the top returns to its initial state only on completion of the second precession under the following conditions:

$$\frac{I_x}{I_z} = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \text{ etc. as } \theta \rightarrow 0 \quad (4.7)$$

Figures 1 to 4 below show graphically the return to the initial state of the symmetric spinning top with a square footprint after one precession. The initial state is shown on the extreme right of each diagram. The condition for this return to occur is that  $\Omega_{\zeta}/\Omega_{Pr} = n$ , where  $n$  is 0, 1, 2, 3, etc. For the angle  $\theta \rightarrow 0$  this is equivalent to the condition  $I_x/I_z = 1, 2, 3, 4$ , etc.




Figure 1.  $\Omega_{\zeta}/\Omega_{Pr} = 0$




Figure 2.  $\Omega_{\zeta}/\Omega_{Pr} = 1$




Figure 3.  $\Omega_{\zeta}/\Omega_{Pr} = 2$






Figure 4.  $\Omega_{\zeta}/\Omega_{Pr} = 3$

The diagrams show the movement of the spinning top in the XY plane of the inertial frame, where the axis Z is aligned with the angular momentum  $M$ . The top spins anticlockwise and the precession is in the same direction, as indicated by the double arrows. The arrows inside the squares show the orientation of the observer on the spinning top and the bold arrows show when the observer faces directly vector  $M$ .


The figures 5 to 8 below show graphically the return to the initial state of the symmetric spinning top with a square footprint after two precessions. The condition for this to occur is that  $2\Omega_c/\Omega_{Pr} = n$ , where  $n$  is  $-1, 1, 3, 5$ , etc. For the angle  $\theta \rightarrow 0$  this is equivalent to the condition  $I_x/I_z = 1/2, 3/2, 5/2, 7/2$ , etc.




**Figure 5.**  $2\Omega_c/\Omega_{Pr} = -1$



**Figure 6.**  $2\Omega_c/\Omega_{Pr} = 1$



**Figure 7.**  $2\Omega_c/\Omega_{Pr} = 3$




**Figure 8.**  $\Omega_c/\Omega_{Pr} = 5$

The top spins anticlockwise, except for figure 5, and the precession is also anticlockwise as indicated by the double arrows. The diagrams show only the first precession, but we can visualise the second precession by turning the arrows in the boxes to face the opposite direction. The arrows in red show the state where the observer on the spinning top faces directly vector  $M$  on the second precession.

### 5. Slightly asymmetric almost vertical spinning top

A spinning top is called asymmetrical when  $I_x \neq I_y$ , it may for example have an elliptical footprint in the  $xy$  plane. It does not mean that there is asymmetry between the upper and lower halves. An asymmetrical spinning top never returns to its initial state [16]. The angle  $\theta$  and all angular velocities  $\Omega$  vary periodically in time. As  $\theta$  increases, vectors  $\Omega$  decrease and change not only their magnitudes but also their directions.

The unit vector  $\mathbf{z}$  that starts in the centre of the spinning top is aligned with the spin vector  $\mathbf{s}$  and moves on the surface of a sphere between two concentric circles, lying in two parallel planes perpendicular to the vector  $\mathbf{M}$  aligned with axis  $\mathbf{Z}$  [17]. This wavy circular movement, known as nutation, is shown in figure 9.



**Figure 9.** The trajectory of the spin vector  $\mathbf{s}$ .

Looking from above we see the trajectory of the end of the spin vector  $\mathbf{s}$  as a periodic wave confined between two concentric circles. The trajectory touches the inner circle when  $\theta$  is at its minimum and the outer circle when  $\theta$  is at its maximum. The variation  $\Delta\theta = \frac{1}{2}(\max \theta - \min \theta)$  decreases with  $\Delta I = |I_x - I_y|$  and disappears when  $I_x = I_y$ .

Formula (4.1) for  $\Omega_\zeta$  still holds at any instant of time, i.e. only for the instantaneous angular velocities. When the spinning top is almost vertical ( $\theta \rightarrow 0$ ) then the formula is almost linear and taking the average of the angular velocities over one precession we get

$$\overline{\Omega_\zeta} = \overline{\Omega_z} - \overline{\Omega_{Pr} \cos \theta} \quad (5.1)$$

The average  $\Omega_\zeta$  may be different for each precession, but only very slightly when  $I_x \approx I_y$ . Thus the equations (4.2) to (4.7) for the return of symmetrical spinning tops to the initial state hold approximately also for slightly asymmetric and almost vertical spinning tops. For the return after one or two precession we have

$$\frac{I_x}{I_z} = 1, 2, 3, \text{ etc.} \quad \frac{I_x}{I_z} = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \text{ etc.} \quad \text{for } I_x \approx I_y \text{ and } \theta \rightarrow 0 \quad (5.2)$$

Only ideal symmetric spinning tops ( $I_x = I_y$ ) can return to their initial state. Real spinning tops, represented in figures 10 to 13 by a top with a rectangular footprint, never return to the initial state. The position of a top at 0, 360 degrees and multiples thereof (in polar coordinates) are different. This is indicated by a gap in the trajectory of the spin axis close to 360 degrees. However, such spinning tops can come close to the initial state if  $I_x \approx I_y$  and  $\theta \rightarrow 0$ . The condition for this to occur after one precession is  $I_x/I_z = 1, 2, 3, \text{ etc.}$




Figure 10.  $\Omega_\zeta/\Omega_{Pr} = 0$




Figure 11.  $\Omega_\zeta/\Omega_{Pr} = 1$




Figure 12.  $\Omega_\zeta/\Omega_{Pr} = 2$





Figure 13.  $\Omega_\zeta/\Omega_{Pr} = 3$

As with the symmetrical spinning tops the arrows inside the rectangles show the orientation of the observer on the spinning top and the bold arrows show when the observer faces directly vector  $M$ .

Figures 14 to 17 show the trajectory of the end of the spin vector  $s$  and the movement of the spinning top. The condition for the top returning to the initial state only after two precessions is  $I_x/I_z = 1/2, 3/2, 5/2, \text{ etc.}$  Again the gap in the trajectory of the spin vector close to 360 degrees indicates that this is only approximate.

$$2\Omega_\zeta/\Omega_{Pr} = -1$$


$$I_x/I_z = 1/2$$



**Figure 14.**  $2\Omega_\zeta/\Omega_{Pr} = -1$

$$2\Omega_\zeta/\Omega_{Pr} = 1$$


$$I_x/I_z = 3/2$$



**Figure 15.**  $2\Omega_\zeta/\Omega_{Pr} = 1$

$$2\Omega_\zeta/\Omega_{Pr} = 3$$


$$I_x/I_z = 5/2$$



**Figure 16.**  $2\Omega_\zeta/\Omega_{Pr} = 3$

$$2\Omega_\zeta/\Omega_{Pr} = 5$$

$$I_x/I_z = 7/2$$



**Figure 17.**  $\Omega_\zeta/\Omega_{Pr} = 5$

The top spins anticlockwise, except for figure 14, and the precession is also anticlockwise as indicated by the double arrows. The diagrams show only the first precession, but we can visualise the second precession by turning the arrows in the boxes to face the opposite direction. The arrows in red show the state where the observer on the spinning top faces directly vector  $M$  on the second precession.

## 6. Elementary particles as stationary circular waves

Euler's equations of motion do not require that the body must be rigid, i.e. that parts of the body remain in constant distance from each other, the conditions are much more relaxed. In the absence of external forces, any internal forces cancel out and the centre of mass of any body moves at a constant velocity. The angular momentum  $\mathbf{M}$  also remains constant. The moments of inertia remain constant not only for rigid bodies, but for a varied length of time also for deformable bodies. Hurricanes retain their shape for days or at least hours, tornados for minutes and dust devils for tens of seconds. During this time Euler's equations apply.

From formulas (4.2) and (4.3) we know that a symmetrical top ( $I_x = I_y$ ) returns to its initial state after one or two precessions if the following equivalent conditions are satisfied

$$\frac{\Omega_\zeta}{\Omega_{Pr}} = \frac{\Omega_z}{\Omega_{Pr}} - \cos \theta = \left( \frac{I_x}{I_z} - 1 \right) \cos \theta = -\frac{1}{2}, 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \text{etc...} \quad (6.1)$$

However, for a perfect circular top the axis  $z$  follows a perfect circular cone and in the absence of friction its movement does not generate any waves in its environment. If such an absolutely perfect top exists in the relativistic ether then it cannot be detected as a wave and thus it is not a suitable model of an elementary particle. An alternative, and a more appropriate description of elementary particles, is provided by considering asymmetrical tops ( $I_x \neq I_y$ ). We shall assume that in nature there is always some asymmetry and consider particles as near symmetrical ( $I_x$  is close to  $I_y$ ) and near vertical ( $\theta$  is close to 0) spinning tops.

For an asymmetrical top the axis  $z$  follows a periodic wavy trajectory so that the top generates circular waves in its environment. If the top is not rigid then the waves that it generates interfere with the top itself. The top effectively becomes a circular wave that is chasing its tail, as shown in figure 9. The head interferes with the tail in a positive or a negative way depending on the phase difference. If the head and tail are in opposite phases after one precession then the top destroys itself. If there were no phase difference at all we would get a stationary circular wave.

Using formula (6.1) we find that a slightly asymmetric almost vertical spinning top almost returns to its initial state after one or two precessions if the following equivalent conditions are approximately satisfied

$$\frac{\Omega_\zeta}{\Omega_{Pr}} + 1 = \frac{\Omega_z}{\Omega_{Pr}} = \frac{I_x}{I_z} \quad \rightarrow \quad \frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}, 3, \text{etc...} \quad \text{for } I_x \rightarrow I_y \text{ and } \theta \rightarrow 0 \quad (6.2)$$

It may be tempting to think that if the ratio  $I_x/I_z$  were slightly greater, to compensate for  $\cos(\theta)$  being less than one, then the formula above would result in  $\Omega_\zeta/\Omega_{Pr}$  becoming an integer or half-integer. This would mean that asymmetric spinning tops would return to their initial state, which we know to be impossible. However, an asymmetric top can almost return to its initial state and create an almost stationary wave. The closer it returns to the initial state the longer it will take for the top to destroy itself.

## 7. Spin of particles

The spin of a particle  $M_z$  is the  $z$  component of the total angular momentum  $\mathbf{M}$  for which we have

$$M_z = I_z \Omega_z \quad (7.1)$$

where  $I_z$  is the moment of inertia about  $z$  axis and  $\Omega_z$  the angular velocity about  $z$  axis. Since the angular velocity of precession  $\Omega_{Pr}$  is never zero we can write the above formula as

$$M_z = I_z \Omega_{Pr} \left( \frac{\Omega_z}{\Omega_{Pr}} \right) \quad (7.2)$$

In Quantum Mechanics for the spin of a particle we have the following formula involving the Heisenberg constant

$$M_z = \hbar \sigma \quad \sigma = \text{integer or half-integer} \quad (7.3)$$

where  $\sigma$  is an integer or half-integer. Let us assume that the following relationship between  $I_z$  and  $\Omega_z$  holds

$$\hbar = I_z \Omega_{Pr} \quad (7.4)$$

This simply says that a spinning top with a small resistance to spinning (small  $I_z$ ) precesses faster than a spinning top with a high resistance to spinning (large  $I_z$ ) and that this relationship is inversely proportional. On the assumption (7.4) the formula (7.2) for the spin of a particle becomes

$$M_z = \hbar \left( \frac{\Omega_z}{\Omega_{Pr}} \right) \quad (7.5)$$

Formula (6.2) for the possible values of the ratio  $\Omega_z/\Omega_{Pr}$  tells us

$$\frac{\Omega_z}{\Omega_{Pr}} \rightarrow \frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}, 3, \text{etc...} \quad \text{for } I_x \rightarrow I_y \text{ and } \theta \rightarrow 0 \quad (7.6)$$

Thus we arrive at an explanation why the spin of elementary particles  $\sigma$  is an integer or a half-integer. Particles are disturbances in the relativistic ether that last long enough for us to observe. For this they have to be almost stationary circular waves and hence have  $\Omega_z/\Omega_{Pr}$  that is equal to an integer or a half-integer, which according to (6.2) is equivalent to  $I_x/I_z$  being an integer or a half-integer.

## 8. Baryons, mesons and their mass

The spin values  $I_x/I_z$  in (6.1) do not include 0 and for  $\frac{1}{2}$  we have a disc that has virtually no mass and hence cannot be a hadron. We address this below and in the process explain the structure of baryons and mesons. The mass formula (2.8) can be modified to take into account the structure of particles. Figures 18 and 19 show a baryon with  $\sigma = 2.5$  and a meson with  $\sigma = 3$ . The parameter  $\sigma$  is the spin of a particle when the entire particle rotates in the same direction. The horizontal lines show the heights for baryons with  $\sigma = 0.5, 1.5, 2.5$  and mesons with  $\sigma = 1, 2$  and 3. The line through the centre of the baryon is a disc with spin 0.5.




Figure 18. Baryon




Figure 19. Meson

To obtain a baryon with spin 0.5 we take the height and mass of  $\sigma = 1.5$ , but with the disc rotating in the opposite direction. In the absence of the disc the spin would be reduced by 0.5 and with the disc rotating in the opposite direction the spin is reduced by another 0.5. We find a similar phenomenon to the baryon disc in hurricanes where the visible part (warm air) rises up from the ground (or sea) rotating anticlockwise, but when it ascends above the hurricane it suddenly turns and rotates clockwise. We can extend this idea, by rotating not only the disc, but also its adjacent parts in the direction opposite direction to the end parts. This way we can obtain any half integer spin  $s$  smaller than  $\sigma$  with the mass of  $\sigma$ .

We can do the same for mesons, although in the absence of the baryon disc we have to start with an even  $\sigma$  to obtain spin 0, if we want a symmetrical arrangement of the top and bottom. If we are prepared to accept asymmetrical particles, for example the top rotating in the opposite direction to the bottom, then we can start with any  $\sigma$ .

The horizontal lines in figures 1 and 2 separate the particles into “slices”, each contributing with spin  $\frac{1}{2}$ . The slices that rotate in the opposite direction to the particle with spin  $s$  are called “contras” and the number of contras is denoted by the variable  $c$ . Thus the relationship between the variables  $\sigma$ ,  $s$  and  $c$  is as follows:  $s = \sigma - c$ , i.e.  $\sigma = s + c$ . Substituting this into the mass formula (2.8) we obtain

$$m = b(2s + 2c - 1)^\beta \quad (8.1)$$

The formula above allows one family of particles to have particles with the same mass but different spin and vice-versa, for a given spin there may be particles with different masses.

## 9. Electric charge of particles

In Special Relativity the electro-magnetic field is transformed between two inertial observers using Lorentz transformations. If the relative velocity  $\mathbf{v}$  between the observers is along their  $z$  axis then the electric field  $E_z$  and the magnetic field  $H_z$  are the same for both observers, but the  $x$  and  $y$  components are different [12].

$$E_x = \frac{E'_x + \frac{v}{c} H'_y}{\sqrt{1 - \frac{v^2}{c^2}}} \quad E_y = \frac{E'_y - \frac{v}{c} H'_x}{\sqrt{1 - \frac{v^2}{c^2}}} \quad E_z = E'_z \quad (9.1)$$

The same formulas are for the magnetic field  $H$ , but the  $+$  and  $-$  signs in the nominator are swapped round. Let us assume that in a “stationary frame” the values of  $E_x'$ ,  $E_y'$  and  $E_z'$  are zero. Then the frame moving in the direction of axis  $z$  acquires electric field that demonstrates itself in non-zero  $E_x$  and  $E_y$ .

Lorentz transformations are also used for the space-time coordinates, with the following formulas [13]

$$z = \frac{z' + vt'}{\sqrt{1 - \frac{v^2}{c^2}}} \quad ct = \frac{ct' + (v/c)z'}{\sqrt{1 - \frac{v^2}{c^2}}} \quad x = x' \quad y = y' \quad (9.2)$$

These transformations can also be expressed using the hyperbolic functions  $\sinh$  and  $\cosh$  [13]

$$z = z' \cosh(\omega) + ct' \sinh(\omega) \quad ct = z' \sinh(\omega) + ct' \cosh(\omega) \quad (9.3)$$

The variable  $\omega$  is related to velocity  $v$  as shown below [13], [18] thus  $\omega = 0$  for  $v = 0$  and  $\omega = \infty$  for  $v = c$

$$\tanh(\omega) = v/c \quad \omega = \frac{1}{2} \ln \frac{1 + v/c}{1 - v/c} \quad (9.4)$$

If  $v$  is much smaller than  $c$  then we can expand the above formula for  $\omega$  in terms of  $v/c$  as follows

$$\omega \cong \frac{v}{c} + \frac{1}{3} \left( \frac{v}{c} \right)^2 + ect \quad \text{i.e.} \quad \omega \cong \frac{v}{c} \quad \text{for } v \ll c \quad (9.5)$$

Yet another way to express the transformations is by geometric functions with imaginary argument  $i\omega$  [19]


$$z = z' \cos(i\omega) - ict' \sin(i\omega) \quad ct = -iz' \sin(i\omega) + ct' \cos(i\omega) \quad (9.6)$$

In this form Lorentz transformations may be considered as rotations in  $xt$  plane. We have now explored all possible rotations in space time and we will show on the next page that they can provide an explanation not only for the spin of particles which was already explained, but also explain the electric charge.


Let us consider vortices and in particular the structure and the air flow of a hurricane. The hot air produced by the heat from land or sea rises up in a spiral twisting about the hurricane's eye. The cold air found at high altitudes spirals just above the visible hurricane cloud in an almost flat invisible disc. When it reaches the hurricane's eye it suddenly falls down at almost a constant stream to fill the void left by the hot air.

Let us now consider particles. In Lorentz transformations the electric field can be created only if the magnetic field is already present. Thus, let us assume that in the case of particles the spiral flow of the relativistic ether in  $xy$  plane creates the magnetic field and then the linear constant flow of the ether along the spin axis  $z$  creates the electric charge. More accurately perhaps, the two distinctly different flows of ether together create the electro-magnetic field and the electric charge.

Figures 20 and 21 show two man-made water vortices. One vortex is made by rotating a disc with a hole in its centre close to the bottom of a water container. The other vortex uses a couple of bottles firmly connected together at their necks. The bottles are half full with water and are given some angular momentum by a circular movement using hands. Then the coupled bottles are placed in a holder to keep them steady.



**Figure 20.** Air flows down the eye



**Figure 21.** Air flows up the eye

The main difference between the two vortices is that in figure 20 the air comes down the vortex eye as in the case of a hurricane. This can be seen by the bubbles rising from the bottom of the container where the disc rotates. On the other hand in figure 21 the air comes up the vortex eye as the water is filling the bottom bottle and displaces the air within it. This is not what is observed on water vortices in nature, but the experiment shows that it is possible.


Thus we assume that a particle with the ether coming out from the narrow ends has a positive electric charge of 1 unit, while a particle where the ether comes in through the narrow ends has a negative charge -1. If the ether comes in from one end and goes out from the other end then the particle has zero charge. In addition to the vertical flow of ether there is a spiral flow of ether where the twin vortices are joined at their wide ends. Thus the ether that comes in through the wide end goes out through the narrow end and vice-versa.

This scheme satisfies the fundamental condition regarding anti-particles, namely that they can be described by the same equations as the particles, but with the reversed time axis. We can imagine this by filming the whole process and then playing the film backwards.


## 10. Hurricanes and Typhoons

We will now take a detailed look at the basic building blocks of particles and represent them by hurricanes and typhoons. For simplicity consider that hurricanes are confined to the northern hemisphere and typhoons to the southern hemisphere, rotating the opposite way. A hurricane has an eye that opens up to a wide spiral, as shown in figure 23. This spiralling cloud is fed by the hot air rising up from the ground or the sea and twisting round the eye, as shown in figure 22. The whole visible cloud rotates anti-clock wise.

In addition not seen on the photographs is the cold air falling down the eye. Also not seen on the photographs, there is a transparent accretion disc just above the clouds that brings the cold air to the edge of the eye. The disc rotates in the opposite direction to the cloud, i.e. clockwise, as shown in figure 23. Since the cold air is not visible its precise circulation is not known.




**Figure 22.** Hurricane top view




**Figure 23.** Hurricane side view

Schematically we shall represent hurricanes and typhoons by drawings shown in figures 22 and 23.



**Figure 24.** Hurricane diagram.




**Figure 25.** Typhoon diagram.

The wall of the hurricane eye may be thought of as a right-handed screw. If we hold the screw still pressing down on a piece of wood and rotate the wood, it will go up the screw. The hot air moves by a right hand rule, with the thumb up and fingers bent pointing anti-clockwise. Similarly, we can think of the typhoon wall as a left-handed screw (seldom used in practice). The hot air twists up the wall, but this time clockwise.


Anti-hurricanes do not occur on Earth, but we can imagine them as a film of a hurricane played backwards in time. In an anti-hurricane the eye is still while the hot air twists clockwise and moves down. At the same time a column of cold air is rising up through the middle of the eye. Similar time reversal takes place in anti-typhoons. Anti-hurricane is not a typhoon and anti-typhoon is not a hurricane.

### 11. Diagrams of h and t vortices


The schematic drawings of hurricanes and typhoons are shown in figures 26 to 33, where the subscripts “c” and “a” stand for clock-wise and anti-clock-wise. In reality the eye has more or less the same height as the width and flattens out much more suddenly, as given by formula (2.1). The vortices spinning anti-clockwise have the spin  $s$  of  $+1/2$  and the those spinning clockwise have the spin of  $-1/2$ . The electric charge  $q$  is  $+1/2$  if the stream through the eye exits via the narrow end, and  $-1/2$  if it enters at the narrow end.




**Figure 26.**  $h_c \quad s = -1/2 \quad q = 1/2$



**Figure 27.**  $t_a \quad s = 1/2 \quad q = 1/2$




**Figure 28.**  $h_a \quad s = 1/2 \quad q = 1/2$




**Figure 29.**  $t_c \quad s = -1/2 \quad q = 1/2$


The vortices h, t (above) have the electric charge  $+1/2$ , while anti-h, anti-t (below) have the charge  $-1/2$ .




**Figure 30.**  $anti-h_a \quad s = 1/2 \quad q = -1/2$



**Figure 31.**  $anti-t_c \quad s = -1/2 \quad q = -1/2$




**Figure 32.**  $anti-h_c \quad s = -1/2 \quad q = -1/2$




**Figure 33.**  $anti-t_a \quad s = 1/2 \quad q = -1/2$

**12. Diagrams of the baryon disc**


The images below are the invisible discs of hurricanes, typhoons and their imaginary anti-partners. For hurricanes and typhoons the discs spin in the opposite direction to the visible clouds of the hot air and bring the cold air to the vortex eye, where the air falls down. The baryon discs do not carry any electric charge. The discs are the spiral inlets of ether for the positive particles and the outlets for the negative particles. The discs  $i_c$  and  $o_a$  ( $i$  = inlet,  $o$  = outlet) are anti-vortices annihilating each other, as are  $o_c$  and  $i_a$ .




**Figure 34.** Inlet  $i_c$   $s = -1/2$



**Figure 35.** Outlet  $o_a$   $s = +1/2$




**Figure 36.** Outlet  $o_c$   $s = -1/2$




**Figure 37.** Inlet  $i_a$   $s = +1/2$

For the neutral particles one side of the disc is an outlet of ether and the other side an inlet of ether. Both disc sides spin in the same direction as one unit with spin  $\pm 1/2$ , there is no net flow of ether. The discs  $io_a$  and  $oi_c$  ( $io$  = inlet-outlet) are anti-vortices annihilating each other, as are  $oi_a$  and  $io_c$ .



**Figure 38.** In-Out $_c$  and Out-In $_c$




**Figure 39.** In-Out $_a$  and Out-In $_a$

This scheme satisfies the fundamental condition regarding anti-particles, namely that they can be described by the same equations as the particles, but with the reversed time axis. We can imagine this by filming the whole process and then playing the film backwards.


**13. Diagrams of proton and neutron**

The notation for baryons is a column containing three entries, names of the vortices, as shown in (13.1). This reflects the structure of a proton and an anti-proton in figures 40 and 41. A proton spinning anti-clockwise consist of vortex  $h_a$  in the down entry,  $t_a$  in the up entry and  $i_c$  in the middle. The anti-proton is shown to spin the opposite way and consists of anti- $h_c$  down, anti- $t_c$  up and anti- $i_a$  in the middle, that may be also denoted as  $o_a$ . Figures 44 and 43 show the structure of the discs from the bird's eye view.


$$p_a = \begin{bmatrix} t_a \\ i_c \\ h_a \end{bmatrix} \qquad \bar{p}_c = \begin{bmatrix} \bar{t}_c \\ \bar{i}_a \\ \bar{h}_c \end{bmatrix} \qquad (13.1)$$




**Figure 40.** Proton



**Figure 41.** Anti-proton



**Figure 42.** Proton inlet disc  $i_c$



**Figure 43.** Anti-proton outlet disc  $o_a$

On the diagrams the spin of the proton is  $+1/2$  and that of the anti-proton is  $-1/2$ . Turning them upside down each would spin in the opposite direction. In a collision of these two particles the magnetic field that they generate make them aligned in such a way that they spin in the opposite direction and annihilate each other.

There are two varieties of neutrons, one consisting of vortex h and anti-h, the other of vortex t and anti-t. The notation for the neutron shown in (13.2) is for the h variety. This reflects the structure of a neutron h and an anti-neutron h in figures 44 and 45. A neutron h spinning anti-clockwise consist of vortex  $h_a$  in the down position,  $\bar{h}_a$  in the up position and  $oi_c$  in the middle. The anti-neutron h is shown to spin the opposite way, i.e. clockwise and consists of vortex  $\bar{h}_c$  down,  $h_c$  up and  $io_a$  in the middle.

$$nh_a = \begin{bmatrix} \bar{h}_a \\ oi_c \\ h_a \end{bmatrix}$$

$$\bar{nh}_c = \begin{bmatrix} h_c \\ io_a \\ \bar{h}_c \end{bmatrix} \quad (13.2)$$




Figure 44. Neutron h




Figure 45. Anti-neutron h

Figures 46 and 47 show the structure of the discs from the bird's eye view. The baryon disc  $io_a$  is a part of a neutral particle and is equivalent to anti- $oi_c$ .




Figure 46. Neutron h disc  $oi_c$




Figure 47. Anti-neutron h disc  $io_a$

The spin of the neutron h is  $+1/2$  and the spin of the anti-neutron h is  $-1/2$ . However, turning either of them upside down they would spin in the opposite direction. In a collision of these two particles they align in such a way that they spin in the opposite direction and annihilate each other.

The notation for the neutron shown in (13.3) is for the t variety. This reflects the structure of a neutron t and an anti-neutron t in figures 48 and 49. A neutron t spinning clockwise consist of vortex  $t_c$  in the down position, anti- $t_c$  in the up position and  $oi_a$  in the middle. The anti-neutron t is shown to spin the opposite way, i.e. anti-clockwise and consists of vortex anti- $t_a$  down,  $t_a$  up and  $io_c$  in the middle.

$$nt_c = \begin{bmatrix} \bar{t}_c \\ oi_a \\ t_c \end{bmatrix} \qquad \bar{nt}_a = \begin{bmatrix} t_a \\ io_c \\ \bar{t}_a \end{bmatrix} \qquad (13.3)$$




Figure 48. Neutron t




Figure 49. Anti-neutron t

Figures 50 and 51 show the structure of the discs from the bird's eye view. The baryon disc  $io_c$  is a part of a neutral particle and is equivalent to anti- $oi_a$ .




Figure 50. Neutron t disc  $oi_a$





Figure 51. Anti-neutron t disc  $io_c$

The spin of the neutron t is  $-1/2$  and the spin of the anti-neutron t is  $+1/2$ . This is opposite to the neutron h and anti-neutron h in figures 46 and 47. For both neutrons h and t the ether flow is down, hence they have different chirality, one is right handed and the other left handed.


If we remove the baryon discs for all the drawings of protons and neutrons we obtain two mesons, one positive and one neutral, say  $\rho^+$  and  $\rho^0$ . Thus there will be two versions of  $\rho^0$ , right handed and left handed.

**14. CPT invariance of Particles**

CPT stands for charge, parity and time transformation of particles. It always returns to the original particle. It is the most fundamental conservation law observed in particle physics, but there is no explanation for this in terms of quarks. We shall now use proton to demonstrate its invariance under CPT transformations.




**Figure 52.** Proton anti-clock-wise ( $p_a$ )




**Figure 53.** Inlet disc clock-wise

To apply C transformation we change the direction of the ether flow, keeping the directions of all rotations.




**Figure 54.**  $C(p_a)$  = anti-proton anti-clock-wise




**Figure 55.** Outlet disc clock-wise

To apply P transformation we change the handedness of particles, i.e. rotate them by 180 degrees about z axis and take the mirror image in yz plane. Rotation is not needed, since particles are symmetrical about z.




**Figure 56.**  $CP(p_a)$  = anti-proton clock-wise



**Figure 57.** Outlet disc anti-clock-wise

To apply T transformation we reverse the time, i.e. reverse all the arrows, leaving the rest untouched. This way the last images return back to the original proton images above, and therefore proton is CPT invariant.


Next we shall use neutron h to demonstrate its invariance under CPT transformations.



**Figure 58.** Neutron h anti-clock-wise ( $nh_a$ )

**Figure 59.** Neutron h disc out-in-c


To apply C transformation we change the direction of the ether flow, keeping the directions of all rotations.



**Figure 60.**  $C(nh_a) =$  anti-neutron h anti-clock-wise

**Figure 61.** Neutron h disc in-out-c

To apply P transformation we change the handedness of particles, i.e. rotate them by 180 degrees about z axis and take the mirror image in yz plane. Rotation is not needed, since particles are symmetrical about z.




**Figure 62.**  $CP(nh_a) =$  anti-neutron h clock-wise

**Figure 63.** Neutron h disc in-out-a

To apply T transformation we reverse the time, i.e. reverse all the arrows, leaving the rest untouched. This way the last images return back to the original images above, and therefore neutron h is CPT invariant. For neutron t the figure headings above are identical, there is no need to demonstrate this separately. For the neutral mesons we simply remove the baryon disc, there is no need to demonstrate this graphically either.


## 15. Matter and anti-matter

All stable matter known to us is made of atoms and they in turn consist of protons, neutrons and electrons. Although there are anti-protons, anti-neutrons and anti-electrons (positrons) there are no stable anti-atoms, i.e. anti-matter. The conventional thinking is that this is due to some asymmetry that occurred very soon after the big-bang. We will now propose an alternative using the diagrams in figure 64 and 65 below.



**Figure 64.** Matter

The protons above are in blue, the neutrons in black. The inner jets from the protons push the ether to the centre of the atom, while the neutrons suck it out. All outer jets face away from the centre, pushing all particles to the centre and keeping the atom together and stable.



**Figure 65.** Anti-matter

The anti-protons above are in red, the anti-neutrons in black. The inner jets from the anti-proton suck out the ether from the centre and the anti-neutron replenish it. All outer jets face towards the centre, pulling all particles away from the centre and the anti-atom does not stay together.

## 16. Summary and conclusions

It has been shown by solving the Euler equations of motion for rotating bodies that the motion of a Vir can result in a circular wave, where the head chases its tail. Such a wave destroys itself after one precession if the head and tail are out of phase, but results in a stationary circular wave if they are in phase.

If a spinning top has a perfectly circular footprint then its spin axis precesses in a perfect circle. If the top has an elliptical footprint then its spin axis follows a wavy circular motion that the top never returns to its initial state. However, if the top is only slightly elliptical and its axis almost coincides with the total angular momentum vector then its motion can produce an almost standing circular wave.

The standing circular wave occurs if the ratio of the Vir moments of inertia  $I_x/I_z$  is an integer or half-integer. If the ratio is an integer then the top (almost) returns to its initial state after one precession and if it is a half-integer then after two precessions. This restricts particles to the spin that we observe.

Electric and magnetic fields for two inertial observers moving with relative velocity  $v$  are related by Lorentz transformations used in Special Relativity. These transformations can be expressed by hyperbolic functions of a real argument  $\omega$  or by an imaginary argument  $i\omega$  with trigonometric functions. Thus the transformations can be considered as rotations in  $zt$  plane, where axis  $z$  is aligned with the velocity of the moving observer. Thus the space rotation of a hurricane about its spin axis is caused by a spiral movement of the air, the rotation in a space-time plane is caused by a linear movement of the relativistic ether along Vir spin axis.

This completes the description of all Vir rotations in space and time and reveals the origin and nature of the electric charge in elementary particles. If the rotation in  $zt$  plane is positive the ether flows out from the ends of the Vir and we have a positive particle. If the ether flows in through the ends of the Vir we have a negative particle. If the ether flows in from one end and out from the other then we have a neutral particle.

CPT is the most fundamental conservation law observed in particle physics, but there is no explanation for this in terms of quarks. The time transformation  $T$  reverses all motion and transforms a particle to an anti-particle. Using this concept of anti-particles Vir Theory shows that CPT transformation must be invariant.

Although there are anti-protons, anti-neutrons and anti-electrons (positrons) there are no stable anti-atoms, i.e. anti-matter. The conventional thinking is that this may be due to some asymmetry that occurred very soon after the big-bang. Vir Theory explains how this comes about without resorting to the big-bang.

When considering black holes they are described by only three quantities: mass, spin and electric charge. These are the essential properties also for elementary particles. Like particles black holes have accretion disc and two opposite jets perpendicular to the disc. It appears that the smallest and the most massive objects known to us may have the same nature.

The integrals for the mass and the moments of inertia for the Vir can be evaluated algebraically and using these expressions one can obtain a formula for the relationship between the spin and mass of Vir. It is a formula that allows several masses for a given spin and vice-versa. This formula has been tested very successfully using the existing particle data, as will be shown shortly in the paper entitled "Mendeleev-like Tables of Elementary Particles".

### Acknowledgements

I would like to thank my wife Jo for her loving care without which this work would not have been possible. I thank God for everything.

### Sources of Figures

Figure 22. Image courtesy NASA 11 Sep 2003 *Hurricane Isabel*  
<http://www.hurricanetrackinfo.com/>

Figure 23. Image courtesy NOAA/NASA 22 Aug 2005 *Hurricane Anatomy*  
[http://earthobservatory.nasa.gov/Library/Hurricanes/Images/hurricane\\_structure.jpg](http://earthobservatory.nasa.gov/Library/Hurricanes/Images/hurricane_structure.jpg)

All other images are copyright of the author. All URLs given above were active in Nov 2007.

### References

- [1] Troutman J 1983 *Variational Calculus with Elementary Convexity* (Springer Verlag) p.5-6, 230-232, 266-267
- [2] Courant R & Hilbert D 1989 *Methods of Mathematical Physics* (John Wiley & Sons) p.164
- [3] Meyer-Ardent J R 1972 *Introduction to Classical and modern Optics* (Prentice-Hall) p.14
- [4] Landau L D and Lifshitz E M 1976 *Mechanics* (Pergamon Press) p.2
- [5] Landau L D and Lifshitz E M 1975 *The Classical Theory of fields* (Pergamon Press) p.24
- [6] Landau L D and Lifshitz E M 1975 *The Classical Theory of fields* (Pergamon Press) p.245
- [7] Landau L D and Lifshitz E M 1965 *Quantum Mechanics* (Pergamon Press) p.20, 25
- [8] Landau L D and Lifshitz E M 1976 *Mechanics* (Pergamon Press) p.114
- [9] Brown H R 2007 *Physical Relativity* (OUP)
- [10] Germuska M J, Germuska R B, Germuska M A 2016 Shapes with the Minimum Moment of Inertia *viXra:1612.0233 [General Mathematics]*
- [11] Germuska M J 2016 The Shape of Water and Air Vortices *viXra:1612.0211 [Classical Physics]*
- [12] Landau L D and Lifshitz E M 1975 *The Classical Theory of fields* (Pergamon Press) p.62
- [13] Landau L D and Lifshitz E M 1975 *The Classical Theory of fields* (Pergamon Press) p.10, 11
- [14] Landau L D and Lifshitz E M 1976 *Mechanics* (Pergamon Press) p.115
- [15] Landau L D and Lifshitz E M 1976 *Mechanics* (Pergamon Press) p.116

13 Dec 2016

[16] Landau L D and Lifshitz E M 1976 *Mechanics* (Pergamon Press) p.120

[17] Landau L D and Lifshitz E M 1976 *Mechanics* (Pergamon Press) p.113

[18] Churchill R V 1960 *Complex Variables and Applications* (McGraw-Hill) p.63

[19] Churchill R V 1960 *Complex Variables and Applications* (McGraw-Hill) p.54