

Dynamic Geometry Waves Theory Foundation

Calin Vasilescu

January 1, 2017

Abstract

The starting point for the concept is the idea that what we use as clocks don't measure the true time. Based on this idea, we can define time as absolute and think that clocks don't measure the absolute time, but they have a tick rate that depends on the speed relative to the flat absolute space. The idea is what I think is a natural interpretation of what happens in the very popular thought experiment of Einstein's relativistic train. If the light beam is a light clock, it becomes clear that the trajectory of the light beam is the only thing that makes the tick rate change. The idea is that the same thing happens with all the clocks we use. Following this idea we can construct a whole theory closely following the empirical evidence we already have. This model suggests that space doesn't have a variable geometry, instead complex particles inner geometry changes what we call time and space.

This concept is only intended to be a starting point for a proper theory of space. It only contains few principles that are intended to give a better and more natural explanation how the universe works. I will try to explain simple ideas about time, space and fields.

1. Space geometry

For this concept to work, we need to define a **flat space geometry**. This flat geometry space-time cannot be changed by definition. We will define an absolute reference frame with an origin for x, y and z axis. We will refer to this frame with flat space as the **alpha frame (AF)**. AF will use alpha meters and time. This AF can be real if we can prove the electromagnetic waves need an Aether to propagate, only the origins are arbitrarily chosen. Otherwise, empty space is an absolute void and flat geometry is the way to tell that the structure of space and time does not change.

2. Time and space definition

Before Special Relativity it was thought space and time were absolute and we also measured them as absolute. Based on this, units of time and space were defined. Once Special Relativity (SR) was accepted by the mainstream, these definitions remained the same but they were not absolute anymore. Instead, speed of light became absolute which apparently made things work surprisingly well and equations had a remarkable symmetry. Moreover, since the idea of an Aether was not necessary anymore, only the speed of light could remain absolute and constant in every

reference frame. This points to the idea of a space that has a dynamic structure and it is real. However, we can define an absolute space and time and speed of light will be observer dependent in the AF frame. A medium for waves like the electromagnetic waves to propagate is no longer necessary in this context but the concept doesn't exclude it.

When time was viewed as absolute, clocks were made and thought to measure this time (absolute). Clocks accuracy improved over time, but they kept measuring the same thing, namely relative time. We though clocks were measuring absolute time, but they were measuring the relative time. If we define **alpha time** as absolute, we can see that our clocks will not be able to measure measure alpha time. Their tick rate will indicate **alpha speed of the clock**.

An object like an atom, for example, that is stationary in **AF** will have the highest oscillating frequency compared to the frequency in any other reference frame we choose. In other words, **a clock at rest in AF will have zero time dilation factor**. Any other reference frame we choose will have clocks at rest at a lower tick rate.

Another property of time is that along the time axis the total amount of information in the universe should be conserved. On any other dimension that doesn't happen. This is the essence of time and it is how true time should be defined.

3. Alpha dimensions

In the **alpha reference frame** we will define alpha time and alpha length and relative alpha speed of light (observer dependent):

$$x_\alpha, t_\alpha, c_{r\alpha}$$

$c_{r\alpha}$ is a relative speed of light seen in *AF*, as the difference between c and object speed in *AF*.

In any other reference frame, c will be measured will not vary because those frames use clocks that have variable tick rates in *AF*.

Figure 1, represents 2 regions of space viewed from *alpha frame*.




Fig. 1. Alpha frame 2d space




Fig. 2. Simple SR diagram for AF , the arrows represent flashes of light travelling from the floor to the ceiling and back. Notice that the box needs to expand on x axis to explain the experimental evidence. This is explained in chapter 6.

4. From SR to Alpha Space-time

Figure 1, shows a basic SR diagram. An observer at rest in the AF (obviously it works for any reference frame we choose), will measure light as constant no matter how fast the box is moving. If trying to measure the speed of light (two way) in the moving box you get the same value for c . SR conclusion is that something must be happening with time in the moving box and that is time dilation. However, there is another possibility I think can answer more questions. We assume that clocks measure time as absolute, then SR demonstrated time was relative. If we define time as absolute then the new idea is that our clocks we use don't measure time but their tick rate depend on speed. In other words, they don't measure alpha time, but relative time. A clock in the moving box (fig. 2), is in fact the light clock. The light clock will complete a cycle in a longer time viewed for AF. The tick rate is reduced by Lorentz factor γ . Any other clock we used would do the same thing. It is as if time dilates but the true alpha time doesn't change. If we put the problem this way, we will be able to explain the mechanics contained in this paper. For the idea to work we need to take into account the length extension (not contraction) as shown in chapter 9.

5. The illusion of Matter

If the mechanism of clocks (which applies to the most accurate atomic clocks available), hold to any clock, it means the theory need to unify all the fields we know, including gravitational into a single fundamental field. This points to the idea that, everything in the universe is only made only of waves in the electromagnetic field. Nicola Tesla said “Everything is light” and I agree. The waves corresponding to this field interacts in such a manner that they create structures like particles we identify as matter and also all force effects. In figure 2, I've used a light flash as a clock. If you sent a beam of electrons instead, it wouldn't work correctly . **It becomes clear that the particle geometry matters.** If we take for example quarks that are supposed to consist of different matter, despite all the evidence, no one has actually seen a bare quark. Instead, we observe clusters of known particles with $\frac{1}{2}$ spin origin. The spin indicates a complex geometry particle (non zero rest mass).

This model suggests that **space doesn't have a variable geometry (the term is usually curved geometry)**. Complex particles inner geometry changes what we call time and space. An atom moving faster changes geometry and its internal tick rate changes. Because the same thing happens to electrons, it means they also have an internal cycle, you could use as a clock. When travelling faster the geometry of the electron changes and makes it complete the cycles in a longer time. Most probably it is a spiral (when viewed relatively) or helix motion. But this is more of a speculation. Careful studies must be carried to confirm the geometry. What is obvious is that **moving at c transforms any geometry trajectory into a single line trajectory** (in

reality perfect straight trajectories do not exist because of all sorts of interactions). At rest the wave orbits around its centre. Orbits can have a precession as well or can stabilize. However, particles are never at rest in AF. The electron particle also has a wavelength associated with it (de Broglie). This wavelength experiences Doppler effect. Hence the moving electron will be measured with higher frequency / lower wavelength. However the Doppler effect is a different aspect. The clock tick rate is not the same thing as the associated wavelength. This has only to do with what a wave detector reads. The Doppler effect is like an illusion. It is like when you move toward pulses they appear to increase frequency. If a particle is accelerated the deBroglie frequency increases (whether is only relatively or in AF needs to be demonstrated) and the orbits frequency is reduced in AF . This is the advantage of using the absolute reference frame. There are many other advantages. The reality might not be possible to describe correctly without an absolute space and time.

To summarise, it is the complex particles geometry (this refers to any particle that has mass) that give the illusion of changing the geometry of spacetime.

6. Zero versus non zero mass particles

The photons as electromagnetic waves, travel in straight line always. All other particles are structures of values that describe circular or spiral patterns, thus the lead wave speed is reduced. Fig. 3. “Mass” is a question of frequency and length of the wave like radiation (photons) does.




Fig. 3. Electron vs. Photon

The electron goes on a different trajectory because the fields disturbances that generates them could not be the same (the gravity explanation might suggest they are almost the same). In the case of electron the propagation happens in such a way it forms a circular trajectory. In AF the trajectory it can form a helix (Fig.4) but viewed from a moving reference frame it can be seen as a 2d moving spiral.

Parametric plot:




Fig.4 .An electron trajectory in AF .
The electron is moving in the z direction,
but the waves follow a helix




Fig.5 .A free electron and a photon
trajectory in a reference frame attached
to a moving object in AF . In this diagram
the electron is travelling at approximately $0.7c$
*frequencies are not representative.
The idea is the electron wave frequency is higher than a frequency
of a sub gamma photon. The Doppler shift is not taken into consideration here.

There seems to be a consistency between my idea and de Broglie discoveries and opinions.

“Thus to describe the properties of matter as well as those of light, waves and corpuscles have to be referred to at one and the same time. The electron can no longer be conceived as a single, small granule of electricity; it must be associated with a wave and this wave is no myth; its wavelength can be measured and its interferences predicted. It has thus been possible to predict a whole group of phenomena without their actually having been discovered. And it is on this concept of the duality of waves and corpuscles in Nature, expressed in a more or less abstract form, that the whole recent development of theoretical physics has been founded and that all future development of this science will apparently have to be founded.” De Broglie

Helix or spiral motion are speculation only, but the idea is some kind a loop motion must exist in order to experience these effects. However, these patterns seem the most likely to occur.

Particles like electrons at rest they can be viewed as circles or loops. The wave still travels at c . That is how they can have rest mass. The same effect can be applied to more complex entities like atoms, since they are composed of elements like electrons and quarks since these follow spiral trajectories.

7. Gravitational effect of particles

Looking at what happens to an electron while moving, the idea of curved spacetime is not very convincing anymore. The gravitational field effect has a unique property. It is always attractive.

Analysing the studies made on gravitational effects and experimental data available, we can realise that the gravitational effects cannot simply be attributed to a simple gravitational field around the matter. The spacetime concept is a step forward from newtonian gravity, but it seems to me it cannot explain all the effect that appear in the real world.

For understanding gravity, in this context, we will use the term particle will describe to a wave that is **either a straight or a spiral / helix like trajectory and not spheres or point like entities (singularity is not accepted in this context)**.

The gravitational effect, I think, is best described using a about what photons are and how the gravity between them is generated. That is because this alpha space concept treats matter as field waves of different trajectory patterns. Understanding how gravity works for photons is crucial. The original paper is **Tolman, R.C., Ehrenfest, P., and Podolsky, B. Phys. Rev. (1931) 37, 602**. The idea was studied in the paper “Gravitational interaction for light-like motion in classical and quantum theory” Nikolai V. Mitskievich. **The study concludes that two photons moving in parallel will not experience any gravitational effect**. If they travel antiparallel they experience a gravitational effect, but twice as big as it would be judging the relativistic masses (using a quasinewtonian model).

Using the ideas expressed in this paper, we can analyze a case of two

hypothetical particles, as in figure 6.




Fig.6. This diagram represents two hypothetical particles at rest in *AF* and two wave beams of the same properties and the same length as the H particles

In the case of the anti-parallel wave beams the gravitational effect should be maximum. For the H particles, it is clear that the gravitational effect will be smaller. For simplification instead of a circular trajectory I have used a squared pattern, but the gravitational total effect should be the same. We can see that in the case of H particles, only **half** of the fundamental waves travel anti-parallel. Those are the vertical lines. This explains the effects seen in Tolman experiment.

An interesting fact that can be seen looking at this diagram is that a gravitational effect will be produced within the particle itself as long as the absolute speed in *AF* is not comparable to c (the effect should be reduced at 'relativistic' speeds). The magnitude of the gravitational effect is twice as big as it is between the H particles. If the radius of the particle is small, the gravitational effect will be considerable and could help keeping the integrity of the particle.

The most important idea is that in order to understand gravity, **we need to understand how gravitational effects occur between photons only**. Then we can extrapolate it to all particles.

8. Rest mass principle

Looking at the electron model explained by this concept the idea of kinetic energy doesn't have the same importance as in classical theories like newtonian mechanics. On fig. 5 The electron size in AF should be thought in terms of the length of the spiral. We can see that the energy of the electron in alpha frame doesn't change. However an observer that uses a normal clock will measure the electron moving at different speeds thinking it must have different speeds, thus different kinetic energy.

I will use electron as a mass particle. The relativistic mass formula is $m = m_0 \gamma$. Combined with the model described in fig.5, this means, for an electron to get more mass, it needs an additional length of the spiral. If you give energy to an electron, that energy adds up at one the end of the spiral. This in reality is probably viewed as an increase in the number of particles rather than increasing the energy of the electron especially for electrons attached to an atom. The increased mass m_{add} could have the formula $m_{\text{add}} = m_0 \cdot dl$ at constant deBroglie frequency, where dl is the increase in the length measured **along the spiral**. This is a way of increasing true energy of the electron but it is not what happens during acceleration.

The absolute mass for a fixed length (multiple of wavelength) along the complex trajectory would be :

$$m_a = m_e n, \quad \text{where } n = x_a / \lambda_a.$$

x_a is a length dimension in AF

λ_a is the wavelength of the electron in AF

m_e is the mass generated by a single oscillation

Notice that m_e is thought to be the absolute rest mass of a single oscillation of an electron (along the spiral). The oscillation refers to the internal frequency of the electron.

However, this way of calculating the absolute mass is wrong because as explained in chapter 7, gravitational effects do not work this way and there is no meaning of an absolute mass of an object in the absence of other objects.

An apparent increase of energy can happen due to observer speed. In a moving reference frame, when the speed of the electron increases, the internal frequency will increase due to Doppler shift. Although in the case of moving observer the energy of the wave remains constant in AF , the effect on the observer is the same as an increase of energy due to source motion in AF because of the frequency at which the waves hit the observer. A clock at rest in the moving reference frame will also be influenced by the relative speed. Fig.6




Fig.6. Electron red-shift from a moving reference frame

The red region represents a redshift and the blue portion a blue shift as a Doppler shift effect. The equation is different from relativistic formula and has a greater effect. Otherwise in case of a moving observer, if the clocks were slowed down in the case of the redshift by the same amount the effect would've been null. The equation for Doppler shift for a source travelling towards the observer at rest in AF is:

$$f = \frac{c}{c - v_a} f_a$$

f is the frequency perceived and f_a is the absolute frequency.

If the observer is moving the equation (corresponding to a moving observer) should be corrected with the Lorentz factor.

For redshift, in the classical Doppler effect, the same thing happens, the frequency of the source is not modified, but the recessional motion causes the illusion of a lower frequency.

The equation of gravitational redshift / blueshift is not going to be covered in this paper and requires more studies to be done.

Although we can have an equivalence between absolute mass and the mass seen from a reference frame, the idea of mass is not very useful for this concept. That is because of the way particles create gravitational effects (see chapter 7). A single photon travelling through space doesn't create any gravitational effect until a second photon that has a velocity which is not parallel to the first photon, appears.

If we look at a proton, the constituents (quarks and electrons) obey the same rule in my opinion. The mass of an electron is generated relativistically. For example an orbiting photon as a hypothetical particle (H) can have a huge relativistic "mass" at rest if we consider the centre of the orbit as the particle position. I think this is the principle of rest mass.

A quark at rest has a small mass. However, within a proton, if quarks move at a higher speed the relativistic mass increases up to the total mass of the proton. Gravitational effect cannot hold the proton together though. Strong nuclear force must be an effect of the way quarks interact with each other.

Mass doesn't seem to have a meaning as an intrinsic property of a wave. We can measure mass through gravity force or inertia, but can a body have mass without exerting any gravity force? I think not. If we can understand how light waves interact with other waves, creating a trajectory deviation we call gravity, then we can understand what gravity is. I think we should focus on this.

The proton mass is not composed of its constituents rest mass. That proves or at least suggests mass is only a relativistic effect. String theory also treats particle as vibrating strings. That means particles constituents are never at rest, but travel at c . It is obvious since electromagnetic waves travel at c , electron waves excite the same fields and must travel at the same speed.

A black hole is an object that behaves like a particle. It is possible to simulate a black hole made entirely of light waves. If BH were tiny as atoms we would think of them as new particles with rest mass. Simply because they are big, they can capture all sorts of particles and extreme amounts of energy.

Energetically if we compare a photon and the hypothetical particle, when you push energy into a photon, it increases its frequency. The H particle will do the same as the centre of the orbit accelerates. The speed limit for the H particle is obvious. The frequency increase of the H particle is explained by the photon frequency increase, although this is not very clear to me other than energetically (for a fixed amplitude of the EM wave).

A scenario of accelerating particles at c in the context of my concept

We can start with a pair of an electron and a positron that fall in a gravitational field. The deBroglie wavelength will gradually reduce as they gain energy and speed. We can monitor a similar pair made of two photons of the same initial wavelength as the e^- / e^+ pair. As the e^- / e^+ pair descends the speed increases, thus the orbital period decreases and the deBroglie frequency increases. For the photon pair only the frequency increases obviously at the same rate. No matter how long they fall, the orbital period will reduce but never reaches zero, but the deBroglie frequency will go extreme. If the pairs are on a slightly convergent way at some point the e^- / e^+ pair will collide and decay into two pair of photons. These photons will be the same frequency as the pair of photons that started the journey with the electrons and will have the same energy. This is the electromagnetic wave - matter equivalence.

9. Equations for transition between AF and a reference frame

Since by definition clock we use don't measure absolute time means in AF a normal second, cannot be compared to alpha seconds. In other words, alpha time is not measured in seconds but in a different unit of measurement we can call alpha second.

We will consider an example where an electron has $v_\alpha = I m_\alpha / s_\alpha$

For explaining the principles of conversion we will define a hypothetical non zero mass (spiral trajectory pattern) like particle called H particle that has certain properties.

For the alpha second definition we will use the time for a free, **stable (integer**

number of wavelengths) H particle to complete a single loop when stationary in AF . The loop will be of a radius of $l m_\alpha$ in alpha space. Hence the definition of an alpha meter will be the radius of a H particle.

The wavelength of an H particle in AF will be defined as $\lambda_H = l nm$. We can now say that the H particle number of oscillations per a complete loop (if you take a snapshot of the particle) is:

$$n_H = 2\pi \frac{r}{\lambda_H} = 2\pi \cdot 10^9$$

No matter how particle moves or how the observer is seeing it, this number will remain constant.

The energy of the particle can have the form:

$$E_H = n_H \frac{c}{\lambda_H} h = 6.54 \cdot 10^{-7} J_\alpha$$

where h is the Planck constant and c is the speed of light. Notice that the joule unit of measurement it only applies to absolute space-time (AF).

To simplify the model further instead of a circle we can think of the H particle as oscillating on a single axis. When moving perpendicular to the axis it makes a zig zag pattern like in fig.3. For this the width of the particle will be:

$$l_H = \pi r = \pi [m_\alpha] , \text{ no matter the speed.}$$

That means the speed of light is:

$$c = \pi [m_\alpha / s_\alpha].$$

If the particle is moving a v_H in AF , then we can measure the length of the particle in the travelling direction:

$$x_v = l \cdot v_H$$

An observer will see a moving particle or extrapolating this, an object, in AF , increasing its size in the direction of travel.

If the absolute time increase for a particle to complete a loop is t at rest.

If the H particle has a length x_H at speed v_H and a length x_H' at speed v_H' the relation between them is the following:

$$x_{H'} = x_H \cdot \sqrt{\frac{1 - \frac{v_H^2}{c^2}}{1 - \frac{v_{H'}^2}{c^2}}}$$

The equation shows that with increasing speed the length in the absolute space-time (AF) is increasing and **not contracting as SR says**.

If the oscillation period of the H particle (also measured as tick rate in case of an atomic clock) is T_H at v_H and a period of $T_{H'}$ at a speed $v_{H'}$ the relation between them in the following equation:

$$T_{H'} = T_H \frac{v_H}{v_{H'}} \sqrt{\frac{1 - \frac{v_H^2}{c^2}}{1 - \frac{v_{H'}^2}{c^2}}}$$

As speed in AF increases the time for a fundamental wave within a particle to complete an orbit increases. That is what we take as the tick rate of the clock or tick rate of a clock. It is clear that time itself does not change.

10. A possible dark matter scenario

The idea is, in a classical world the only thing it can make elements like photons or electrons have a fixed value of energy for a fixed wavelength is a multiple of a wavelength. In the case of an electron, when it is attached to an atom its energy level stabilizes at number multiple of its wavelength. But this doesn't necessarily mean that when it is travelling freely after other interactions it keeps the same number. It can be any number, but when is absorbed the atom gets the length it can accommodate. The same thing may happen to photons. If a wave of a certain length hits the electron wave, it gives its energy to the electron. Part of the light wave can still travel further if it is not entirely absorbed.

This seems to contradict the photoelectric effect. However, since the source of photons is the electron transition between orbitals, a possible explanation is, they are usually (if not always) emitted with the same length (same number of oscillations) for the same wavelength.

The truncated parts of photons or possibly other particles can manifest as black matter / energy and cannot be detected directly.

Another possible source of DM can be related to the electromagnetic interferences. Waves that cancel each other the E or H field, might become undetectable.

In any case, any source of dark matter should have an electromagnetic origin.

11. Two way speed of light

For this theory to work speed of light needs to be constant only when measured between a point A and a point B and back. The speed from A to B will not be equal to the speed from B to A in a particular reference frame we chose to measure it.

It is interesting to see what we should actually measure if one way speed of light could be measured.

Consider a box within the measurement of the speed of light is done. A light beam is sent from a source to a mirror and then it comes back. If we ignore the length extension I've mentioned in chapter 9, we get

$$c_{mF} = (c-v) \cdot \gamma$$

$$c_{mR} = (c+v) \cdot \gamma$$

where c_{mF} is the speed measured forward and c_{mR} is the speed measured when the light returns from the mirror. γ is the Lorentz factor.

$$c_{mF} = (c-v) \cdot \gamma = \frac{d'}{t_F} = \frac{d \cdot \gamma}{t_F}$$

$$\frac{d}{t_F} = c-v = c'_{mF}$$

However the box will increase in size by Lorentz factor and we get:

c'_{mF} is the real measurement after length correction.

Hence the real measurements should show:

$$c'_{mF} = c-v \quad \text{and} \quad c'_{mR} = c+v$$

These seems to confirm the conclusions of *Stephan J.G in his paper, **GPS and the One-Way Speed of Light***.

This concept is supported by the evidence we have about the speed of light as a constant when measured both ways just like relativity. However the only experiment claiming to have achieved the correct measurement for the one way speed of light supports this concept and indicate the same results.

Reginald T. Cahill in his paper, *One Way Speed of Light Measurements Without Clock Synchronisation*, concludes:

“The absence of the Fresnel drag in RF coaxial cables enables 1st order in v/c measurements of the anisotropy of the speed of light. “

12. A photon classical model

I'm going to make a very synthetic description of a model of photon, that seems to be consistent with QM and experimental evidence. It might not be very precise, but it will cover the essential ideas. A photon is an energy pulse that contains a packet of electromagnetic waves of approximately the same frequency. The structure of the waves within the photon is generated by electron transitions. That is the reason why a photon is not a simple EM wave, but a collection of waves in a single pulse.

In Quantum Mechanics, the square root of the orbital angular momentum of a photon is zero in the $m = 0$ state. The orbital angular momentum itself has strong fluctuations and it is zero only on average. Just like in QM where a photon is in a superposition of all possible states, the photon wave contains an infinity of electric field vectors (E), disposed on all directions perpendicular to the direction of travel at once.




Fig. 6. Photon OAM, $m=0$,
source:wikipedia.com

For $|m| > 0$, these vectors rotate, all at the same time. The helical modes are characterized by an integer number m , positive or negative.




Fig. 7. A reduced model of a classical representation of a clockwise spin photon electric field. The w_i symbols are separate electric field vectors corresponding to 5 waves

In figure 7, we have a representation of photon electric field vectors that compose the particle. The diagram only shows 5 electromagnetic waves rotating

clockwise (CW), but in reality a photon should contain an infinity of E vectors, covering half a circle, thus, 180° directions, rotating at the same time. You can see that the photon just like in QM, contains all directions at the same time in a simple classical superposition.

We can see that a photon is a little bit more than a simple EM wave. However, each wave it contains (as a classical superposition) has exactly the same behaviour as a classical radio wave. That is very important because in that case, two photons can combine amplitudes while travelling and form a photon with inner EM waves of higher amplitudes. However, when the new photon is absorbed by an electron, the electron only absorbs energies corresponding to the energies the electron can emit. These energies are related to the orbital numbers when the electron is attached to an atom. This means part of the photon can pass through. For a photon to be absorbed it needs a frequency threshold, but also an amplitude threshold. If the amplitude threshold is not met the energy will be re-emitted as the electron will be unable to change orbital state. If the frequency threshold is not met the photon will not interact and simply pass through.

In figure 8, there is a simple representation of the electric field within a high amplitude photon when passing through a vertical filter. The high amplitude photon is a combination of amplitudes of a limited amount of photons. This corresponds to a thin, light beam. The circle is in the $x-z$ axis and motion is in y axis. The diagram shows a slice through the y axis. It is clear that, if we place another V filter nothing changes in that pattern. The diagram doesn't follow natural proportions. The area of the petal of vectors should be $\pi r^2/2$ (half of a circle area). The total amplitude of the photon will be half the original amplitude. The combined magnitude of the transmitted amplitude will be $E_o/2$.




Fig.8. A photon passing through a linear polarization filter. E represents the E_i vectors summation

If we then let the linearly polarized photon pass through a CW circular polarizing filter (fig. 7) , the waves get arranged so that the electric fields will point in all 360° directions again, but also spinning CW all at the same time. There will be a 90 degrees phase difference from B field. A better picture is in fact the original diagram in figure 7, with the only difference that the new photon is now half the amplitude of the linearly polarized photon.

Now it becomes clear that if placing a horizontal filter will allow 50% of the amplitude of the photon to pass through. Also, it is clear that if we remove the circular filter, the light beam will be almost completely blocked. How much it is blocked depend on the material used. A perfect filter should be impossible to build (Fig.10).




Fig.9 Circular filter. All vectors are rotating CW




Fig. 10. The diagram shows a circularly polarized photon passing through a linear filter. Only a small fraction of the amplitude of the photon will be able to pass through. The red line is the petal corresponding to the horizontal filter, $E_r \ll E_0$
 Note. The diagram doesn't respect proportions, it only presents the idea.

If we emit a single photon, it will not be detected because the combined amplitudes of the EM waves that composes it will be less than the threshold amplitude for an electron transition. However a high amplitude photon will be able to pass through.

In QM you can't combine photons or split them. However, nothing prevents that in reality. Only emissions and absorptions (after stabilization on an orbital) are quantized. Classical EM waves can easily do that.

13. Paradoxes in SR and GR

General Relativity limits the particle speed justifying that to accelerate a particle to c you need infinite energy and sometimes people ask what would happen if you exceed this limit. Some even think that would mean going back in time. My model seems more natural to me. Exceeding c is clearly impossible and also there is no reason to think of going back in time.

Because in AF time is absolute, there are no paradoxes, no problems with the moment "now". What is *now* here it is *now* everywhere in AF. The popular twin paradox is not a matter of time any more. The twin that returns younger than his brother does that because the particles he is made of experiences slower cycles.

Time travelling is not possible here because it is in contradiction to the way alpha time is defined.

This theory allows the possibility of taking a snapshot of the universe (a single line through the space axis) and containing all the information in the universe. Thus, the next frame can be generated. However, quantum effects, could prevent knowing the hypothetical “next” frame and an infinity of possible next frame can exist, the one that follows is not known until it happens. If we imagine a random number RN that is either 1 or 2 and an event E that can generate two possible outcomes O1 and O2, then these outcomes are equally valid. Which one follows is impossible to know. This means the universe will certainly generate only one of these outcomes. The generation of a random outcome is subject of controversy.

There is a thought experiment we can think of. Imagine two circles of equal radius approaching each other from opposite directions. The path they describe is a straight line and the circles move so as the centres of the both circles are on the line. After they collide, we assume they will bounce in opposite directions. The question is if the trajectory of both circles will not follow the same line, how can we predict which side of the line they will go ? No matter how much we zoom in we will always see continuous lines and not points. But the circles are made of points and spaces between points that are also infinitesimal. That is a sort of a mechanical RN generator. If we can zoom in and ultimately we see the points, then nothing is random here.

14. Black Holes Dynamics Interpretation

First of all this model does not allow any singularity neither point particles.

It is very likely that strong / weak forces to have a fundamentally electromagnetic origin. In that case it is a single shell of the BH, comprising all sorts particles orbiting at near c (elongated helix-es). Most of these particles are turned into various extreme high frequency EM waves (photon geometry). This is the optimum way to compress energy. This way the shell becomes very thin and gravitational effect is maximum.

The gravity produced is the same gravity planets produce, but different structure bodies produce different effects on close objects trajectories.

What differs from a planet in case of the BH are the “relativistic” and a stronger frame dragging effects.

I think it is important to take into account “relativistic” increase in “masses”. The phenomenon that can also be explained by relativity (however, only if using infinitesimal mass particles instead of photons) is that photons moving in parallel do not attract but those traveling anti-parallel do. This has been confirmed experimentally. In case of other particles a similar thing happens. If moving parallel the gravity does not increase.

All particles approaching the black hole horizon are accelerated by extreme frame dragging at near c , on the outer shell and begin to orbit along with the orbiting

radiation.

Most probably the greatest amount of gravity is produced by the relativistic motion of electro-magnetic waves while orbiting and not by their value of rest mass.

There is not reason for any particles inside the black hole. All particles and photons are concentrated on the shell that corresponds to what is called event horizon in General Relativity.

Fast jets and fireworks

By analyzing the model, it is clear that particles or radiation cannot escape tangentially to the surface of rotation. In order to escape, light or extreme speed particles should precess and exit through poles transforming the orbital trajectory into an elongated spiral trajectory. The idea is that light cannot be stopped and can only be deviated. The poles are the only way to escape without being deviated back into the BH.

According to data available, BH are relatively stable, but especially during collisions with massive objects, part of the particles are destabilized and can escape the BH.

15. The universe boundaries

Presumably the universe is not infinite, we can use this model to predict how the universe boundaries would look like. These will be similar to a black hole boundary. Gravitational effects will keep the waves in an orbit around the universe. Eventually all particles will establish on an orbit around the centre of symmetry.

16. The electron as a helically-circulating spin-1/2 charged photon generating the de Broglie wavelength

This model is found in the paper *The electron is a helically-circulating spin-1/2 charged photon generating the de Broglie wavelength* written by Gauthier R. Although the author says his approach is relativistic, it perfectly fits my concept.

“My approach [1] is to model the relativistic electron as a helically-circulating double-looping charged photon. The photon carries the electron’s charge and has spin $S_z = \pm\hbar/2$, the same as that of an electron, rather than spin $S_z = \pm 1 \hbar$

of an uncharged photon. By equating the moving electron’s relativistic energy $E = \gamma mc^2$ with the photon’s energy $E = h\nu$, the charged photon is found to have a relativistic frequency $\nu = \gamma mc^2 / h$ and a relativistic wavelength $\lambda = h / \gamma mc$. While this relativistic frequency $\nu = \gamma mc^2 / h$ was used by de Broglie to derive the electron’s de Broglie wavelength, the relativistic wavelength $\lambda = h / \gamma mc$ of a hypothesized photon that models a relativistic electron has never to my knowledge previously been utilized in describing the electron, neither by de Broglie nor by others (including other electron modelers.)”Gauthier R.

“This spin 1/2 charged photon model of the electron is a generic model of the photon because it does not present a more detailed model of a photon that may compose an electron. This generic model of the charged photon could be used with a variety of more detailed photon models that also

have the basic photon characteristics of light speed, frequency, wavelength, energy and momentum listed above.

In a more detailed charged photon model, the charged photon's spin $S_z = \pm \hbar / 2$ must remain constant at all velocities to give the electron model a spin of $\hbar / 2$ at all velocities. " R.Gauthier




Figure 1. The electron's de Broglie wavelength is generated from the spin 1/2 charged photon model of electron. The electron is moving to the right with speed v . The charged photon modeling the electron is moving helically to the right with speed c along its helical trajectory. The helical trajectory for one wavelength of the double-looping charged photon is mathematically unrolled to show the geometrical relations more clearly. Each wavelength of the double-looping charged photon along its helical trajectory generates the electron's de Broglie wavelength along the helical axis of the charged photon. As the electron moves to the right with speed v , the de Broglie waves move to the right with speed $v_{phase} = c^2 / v$.

Fig.11. A diagram in R. Gauthier paper showing the model of an electron.

"An objection to the spin 1/2 charged photon model of the electron is that an electron has spin $S_z = \pm \hbar / 2$ and is a fermion while a photon has spin $S_z = \pm \hbar$ and is a boson, so an electron cannot be a charged photon. But if a circulating photon carries the electron's charge and has spin 1/2, it is not a boson but a fermion. In other words, I am proposing that photons may be of two varieties: 1) uncharged and moving linearly with spin 1 (a boson) with no rest mass, and 2) charged and moving helically with spin 1/2 (a fermion) and having a rest mass. " Gauthier R.

In this paper Gauthier is using a relativistic formula. However if we assume the photon moves in absolute space or more precisely in an electromagnetic medium then the relativistic effects occur naturally. It is clear how the electron speed is limited by c . As the electron accelerates, the helix pitch is increasing. That is what in fact happens to clocks, the tick rate is given by the pitch of the helix and an increase in pitch corresponds to a time dilation in Relativity. In reality clock tick rates are reduced. This happens to any particle. Obviously biological clocks, for example, will follow the same principles.

Regarding the classical way of charge generation, there is a mechanism that can explain it. A better explanation is made by Ph.D. H. Shantz in chapter 16.

Sending two electro-magnetic waves one toward each other, depending on the phase, the electric and magnetic fields will interact. If the amplitudes overlap the resultant will be doubled. If the electric fields values cancel each other, the magnetic field values will double and vice-versa. This way it becomes clear that the energy is conserved. However if you think about sending two waves in the same direction, in that case the waves would be cancelled completely. However this case is equivalent to not generating the wave at all. In mechanics, this is equivalent to trying to pitch a string from both sides. Obviously there will be no release of energy.

By overlapping two electromagnetic waves it becomes possible to create a charge. For example, if two electromagnetic waves are forced to travel in a circle, in opposite directions with a phase shift of 180 degrees between electric field amplitudes, the electric field will be cancelled. This way it will become a negatively charged particle. If the magnetic fields cancel, the particle created will possess a positive charge.

If we look at an e^+ and an e^- pair and assume that the electron has a double amplitude magnetic field and the positron has a double amplitude electric field, when they interact they can create two simple electromagnetic waves that obviously carry the same energy. This is a matter – energy equivalence that also demonstrates energy conservation. It also means that all energy in the universe is fundamentally kinetic energy.

17. Electromagnetic waves vector summation

For explaining the electromagnetic interference I will use the *H.G. Schantz's* paper as a reference. He manages to explain this in the most elegant and logical way which reveals a perfect energy / information conservation principle.

Dirac concluded in his paper, that two photons can never interfere with each other (P.A.M. Dirac, *The Principles of Quantum Mechanics*, 4th ed. (revised), 1967, p. 9).

“ I believe Dirac was wrong.

Interference is defined with respect to the electric field. But when two photons interfere, a very interesting thing happens. Electric energy transforms to magnetic energy, but conservation of total energy holds. The figure below illustrates how this works for constructive interference (left) and destructive interference (right).

In “destructive interference,” magnetic fields (H) add while electric fields (E) cancel. In “constructive” interference, electric fields add while magnetic fields cancel. Energy is always conserved, merely transitioning between the electric and magnetic forms. The “S” is showing the Poynting vector, the local flux of EM energy, given by $S = E \times H$.

The truly fascinating aspect of this is that when one field or the other goes to zero, the energy comes to a rest, even though the waves keep propagating through each other. Some of the energy is exchanged between the two, so some of the forward propagating energy becomes backwards propagating energy and vice versa.

I discuss this at greater length in a blog post, "[Dirac's Big Mistake: What EM Tells us About QM.](#)"

So in summary, in a destructive interference, energy is conserved because the cancelled E-Field generates an enhanced H-Field and the "missing" electric energy transforms to magnetic energy.

You'll find Princeton physics professor Kirk McDonald offers an excellent analysis here, as well: "[Does Destructive Interference Destroy Energy?](#)" H.G. Schantz

If this explanation for waves interference hold, then it can explain how it is possible to create apparent charges using electromagnetic waves only, as seen in chapter 16.

18. The electro-magnetic medium

If you can explain the electromagnetic waves propagation without using a medium, then waves propagate in an absolute frame of reference. However, if the medium is necessary to provide this propagation then I can say that this medium needs to have constant permittivity/permeability. There is no definite motion of this medium. Even if it had, it wouldn't have absolutely any effect on the propagation of waves.

19. Time definition aspects

*In his paper, **Time Invariance of the Fundamental Physical Constants**, the author Muger B. Răuț concludes:*

This paper shows that the variation of certain fundamental constants is practically impossible in a physical time frame of reference. We can have as many time frames of reference we want but when we transform them all into physical time frames of reference, with time as a measure of movement, physical equations retain their form and meaning and values of certain physical quantities and fundamental constants are the same. Therefore the question of variation of certain fundamental constants is only possible for those frames of reference other than physical time.

I've defined space and time as absolute, so that they can become frame independent. The absolute time is not physical. By proposing a special particle geometry, I have shown that causality is constant in absolute time and space (which is not physical) but the measured time and lengths vary from frame to frame. This way there seems to be no contradiction between my model and the ideas expressed in the M. Raut's paper.

20. A classical view on the photoelectric effect

In the double slit experiments, in terms of Quantum Mechanics (QM), it is often said that the measurement collapses the wavefunction or more exactly, the measurement destroys the interference. However, we can also think, that is not the measurement that destroys the interference (except for some experiments). The delayed choice quantum eraser can be used as a proof. People developed the DCQE (Delayed Choice Quantum Eraser) experiment, where there is no measurement involved at the slits, but they went forward with QM believing that even simply watching the results destroys the interference. If QM is all correct, this is a rational conclusion. However, I think it is not the case. The theory is probably wrong. Apparently, the detectors cannot detect the energy of a photon, if the energy that reaches the detector is by a certain degree less than the energy released by the atom. QM says, that there is a significant probability of detection when the wave passes through both slits. That must be wrong. The probability is **almost** zero. Unless the other part of the wave hits the excited atom in a very short time, the atom will not release the electron.

Let's suppose we have a photon with energy E_γ . After it passes the slits we get two waves of $E_{\gamma L}$ and $E_{\gamma R}$. Say, the energy required for an atom to release an electron is E_e . If a wave with energy E_L hits the atom, it will excite it, but the atom will not release the electron yet. Since the excited state does not correspond to a stable electron orbital, it will quickly lose energy. However, if the wave E_R comes quickly enough, there will still be enough energy left in the excited atom and $E_R + E_L > E_e$. Hence the electron will be released.

Notice, that the energy of a photon when it was just released will always have more energy than E_e otherwise the frequency of the wave is not enough to trigger a photo-electron release.

We can, then say that this mechanism should work at any frequency of the photon. More exactly, a wave of any frequency could trigger an electron release, because photons can add up energies to the atom and excite it until releasing the electrons. It is clear that it doesn't since the experiments show that the frequency matters. However, we can then understand that for a photon to interfere with an electron, it needs a minimum frequency. It is possible to explain how this happens. A low frequency wave interferes with an electron attached to an atom as well, just like a high frequency does. However the excited state of the atom, loses energy quicker than the low frequency wave is able to provide the energy for releasing an electron.

Supposing we have a Hydrogen atom. We can write:

$$E_A(t) = E_{A0} + E_\gamma - P_A * t$$

where, E_{A0} is the energy of the atom when the electron is a stable configuration, like say first orbital

t is measured in absolute seconds, according to this concept. All symbols for time denote absolute time

$E(t)$ is the energy stored in the orbiting electron, which is a function of time.

E_A is the energy received from an electromagnetic wave.

P_A is the radiation power of the atom in the unstable state.

$$E_{\gamma} = P_{\gamma} t_A$$

t_A is the emission time of an electromagnetic wave.

E_{γ} is the energy produced by an electromagnetic wave.

We can also write:

$$E_{\gamma}(t) = P_{\gamma} t$$

P_{γ} - power produced by an electromagnetic wave

This power depends on the frequency:

$$P_{\gamma} = k\nu$$

k is a constant and ν is the frequency of the wave.

We can now write:

$$E_{\gamma}(t) = k\nu t$$

To produce a photo-electron, the energy E_A needs to reach E_{e^-} .

$$E_A(t) = E_{A0} + k\nu t - P_A t$$

In conclusion, to produce photo-electron, the electromagnetic wave must have a frequency that satisfies the following inequality:

$$\nu \geq P_A/k$$

These equations need to be expressed in absolute time. Applying relativistic equations to them will generate wrong results in certain situations, because the relativistic effect are generated automatically by the wave behavior of all particles.

Conclusions:

The paper shows that the propagation of values (waves) of electromagnetic field happens at the same constant speed in absolute space and time as defined. Light waves travel straight and thus the forward speed is c . Particles with mass travel in a helix / spiral and thus the forward speed is reduced, but the internal field propagation is at the same constant speed c . Thus, c limit becomes obvious for non zero mass particles. The c limit of light is the natural propagation speed of EM waves and is not a limit but the speed the propagation it is always happening. It feels natural for fields to travel at a certain speed rather than infinite. Infinite speed would be unimaginable and a universe like that wouldn't work. The value $2.999 \cdot 10^8$ m/s is because the conventions we use when defining dimensions. This constant is dictated by the permittivity and permeability properties of the vacuum. The only important thing is that is constant and non infinite.

The reason why c is not variable is because the propagation environment and propagation mechanics don't change. There seems to be no reason for the vacuum properties to change. The fundamental waves that compose other particles propagate the same way. In other words, there is a single fundamental speed in the universe. Speeds below c , are apparent. If you send a light beam through a channel with mirrors and it goes in a zigzag pattern, it reaches the observer slower but the wave has travelled the same speed. The apparent speed is lower. Otherwise, you would say only c is constant and other particles travel a various speeds. That is not the case, according to this concept.

GR limits the particle speed justifying that to accelerate a particle to c , you need infinite energy and sometimes people ask what would happen if you exceed this limit. Some even think that would mean going back in time. My model seems more natural. Exceeding c is clearly impossible and also there is no reason to think of going back in time.

Usually theories start from simple concepts. If the concept is wrong, then mathematics can't do anything to fix it, unless you accept the mathematical predictions are irrational. Mathematics can be used to further develop the theory and to make sophisticated predictions. My opinion is, both Quantum Mechanics and General Relativity theories, at some point make irrational predictions.

These simple ideas expressed in this paper could represent the foundation for a new theory if proved to be correct.

Acknowledgments

I gratefully acknowledge the users on Physics Forums of Bernhardt Media LLC for getting qualified answers and The Naked Scientists Forum of The University of Cambridge for answering various questions that helped me identify certain issues and develop this paper.

References

- [1]. *Ether and the Theory of Relativity*, published by Methuen & Co. Ltd, London, in 1922.
http://www-history.mcs.st-andrews.ac.uk/Extras/Einstein_ether.html
- [2]. *The wave nature of the electron*, LOUIS DE BROGLIE Nobel Lecture, December 12, 1929. https://www.nobelprize.org/nobel_prizes/physics/laureates/1929/broglie-lecture.pdf
- [3]. *Development of our conception of the nature and constitution of radiation* by Albert Einstein, *Pysikalische Zeitschrift* 22, 1909
<http://photontheory.com/Einstein/Einstein06.html#Ziegler>
- [4]. *Gravitational interaction for light-like motion in classical and quantum theory* Nikolai V. Mitskievich
- [5]. <https://www.physicsforums.com/threads/plausibility-arguments-for-noninteraction-of-parallel-beams-of-light.514429/>
- [6]. *GPS and the One-Way Speed of Light*, Stephan J.G. Gift Additional information is available at the end of the chapter <http://dx.doi.org/10.5772/50670>
- [7]. *Quarks*, Darin Acosta <http://marge.phys.washington.edu/Library/quarks.pdf>
- [8]. <http://www.nature.com/news/mystery-of-black-hole-fireworks-solved-1.17656>
- [9]. https://en.wikipedia.org/wiki/Redshift#Doppler_effect
- [10]. https://en.wikipedia.org/wiki/Orbital_angular_momentum_of_light
- [11]. https://www.researchgate.net/publication/281322004_The_electron_is_a_helicily-circulating_spin-12_charged_photon_generating_the_de_Broglie_wavelength
- [12]. *The art and science of UWB antennas*, H.G. Schantz, <http://aetherczar.com/?p=3667>
- [13]. *One-WaySpeedofLightMeasurementsWithoutClockSynchronisation* Reginald T. Cahill, http://www.ptep-online.com/index_files/2012/PP-30-08.PDF
- [14]. *Time Invariance of the Fundamental Physical Constants*, Mugur B. Răuț, <http://vixra.org/abs/1612.0384>