

Part D.**VBGC 1.2 - The extension and generalization of BGC as applied on o-primeths (${}^o\mathcal{P}^*$)**

VBGC 1.2 (version 1.2, the same with the version of this article) – main statement:

1. Defining o-primeths as: ${}^0P_x = P\left(\frac{x}{\text{0iterations}}\right)$, ${}^1P_x = P\left(\frac{P(x)}{\text{1iteration}}\right)$,

$${}^2P_x = P\left(\frac{P(P(x))}{\text{2iterations}}\right) \dots {}^oP_x = P\left(\frac{P(P(\dots P(x)))}{\text{oiterations}}\right), \text{ with } P(x) \text{ being the } x\text{-th prime in the set of}$$

standard primes (usually denoted as $P(x)$ or P_x and equivalent to 0P_x alias “0-primeths”) and the generic oP_x being named the generic set of o-primeths (with “o” being the number of “order” of iterations).

a. I have used the notation 0P_x and oP_x instead of the standard notation

$$P^1(x) = P(x) \left[= {}^0P_x \right] \text{ and } P^n(x) = \frac{P(P..P(x))}{\text{nnested functions } P} \left[= {}^{n-1}P_x \right], \text{ so that to strictly measure}$$

the number of recursive steps (iterations) to produce a generic set oP from ${}^{(o-1)}P$ by applying one additional recursive step and also to not generate the confusion between

$$P^n(x) = \frac{P(P..P(x))}{\text{nnested functions } P} \text{ and the exponential product } \left[P(x) \right]^n = \frac{P(x) \cdot P(x) \cdot \dots P(x)}{\text{ntimes}}.$$

b. It is also true that producing the elements of the (prime) function $P(x)$ from the natural set \mathbb{N}^* is also like selecting just the naturals with prime indexes from \mathbb{N}^* , so that 0P can be identified with \mathbb{N}^* and the set of primes \mathcal{P}^* can be identified with 1P : however, \mathbb{N}^* is not a set of primes and that is why I have avoided to note \mathbb{N}^* with 0P AND ALSO decided to count the sets of o-primeths starting from 0 (so that ${}^0P_x = P(x)$) in the purpose to strictly measure the number of iterations starting from 1, so that

$${}^1P_x = P\left(\frac{P(x)}{\text{1iteration}}\right).$$

2. The inductive variant of VBGC states that: “Any even positive integer

$$\boxed{2m > 2 \cdot 2^{(a+1)(b+2)(a+b+1)}} \text{ can be written as the sum of at least one pair of distinct o-primeths$$

$$\boxed{{}^aP_x > {}^bP_y}, \text{ with the positive integers pair } \boxed{(a, b), \text{ with } a \geq b \geq 0} \text{ defining the (recursive) orders$$

$$\text{of each of those o-primeths AND the pair of distinct positive integers } \boxed{(x, y), \text{ with } x > y > 1}$$

defining the indexes of each of those o-primeths.”.

3. Alternative formulation for the inductive variant of VBGC, using the standard notation

$$P^1(x) = P(x) = {}^0P_x, P^2(x) = P(P(x)) = {}^1P_x \text{ and } P^a(x) = {}^{(a-1)}P_x: \text{ “Any even positive integer”}$$

$$\boxed{2m > 2 \cdot 2^{a(b+1)(a+b-1)}} \text{ can be written as the sum of at least one pair of distinct o-primeths$$

$$\boxed{P^a(x) > P^b(y)}, \text{ with the positive integers pair } \boxed{(a, b), \text{ with } a \geq b \geq 0} \text{ defining the (recursive)$$

$$\text{orders of each of those o-primeths } P^a(x) \text{ and } P^b(y) \text{ AND the pair of distinct positive$$

$$\text{integers } \boxed{(x, y), \text{ with } x > y > 1} \text{ defining the indexes of each of those o-primeths.”.$$

4. **The analytic variant of VBGC (from which the inductive VBGC can be intuitively inducted) states that:** “For any pair of finite positive integers (a, b) , with $a \geq b \geq 0$ defining the (recursive) orders of an a-primeth $({}^aP)$ and a b-primeth respectively $({}^bP)$, there will always exist a single finite positive integer $(n_{a,b} = n_{b,a}) \geq 3$ so that, for any positive integer $m > n_{a,b}$ it will always exist at least one pair of finite *distinct* positive integers (x, y) , with $x > y > 1$ (indexes of distinct odd o-primeths) so that: ${}^aP_x + {}^bP_y = 2m$ AND ${}^aP_x > {}^bP_y$ AND the function $f(a, b) = f(b, a) = (n_{a,b} = n_{b,a}) \geq 3$ has a finite positive integer value for any combination of finite positive integers (a, b) , without any catastrophic-like infinities for any (a, b) pair of finites positive integers.

- a. **Important note.** I have chosen the additional conditions $(a \geq b \geq 0) \wedge (x > y > 1) \Leftrightarrow {}^aP_x > {}^bP_y$ so that to lower the nof. lines per each GM and to simplify the algorithm of searching $({}^aP_x, {}^bP_y)$ pairs, as the set aP is much less dense that the set bP for $a > b$ AND the sieve using aP (which searches an aP starting from $2m$ to 3) finds a $({}^aP_x, {}^bP_y)$ pair much more quicker than a sieve using bP (if $a > b$).
- b. $f(0, 0) = (n_{0,0}) = 3$
- c. $f(1, 0) = f(0, 1) = (n_{1,0} = n_{0,1}) = 3$
- d. $f(2, 0) = f(0, 2) = (n_{2,0} = n_{0,2}) = 2564$
- e. $f(1, 1) = (n_{1,1}) = 40\ 306$
- f. $f(2, 1) = f(1, 2) = (n_{2,1} = n_{1,2}) = 1\ 765\ 126$
- g. $f(2, 2) = (n_{2,2}) = 161\ 352\ 166$
- h. $f(3, 0) = f(0, 3) = (n_{3,0} = n_{0,3}) = ?$ [working in progress on this function value]
- i. $f(3, 1) = f(1, 3) = (n_{3,1} = n_{1,3}) = ?$ [working in progress on this function value]
- j. $f(3, 2) = f(2, 3) = (n_{3,2} = n_{2,3}) = ?$ [working in progress on this function value]
- k. $f(3, 3) = (n_{3,3}) = ?$ [working in progress on this function value]
- l. ...[working progress on other higher indexes function values]

m. Interestingly, $f(a,b)$ applied on $a \in \{0,1,2\}$ and $b \in \{0,1,2\}$ has its value in the set


$$F = \{3, 3, 2564, 40306, 1\,765\,126, 161\,352\,166\}$$
 which has an exponential pattern

such as: $E_F = \{\approx 1.1, \approx 1.1, \approx 7.8, \approx 10.6, \approx 14.4, \approx 18.9\}$, with a relatively constant geometric progression between its last 4 elements so that

$$\left(\frac{\approx 18.9}{\approx 14.4}\right) \cong \left(\frac{\approx 14.4}{\approx 10.6}\right) \cong \left(\frac{\approx 10.6}{\approx 7.8}\right) \cong 1.32$$

. The gap between the exponents ≈ 1.1 and ≈ 7.8 may be possibly filled by $\ln[f(3,0) = f(0,3)]$,

$\ln[f(4,0) = f(0,4)]$... which are still in work to compute in the near future (see the next figures).




n. $F = \{3, 3, 2564, 40306, 1\,765\,126, 161\,352\,166\}$ also has a correspondent matrix

$$M_{f(a,b)} = \begin{pmatrix} n_{0,0} & n_{1,0} = n_{0,1} & n_{2,0} = n_{0,2} \\ n_{0,1} = n_{1,0} & n_{1,1} & n_{2,1} = n_{1,2} \\ n_{0,2} = n_{2,0} & n_{1,2} = n_{2,1} & n_{2,2} \end{pmatrix} = \begin{pmatrix} 3 & 3 & 2564 \\ 3 & 40306 & 1765126 \\ 2564 & 1765126 & 161352166 \end{pmatrix} \text{ and}$$

a matrix of exponents $ME_{f(a,b)} = \begin{pmatrix} \ln(n_{0,0}) & \ln(n_{1,0} = n_{0,1}) & \ln(n_{2,0} = n_{0,2}) \\ \ln(n_{0,1} = n_{1,0}) & \ln(n_{1,1}) & \ln(n_{2,1} = n_{1,2}) \\ \ln(n_{0,2} = n_{2,0}) & \ln(n_{1,2} = n_{2,1}) & \ln(n_{2,2}) \end{pmatrix},$

$$ME_{f(a,b)} = \begin{pmatrix} 1.1 & 1.1 & 7.85 \\ 1.1 & 10.6 & 14.38 \\ 7.85 & 14.38 & 18.9 \end{pmatrix} \text{ which can both be graphed as a surfaces (see the}$$

next figure).



- o. More interestingly, the function $f_x(a,b) = 2^{(a+1)(b+2)(a+b+1)}$ generates positive integer values that are relatively close BUT strictly larger than the values of $f(a,b)$ for $a \in \{0,1,2\}$ and $b \in \{0,1,2\}$, so that the author proposes a variant of inductive VBGC stating that:

“Any even positive integer $2m > 2 \cdot 2^{(a+1)(b+2)(a+b+1)}$ can be written as the sum of at least one pair of distinct o-primeths ${}^a P_x > {}^b P_y$, with the positive integers pair (a,b) , with $a \geq b \geq 0$ defining the (recursive) orders of each of those o-primeths AND the pair of distinct positive integers (x,y) , with $x > y > 1$ defining the indexes of each of those o-primeths.”

- p. The function $f_x(a,b) = 2^{(a+1)(b+2)(a+b+1)}$ has its values in the matrix

$$M_{f_x(a,b)} = \begin{pmatrix} 4 & 64 & 4096 \\ 256 & \cong 2.621 \times 10^5 & \cong 4.295 \times 10^9 \\ \cong 2.621 \times 10^5 & \cong 6.872 \times 10^{10} & \cong 1.153 \times 10^{18} \end{pmatrix} \text{ in which each element is}$$

larger than its correspondent element from $M_{f(a,b)} = \begin{pmatrix} 3 & 3 & 2564 \\ 3 & 40306 & 1765126 \\ 2564 & 1765126 & 161352166 \end{pmatrix}$

5. AND

- a. for $(a,b) = (1,0)$ AND $m > 28$, it will always exist at least one pair of finite distinct positive integers (x,y) , with $x > y > 1$ AND ${}^1 P_x + {}^0 P_y = 2m$ AND x (or y) in the double-open interval $(\ln(2m), 2m / \ln(2m))$.

- i. **Important note:** VBGC is much “stronger” and general than BGC and proposes a much more rapid and efficient (at-least-one-GIP)-sieve than the GKRC. The GM of GIPs generated by VBGC has a smaller nof. lines than the GM of GIPs generated by GKRC. VBGC is a useful optimized sieve to push forward the limit

$4 \cdot 10^{18}$ to which BGC was verified to hold [53]. When verifying BGC for a very large number N , one can use the VBGC(a,b) with a minimal positive value for the difference $N - f(a,b)$.

6. Important note: VBGC essentially (and alternatively) states that there is an infinite number of conjectures indexable as VBGC(a,b), all stronger than BGC, EACH of if associated with a pair (a,b) , with $a \geq b > 0$ AND a finite positive integer $n_{a,b} = f(a,b)$.

a. VBGC(0,0) is in fact ntBGC.

VBGC 1.2 – secondary statements (also part of VBGC):

1. The different special cases of VBGC can be named after the pair (a,b) [VBGC(a,b)] AND:

a. **VBGC(0,0)** is in fact ntBGC (defined in the Part B of this article)

b. **VBGC(1,0)**^[1] is a GLC stronger and more elegant than ntBGC, as it acts on a limit $2f(1,0) = 6$ identical to ntBGC inferior limit (which is $2f(0,0) = 6$) BUT the associated $G_{1,0}(m)$ (which counts the number of pairs of possible GIPs for any even integer $m > 3$) has significantly smaller values than the function $G_{0,0}(m)$ of ntBGC [which is VBGC(0,0)]

c. **VBGC(2,0)** is obviously a stronger GLC than VBGC(1,0) is AND ALSO $G_{2,0}(m)$ has smaller non-0 values than $G_{1,1}(m)$ for $m \in (f(2,0), \infty)$

d. **VBGC(1,1)** (anticipated by my discovery of **VBGC(1,0) from 2007 and officially registered in 2012 at OSIM**^[1]) is an obviously stronger GLC than VBGC(1,0) and is equivalent to Bayless-Klyve-Oliveira e Silva Goldbach-like Conjecture (BKOS-GLC) published in Oct. 2013 [54] alias “Conjecture 9.1” (rephrased) (tested by these authors up to $2m = 10^9$): all even integers $2m > [2 \cdot 40306 (= 2f(1,1))]$ can be expressed as the sum of at least one pair of prime-indexed primes [PIPs] ($1P_x$ and $1P_y$). This article of Bayless, Klyve and Oliveira (2012, 2013) was based on a previous article by Barnett and Broughan (published in 2009) [55], but BKOS-GLC was an additional result to this 2009 article. Mr. George Anescu (a friend and collaborator) have also helped me to retest VBGC(1,1) up to $2m = 10^{10}$, but also helped me verifying all VBGC(a,b) for all pairs $(a,b) \in \{(1,0), (1,1), (2,0), (2,1), (2,2)\}$ ^[6].

2. When $a \rightarrow \infty, b \rightarrow \infty$ and $m \rightarrow \infty$, $G_{a,b}(f(a,b)+1) \rightarrow 1$ and the “comets” of VBGC(a,b) tend to narrow progressively for each pair of positive integers (a_2, b_2) , with $a_2 > a_1$ and $b_2 > b_1$.

3. All VBGC(a>0,b≥0) can be used to produce more rapid algorithms for the experimental verification of ntBGC for very large positive integers

a. For VBGC(1,0), the average number of attempts (ANA) to find the first pair (x,y) for each integer m, in the interval $[3, 2m]$ tends asymptotically to $\ln(\sqrt{n}) = \ln(n)/2$ when searching

[6] The code-source (written by Mr. George Anescu in Microsoft Visual C++ language/environment using parallel processing) that was used to test BKOS-GLC up to $n=10^{10}$ (using a laptop PC with an Intel^R CoreTM processor i7-3630 QM CPU at 2.4 GHz with 4 processors (8 hyper-threads), can be found at this URL: dragoi.com/test_primes.rar

just the 1-primeths subset in descending manner, starting from the largest 1-primeth $\leq 2m-1$ and verifying if $(2m - {}^1P_x)$ is a 0-primeth

Conclusions on VBGC 1.2:

1. VBGC(a,b) is essentially an extension and generalization of BGC as applied on (the extended and generalized concept of) all ${}^o\wp^*$ subsets of o-primeths.
2. VBGC distinguishes as a very important (unified) conjecture of primes and a very special self-similar propriety of the primes as the rarefied ${}^o\wp^*$ is self-similar to the more dense ${}^{(o-1)}\wp^*$ in respect to the ntBGC. In other words, each of the o-primeths sets behaves as a “summary of” the 0-primeths set in respect to the ntBGC: this is a (quasi)fractal-like BGC-related behavior of the infinite number of the o-primeths sets (Batchko R.G. has also reported other quasi-fractal/quasi-self-similar structure in the distribution of the prime-indexed primes [56]: Batchko also used a similar general definition for primes with recursive prime indexes, briefly named in my article as “o-primeths”). **Essentially, VBGC conjectures that ntBGC is a common propriety of all the o-primeths sets (for any positive integer order o), differing just by the inferior limit of each VBGC(a,b) defined by the function $f(a,b)$. I have called VBGC as “vertical” motivated by the fact that VBGC is a “vertical” (recursive) generalization of the ntBGC on the infinite super-set of o-primeths sets.**
 - a. The set of values of $f(a,b)$ is a set of critical density thresholds/points of each o-primeths set in respect to the set VBGC(a,b) conjectures.

Future challenges for VBGC (to be also approached in the next versions of this article):

1. To calculate the values of the function $f(a,b) = n_{a,b} = n_{b,a}$ and test/verify VBGC(a,b) for large positive integers pairs $(a > 2, b > 2)$ (a,b), but also for the pairs (a,b) with large $(a-b)$ differences.

Potential applications of VBGC (to also be created in the next versions of this article):

1. VBGC can offer a potential infinite set of Goldbach Comets, one for each sub-VBGC applied on each order of o-primeths
2. VBGC can be used to optimize the algorithms of finding/verifying very large primes (o-primeths)/potential primes (o-primeths)
3. VBGC can be used as a model to also formulate a Vertical (generalization) of the Ternary Goldbach Conjecture/Theorem (VTGC)
4. VBGC can be theoretically used to optimize the algorithms of [prime/integer factorization](#)^[URL2,URL3] (the main tool of [cryptography](#))
5. VBGC can offer a rule of decomposition of [Euclidean](#)^[URL2,URL3,URL4]/[non-Euclidean](#)^[URL2] spaces/volumes with a finite $2N$ (positive) integer number of dimensions into pair of spaces, both with a (positive) o-primeth number of dimensions
6. VBGC can be used in [M-Theory](#) to simulate decompositions of $2N$ -branes (with a finite $2N$ [positive] integer number of dimensions) into pair of branes both with a (positive) o-primeth number of dimensions
7. VBGC can be also used to predict possible symmetries/asymmetries in [crystallography](#), as based on o-primeths.

Acknowledgements

I would like to express all my sincere gratitude and appreciation to all my mathematics, physics, chemistry and medicine teachers for their support and fellowship throughout the years, which provided substantial and profound inner motivation for the redaction and completion of this manuscript. I would also like to emphasize my friendship with George Anescu (physicist and mathematician) who helped me verify the VBGC up to $n=10^{10}$.

My special thanks to professor Toma Albu^[7] who had the patience to read and listen my weak voice in mathematics as a hobbyist. Also my sincere gratitude to professor Șerban-Valentin Strătilă^[8] that advised me on the first special case of VBGC discovered in 2007 and he urged me to look for a more general conjecture based on my first observation.

Competing interests

Author has declared that no competing interests exist.

Addendum

Method for verifying VBGC. We have used Microsoft Visual C++. First, we have created (and stored on hard-disk) a set of “.bin” files containing all the standard primes (alias 0-primeths) (a file of ~3.6GigaBytes), the 1-primeths and the 2-primeths respectively, all in the double-open interval $(1, 10^{10})$.

For every (a, b) pair with $a \geq b$, we have verified each ${}^a P_x (> {}^b P_x)$ from the (less) dense subset of ${}^a P$ superposing the double-open interval $(2, 2m \geq 6)$ (starting from that ${}^a P_x$ which was the closest to $2m - 1$ in descending order): we have then verified if the difference $(2m - {}^a P_x)$ is an element in the (more) dense set ${}^b P$ by using binary section method.

We have then computed each value of $f(a, b)$ (with the additional condition ${}^a P_x \neq {}^b P_y \Leftrightarrow {}^a P_x > {}^b P_y$ in at least one Goldbach partition for any $m > f(a, b)$, with ${}^a P_x + {}^b P_y = 2m$). The computing time for determining and verifying $f(2, 1) = f(1, 2) = (n_{2,1} = n_{1,2}) = 1\ 765\ 126$ and

$f(2, 2) = (n_{2,2}) = 161\ 352\ 166$ was about 30 hours.

[7] The CV of Professor Albu T. is also available online ([URL](#))

[8] The CV of Professor Strătilă Ș-V. is also available online ([URL](#))

ENDNOTE ADDITIONAL REFERENCES (in order of citation in this article)

- [1] Weisstein E. W. (1999-2014). "Goldbach Conjecture", web article From MathWorld—A Wolfram Web Resource. ([URL1](#), [URL2](#); [URL3](#))
- [2] Caldwell C.K. (1999-2015). "Goldbach's conjecture (another Prime Pages' Glossary entries)", web article. ([URL](#))
- [3] Oliveira e Silva T. (2014). "Goldbach conjecture verification", web article. ([URL](#))
- [4] Ye J. D., Liu C. (2013). "A Study of Goldbach's conjecture and Polignac's conjecture equivalence issues", IACR Cryptology ePrint Archive, Volume 2013 ([URL](#))
- [5] Ye J. D., Liu C. (2014). "A Study of Relationship of RSA with Goldbach's Conjecture and Its Properties" ([URL1](#), [URL2](#))
- [6] Liu C. (2015). "A Study of Relationship Among Goldbach Conjecture, Twin Prime and Fibonacci Number" ([URL1](#), [URL2](#))
- [7] Liu C., Chang C-C., Wu Z-P., Ye S-L (2015). "A Study of Relationship between RSA Public Key Cryptosystem and Goldbach's Conjecture Properties", International Journal of Network Security, Vol.17, No.4, PP.445-453, July 2015 ([URL](#))
- [8] Fernandez N. (1999). "The Exploring Primeness Project", website. ([URL1](#), [URL2](#), [URL3-OIES page](#), [URL4-OIES page](#))
- [9] Weisstein E. W. (1999-2015). "Prime Number", web article From MathWorld—A Wolfram Web Resource. ([URL](#))
- [10] Wells, D (1986). "The Penguin Dictionary of Curious and Interesting Numbers", Middlesex, England: Penguin Books, 1986, p. 31. ([URL](#))
- [11] Weisstein E. W. (1999-2015). "Goldbach Partition", web article ([URL](#)); Some JavaScript/ Wolfram Language online calculator of Goldbach partitions can be found at: [URL1](#), [URL2](#), [URL3](#))
- [12] Granville A. (1993). "Harald Cramér and the distribution of prime numbers" (based on a lecture presented on 24th September 1993 at the Cramér symposium in Stockholm. ([URL](#))
- [13] Granville A. (2009). "Different Approaches to the Distribution of Primes", Milan Journal of Mathematic vol. 78 (2009), p. 1–25 ([URL](#))
- [14] Soundararajan K.(2006). "The distribution of prime numbers" ([URL](#))
- [15] Diamond H.G.(1982). "Elementary methods in the study of the distribution of prime numbers", Bull. Amer. Math. Soc. (N.S.), Volume 7, Number 3 (1982), p. 553-589. ([URL](#))
- [16] Liang W.,Yan H., Zhi-cheng D. (2006). "Fractal in the statistics of Goldbach partition" ([URL](#))
- [17] Weisstein E. W. (1999-2014). "Euclid's Theorems", web article From MathWorld—A Wolfram Web Resource. ([URL](#))
- [18] Weisstein E. W. (1999-2014). "Prime Number Theorem", web article From MathWorld—A Wolfram Web Resource. ([URL](#))
- [19] Pakianathan J. and Winfree T. (2011). "Quota Complexes, Persistent Homology and the Goldbach Conjecture", pages 9-10 ([URL](#))
- [20] Zhang S. (2008, 2010). "Goldbach Conjecture and the least prime number in an arithmetic progression", page 2 ([URL](#))
- [21] Wikipedia contributors (last update: 2 Aug. 2015). "Goldbach's weak conjecture", Wikipedia, The Free Encyclopedia ([URL](#) accessed on 13 December 2015)
- [22] Helfgott H.A. (2013). "The ternary Goldbach conjecture is true"* ([URL1](#), [URL2](#), [URL3](#)) (*although it still has to go through the formalities of publication, Helfgott's preprint is endorsed and believed to be true by top mathematicians, including the Fields medalist Terence Tao who showed in 2012 that any odd integer is the sum of at most 5 primes, as can be found at: [URL1](#), [URL2](#))
- [23] Helfgott H.A. (2014, 2015). "The ternary Goldbach problem", Snapshots of modern mathematics from Oberwolfach, No. 3/2014 ([URL1](#); [URL2](#), [URL3](#))
- [24] Platt D.J. (2014). "Proving Goldbach's Weak Conjecture" ([URL](#))
- [25] Lemoine E.(1894). "L'intermédiaire des mathématiciens", 1 (1894), 179; *ibid* 3 (1896), page 151
- [26] Wikipedia contributors (last update: 25 Nov 2014). "Lemoine's conjecture", Wikipedia, The Free Encyclopedia ([URL](#) accessed on 4 Jan 2016)
- [27] Levy H.(1963). "On Goldbach's Conjecture", Math. Gaz. 47 (1963): page 274
- [28] Weisstein E. W. (1999-2014). "Levy's Conjecture", web article From MathWorld—A Wolfram Web Resource. ([URL](#))
- [29] Kiltinen J. and Young P. (September 1984). "Goldbach, Lemoine, and a Know/Don't Know Problem", Mathematics Magazine (Mathematical Association of America) 58 (4): pages 195–203 ([URL1](#), [URL2](#))
- [30] Smarandache F.(1999, 2000[republished], 2007[republished]). "Conjectures on partitions of integers as summations of primes", "Math Power", Pima Community College, Tucson, AZ, USA, Vol. 5, No. 9, pp. 2-4, September 1999; ([URL](#))
- [31] Weisstein E. W. (1999-2014). "Twin Prime Conjecture", web article From MathWorld—A Wolfram Web Resource. ([URL](#))
- [32] Ikorong G.A.N (2007). "Around the twin primes conjecture and the Goldbach conjecture I", Analele Științifice ale Universității "Al. I. Cuza" din Iași (S.N.), Matematică, Tomul LIII, 2007, f.1 ([URL](#))
- [33] Ikorong G.A.N (2008). "Playing with the Twin Primes Conjecture and the Goldbach Conjecture", Alabama Journal of Mathematics, Spring/Fall 2008, pages 45-52 ([URL](#))
- [34] Gerstein L.J. (1993). "A Reformulation of the Goldbach Conjecture", Mathematics Magazine vol. 66, no.1, February 1993. pages 44-45 ([URL](#))
- [35] Polymath D.H.J. (2014). "The "bounded gaps between primes" Polymath project - a retrospective" ([URL](#))
- [36] Chen, J.R. (1966). "On the representation of a large even integer as the sum of a prime and the product of at most two primes", Kexue Tongbao 11 (9): pages 385–386
- [37] Chen, J.R. (1973). "On the representation of a larger even integer as the sum of a prime and the product of at most two primes", Scientia Sinica 16: pages 157–176
- [38] Cheng-Dong P., Xia-Xi D., Yuan W. (1975). "On the representation of every large even integer as a sum of a prime and an almost prime", Scientia Sinica Vol. XVIII No.5 Sept.-Oct. 1975: pages 599–610 ([URL](#))
- [39] Cai, Y.C. (2002). "Chen's Theorem with Small Primes", Acta Mathematica Sinica 18 (3): pages 597–604 ([URL](#))
- [40] Cai, Y.C. (2008). "On Chen's theorem (II)", Journal of Number Theory, Volume 128, Issue 5, May 2008, pages: 1336–1357 ([URL](#))

-
- [41] [Rivera C. \(1999-2001\)](#). “Conjecture 22. A stronger version of the Goldbach Conjecture (by Mr. Rudolf Knjzek, from Austria)”, web article from Prime Problems & Puzzles. ([URL](#))
- [42] [Rivera C. \(1999-2001\)](#). “Conjecture 22. A stronger version of the Goldbach Conjecture (by Mr. Rudolf Knjzek, from Austria and narrowed by Rivera C.)”, web article from Prime Problems & Puzzles. ([URL](#))
- [43] [Woon M.S.C. \(2000\)](#). “On Partitions of Goldbach’s Conjecture” ([URL](#))
- [44] [Sun Z-W. \(2013, 2014\)](#). Chapter “Problems on combinatorial properties of primes” (19 pages) in “Number Theory: Plowing and Starring Through High Wave Forms: Proceedings of the 7th China-Japan Seminar” (edited by Kaneko M., Kanemitsu S. and Liu J.), pages: 169 – 188 ([URL1-book excerpt](#), [URL2- full book](#))
- [45] [Sun Z-W. \(2014\)](#). “Towards the Twin Prime Conjecture”, A talk given at: NCTS (Hsinchu, Taiwan, August 6, 2014), Northwest University (Xi’an, October 26, 2014) and at Center for Combinatorics, Nankai University (Tianjin, Nov. 3, 2014) ([URL](#))
- [46] See also Sun’s Z-W. personal web page on which all conjectures are presented in detail ([URL](#))
- [47] See also the first announcement of this conjecture made by Sun Z-W. himself on 6 Feb 2014) ([URL](#))
- [48] See also the sequence [A218829](#) on [OEIS.org](#) proposed by Sun Z-W. ([URL1](#), [URL2](#))
- [49] Alternative terms for “primeths”: “higher-order prime numbers”, “superprime numbers”, “super-prime numbers”, ”super-primes”, ” superprimes” or “prime-indexed primes[PIPs]” ([URL-OIES page](#))
- [50] [Murthy A. \(2005\)](#). “Generalized Partitions and New Ideas on Number Theory and Smarandache Sequences” (book), page 91 ([URL1-book](#), [URL2 – page 181](#))
- [51] [Seleacu V. and Bălăcenoiu I. \(2000\)](#). “Smarandache Notions, Vol. 11” (book), page181 ([URL](#))
- [52] Primes subset (3, 5, 11, 17, 31, 41, 59, 67, 83, 109, 127, 157, ...), also known as sequence **A006450** in [OEIS](#) ([URL-OIES page](#))
- [53] [Oliveira e Silva T. \(30 Dec. 2016\)](#). “Goldbach conjecture verification” (web article) ([URL](#))
- [54] [Bayless J., Klyve D. and Oliveira e Silva T. \(2012, 2013\)](#). “New bounds and computations on prime-indexed primes” (23 pages article,), *Integers: Annual Volume 13* (2013), page 17 ([URL1](#), [URL2](#), [URL3](#))
- [55] [Broughan K.A., Ross Barnett A. \(2009\)](#). “On the Subsequence of Primes Having Prime Subscripts” (10 pages), Article 09.2.3 from the *Journal of Integer Sequences*, Vol. 12 (2009) ([URL1](#), [URL2](#), [URL3](#), [URL4](#))
- [56] [Batchko R.G. \(2014\)](#). “A prime fractal and global quasi-self-similar structure in the distribution of prime-indexed primes”, ArXiv article, submitted on 10 May 2014 (v1), last revised 17 May 2014 (this version, v2) ([URL1](#), [URL2](#))