

# Gravity as the cause for nuclear force and cosmological acceleration

Henok Tadesse, Electrical Engineer, BSc.

Ethiopia, Debrezeit, P.O Box 412

Mobile: +251 910 751339; email [entkidmt@yahoo.com](mailto:entkidmt@yahoo.com) or [wchmar@gmail.com](mailto:wchmar@gmail.com)

21 January 2017

## Abstract

The origin of the force holding protons and neutrons together in the nucleus is one of the daunting puzzles of physics, regardless of the Standard Model explanation. One possible consideration is the force of gravity as responsible for the stability of the nucleus. However, this idea will be immediately dismissed because gravitational force as we know it is weaker than electromagnetic force by a factor of about  $8 \times 10^{-37}$ . We know gravitation as was introduced by Newton and have been stuck with that for centuries. This paper reveals a drastically different law of gravitation that ultimately resolves the mystery of nuclear force. This theory is also promising to explain the phenomenon of cosmological redshift.

## Introduction

The reason why the nucleus doesn't fly apart under the electrostatic repulsion forces of its protons, packed within an extremely small space ( the diameter of the nucleus is of the order of  $1 \times 10^{-14}$  m ), was one of the long standing mysteries of physics. The origin of the force holding protons and neutrons together in the nucleus is explained in the Standard Model by the interaction of elementary particles called Quarks and Gluons. In this paper, we propose that both the stability of the nucleus and cosmological acceleration phenomenon may be due to the force of gravitation.

## Gravity

In a previous paper[1] I have proposed that gravity is a difference between electrostatic attraction and repulsion forces. This idea was also supported by a compelling theory ( Apparent Source Theory) and experimental and observational evidences[1].

I restated Coulomb's law as [1]:

$$F_{att} = \frac{1}{4\pi\epsilon_{att}} \frac{Q1 \cdot Q2}{r^2}$$
$$F_{rep} = \frac{1}{4\pi\epsilon_{rep}} \frac{Q1 \cdot Q2}{r^2}$$

where  $\epsilon_{att}$  is the permittivity for opposite charges and  $\epsilon_{rep}$  is for similar charges.

The idea that gravity is a difference between electrostatic attraction and repulsion forces was/is a very compelling one.

## Nuclear force

The mystery regarding nuclear force can be stated as:

*Why does the nucleus not fly apart under the electrostatic repulsion force of the protons?*

*And what holds the protons together in the nucleus ? i.e. the protons and neutrons would drift away from each other if there is no some kind of binding force ?*

We propose here that nuclear force is in fact gravitational. Since gravitational force as we know it cannot account for nuclear stability, we have to re-write our understanding of it.

The force of gravity is a difference between the attractive and repulsive electrostatic forces. This difference results from different expressions ( formulas ) for attractive and repulsive electrostatic forces.

$$F_{att} = f(r) \quad \text{and} \quad F_{rep} = g(r)$$

where

$F_{att}$  is the electrostatic attraction force,

$F_{rep}$  is the electrostatic repulsion force,

$f(r)$  is the expression for distance dependence of electrostatic attractive force,

$g(r)$  is the expression for distance dependence of electrostatic repulsive force and


$r$  is the distance between the two charges

For centuries, we have been stuck with the inverse squared distance Newton's law of gravitational force and that may have been the root problem.

Now we put down the requirements for the new laws of electrostatic attraction, electrostatic repulsion and gravitational forces.

1. At extremely small distances, as in the distance between protons in the nucleus, the attractive force should be greater than the repulsive force in such a way that the gravitational force, which is  $F_{att} - F_{rep}$ , should be greater than the repulsion force  $F_{rep}$ , by a factor of about 137 because it is known that the strong nuclear force of the Standard Model is greater than electromagnetic force by this factor.
2. This gravitational force should diminish to nearly zero at distances of the order of the diameter of atoms and beyond. At macroscopic distances, the attraction and repulsion forces should essentially follow inverse squared distance dependence of Coulomb's law.

Graphically, the attractive and repulsive electrostatic forces look like as follows, qualitatively.



We see from the above curves that, at a distance of  $1 \times 10^{-14} \text{ m}$ , the difference between the attractive and repulsive electrostatic forces, which is the gravitational force, is greater than the electrostatic repulsion force. At large distances, the two formulae essentially follow the inverse squared distance law with which we are familiar.

The above is just a qualitative graphical representation of the forces. The exact formulae for the electrostatic attraction and repulsion forces and for the gravitational force should be revealed by further research. We simply make a heuristic attempt in this paper.

The electrostatic force can be written as :

$$F_{att} = \frac{1}{4\pi\epsilon_{att}} \frac{Q1 \cdot Q2}{r^2}$$

$$F_{rep} = \frac{1}{4\pi\epsilon_{rep}} \frac{Q1 \cdot Q2}{r^2}$$

We introduce an additional factor for each formula that will make the electrostatic attraction force greater than the electrostatic repulsion force at distances of the order of the diameter of the nucleus and essentially reduce to 1 at distances comparable to the diameter of the atom. Note that we are just trying to find an example of a factor that fulfills the requirement we stated . We don't mean that this is the exact expression.

$$F_{att} = \frac{1}{4\pi\epsilon_0} \left( 1 + \frac{1}{K_{att} r^2} \right) \frac{Q1 \cdot Q2}{r^2}$$

$$F_{rep} = \frac{1}{4\pi\epsilon_0} \left( 1 + \frac{1}{K_{rep} r^2} \right) \frac{Q1 \cdot Q2}{r^2}$$

For the factors to diminish to nearly 1 for distances of about  $10^{-10}$  m (the diameter of atoms) , the constants  $K_{att}$  and  $K_{rep}$  should be very large numbers.

For example, if  $r = 10^{-11}$  m , the factor  $K_{rep}$  should be about  $10^{30}$  for  $\epsilon_{rep}$  to be  $1.00000001 * (1/(4\pi\epsilon_0))$  ).

Now let us estimate the values of  $K_{att}$  and  $K_{rep}$  so that gravitational force in the nucleus is about 137 times the electrostatic repulsion force. i.e.

$$\frac{F_{att} - F_{rep}}{F_{rep}} = 137 \Rightarrow \frac{\frac{1}{K_{att} r^2} - \frac{1}{K_{rep} r^2}}{1 + \frac{1}{K_{rep} r^2}} = 137 \Rightarrow \frac{\frac{1}{K_{att}} - \frac{1}{K_{rep}}}{r^2 + \frac{1}{K_{rep}}} = 137$$

For example, if we assume  $K_{rep} = 10^{30}$  ,  $r = 10^{-14}$  m , then  $K_{att}$  will be:

$$K_{att} = 7.22647781471311 \times 10^{25}$$


### The nucleus as a quantum system

Since gravitational force is greater than the repulsion force of the protons, the protons and the neutrons will be attracted to the center of the nucleus, in the same way that an electron in an atom is attracted towards its nucleus. Therefore, the protons and neutrons should revolve around the center of the nucleus to avoid falling into the center. From this follows allowable orbitals of quantum mechanics in the nucleus. Therefore, a nucleus, like the atom, is a quantum system and will have only discrete states.

## Cosmological acceleration

We can even add another requirement for the formulas expressing the two forces. That is, the formulas should also be such that the electrostatic repulsion force becomes greater than the electrostatic attraction force beyond some astronomical distance, so that gravity turns from attractive force to repulsive force beyond a certain astronomical distance. This may explain why the universe doesn't collapse and why galaxies are moving away from us and from each other ( i.e. 'expanding universe' ).

All we need is introduce yet another factor. Graphically the forces look like as follows in astronomical scales, qualitatively.



The forces with the additional factors are given by:

$$F_{att} = \frac{1}{4\pi\epsilon_0} \left( 1 + \frac{1}{K_{att} r^2} \right) ( 1 + K_{astatt} r ) \frac{Q1 \cdot Q2}{r^2}$$

$$F_{rep} = \frac{1}{4\pi\epsilon_0} \left( 1 + \frac{1}{K_{rep} r^2} \right) ( 1 + K_{astrep} r ) \frac{Q1 \cdot Q2}{r^2}$$

$K_{astatt}$  and  $K_{astrep}$  can be calculated from astronomical observations.

## Conclusion

In this paper, the mystery of nuclear force has been revealed. The nuclear force is basically a gravitational force, but not gravitational force as we know it. The laws of electrostatic force and gravitation have been modified to explain nuclear force. This theory is also promising to explain the cause of cosmological redshift. There is only one fundamental force in the universe: electromagnetic force.

Thanks to God and His Mother Our Lady Saint Virgin Mary

## References

1. Absolute/Relative Motion and the Speed of Light, Electromagnetism, Inertia and Universal Speed Limit  $c$  - an Alternative Interpretation and Theoretical Framework, Henok Tadesse, Vixra

<http://vixra.org/pdf/1508.0178v9.pdf>

## Bibliography

1. The Cult of the Nuclear Force

<https://nohiggs.wordpress.com/2011/10/14/when-science-and-reality-do-not-agree/>

2. The Conventional Concept of a Nuclear Strong Force is a Colossal Blunder,

<http://www.sjsu.edu/faculty/watkins/strongforce.htm>

3. The Farce of Modern Physics, David Pratt, 2008

<http://www.davidpratt.info/farce.htm>

4. Strong interaction, Wikipedia

[https://en.wikipedia.org/wiki/Strong\\_interaction](https://en.wikipedia.org/wiki/Strong_interaction)

5. Nuclear Forces

<http://www.alternativephysics.org/book/NuclearForces.htm>