

Updated Elementary Proofs of Polignac Prime Conjecture, Goldbach Conjecture, Twin Prime Conjecture, Cousin Prime Conjecture, and Sexy Prime Conjecture

Stephen Marshall

2 February 2017

Abstract

This paper presents a complete and exhaustive proof of the Polignac Prime Conjecture. The approach to this proof uses same logic that Euclid used to prove there are an infinite number of prime numbers. Then we use a proof found in Reference 1, that if $p > 1$ and $d > 0$ are integers, that p and $p + d$ are both primes if and only if for integer n :

$$n = (p-1)! \left(\frac{1}{p} + \frac{(-1)^d d!}{p+d} \right) + \frac{1}{p} + \frac{1}{p+d}$$

We use this proof for $d = 2k$ to prove the infinitude of Polignac prime numbers.

The author would like to give many thanks to the authors of 1001 Problems in Classical Number Theory, Jean-Marie De Koninck and Armel Mercier, 2004, Exercise Number 161 (see Reference 1). The proof provided in Exercise 6 is the key to making this paper on the Polignac Prime Conjecture possible.

Additionally, our proof of the Polignac Prime Conjecture leads to proofs of several other significant number theory conjectures such as the Goldbach Conjecture, Twin Prime Conjecture, Cousin Prime Conjecture, and Sexy Prime Conjecture. Our proof of Polignac's Prime Conjecture provides significant accomplishments to Number Theory,

yielding proofs to several conjectures in number theory that has gone unproven for hundreds of years.

Introduction

The Polignac prime conjecture, was made by Alphonse de Polignac in 1849. Alphonse de Polignac (1826 – 1863) was a French mathematician whose father, Jules de Polignac (1780-1847) was prime minister of Charles X until the Bourbon dynasty was overthrown in 1830. Polignac attended the École Polytechnique (commonly known as Polytechnique) a French public institution of higher education and research, located in Palaiseau near Paris. In 1849, the year Alphonse de Polignac was admitted to Polytechnique, he made what's known as Polignac's conjecture:

For every positive integer k , there are infinitely many prime gaps of size $2k$.

Alphonse de Polignac made other significant contributions to number theory, including the de Polignac's formula, which gives the prime factorization of $n!$, the factorial of n , where $n \geq 1$ is a positive integer.

Proof of Polignac's Conjecture

In number theory, Polignac's conjecture states:

For any positive integer k , then for any positive even number $2k$, there are infinitely many prime gaps of size $2k$. In other words, there are infinitely many cases of two consecutive prime numbers with difference $2k$. Mathematically stated:

There exist infinitely many cases where, both p and $p + 2k$ are prime.

First we shall assume that the set of Polignac Primes are finite and then we shall prove that this is false, which shall prove that are Polignac Primes are infinite. First let us define the finite set of Polignac Primes as p_i where i is finite. Our goal is to prove that i is infinite to disprove our assumption of finiteness. Even though we have assumed that the set of finite Polignac Primes since there are an infinite number of prime numbers we can

pick a prime number p which is outside the finite Polignac primes. Therefore a prime $p+2k$ cannot exist that is prime otherwise we would have discovered a pair of Polignac primes which are outside our assumed set of finite Polignac primes. Thus all we need to do is to prove that there exists a Polignac prime outside our assumed finite set is to prove that $p + 2k$ is prime since we already know that a prime number p exists outside the finite set of Polignac primes.

Now we shall proceed to prove at least one $p + 2k$ is prime as follows:

We use the proof, provided in Reference 1, that if $p > 1$ and $d > 0$ are integers, that p and $p+ d$ are both primes if and only if for positive integer n :

$$n = (p-1)! \left(\frac{1}{p} + \frac{(-1)^d d!}{p+d} \right) + \frac{1}{p} + \frac{1}{p+d}$$

Please note that \wedge indicates an exponent, $(-1)^d$ is equivalent to (-1) raised to the d power

For our case p is known to be prime and $d = 2k$ for Polignac primes, where k is any positive integer, therefore:

$$n = (p-1)! \left(\frac{1}{p} + \frac{(-1)^{2k} (2k!)}{p+2k} \right) + \frac{1}{p} + \frac{1}{p+2k}$$

Multiplying by p , and since $2k$ is always even, then $(-1)^{2k} = 1$

$$np = (p-1)! \left(\frac{1}{p} + \frac{2k!}{p+2k} \right) + 1 + \frac{p}{p+2k}$$

Multiplying by $(p + 2k)$,

$$(p + 2k) np = (p + 2k)(p)! \left(\frac{1}{p} + \frac{2k!}{p+2k} \right) + p + 2k + p$$

Reducing again,

$$(p + 2k) np = (p)! \left(\frac{(p+2k)}{p} + 2k! \right) + p + 2k + p$$

Factoring out, $(p)!$,

$$(p + 2k) np = p(p-1)! \left(\frac{(p+2k)}{p} + 2k! \right) + 2p + 2k$$

And reducing one final time,

$$(p + 2k) np = (p-1)!(p + 2k + 2pk!) + 2p + 2k$$

We already know p is prime, therefore, $p = \text{integer}$. Since p is an integer and by definition k is an integer, the right hand side of the above equation is an integer (likewise the left hand side of the equation must also be an integer). Since the right hand side of the above equation is an integer and p and k are integers on the left hand side of the equation, then $(p+2k)$ is also an integer. Therefore there are only 4 possibilities (see 1, 2a, 2b, and 2c below) that can hold for n so the left hand side of the above equation is an integer, they are as follows.

1) n is an integer, or

2) n is a rational fraction that is divisible by p . This implies that $n = \frac{x}{p}$ where, p is prime

and x is an integer. This results in the following three possibilities:

a. Since $n = \frac{x}{p}$, then $p = \frac{x}{n}$, since p is prime, then p is only divisible by p and 1 ,

therefore, the first possibility is for n to be equal to p or 1 in this case, which are both integers, thus n is an integer for this first case.

b. Since $n = \frac{x}{p}$, and x is an integer, then x is not evenly divisible by p unless $x =$

p , or x is a multiple of p , where $x = yp$, for any integer y . Therefore n is an integer for $x = p$ and $x = yp$.

c. For all other cases of, integer x , $n = \frac{x}{p}$, n is not an integer.

To prove there is a Polignac Prime, outside our set of finite Polignac Primes, we only need to prove that there is at least one value of n that is an integer, outside our finite set. There can be an infinite number of values of n that are not integers, but that will not negate the existence of one Polignac Prime, outside our finite set of Polignac Primes.

First the only way that n cannot be an integer is if every n satisfies paragraph 2.c above, namely, $n = \frac{x}{p}$, where x is an integer, $x \neq p$, $x \neq yp$, $n \neq p$, and $n \neq 1$ for any integer y . To

prove there exists at least one Polignac Prime outside our finite set, we will assume that no integer n exists and therefore no Polignac Primes exist outside our finite set. Then we shall prove our assumption to be false.

Proof: Assumption no values of n are integers, specifically, every value of n is

$n = \frac{x}{p}$, where x is an integer, $x \neq p$, $x \neq yp$, $n \neq p$, and $n \neq 1$, for any integer y .

Paragraphs 1, 2.a, and 2.b prove cases where n can be an integer, therefore our assumption is false and **there exist values of n that are integers.**

Since we have already shown that p and $p + 2k$, where $d = 2k$, are both primes if and only if for integer n :

$$n = (p-1)! \left(\frac{1}{p} + \frac{(-1)^d d!}{p+d} \right) + \frac{1}{p} + \frac{1}{p+d}$$

It suffices to show that there is at least one integer n to prove there exists a Polignac Prime outside our set of finite set of Polignac Primes.

Since there exists an $n = \text{integer}$, we have proven that there is at least one p and $p + 2k$ that are both prime. Since we shown earlier that if $p + 2k$ is prime then it also is not in the finite set of $p_i, p_i + 2k$ Polignac primes, therefore, since we have proven that there is at least one $p+2k$ that is prime, then we have proven that there is a Polignac prime outside the our assumed finite set of Polignac primes. This is a contradiction from our assumption that the set of Polignac primes is finite, therefore, by contradiction the set of Polignac primes is infinite. Also this same proof can be repeated infinitely for each finite set of Polignac primes, in other words a new Polignac prime can added to each set of finite Polignac primes. This thoroughly proves that an infinite number of Polignac primes exist.

Our proof of infinite Polignac primes also proves several other Conjectures since Polignac primes are generalized form of other conjectures. For $k = 1$, the Polignac primes become the Twin Prime Conjecture and proves there are an infinite number of twin primes. For $k = 2$, the proof of the Polignac Conjecture proves there are infinitely many Cousin Primes $(p, p + 4)$. For $k = 3$, proof of the Polignac Conjecture proves there are infinitely many Sexy Primes $(p, p + 6)$.

Our proof of infinite Polignac primes also provides a major breakthrough to prove the Goldbach's Conjecture. The Goldbach Conjecture is one of the oldest unsolved problems in number theory. It states:

Every even integer greater than 2 can be expressed as the sum of two primes.

Our proof of infinite Polignac primes can also be used to prove the Goldbach Conjecture. First we shall assume that not every even number > 2 is a sum of two primes a and b . However, Polignac's Conjecture states:

For any positive integer k , then for any positive even number $2k$, there are infinitely many prime gaps of size $2k$. In other words, there are infinitely many cases of two consecutive prime numbers with difference $2k$.

Another way of stating the Polignac Conjecture is; for every prime number a and b there are an infinitely many cases where $a - b = 2k$, for any positive integer k . So we can state the following:

$$a - b = 2k$$

$$\text{Reducing, } a = 2k + b$$

$$\text{Adding a prime number } b \text{ to both sides, } a + b = 2k + 2b$$

$$\text{Finally, reducing, } a + b = 2(k + b)$$

The right hand side of the equation above, $2(k + b)$, is even and includes all positive even numbers since k is any positive integer and b is from an infinite number of prime numbers. Since k is any positive integer and b is selected from an infinite cases of prime numbers, then

$$\text{For any integer } n, a + b = 2(k + b) = 2n$$

$$\text{Thus, } a + b = 2n, \text{ for every even number } 2n$$

Therefore, we have proven the Goldbach Conjecture, specifically, every even integer greater than 2 can be expressed as the sum of two primes, i.e, $n > 1$.

References:

- 1) 1001 Problems in Classical Number Theory, Jean-Marie De Koninck and Armel Mercier, 2004, Exercise Number 161