

# Superconducting Josephson Junction Laser

*A team of researchers led by Leo Kouwenhoven at TU Delft has demonstrated an on-chip microwave laser based on a fundamental property of superconductivity, the ac Josephson effect. They embedded a small section of an interrupted superconductor, a Josephson junction, in a carefully engineered on-chip cavity. Such a device opens the door to many applications in which microwave radiation with minimal dissipation is key, for example in controlling qubits in a scalable quantum computer. [8]*

*Optical scientists from the Warsaw Laser Centre of the Institute of Physical Chemistry of the Polish Academy of Sciences and the Faculty of Physics of the University of Warsaw have generated ultrashort laser pulses in an optical fiber with a method previously considered to be physically impossible. [7]*

*Researchers at the Max Planck Institute for the Science of Light in Erlangen have discovered a new mechanism for guiding light in photonic crystal fiber (PCF). [6]*

*Scientists behind a theory that the speed of light is variable - and not constant as Einstein suggested - have made a prediction that could be tested. [5]*

*Physicists' greatest hope for 2015, then, is that one of these experiments will show where Einstein got off track, so someone else can jump in and get closer to his long-sought "theory of everything."*

*This article is part of our annual "Year In Ideas" package, which looks forward to the most important science stories we can expect in the coming year. It was originally published in the January 2015 issue of Popular Science. [4]*

*The self maintained electric potential of the accelerating charges equivalent with the General Relativity space-time curvature, and since it is true on the quantum level also, gives the base of the Quantum Gravity.*

*The magnetic induction creates a negative electric field, causing an electromagnetic inertia responsible for the relativistic mass change; it is the mysterious Higgs Field giving mass to the particles.*

*The Planck Distribution Law of the electromagnetic oscillators explains the electron/proton mass rate by the diffraction patterns. The accelerating charges explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the*

*electron's spin also, building the bridge between the Classical and Relativistic Quantum Theories.*

## Contents

Preface.....	3
Popular questions about the Higgs Field and General Relativity:.....	3
How can we answer these questions? .....	3
Researchers demonstrate new type of laser .....	4
Superconducting Josephson junction .....	4
Josephson junction laser.....	4
Low-loss quantum control.....	5
Researchers develop surprising technique for ultrashort laser pulses .....	5
Impossible (but working!) recipe for ultrashort laser pulses .....	6
Optical fibre with Einstein effect .....	7
Photonic crystals give butterflies their colour and can also guide light .....	7
The holes define the space in a photonic crystal fibre.....	8
In a twisted PCF, the light follows the shortest path in the interior of the fibre .....	8
A twisted PCF as a large-scale environmental sensor.....	8
Theory that challenges Einstein's physics could soon be put to the test .....	9
RADICAL IDEA .....	9
THE HORIZON PROBLEM.....	10
The Test in 2015 .....	10
The Classical Relativistic effect .....	11
The Relativistic Quantum Mechanics .....	11
The Heisenberg Uncertainty Relation.....	11
The General Relativity - Electromagnetic inertia and mass.....	11
Electromagnetic Induction .....	11
Relativistic change of mass.....	11
The frequency dependence of mass .....	12
Electron – Proton mass rate .....	12
The weak interaction .....	12
The General Weak Interaction .....	13
The Higgs boson .....	14
Higgs mechanism .....	14

Gravity from the point of view of quantum physics .....	15
What is the Spin? .....	15
The Graviton .....	15
Conclusions .....	15
References .....	16

Author: George Rajna

## Preface

### **Popular questions about the Higgs Field and General Relativity:**

- 1.) If the Higgs field is responsible for imbuing particles with mass, and mass is responsible for gravity, is it possible that the Higgs field will provide the missing link between general relativity and quantum mechanics i.e. could the Higgs field be the basis of a quantum theory of gravity?
- 2.) Can the theoretical Higgs Field be used as the “cause” of relativistic momentum or relativistic kinetic energy of a moving body?
- 3.) Does Einstein's General Relativity need to be adjusted for the Higgs field?
- 4.) Since the Higgs field gives most particles mass, and permeates all space, then GR needs the Higgs field to be a theory of space?
- 5.) So where GR is highly curved, the Higgs field is also curved? And does a highly curved Higgs field affect the way particles acquire mass? For that matter, a curved space-time would also curve electromagnetic field?

### **How can we answer these questions?**

Discovering the magnetic effect of the electric current from the observed effects of the accelerating electrons - causing naturally the experienced changes of the electric field potential along the electric wire - the accelerating electrons explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the electron's spin also, building the bridge between the Classical and Quantum Theories. [1]

Another bridge between the classical and quantum mechanics in the realm of relativity is that the charge distribution is lowering in the reference frame of the accelerating charges linearly:  $ds/dt = at$  (time coordinate), but in the reference frame of the current it is parabolic:  $s = a/2 t^2$  (geometric coordinate).

One origin of the Quantum Physics is the Planck Distribution Law of the electromagnetic oscillators, giving equal intensity for 2 different wavelengths on any temperature. Any of these two wavelengths will give equal intensity diffraction patterns, building different asymmetric constructions, for example proton - electron structures (atoms), molecules, etc. Since the particles are centers of diffraction patterns they also have particle – wave duality as the electromagnetic waves have. [2]

The Electroweak Interaction shows that the Weak Interaction is basically electromagnetic in nature. The arrow of time shows the entropy grows by changing the temperature dependent diffraction patterns of the electromagnetic oscillators.

## Researchers demonstrate new type of laser

Lasers are everywhere nowadays: Doctors use them to correct eyesight, cashiers to scan your groceries, and quantum scientist to control qubits in the future quantum computer. For most applications, the current bulky, energy-inefficient lasers are fine, but quantum scientist work at extremely low temperatures and on very small scales. For over 40 years, they have been searching for efficient and precise microwave lasers that will not disturb the very cold environment in which quantum technology works.

A team of researchers led by Leo Kouwenhoven at TU Delft has demonstrated an on-chip microwave laser based on a fundamental property of superconductivity, the ac Josephson effect. They embedded a small section of an interrupted superconductor, a Josephson junction, in a carefully engineered on-chip cavity. Such a device opens the door to many applications in which microwave radiation with minimal dissipation is key, for example in controlling qubits in a scalable quantum computer.

The scientists have published their work in Science on the 3rd of March.

Lasers have the unique ability to emit perfectly synchronized, coherent light. This means that the linewidth (corresponding to the color) is very narrow. Typically lasers are made from a large number of emitters (atoms, molecules, or semiconducting carriers) inside a cavity. These conventional lasers are often inefficient, and dissipate a lot of heat while lasing. This makes them difficult to operate in cryogenic environments, such as what is required for operating a quantum computer.

## Superconducting Josephson junction

In 1911, the Dutch physicist Heike Kamerlingh Onnes discovered that some materials transition to a superconducting state at very low temperatures, allowing electrical current to flow without any loss of energy. One of the most important applications of superconductivity is the Josephson effect: if a very short barrier interrupts a piece of superconductor, the electrical carriers tunnel through this non-superconducting material by the laws of quantum mechanics. Moreover, they do so at a very characteristic frequency, which can be varied by an externally applied DC voltage. The Josephson junction is therefore a perfect voltage to light (frequency) converter.

## Josephson junction laser

The scientists at QuTech coupled such a single Josephson junction to a high-quality factor superconducting micro-cavity, no bigger than an ant. The Josephson junction acts like a single atom, while the cavity can be seen as two mirrors for microwave light. When a small DC voltage is applied to this Josephson junction, it emits microwave photons that are on resonance with the cavity frequency. The photons bounce back and forth between two superconducting mirrors, and force the Josephson junction to emit more photons synchronized with the photons in the cavity. By cooling the device down to ultra-low temperatures (< 1 Kelvin) and applying a small DC voltage to the Josephson junction, the researchers observe a coherent beam of microwave photons emitted at the output of the cavity. Because the on-chip laser is made entirely from superconductors, it is very

energy efficient and more stable than previously demonstrated semiconductor-based lasers. It uses less than a picoWatt of power to run, more than 100 billion times less than a light globe.

### Low-loss quantum control

Efficient sources of high quality coherent microwave light are essential in all current designs of the future quantum computer. Microwave bursts are used to read out and transfer information, correct errors and access and control the individual quantum components. While current microwave sources are expensive and inefficient, the Josephson junction laser created at QuTech is energy efficient and offers an on-chip solution that is easy to control and modify. The group is extending their design to use tunable Josephson junctions made from nanowires to allow for microwave burst for fast control of multiple quantum components. In the future, such a device may be able to generate so-called "amplitude-squeezed" light with smaller intensity fluctuations compared to conventional lasers, this is essential in most quantum communication protocols. This work marks an important step towards the control of large quantum systems for quantum computing. [8]

### Researchers develop surprising technique for ultrashort laser pulses

Pulse lasers built entirely on optical fibers are increasingly used by industry. Optical scientists from the Warsaw Laser Centre of the Institute of Physical Chemistry of the Polish Academy of Sciences and the Faculty of Physics of the University of Warsaw have generated ultrashort laser pulses in an optical fiber with a method previously considered to be physically impossible. Their solution is not only useful, but also surprisingly simple.

An innovative fiber laser has been developed at the Laser Centre of the Institute of Physical Chemistry of the Polish Academy of Sciences (IPC PAS) and the Faculty of Physics of the University of Warsaw. Using a simple solution, the Warsaw optical scientists have "forced" one of the types of optical fiber lasers to generate ultrashort, high-energy pulses. The new laser is devoid of any mechanically sensitive external parts, which seems to be especially interesting for future applications. The invention greatly expedites the processing of materials in industrial laser machines.

"Fiber lasers can be built so that all the processes important for the generation and shaping of the ultrashort pulses take place in the fiber itself. Such devices, without any external mechanically sensitive components, operate in a very stable manner, and are ideal for working in difficult conditions," says Dr. Yuriy Stepanenko (IPC PAS).

Laser action in the fiber leads to the generation of a continuous light beam. The release of energy in the shortest possible pulses is, however, much more favourable, since it signifies a great increase of power. Pulses are generated in fiber lasers via a saturable absorber system. When the light intensity is low, the absorber blocks light; when it is high, the absorber lets it through. Since femtosecond pulses have greater intensity than a continuous beam, the parameters of the absorber can be adjusted so that it only admits pulses.

"Up to now, graphene sheets, among others, have been used as the saturable absorbers, in a form of a thin layer deposited on the tip of the fiber. But the diameters of optical fibers are in the order of single microns. Even a little energy crammed in such a small cross-section has a significant density per unit area, affecting the lifetime of the materials. Therefore, if an attempt was made to increase the power of the femtosecond pulses, the graphene on the tip of the connector was destroyed.

Other absorbers, such as carbon nanotubes, may also undergo degradation," explains Jan Szczepanek, a PhD student from the Faculty of Physics of the University of Warsaw.

In order to generate higher energy femtosecond pulses in the optical fiber, the Warsaw physicists decided to improve saturable absorbers of a different type, via the clever use of optical phenomena such as nonlinear effects that cause a change in the refractive index of glass.

### **Impossible (but working!) recipe for ultrashort laser pulses**

A nonlinear artificial saturable absorber works as follows. The plane of polarization of low intensity light beam does not change in the absorber and the output polarizer blocks the light (images at the bottom). At a high enough intensity, ...more

Electric and magnetic fields of light usually oscillate in random, mutually perpendicular directions. When the fields oscillate all the time in the same plane, the wave is called linearly polarized. In classical optics, it is assumed that when such a wave passes through a medium, it experiences a constant refractive index, regardless of the light intensity. In nonlinear optics this is different: At a sufficiently high light intensity, the refractive index begins to increase slightly.

A nonlinear artificial saturable absorber works as follows. At the input, the linearly polarized light is divided into a beam with a low intensity and a beam with a high intensity. The medium of the absorber can be chosen for both light beams to experience a slightly different refractive index in order for them to travel at slightly different (phase) velocities. As a result of the velocity difference, the plane of polarization starts to rotate. At the output of the absorber, there is a polarization filter that only lets through waves oscillating perpendicularly to the plane of polarization of the incoming light. When the laser is operating in continuous mode, the light in the beam is of a relatively low intensity, an optical path difference does not occur, the polarization does not change, and the output filter blocks the light. At a high enough intensity, typical for femtosecond pulses, the rotation of polarization causes the pulse to pass through the polarizer.

For the saturable absorber with polarization rotation to work, the fibre must have different refractive indices in different directions (thus it has to be birefringent), and both indices should also be stable. The problem is that in ordinary optical fibers, birefringence occurs accidentally, e.g. due to stress caused by the touch of a finger. Lasers built in this manner are extremely sensitive to external factors. In turn, birefringence of the polarization preserving fibers is so large that the light propagates in only one direction, and the construction of artificial saturable absorbers becomes physically impossible.

"Birefringent optical fibers retaining the polarization state of the entering light are already in production. We are the first to demonstrate how they can be used to construct a saturable absorber: We cut the optical fiber into segments of an appropriate length and then reconnect them, rotating each successive segment 90 degrees in relation to its predecessor," says Ph.D. student Szczepanek.

"Rotation means that if in one segment a pulse with, shall we say, vertical polarization travels slowly, in the next, it will run faster and catch up with the second pulse, polarized perpendicularly. A simple procedure has therefore allowed us to eliminate the main obstacle to increasing the energy, that is,

the great difference in velocities between pulses of different polarities, typical for all polarization preserving fibres," explains Dr. Stepanenko.

The more rotated segments there are, the better the quality of the pulses generated in the fiber. In the laser built in the Warsaw laboratory, the saturable absorber consisted of a fiber with a length of approximately 3 m, divided into three segments, and a filtering polarizer. The potential number of rotated segments can be increased up to even a dozen or so.

The new laser produces high-quality femtosecond pulses, and their energy can be up to 1000 times larger than typical for lasers with material absorbers. In comparison to the devices with artificial absorbers, the laser made by Warsaw scientists has a much simpler construction and therefore its reliability is significantly greater. [7]

## **Optical fibre with Einstein effect**

Researchers at the Max Planck Institute for the Science of Light in Erlangen have discovered a new mechanism for guiding light in photonic crystal fibre (PCF). PCF is a hair-thin glass fibre with a regular array of hollow channels running along its length. When helically twisted, this spiralling array of hollow channels acts on light rays in an analogous manner to the bending of light rays when they travel through the gravitationally curved space around a star, as described by the general theory of relativity.

Optical fibres act as pipes for light. And just as the inside of a pipe is enclosed by a wall, optical fibres normally have a light-guiding core, whose glass has a higher refractive index than the glass of the enclosing outer cladding. The difference in the refractive index causes the light to be reflected at the cladding interface and trapped in the core like water in a pipe. A team headed by Philip Russell, Director at the Max Planck Institute for the Science of Light, is the first to succeed in guiding light in a PCF with no core.

## **Photonic crystals give butterflies their colour and can also guide light**

A typical photonic crystal consists of a piece of glass with holes arranged in regular periodic pattern throughout its volume. Since glass and the air have different refractive indices, the refractive index has a periodic structure. This is the reason these materials are called crystals—their atoms form an ordered, three-dimensional lattice as found in crystalline salt or silicon, for example. In a conventional crystal, the precise design of the 3-D structure determines the behaviour of electrons, resulting for example in electrical insulators, conductors and semiconductors.

In a similar manner, the optical properties of a photonic crystal depend on the periodic 3-D microstructure, which is responsible for the shimmering colours of some butterfly wings, for example. Being able to control the optical properties of materials is useful in a wide variety of applications. The photonic crystal fibres developed by Philip Russell and his team at the Erlangen-based Max Planck Institute can be used to filter specific wavelengths out of the visible spectrum or to produce very white light, for example.

As is the case with all optical fibres used in telecommunications, all conventional photonic crystal fibres have a core and cladding each with different refractive indices or optical properties. In PCF,

the air-filled channels already give the glass a refractive index different from the one it would have if completely solid.

### The holes define the space in a photonic crystal fibre

"We are the first to succeed in guiding light through a coreless fibre," says Gordon Wong from the Max Planck Institute for the Science of Light in Erlangen. The researchers working in Philip Russell's team have fabricated a photonic crystal fibre whose complete cross-section is closely packed with a large number of air-filled channels, each around one thousandth of a millimetre in diameter, which extend along its whole length.

While the core of a conventional PCF is solid glass, the cross-sectional view of the new optical fibre resembles a sieve. The holes have regular separations and are arranged so that every hole is surrounded by a regular hexagon of neighbouring holes. "This structure defines the space in the fibre," explains Ramin Beravat, lead author of the publication. The holes can be thought of as distance markers. The interior of the fibre then has a kind of artificial spatial structure which is formed by the regular lattice of holes.

"We have now fabricated the fibre in a twisted form," continues Beravat. The twisting causes the hollow channels to wind around the length of the fibre in helical lines. The researchers then transmitted laser light through the fibre. In the case of the regular, coreless cross-section, one would actually expect the light to distribute itself between the holes of the sieve as evenly as their pattern determines, i.e. at the edge just as much as in the centre. Instead, the physicists discovered something surprising: the light was concentrated in the central region, where the core of a conventional optical fibre is located.

### In a twisted PCF, the light follows the shortest path in the interior of the fibre

"The effect is analogous to the curvature of space in Einstein's general theory of relativity," explains Wong. This predicts that a heavy mass such as the Sun will distort the space surrounding it – or more precisely, distort spacetime, i.e. the combination of the three spatial dimensions with the fourth dimension, time – like a sheet of rubber into which a lead sphere is placed. Light follows this curvature. The shortest path between two points is then no longer a straight line, but a curve. During a solar eclipse, stars which should really be hidden behind the Sun thus become visible. Physicists call these shortest connecting paths "geodesics".

"By twisting the fibre, the 'space' in our photonic crystal fibre becomes twisted as well," says Wong. This leads to helical geodesic lines along which light travels. This can be intuitively understood by taking into account the fact that light always takes the shortest route through a medium. The glass strands between the air-filled channels describe spirals, which define possible paths for the light rays. The path through the wide spirals at the edge of the fibre is longer than that through the more closely wound spirals in its centre, however, resulting in curved ray paths that at a certain radius are reflected by a photonic crystal effect back towards the fibre axis.

### A twisted PCF as a large-scale environmental sensor

The more the fibre is twisted, the narrower is the space within which the light concentrated. In analogy to Einstein's theory, this corresponds to a stronger gravitational force and thus a greater

deflection of the light. The Erlangen-based researchers write that they have created a "topological channel" for the light (topology is concerned with the properties of space which are conserved under continuous distortion).

The researchers emphasize that their work is basic research. They are one of the very few research groups working in this field anywhere in the world. Nevertheless, they can think of several applications for their discovery. A twisted fibre which is less twisted at certain intervals, for example, will allow a portion of the light to escape to the outside. Light could then interact with the environment at these defined locations. "This could be used for sensors which measure the absorption of a medium, for instance." A network of these fibres could collect data over large areas as an environmental sensor. [6]

## **Theory that challenges Einstein's physics could soon be put to the test**

Scientists behind a theory that the speed of light is variable - and not constant as Einstein suggested - have made a prediction that could be tested.

Einstein observed that the speed of light remains the same in any situation, and this meant that space and time could be different in different situations.

The assumption that the speed of light is constant, and always has been, underpins many theories in physics, such as Einstein's theory of general relativity. In particular, it plays a role in models of what happened in the very early universe, seconds after the Big Bang.

But some researchers have suggested that the speed of light could have been much higher in this early universe. Now, one of this theory's originators, Professor João Magueijo from Imperial College London, working with Dr Niayesh Afshordi at the Perimeter Institute in Canada, has made a prediction that could be used to test the theory's validity.

Structures in the universe, for example galaxies, all formed from fluctuations in the early universe – tiny differences in density from one region to another. A record of these early fluctuations is imprinted on the cosmic microwave background – a map of the oldest light in the universe – in the form of a 'spectral index'.

Working with their theory that the fluctuations were influenced by a varying speed of light in the early universe, Professor Magueijo and Dr Afshordi have now used a model to put an exact figure on the spectral index. The predicted figure and the model it is based on are published in the journal Physical Review D.

Cosmologists are currently getting ever more precise readings of this figure, so that prediction could soon be tested – either confirming or ruling out the team's model of the early universe. Their figure is a very precise 0.96478. This is close to the current estimate of readings of the cosmic microwave background, which puts it around 0.968, with some margin of error.

## **RADICAL IDEA**

Professor Magueijo said: "The theory, which we first proposed in the late-1990s, has now reached a maturity point – it has produced a testable prediction. If observations in the near future do find this number to be accurate, it could lead to a modification of Einstein's theory of gravity."

"The idea that the speed of light could be variable was radical when first proposed, but with a numerical prediction, it becomes something physicists can actually test. If true, it would mean that the laws of nature were not always the same as they are today."

The testability of the varying speed of light theory sets it apart from the more mainstream rival theory: inflation. Inflation says that the early universe went through an extremely rapid expansion phase, much faster than the current rate of expansion of the universe.

## THE HORIZON PROBLEM

These theories are necessary to overcome what physicists call the 'horizon problem'. The universe as we see it today appears to be everywhere broadly the same, for example it has a relatively homogenous density.

This could only be true if all regions of the universe were able to influence each other. However, if the speed of light has always been the same, then not enough time has passed for light to have travelled to the edge of the universe, and 'even out' the energy.

As an analogy, to heat up a room evenly, the warm air from radiators at either end has to travel across the room and mix fully. The problem for the universe is that the 'room' – the observed size of the universe – appears to be too large for this to have happened in the time since it was formed.

The varying speed of light theory suggests that the speed of light was much higher in the early universe, allowing the distant edges to be connected as the universe expanded. The speed of light would have then dropped in a predictable way as the density of the universe changed. This variability led the team to the prediction published today.

The alternative theory is inflation, which attempts to solve this problem by saying that the very early universe evened out while incredibly small, and then suddenly expanded, with the uniformity already imprinted on it. While this means the speed of light and the other laws of physics as we know them are preserved, it requires the invention of an 'inflation field' – a set of conditions that only existed at the time.

'Critical geometry of a thermal big bang' by Niayesh Afshordi and João Magueijo is published in Physical Review D. [4]

## The Test in 2015

Over the course of four Thursdays in November 1915, Albert Einstein stood before the Prussian Academy in Berlin and unveiled a set of equations that upended our ideas about space and time. A century later, his grand project remains frustratingly incomplete. Sure, the general theory of relativity has become the foundation of our modern understanding of the big bang, black holes, and gravity itself—but crucial parts of it remain unverified.

One unproved prediction is that an accelerating mass should create gravitational waves, like the ripples from a boat sailing across the surface of a lake. In July, the hunt for those waves will heat up with the launch of a detector called Lisa Pathfinder, which will test technology for a new gravitational wave observatory in space. Starting in 2015, two earthbound experiments, Advanced Ligo and Advanced Virgo, will be brought online. They should be able to pick up gravitational

disturbances from exploding stars. Another test will examine the motions of a triple-star system called PSR J0337+1715 to check if gravity behaves the same toward all kinds of matter, as Einstein believed.

Some scientists suspect other aspects of relativity are just flat-out wrong. For years, cosmologists have observed dramatic, unexplained movements of galaxies. Those motions are normally attributed to hypothetical dark matter and dark energy, but at a series of conferences this year, physicists will explore the possibility that gravity simply doesn't work the way Einstein thought. And then there's the elephant in the lab: General relativity clashes with quantum mechanics. Attempts to reconcile them have so far yielded nothing but endless, inconclusive papers about string theory. [4]

## The Classical Relativistic effect

The moving charges are self maintain the electromagnetic field locally, causing their movement and this is the result of their acceleration under the force of this field.

In the classical physics the charges will distributed along the electric current so that the electric potential lowering along the current, by linearly increasing the way they take every next time period because this accelerated motion.

## The Relativistic Quantum Mechanics

The same thing happens on the atomic scale giving a  $dp$  impulse difference and a  $dx$  way difference between the different part of the not point like particles.

Commonly accepted idea that the relativistic effect on the particle physics it is the fermions' spin - another unresolved problem in the classical concepts. If the electric charges can move only with accelerated motions in the self maintaining electromagnetic field, once upon a time they would reach the velocity of the electromagnetic field. The resolution of this problem is the spinning particle, constantly accelerating and not reaching the velocity of light because the acceleration is radial.

## The Heisenberg Uncertainty Relation

I think that we have a simple bridge between the classical and quantum mechanics by understanding the Heisenberg Uncertainty Relations. It makes clear that the particles are not point like but have a  $dx$  and  $dp$  uncertainty.

## The General Relativity - Electromagnetic inertia and mass

### Electromagnetic Induction

Since the magnetic induction creates a negative electric field as a result of the changing acceleration, it works as an electromagnetic inertia, causing an electromagnetic mass. [1]

### Relativistic change of mass

The increasing mass of the electric charges the result of the increasing inductive electric force acting against the accelerating force. The decreasing mass of the decreasing acceleration is the result of the

inductive electric force acting against the decreasing force. This is the relativistic mass change explanation, especially importantly explaining the mass reduction in case of velocity decrease.

### The frequency dependence of mass

Since  $E = \hbar v$  and  $E = mc^2$ ,  $m = \hbar v / c^2$  that is the  $m$  depends only on the  $v$  frequency. It means that the mass of the proton and electron are electromagnetic and the result of the electromagnetic induction, caused by the changing acceleration of the spinning and moving charge! It could be that the  $m_0$  inertial mass is the result of the spin, since this is the only accelerating motion of the electric charge. Since the accelerating motion has different frequency for the electron in the atom and the proton, they masses are different, also as the wavelengths on both sides of the diffraction pattern, giving equal intensity of radiation.

### Electron – Proton mass rate

The Planck distribution law explains the different frequencies of the proton and electron, giving equal intensity to different lambda wavelengths! Also since the particles are diffraction patterns they have some closeness to each other – can be seen as a gravitational force. [2]

There is an asymmetry between the mass of the electric charges, for example proton and electron, can be understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

### The weak interaction

The weak interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and time reversal symmetry. The Electroweak Interaction shows that the Weak Interaction is basically electromagnetic in nature. The arrow of time shows the entropy grows by changing the temperature dependent diffraction patterns of the electromagnetic oscillators.

Another important issue of the quark model is when one quark changes its flavor such that a linear oscillation transforms into plane oscillation or vice versa, changing the charge value with 1 or -1. This kind of change in the oscillation mode requires not only parity change, but also charge and time changes (CPT symmetry) resulting a right handed anti-neutrino or a left handed neutrino.

The right handed anti-neutrino and the left handed neutrino exist only because changing back the quark flavor could happen only in reverse, because they are different geometrical constructions, the u is 2 dimensional and positively charged and the d is 1 dimensional and negatively charged. It needs also a time reversal, because anti particle (anti neutrino) is involved.

The neutrino is a 1/2spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with  $\frac{1}{2}$  spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction

intensity of the Maxwell–Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The limit of the velocity of particles as the speed of light appropriate only for electrical charged particles, since the accelerated charges are self maintaining locally the accelerating electric force. The neutrinos are CP symmetry breaking particles compensated by time in the CPT symmetry, that is the time coordinate not works as in the electromagnetic interactions, consequently the speed of neutrinos is not limited by the speed of light.

The weak interaction T-asymmetry is in conjunction with the T-asymmetry of the second law of thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes the weak interaction, for example the Hydrogen fusion.

Probably because it is a spin creating movement changing linear oscillation to 2 dimensional oscillation by changing d to u quark and creating anti neutrino going back in time relative to the proton and electron created from the neutron, it seems that the anti neutrino fastest then the velocity of the photons created also in this weak interaction?

A quark flavor changing shows that it is a reflection changes movement and the CP- and T- symmetry breaking!!! This flavor changing oscillation could prove that it could be also on higher level such as atoms, molecules, probably big biological significant molecules and responsible on the aging of the life.

Important to mention that the weak interaction is always contains particles and antiparticles, where the neutrinos (antineutrinos) present the opposite side. It means by Feynman's interpretation that these particles present the backward time and probably because this they seem to move faster than the speed of light in the reference frame of the other side.

Finally since the weak interaction is an electric dipole change with  $\frac{1}{2}$  spin creating; it is limited by the velocity of the electromagnetic wave, so the neutrino's velocity cannot exceed the velocity of light.

## The General Weak Interaction

The Weak Interactions T-asymmetry is in conjunction with the T-asymmetry of the Second Law of Thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes for example the Hydrogen fusion. The arrow of time by the Second Law of Thermodynamics shows the increasing entropy and decreasing information by the Weak Interaction, changing the temperature dependent diffraction patterns. A good example of this is the neutron decay, creating more particles with less known information about them.

The neutrino oscillation of the Weak Interaction shows that it is a general electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. There should be a new research space of the Quantum Information Science the 'general neutrino oscillation' for the greater then subatomic matter structures as an electric dipole change. There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that

is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction.

## The Higgs boson

By March 2013, the particle had been proven to behave, interact and decay in many of the expected ways predicted by the Standard Model, and was also tentatively confirmed to have + parity and zero spin, two fundamental criteria of a Higgs boson, making it also the first known scalar particle to be discovered in nature, although a number of other properties were not fully proven and some partial results do not yet precisely match those expected; in some cases data is also still awaited or being analyzed.

In my opinion, the best explanation of the Higgs mechanism for a lay audience is the one invented by David Miller. You can find it here: <http://www.strings.ph.qmul.ac.uk/~jmc/epp/higgs3.html> .

The field must come first. The boson is an excitation of the field. So no field, no excitation. On the other hand in quantum field theory it is difficult to separate the field and the excitations.

The Higgs field is what gives particles their mass.

There is a video that gives an idea as to the Higgs field and the boson. It is here:

<http://www.youtube.com/watch?v=RIg1Vh7uPyw> . Note that this analogy isn't as good as the Miller one, but as is usually the case, if you look at all the analogies you'll get the best understanding of the situation.

Since the Higgs boson is necessary to the W and Z bosons, the dipole change of the Weak interaction and the change in the magnetic effect caused gravitation must be conducted. The Wien law is also important to explain the Weak interaction, since it describes the  $T_{\max}$  change and the diffraction patterns change. [2]

## Higgs mechanism

The magnetic induction creates a negative electric field, causing an electromagnetic inertia. Probably it is the mysterious Higgs field giving mass to the charged particles? We can think about the photon as an electron-positron pair, they have mass. The neutral particles are built from negative and positive charges, for example the neutron, decaying to proton and electron. The wave – particle duality makes sure that the particles are oscillating and creating magnetic induction as an inertial mass, explaining also the relativistic mass change. Higher frequency creates stronger magnetic induction, smaller frequency results lesser magnetic induction. It seems to me that the magnetic induction is the secret of the Higgs field.

In particle physics, the Higgs mechanism is a kind of mass generation mechanism, a process that gives mass to elementary particles. According to this theory, particles gain mass by interacting with the Higgs field that permeates all space. More precisely, the Higgs mechanism endows gauge bosons in a gauge theory with mass through absorption of Nambu–Goldstone bosons arising in spontaneous symmetry breaking.

The simplest implementation of the mechanism adds an extra Higgs field to the gauge theory. The spontaneous symmetry breaking of the underlying local symmetry triggers conversion of components of this Higgs field to Goldstone bosons which interact with (at least some of) the other

fields in the theory, so as to produce mass terms for (at least some of) the gauge bosons. This mechanism may also leave behind elementary scalar (spin-0) particles, known as Higgs bosons.

In the Standard Model, the phrase "Higgs mechanism" refers specifically to the generation of masses for the  $W^\pm$ , and Z weak gauge bosons through electroweak symmetry breaking. The Large Hadron Collider at CERN announced results consistent with the Higgs particle on July 4, 2012 but stressed that further testing is needed to confirm the Standard Model.

## Gravity from the point of view of quantum physics

### What is the Spin?

So we know already that the new particle has spin zero or spin two and we could tell which one if we could detect the polarizations of the photons produced. Unfortunately this is difficult and neither ATLAS nor CMS are able to measure polarizations. The only direct and sure way to confirm that the particle is indeed a scalar is to plot the angular distribution of the photons in the rest frame of the centre of mass. A spin zero particles like the Higgs carries no directional information away from the original collision so the distribution will be even in all directions. This test will be possible when a much larger number of events have been observed. In the mean time we can settle for less certain indirect indicators.

### The Graviton

In physics, the graviton is a hypothetical elementary particle that mediates the force of gravitation in the framework of quantum field theory. If it exists, the graviton is expected to be massless (because the gravitational force appears to have unlimited range) and must be a spin-2 boson. The spin follows from the fact that the source of gravitation is the stress-energy tensor, a second-rank tensor (compared to electromagnetism's spin-1 photon, the source of which is the four-current, a first-rank tensor). Additionally, it can be shown that any massless spin-2 field would give rise to a force indistinguishable from gravitation, because a massless spin-2 field must couple to (interact with) the stress-energy tensor in the same way that the gravitational field does. This result suggests that, if a massless spin-2 particle is discovered, it must be the graviton, so that the only experimental verification needed for the graviton may simply be the discovery of a massless spin-2 particle. [3]

## Conclusions

Physicists' greatest hope for 2015, then, is that one of these experiments will show where Einstein got off track, so someone else can jump in and get closer to his long-sought "theory of everything." [4]

The electric currents causing self maintaining electric potential is the source of the special and general relativistic effects. The self maintained electric potential of the accelerating charges equivalent with the General Relativity space-time curvature, and since it is true on the quantum level also, gives the base of the Quantum Gravity. Basing the gravitational force on the accelerating Universe caused magnetic force and the Planck Distribution Law of the electromagnetic waves caused diffraction gives us the basis to build a Unified Theory of the physical interactions.

Since graviton is a tensor field, it has spin = 2, could be 2 photons with spin = 1 together.

## References

[1] The Magnetic field of the Electric current and the Magnetic induction

[http://academia.edu/3833335/The\\_Magnetic\\_field\\_of\\_the\\_Electric\\_current](http://academia.edu/3833335/The_Magnetic_field_of_the_Electric_current)

[2] 3 Dimensional String Theory

[http://academia.edu/3834454/3\\_Dimensional\\_String\\_Theory](http://academia.edu/3834454/3_Dimensional_String_Theory)

[3] Graviton Production By Two Photon and Electron-Photon Processes In Kaluza-Klein Theories With Large Extra Dimensions

<http://arxiv.org/abs/hep-ph/9909392>

[4] WHAT TO EXPECT IN 2015: GENERAL RELATIVITY GETS PUT TO THE TEST

<http://www.popsci.com/general-relativity-gets-put-test>

[5] Theory that challenges Einstein's physics could soon be put to the test

<http://phys.org/news/2016-11-theory-einstein-physics.html>

[6] Optical fibre with Einstein effect

<https://phys.org/news/2017-02-optical-fibre-einstein-effect.html>

[7] Researchers develop surprising technique for ultrashort laser pulses

<https://phys.org/news/2017-02-technique-ultrashort-laser-pulses.html>

[8] Researchers demonstrate new type of laser

<https://phys.org/news/2017-03-laser.html>