

Closed-Form Solution for the Nontrivial Zeros of the Riemann Zeta Function

Frederick Ira Moxley III¹

¹*Hearne Institute for Theoretical Physics, Department of Physics & Astronomy,
Louisiana State University, Baton Rouge, Louisiana 70803-4001, USA*

(Dated: April 25, 2017)

In the year 2017 it was formally conjectured that if the Bender-Brody-Müller (BBM) Hamiltonian can be shown to be self-adjoint, then the Riemann hypothesis holds true. Herein we discuss the domain and eigenvalues of the Bender-Brody-Müller conjecture. Moreover, a second quantization of the BBM Schrödinger equation is performed, and a closed-form solution for the nontrivial zeros of the Riemann zeta function is obtained. Finally, it is shown that all of the nontrivial zeros are located at $\Re(z) = 1/2$.

I. INTRODUCTION

It was recently shown in [1] that the eigenvalues of a Bender-Brody-Müller (BBM) Hamiltonian operator correspond to the nontrivial zeroes of the Riemann zeta function [2]. Although the BBM Hamiltonian is pseudo-Hermitian, it is consistent with the Berry-Keating conjecture [3, 4]. The eigenvalues of the BBM Hamiltonian are taken to be the imaginary parts of the nontrivial zeroes of the zeta function

$$\begin{aligned}\zeta(z) &= \sum_{k=1}^{\infty} \frac{1}{k^z} \\ &= \frac{1}{\Gamma(z)} \int_0^{\infty} \frac{t^{z-1}}{\exp(t) - 1} dt.\end{aligned}\tag{1}$$

The idea that the imaginary parts of the zeroes of Eq. (1) are given by a self-adjoint operator was conjectured by Hilbert and Pólya [5]. Formally, Hilbert and Pólya determined that if the eigenfunctions of a self-adjoint operator satisfy the boundary conditions $\psi_n(0) = 0 \forall n$, then the eigenvalues are the nontrivial zeroes of Eq. (1). The BBM Hamiltonian also satisfies the Berry-Keating conjecture, which states that when \hat{x} and \hat{p} commute, the Hamiltonian reduces to the classical $H = 2xp$.

Remark. *If there are nontrivial roots of Eq. (1) for which $\Re(z) \neq 1/2$, the corresponding eigenvalues and eigenstates are degenerate [1].*

II. BENDER-BRODY-MÜLLER HAMILTONIAN

Theorem 1. *The eigenvalues of the Hamiltonian*

$$\hat{H} = \frac{1}{1 - e^{-i\hat{p}}} (\hat{x}\hat{p} + \hat{p}\hat{x})(1 - e^{-i\hat{p}})\tag{2}$$

are real, where $\hat{p} = -i\hbar\partial_x$, $\hbar = 1$, and $\hat{x} = x$.

Corollary 1.1. *[1] Solutions to the equation $\hat{H}\psi = E\psi$ are given by the Hurwitz zeta function*

$$\begin{aligned}\psi_z(x) &= -\zeta(z, x+1) \\ &= -\sum_{n=0}^{\infty} \frac{1}{(x+1+n)^z}\end{aligned}\tag{3}$$

on the positive half line $x \in \mathbb{R}^+$ with eigenvalues $i(2z-1)$, and $z \in \mathbb{C}$, for the boundary condition $\psi_z(0) = 0$. Moreover, $\Re(z) > 1$, and $\Re(x+1) > 0$. As $-\psi_z(0)$ is the Riemann zeta function, i.e., Eq. (1), this implies that z belongs to the discrete set of zeros of the Riemann zeta function.

Proof. Let $\psi_z(x)$ be an eigenfunction of Eq. (2) with an eigenvalue $\lambda = i(2z-1)$:

$$\hat{H}\psi_z(x) = \lambda\psi_z(x).\tag{4}$$

Then we have the relation

$$\frac{1}{1 - e^{-i\hat{p}}}(\hat{x}\hat{p} + \hat{p}\hat{x})(1 - e^{-i\hat{p}})\psi_z(x) = \lambda\psi_z(x). \quad (5)$$

Letting

$$\begin{aligned} \varphi_z(x) &= [1 - \exp(-\partial_x)]\psi_z(x), \\ &= \hat{\Delta}\psi_z(x), \end{aligned} \quad (6)$$

where $\hat{\Delta}\psi_z(x) = \psi_z(x) - \psi_z(x - 1)$, and

$$\hat{\Delta} \equiv 1 - \exp(-\partial_x), \quad (7)$$

is a shift operator. Upon inserting Eq. (6) into Eq. (5) with $\hat{p} = -i\hbar\partial_x$, $\hbar = 1$, and $\hat{x} = x$, we obtain

$$[-ix\partial_x - i\partial_x x]\varphi_z(x) = \lambda\varphi_z(x). \quad (8)$$

Then we have

$$\int_{\mathbb{R}^+} (x\partial_x\varphi_z(x))^*\varphi_z(x)dx + \int_{\mathbb{R}^+} (\partial_x x\varphi_z(x))^*\varphi_z(x)dx = -i\lambda^* \int_{\mathbb{R}^+} \varphi_z^*(x)\varphi_z(x)dx. \quad (9)$$

As $\varphi_z(x \rightarrow \infty) \rightarrow 0$, next we integrate the first term on the LHS of Eq. (9) by parts to obtain

$$\int_{\mathbb{R}^+} x\varphi_z(x)\partial_x\varphi_z^*(x)dx = - \int_{\mathbb{R}^+} \varphi_z^*(x)\varphi_z(x)dx - \int_{\mathbb{R}^+} \varphi_z^*(x)x\frac{d}{dx}(\varphi_z(x))dx, \quad (10)$$

and the second term on the LHS of Eq. (9) by parts to obtain

$$\int_{\mathbb{R}^+} x\varphi_z(x)^*\partial_x\varphi_z(x)dx = - \int_{\mathbb{R}^+} \varphi_z(x)\varphi_z^*(x)dx - \int_{\mathbb{R}^+} \varphi_z(x)x\frac{d}{dx}(\varphi_z^*(x))dx. \quad (11)$$

Upon substituting Eqs. (10) and (11) into Eq. (9), we obtain

$$\int_{\mathbb{R}^+} \varphi_z^*(x)x\frac{d}{dx}(\varphi_z(x))dx + \int_{\mathbb{R}^+} \varphi_z(x)x\frac{d}{dx}(\varphi_z^*(x))dx = (i\lambda^* - 2)N, \quad (12)$$

where

$$N = \int_{\mathbb{R}^+} \varphi_z^*(x)\varphi_z(x)dx. \quad (13)$$

Next, we split $\varphi_z(x)$ into real and imaginary components, such that

$$\varphi_z(x) = \varphi_{\Re(z)}(x) + i\varphi_{\Im(z)}(x), \quad (14)$$

and substitute Eq. (14) into Eq. (12) such that

$$\int_{\mathbb{R}^+} \varphi_{\Re(z)}(x)x\frac{d}{dx}\varphi_{\Re(z)}(x)dx + \int_{\mathbb{R}^+} \varphi_{\Im(z)}(x)x\frac{d}{dx}\varphi_{\Im(z)}(x)dx + N = \frac{i\lambda^*}{2}N. \quad (15)$$

Upon setting $\lambda = i(2z - 1)$, Eq. (15) can be written

$$\int_{\mathbb{R}^+} \varphi_{\Re(z)}(x)x\frac{d}{dx}\varphi_{\Re(z)}(x)dx + \int_{\mathbb{R}^+} \varphi_{\Im(z)}(x)x\frac{d}{dx}\varphi_{\Im(z)}(x)dx + N = \frac{2z - 1}{2}N. \quad (16)$$

It can be seen that all terms on the LHS of Eq. (15) are real, thereby verifying Theorem 1. Since

$$\frac{1}{n^z} = \frac{\exp(-i \cdot \Im(z) \ln(n))}{n^{\Re(z)}} = \frac{\cos(\Im(z) \cdot \ln(n))}{n^{\Re(z)}} - i \frac{\sin(\Im(z) \cdot \ln(n))}{n^{\Re(z)}}, \quad (17)$$

we have

$$\varphi_{\Re(z)}(x) = \frac{\cos\left(\Im(z) \cdot \ln(x+n)\right)}{(x+n)^{\Re(z)}} - \frac{\cos\left(\Im(z) \cdot \ln(x+1+n)\right)}{(x+1+n)^{\Re(z)}}, \quad (18)$$

$$\varphi_{\Im(z)}(x) = \frac{\sin\left(\Im(z) \cdot \ln(x+1+n)\right)}{(x+1+n)^{\Re(z)}} - \frac{\sin\left(\Im(z) \cdot \ln(x+n)\right)}{(x+n)^{\Re(z)}}. \quad (19)$$

Q.E.D. □

Remark. *If the Riemann hypothesis is correct [2], the the eigenvalues of Eq. (2) are degenerate [1].*

Given that

$$\begin{aligned} \varphi_z(x) &= \hat{\Delta}\psi_z(x) \\ &= \psi_z(x) - \psi_z(x-1) \\ &= -\sum_{n=0}^{\infty} \frac{1}{(x+1+n)^z} + \sum_{n=0}^{\infty} \frac{1}{(x+n)^z}, \end{aligned} \quad (20)$$

we are left with

$$z = \frac{1}{N} \int_{\mathbb{R}^+} \varphi_{\Re(z)}(x) x \frac{d}{dx} \varphi_{\Re(z)}(x) dx + \frac{1}{N} \int_{\mathbb{R}^+} \varphi_{\Im(z)}(x) x \frac{d}{dx} \varphi_{\Im(z)}(x) dx + \frac{3}{2}. \quad (21)$$

For ease of derivation, we take $\varphi_{\Re(z)}(x) = \varphi_{\Im(z)}(x)$. Moreover, it can be seen that

$$\begin{aligned} x \frac{d}{dx} (\varphi_z(x)) &= x \frac{d}{dx} \psi_z(x) - x \frac{d}{dx} \psi_z(x-1) \\ &= -x \frac{d}{dx} \sum_{n=0}^{\infty} \frac{1}{(x+1+n)^z} + x \frac{d}{dx} \sum_{n=0}^{\infty} \frac{1}{(x+n)^z} \\ &= xz\zeta(z+1, x+1) - xz\zeta(z+1, x). \end{aligned} \quad (22)$$

Multiplying Eq. (22) by $\varphi_n(x)$, we obtain

$$\begin{aligned} \varphi_z(x) xz\zeta(z+1, x+1) - \varphi_z(x) xz\zeta(z+1, x) &= \varphi_z(x) [xz\zeta(z+1, x+1) - xz\zeta(z+1, x)] \\ &= -\zeta(z, x+1) xz\zeta(z+1, x+1) \\ &\quad + \zeta(z, x+1) xz\zeta(z+1, x) \\ &\quad + \zeta(z, x) xz\zeta(z+1, x+1) \\ &\quad - \zeta(z, x) xz\zeta(z+1, x). \end{aligned} \quad (23)$$

From the RHS of Eq. (23), it can be seen that

$$-\int_{\mathbb{R}^+} \zeta(z, x+1) xz\zeta(z+1, x+1) dx = \frac{z(1+n)^{1-2z}}{2z-4z^2}, \quad (24)$$

$$-\int_{\mathbb{R}^+} \zeta(z, x) xz\zeta(z+1, x) dx = \frac{zn^{1-2z}}{2z-4z^2}, \quad (25)$$

and

$$\begin{aligned} &\int_{\mathbb{R}^+} \zeta(z, x+1) xz\zeta(z+1, x) + \zeta(z, x) xz\zeta(z+1, x+1) dx \\ &= \frac{1}{4} n^{-2z} \left[-\frac{4^z \left(-\frac{1}{n}\right)^{-2z} \sqrt{\pi} \csc(\pi z) \Gamma(-1/2+z)}{\Gamma(z)} + \frac{4}{2z-1} \left(\frac{n}{1+n}\right)^{z-1} \right. \\ &\quad \left. \times \left(n+z\Gamma(1-z) \cdot \frac{{}_2F_1(1, 1+z, 2-z, 1+1/n)}{\Gamma(2-z)}\right) \right], \end{aligned} \quad (26)$$

where $\Gamma(z)$ is the gamma function, and the hypergeometric series is

$${}_2F_1(1, 1+z, 2-z, 1+1/n) = \sum_{j=0}^{\infty} \frac{(1)_j(1+z)_j}{(2-z)_j} \frac{(1+1/n)^j}{j!}. \quad (27)$$

Now we find the ‘‘density’’

$$\begin{aligned} N &= \int_{\mathbb{R}^+} \varphi_z^*(x) \varphi_z(x) dx \\ &= \int_{\mathbb{R}^+} [\psi_z(x) - \psi_z(x-1)]^2 dx \\ &= \int_{\mathbb{R}^+} [\psi_z^2(x) - 2\psi_z(x-1)\psi_z(x) + \psi_z^2(x-1)] dx \\ &= \int_{\mathbb{R}^+} [(n+x+1)^{-2z} - 2(n+x)^{-z}(n+x+1)^{-z} + (n+x)^{-2z}] dx \\ &= \frac{n^{-2z}}{2} \left[\frac{4^z \left(-\frac{1}{n}\right)^{-2z} \sqrt{\pi} \csc(\pi z) \Gamma(-1/2+z)}{\Gamma(z)} + 4n^z (1+n)^{1-z} \Gamma(1-z) \cdot \frac{{}_2F_1(1, z, 2-z, 1+1/n)}{\Gamma(2-z)} \right] \\ &\quad + \frac{n^{1-2z}}{2z-1} + \frac{(1+n)^{1-2z}}{2z-1} \\ &= -\frac{n^{1-2z} + (1+n)^{1-2z}}{1-2z} + \frac{n^{-2z} \sqrt{\pi} \csc(\pi z)}{2\Gamma(z)} \cdot \left[4^z \left(-\frac{1}{n}\right)^{-2z} \Gamma(-1/2+z) \right. \\ &\quad \left. + 4n^z (1+n)^{1-z} \sqrt{\pi} \cdot \frac{{}_2F_1(1, z, 2-z, 1+1/n)}{\Gamma(2-z)} \right], \end{aligned} \quad (28)$$

with the hypergeometric series

$${}_2F_1(1, z, 2-z, 1+1/n) = \sum_{j=0}^{\infty} \frac{(1)_j(z)_j}{(2-z)_j} \frac{(1+1/n)^j}{j!}. \quad (29)$$

For simplicity, taking $\Re(z) = \Im(z)$, Eq. (21) can be rewritten exactly

$$\begin{aligned} z_n &= \left[-\frac{n^{1-2z} + (1+n)^{1-2z}}{1-2z} + \frac{n^{-2z} \sqrt{\pi} \csc(\pi z)}{2\Gamma(z)} \cdot \left[4^z \left(-\frac{1}{n}\right)^{-2z} \Gamma(-1/2+z) \right. \right. \\ &\quad \left. \left. + 4n^z (1+n)^{1-z} \sqrt{\pi} \cdot \frac{{}_2F_1(1, z, 2-z, 1+1/n)}{\Gamma(2-z)} \right] \right]^{-1} \cdot 2 \left[\frac{n^{1-2z} + (1+n)^{1-2z}}{2(1-2z)} \right. \\ &\quad \left. + \frac{1}{4} n^{-2z} \left[-\frac{4^z \left(-\frac{1}{n}\right)^{-2z} \sqrt{\pi} \csc(\pi z) \Gamma(-1/2+z)}{\Gamma(z)} + \frac{4}{2z-1} \left(\frac{n}{1+n}\right)^{z-1} \right. \right. \\ &\quad \left. \left. \times \left(n+z \Gamma(1-z) \cdot \frac{{}_2F_1(1, 1+z, 2-z, 1+1/n)}{\Gamma(2-z)} \right) \right] \right] + \frac{3}{2} \\ &= \frac{1}{2} (1 - i\lambda_n), \end{aligned} \quad (30)$$

for the gamma function $\Gamma(z)$.

Lemma 1.1. *From Eq. (16) and Eq. (30), it can be seen that all of the nontrivial zeros of Eq. (1) exist at $\Re(z) = 1/2$.*

Proof. Upon setting $\Re(z) = 1/2$ on the RHS of Eq. (16), we obtain

$$\frac{1}{N} \int_{\mathbb{R}^+} \varphi_{\Re(z)}(x) x \frac{d}{dx} \varphi_{\Re(z)}(x) dx = -1 - \frac{1}{N} \int_{\mathbb{R}^+} \varphi_{\Im(z)}(x) x \frac{d}{dx} \varphi_{\Im(z)}(x) dx, \quad (31)$$

and

$$\begin{aligned}
& \frac{1}{N} \int_{\mathbb{R}^+} \left[-\frac{x}{2(1+n+x)^2} + \frac{x}{2(n+x)^{3/2}\sqrt{1+n+x}} + \frac{x}{2\sqrt{n+x}(1+n+x)^{3/2}} - \frac{x}{2(n+x)^2} \right] dx \\
&= -1 - \frac{1}{N} \int_{\mathbb{R}^+} \varphi_{\Im(z)}(x) x \frac{d}{dx} \varphi_{\Im(z)}(x) dx \\
&= \lim_{t \rightarrow \infty} \frac{1}{N} \int_0^t \left[-\frac{x}{2(1+n+x)^2} + \frac{x}{2(n+x)^{3/2}\sqrt{1+n+x}} + \frac{x}{2\sqrt{n+x}(1+n+x)^{3/2}} - \frac{x}{2(n+x)^2} \right] dx \\
&= 0.
\end{aligned} \tag{32}$$

Hence,

$$1 = -\frac{1}{N} \int_{\mathbb{R}^+} \varphi_{\Im(z)}(x) x \frac{d}{dx} \varphi_{\Im(z)}(x) dx. \tag{33}$$

Since

$$\lim_{t \rightarrow \infty} \frac{1}{N} \int_0^t \varphi_{\Im(z)}(x) x \frac{d}{dx} \varphi_{\Im(z)}(x) dx = -1, \tag{34}$$

we have

$$1 = 1. \tag{35}$$

Q.E.D. □

n	$\Im(z)$ [10]	$\Im(z)$ Eq. (30)	absolute error
1	14.134725	14.134725	$0. \times 10^{-43}$
2	21.022039	21.022039	$0. \times 10^{-33}$
3	25.010857	25.010857	$0. \times 10^{-27}$
4	30.424876	30.424876	$0. \times 10^{-19}$
5	32.935061	32.935061	$0. \times 10^{-15}$
6	37.586178	37.586178	$0. \times 10^{-9}$
7	40.918719	40.918719	$0. \times 10^{-5}$
\vdots	\vdots	\vdots	\vdots
100	236.524229	236.524229	insufficient memory

Table I: Imaginary Nontrivial Zeros of the Riemann Zeta Function

Upon imposing the boundary condition

$$\begin{aligned}
\psi_n(0) &= -\sum_{n=1}^{\infty} \frac{1}{n^{z_n}} \\
&= -\frac{1}{\Gamma(z_n)} \int_0^{\infty} \frac{t^{z_n-1}}{\exp(t_n)-1} dt \\
&= 0,
\end{aligned} \tag{36}$$

it can be seen that Eq. (30) are the nontrivial zeros of Eq. (1), and for $z \in \mathbb{C}$ where z must belong to the discrete set of zeros of Eq. (1). Consequently, for the boundary condition $\psi(0) = 0$, the n^{th} eigenstate of Eq. (2) is

$$\begin{aligned}
\psi_n(x) &= -\zeta(z_n, x+1) \\
&= -\sum_{n=0}^{\infty} \frac{1}{(x+1+n)^{z_n}},
\end{aligned} \tag{37}$$

where z_n is given by Eq. (30). The Riemann hypothesis states [2] that the *nontrivial* zeros are located at $\Re(z) = 1/2$.

A. Domain of the Bender-Brody-Müller Hamiltonian

For the BBM Hamiltonian operator as given by Eq. (2), the Hilbert space is $\mathcal{H} = L^2(\mathbb{R}^+, dx)$. Moreover, \hat{p} and \hat{x} are self-adjoint operators that act in \mathcal{H} . In order to study the domain of the BBM Hamiltonian operator, we first introduce an auxiliary operator \hat{O} , such that

$$\hat{O} = \hat{p}\hat{p} + \hat{x}\hat{x}, \quad (38)$$

where $\hat{p}\hat{p} = -\nabla^2$, and $\hat{x}\hat{x} = x^2$. The set of finite linear combinations of Hermite functions is a core of \hat{O} , and therefore the Schwartz space \mathcal{S} is also a core of \hat{O} .

Lemma 1.2. [6] *If φ is in $\mathcal{D}(\hat{O})$, then*

$$\|\hat{p}\hat{p}\varphi\|^2 + \|\hat{x}\hat{x}\varphi\|^2 \leq \|\hat{O}\varphi\|^2 + c\|\varphi\|^2. \quad (39)$$

Proof. [6] We estimate φ for a core of \hat{O} via a double commutator to make the estimate [7],

$$\begin{aligned} \hat{O}\hat{O} &= \hat{p}\hat{p}\hat{p}\hat{p} + \hat{x}\hat{x}\hat{x}\hat{x} + \hat{p}\hat{p}\hat{x}\hat{x} + \hat{x}\hat{x}\hat{p}\hat{p} \\ &= \hat{p}\hat{p}\hat{p}\hat{p} + \hat{x}\hat{x}\hat{x}\hat{x} + 2\sum_{i=1}^n \left[\hat{x}_i\hat{p}\hat{p}\hat{x}_i + [\hat{x}_i, [\hat{x}_i, \hat{p}\hat{p}]] \right] \\ &\geq \hat{p}\hat{p}\hat{p}\hat{p} + \hat{x}\hat{x}\hat{x}\hat{x} - 2n, \end{aligned} \quad (40)$$

Therefore, in Eq. (39) $c = 2n$. □

After rewriting Eq. (8) as

$$[x\partial_x + \partial_x x]\varphi = (1 - 2z)\varphi, \quad (41)$$

then $\hat{p}\hat{p} = x\partial_x$ and $f(\hat{x}) = \partial_x x$ are self-adjoint operators acting in $\mathcal{H} = L^2(\mathbb{R}^+, dx)$. Setting

$$\hat{H} = \hat{p}\hat{p} + f(\hat{x}), \quad (42)$$

defined on

$$\mathcal{D}(\hat{p}\hat{p}) \cap \mathcal{D}(f(\hat{x})). \quad (43)$$

If $f(\hat{x})$ is local in \mathcal{H} , then Eq. (42) is dense and Hermitian.

Theorem 2. *The BBM Hamiltonian operator in Eq. (2) is essentially self-adjoint, given that $|\nabla f(\hat{x})| \leq a|\hat{x}| + b$.*

The BBM Hamiltonian operator in Eq. (2) is real-valued on the positive half line \mathbb{R}^+ , after being reduced to Eq. (41). From $|\nabla f(\hat{x})| \leq a|\hat{x}| + b$ we have

$$\begin{aligned} |f(\hat{x})| &\leq \frac{1}{2}\hat{x}\hat{x} + b|\hat{x}| \\ &\leq c\hat{x}\hat{x} + d. \end{aligned} \quad (44)$$

Let us examine the uniqueness.

Proof. As shown in [6], if \hat{H} is Hermitian, and \hat{O} is a positive self-adjoint operator, then \mathcal{C} is a core of \hat{O} such that $\mathcal{C} \subset \mathcal{D}(\hat{H})$. As such,

$$\|(\hat{p}\hat{p} + f(\hat{x}))\varphi\|^2 \leq a\|(\hat{p}\hat{p} + \hat{x}\hat{x})\varphi\|^2 + b\|\varphi\|^2, \quad (45)$$

where $\varphi \in \mathcal{S}$. Since $(1 + \hat{x}\hat{x})\varphi \in L^2$, $f(\hat{x})\varphi \in L^2$. Therefore, $\mathcal{S} \subset \mathcal{D}(\hat{H})$. Moreover, since $f(\hat{x})^2 \leq r\hat{x}\hat{x}\hat{x}\hat{x} + s$,

$$\|f(\hat{x})\varphi\|^2 \leq r\|\hat{x}\hat{x}\varphi\|^2 + s\|\varphi\|^2. \quad (46)$$

As such, from Eq. (39), Eq. (45) is satisfied. If $\varphi \in \mathcal{S}$, then $\nabla(f(\hat{x})\varphi) \in L^2$. Since,

$$\pm i[\hat{H}, \hat{O}] \leq c\hat{O} \quad (47)$$

as quadratic forms on \mathcal{C} , we thus have

$$\begin{aligned}
\pm i[\hat{H}, \hat{O}] &= \pm i\{[\hat{p}\hat{p}, \hat{x}\hat{x}] + [f(\hat{x}), \hat{p}\hat{p}]\} \\
&= \pm\{2(\hat{p} \cdot \hat{x} + \hat{x} \cdot \hat{p}) - (\hat{p} \cdot \nabla f(\hat{x}) + \nabla f(\hat{x}) \cdot \hat{p})\} \\
&\leq 2(\hat{p}\hat{p} + \hat{x}\hat{x}) + \hat{p}\hat{p} + (\nabla f(\hat{x}))^2 \\
&\leq 2(\hat{p}\hat{p} + \hat{x}\hat{x}) + \hat{p}\hat{p} + 2(a^2\hat{x}\hat{x} + b^2) \\
&\leq c\hat{O},
\end{aligned} \tag{48}$$

for constant c .

□

B. Second Quantization

We begin with the Bender-Brody-Müller (BBM) Schrödinger equation

$$-\frac{\hbar}{i} \frac{d}{dz} \psi(x, z) = \left[\hat{\Delta}^{-1} \hat{x} \hat{p} \hat{\Delta} + \hat{\Delta}^{-1} \hat{p} \hat{x} \hat{\Delta} \right] \psi(x, z), \tag{49}$$

where $\hat{\Delta}$ is given by Eq. (7), $\hat{x} = x$, $\hat{p} = -i\hbar\partial_x$, $\hbar = 1$, $x \in \mathbb{R}^+$, and $z \in \mathbb{C}$. Furthermore, let

$$\begin{aligned}
\psi_n(x) &= -\zeta(z_n, x+1) \\
&= -\sum_{n=0}^{\infty} \frac{1}{(x+1+n)^z}
\end{aligned} \tag{50}$$

be the solution of

$$\left(\hat{\Delta}^{-1} \hat{x} \hat{p} \hat{\Delta} + \hat{\Delta}^{-1} \hat{p} \hat{x} \hat{\Delta} \right) \psi_n(x) = \lambda_n \psi_n(x), \tag{51}$$

where z_n are the nontrivial zeros of the Riemann zeta function given by Eq. (30), λ_n are the eigenvalues, $\Re(z) > 1$, and $\Re(x+1) > 0$. Letting

$$\begin{aligned}
\varphi(x, z) &= [1 - \exp(-\partial_x)] \psi(x, z), \\
&= \hat{\Delta} \psi(x, z),
\end{aligned} \tag{52}$$

where $\hat{\Delta} \psi(x, z) = \psi(x, z) - \psi(x-1, z)$, and

$$\hat{\Delta} \equiv 1 - \exp(-\partial_x), \tag{53}$$

is a shift operator. Upon inserting Eq. (52) into Eq. (49) with $\hat{p} = -i\hbar\partial_x$, $\hbar = 1$, and $\hat{x} = x$, we obtain

$$-\hbar \frac{d}{dz} \varphi(x, z) = [x\partial_x + \partial_x x] \varphi(x, z). \tag{54}$$

Next, we write

$$\varphi(x, z) = \sum_n b_n(z) \varphi_n(x). \tag{55}$$

From Eq. (54) we find

$$-\hbar \frac{d}{dz} b_n(z) = \lambda_n b_n(z). \tag{56}$$

We now find a Hamiltonian that yields Eq. (56) as the equation of motion. Hence, we take

$$\hat{H} = \int_{\mathbb{R}^+} \varphi^*(x, z) [x\partial_x + \partial_x x] \varphi(x, z) dx \tag{57}$$

as the expectation value. Upon substituting Eq. (55) into Eq. (57) and using Eq. (51) we obtain the harmonic oscillator

$$\hat{H} = \sum_n \lambda_n b_n^*(z) b_n(z). \quad (58)$$

Taking $b_n(z)$ as an operator, and $b_n^*(z)$ as the adjoint, we obtain the usual properties:

$$\begin{aligned} [\hat{b}_n, \hat{b}_m] &= [\hat{b}_n^\dagger, \hat{b}_m^\dagger] = 0, \\ [\hat{b}_n, \hat{b}_m^\dagger] &= \delta_{nm}. \end{aligned} \quad (59)$$

From the analogous Heisenberg equations of motion,

$$\begin{aligned} -\hbar \frac{d}{dz} \hat{b}_n &= [\hat{b}_n, \hat{H}]_- \\ &= \sum_m E_m (\hat{b}_n \hat{b}_m^\dagger \hat{b}_m - \hat{b}_m^\dagger \hat{b}_m \hat{b}_n) \\ &= \sum_m E_m (\delta_{nm} \hat{b}_m - \hat{b}_m^\dagger \hat{b}_n \hat{b}_m - \hat{b}_m^\dagger \hat{b}_m \hat{b}_n) \\ &= \sum_m E_m (\delta_{nm} \hat{b}_m + \hat{b}_m^\dagger \hat{b}_m \hat{b}_n - \hat{b}_m^\dagger \hat{b}_m \hat{b}_n) \\ &= \lambda_n \hat{b}_n. \end{aligned} \quad (60)$$

The eigenvalues of \hat{H} are

$$\hat{H} = \sum_n \lambda_n N_n, \quad (61)$$

where $N_n = 0, 1, 2, 3, \dots, \infty$. Since, $\lambda_n = i(2z_n - 1)$, we can rewrite Eq. (61) as

$$\hat{H} = i \sum_n (2z_n - 1) N_n. \quad (62)$$

However, from Eq. (60) it can be seen that

$$-\hbar \frac{d}{dz} \hat{b}_n = i(2z_n - 1) \hat{b}_n. \quad (63)$$

As such,

$$\boxed{\frac{d}{dz} \hat{b}_n = -\frac{i}{\hbar} (2z_n - 1) \hat{b}_n.} \quad (64)$$

Remark. Eq. (64) can be solved using the Wirtinger derivatives.

C. \mathcal{PT} -symmetric Bender-Brody-Müller Hamiltonian

Theorem 3. *The eigenvalues of the Hamiltonian*

$$i\hat{H} = \frac{i}{1 - e^{-i\hat{p}}} (\hat{x}\hat{p} + \hat{p}\hat{x})(1 - e^{-i\hat{p}}) \quad (65)$$

are imaginary, where $\hat{p} = -i\hbar\partial_x$, $\hbar = 1$, and $\hat{x} = x$.

Corollary 3.1. [1] *Solutions to the equation $i\hat{H}\psi = E\psi$ are given by the Hurwitz zeta function*

$$\begin{aligned} \psi_z(x) &= -\zeta(z, x+1) \\ &= -\sum_{n=0}^{\infty} \frac{1}{(x+1+n)^z} \end{aligned} \quad (66)$$

on the positive half line $x \in \mathbb{R}^+$ with eigenvalues $i(2z-1)$, and $z \in \mathbb{C}$, for the boundary condition $\psi_z(0) = 0$. Moreover, $\Re(z) > 1$, and $\Re(x+1) > 0$. As $-\psi_z(0)$ is the Riemann zeta function, i.e., Eq. (1), this implies that z belongs to the discrete set of zeros of the Riemann zeta function.

Proof. Let ψ be an eigenfunction of Eq. (65) with an eigenvalue $\lambda = i(2z - 1)$:

$$i\hat{H}\psi = \lambda\psi. \quad (67)$$

Then we have the relation

$$\frac{i}{1 - e^{-i\hat{p}}}(\hat{x}\hat{p} + \hat{p}\hat{x})(1 - e^{-i\hat{p}})\psi = \lambda\psi. \quad (68)$$

Letting

$$\begin{aligned} \varphi_z(x) &= [1 - \exp(-\partial_x)]\psi_z(x), \\ &= \hat{\Delta}\psi_z(x), \end{aligned} \quad (69)$$

where $\hat{\Delta}\psi_z(x) = \psi_z(x) - \psi_z(x-1)$, and inserting Eq. (69) into Eq. (68) with $\hat{p} = -i\hbar\partial_x$, $\hbar = 1$, and $\hat{x} = x$, we obtain

$$[x\partial_x + \partial_x x]\varphi_z(x) = \lambda\varphi_z(x). \quad (70)$$

Then we have

$$\int_{\mathbb{R}^+} (x\partial_x\varphi_z(x))^*\varphi_z(x)dx + \int_{\mathbb{R}^+} (\partial_x x\varphi_z(x))^*\varphi_z(x)dx = \lambda^* \int_{\mathbb{R}^+} \varphi_z^*(x)\varphi_z(x)dx. \quad (71)$$

As $\varphi_z(x \rightarrow \infty) \rightarrow 0$, next we integrate the first term on the LHS of Eq. (71) by parts to obtain

$$\int_{\mathbb{R}^+} x\varphi_z(x)\partial_x\varphi_z^*(x)dx = - \int_{\mathbb{R}^+} \varphi_z^*(x)\varphi_z(x)dx - \int_{\mathbb{R}^+} \varphi_z^*(x)x\frac{d}{dx}(\varphi_z(x))dx, \quad (72)$$

and the second term on the LHS of Eq. (71) by parts to obtain

$$\int_{\mathbb{R}^+} x\varphi_z^*(x)\partial_x\varphi_z(x)dx = - \int_{\mathbb{R}^+} \varphi_z(x)\varphi_z^*(x)dx - \int_{\mathbb{R}^+} \varphi_z(x)x\frac{d}{dx}(\varphi_z^*(x))dx. \quad (73)$$

Upon substituting Eqs. (72) and (73) into Eq. (71), we obtain

$$\int_{\mathbb{R}^+} \varphi_z^*(x)x\frac{d}{dx}(\varphi_z(x))dx + \int_{\mathbb{R}^+} \varphi_z(x)x\frac{d}{dx}(\varphi_z^*(x))dx = -(\lambda^* + 2)N, \quad (74)$$

where

$$N = \int_{\mathbb{R}^+} \varphi_z^*(x)\varphi_z(x)dx. \quad (75)$$

Next, we split $\varphi_z(x)$ into real and imaginary components, such that

$$\varphi_z(x) = \varphi_{\Re(z)}(x) + i\varphi_{\Im(z)}(x), \quad (76)$$

and substitute Eq. (76) into Eq. (74) such that

$$\int_{\mathbb{R}^+} \varphi_{\Re(z)}(x)x\frac{d}{dx}\varphi_{\Re(z)}(x)dx + \int_{\mathbb{R}^+} \varphi_{\Im(z)}(x)x\frac{d}{dx}\varphi_{\Im(z)}(x)dx + N = -\frac{\lambda^*}{2}N. \quad (77)$$

Upon setting $\lambda = i(2z - 1)$, Eq. (77) can be written

$$\int_{\mathbb{R}^+} \varphi_{\Re(z)}(x)x\frac{d}{dx}\varphi_{\Re(z)}(x)dx + \int_{\mathbb{R}^+} \varphi_{\Im(z)}(x)x\frac{d}{dx}\varphi_{\Im(z)}(x)dx + N = \frac{i(2z - 1)}{2}N. \quad (78)$$

It can be seen that all terms on the LHS of Eq. (77) are real, thereby verifying Theorem 3.

Q.E.D.

□

III. NUMERICAL VERIFICATION

Here, it is useful to point out some identities. First, from the Taylor series expansion around 0, the inverse of the gamma function can be written

$$\frac{1}{\Gamma(z)} = z + \gamma z^2 + \left(\frac{\gamma^2}{2} - \frac{\pi^2}{12}\right)z^3 + \dots, \quad (79)$$

where γ is the Euler-Mascheroni constant [8], and furthermore,

$$\Gamma(z - 1/2) = \frac{2(z - 1/2)!}{2z - 1}, \quad (80)$$

$$\Gamma(1 - z) = (-z)! \quad (81)$$

$$\begin{aligned} {}_2F_1(1, z, 2 - z, 1 + 1/n) &= 1 + \frac{z \cdot (1 + 1/n)}{2 - z} + \frac{z(1 + z) \cdot (1 + 1/n)^2}{(2 - z) \cdot (3 - z)} \\ &\quad + \frac{z(z + 1)(z + 2)(1/n + 1)^3}{(2 - z)(3 - z)(4 - z)} + \dots \end{aligned} \quad (82)$$

$${}_2F_1(1, 1 + z, 2 - z, 1 + 1/n) = 1 + \frac{(1 + z) \cdot (1 + 1/n)}{2 - z} + \frac{(1 + z)(2 + z)(1 + 1/n)^2}{(2 - z)(3 - z)} \quad (83)$$

$$+ \frac{(1 + z)(2 + z)(3 + z)(1 + 1/n)^3}{(2 - z)(3 - z)(4 - z)} + \dots \quad (84)$$

$$\csc(\pi z) = \frac{1}{\pi z} + \frac{\pi z}{6} + \frac{7(\pi z)^3}{360} + \dots \quad (85)$$

Using Eqs. (79), (80), (82), (85) in Eq. (28), we find

$$\begin{aligned} N &\approx -\frac{n^{1-2z} + (1 + n)^{1-2z}}{1 - 2z} + \frac{n^{-2z}\sqrt{\pi}}{2} \cdot \left(\frac{1}{\pi z} + \frac{\pi z}{6} + \frac{7(\pi z)^3}{360} + \dots\right) \\ &\quad \cdot \left(z + \gamma z^2 + \left(\frac{\gamma^2}{2} - \frac{\pi^2}{12}\right)z^3 + \dots\right) \cdot \left[4^z \left(-\frac{1}{n}\right)^{-2z} \cdot \frac{2(z - 1/2)!}{2z - 1}\right. \\ &\quad + 4n^z(1 + n)^{1-z}\sqrt{\pi} \left[1 + \frac{z \cdot (1 + 1/n)}{2 - z} + \frac{z(1 + z) \cdot (1 + 1/n)^2}{(2 - z) \cdot (3 - z)}\right. \\ &\quad \left. \left. + \frac{z(z + 1)(z + 2)(1/n + 1)^3}{(2 - z)(3 - z)(4 - z)} + \dots\right]\right]. \end{aligned} \quad (86)$$

Using Eqs. (79), (80), (81), and (84) in Eq. (26), we also find

$$\begin{aligned} &\int_{\mathbb{R}^+} \zeta(z, x + 1)xz\zeta(z + 1, x) + \zeta(z, x)xz\zeta(z + 1, x + 1)dx \\ &\approx \frac{1}{4}n^{-2z} \left[-4^z \left(-\frac{1}{n}\right)^{-2z} \sqrt{\pi} \left(\frac{1}{\pi z} + \frac{\pi z}{6} + \frac{7(\pi z)^3}{360} + \dots\right) \cdot \left(z + \gamma z^2 + \left(\frac{\gamma^2}{2} - \frac{\pi^2}{12}\right)z^3 + \dots\right) \right. \\ &\quad \cdot \frac{2(z - 1/2)!}{2z - 1} + \frac{4}{2z - 1} \left(\frac{n}{1 + n}\right)^{z-1} \cdot \left(n + z(-z)!\right) \left[1 + \frac{(1 + z) \cdot (1 + 1/n)}{2 - z}\right. \\ &\quad \left. \left. + \frac{(1 + z)(2 + z)(1 + 1/n)^2}{(2 - z)(3 - z)} + \frac{(1 + z)(2 + z)(3 + z)(1 + 1/n)^3}{(2 - z)(3 - z)(4 - z)} + \dots\right]\right]. \end{aligned} \quad (87)$$

Hence, the nontrivial zeros can be written approximately

$$\begin{aligned}
z_n \approx & \left[-\frac{n^{1-2z} + (1+n)^{1-2z}}{1-2z} + \frac{n^{-2z}\sqrt{\pi}}{2} \cdot \left(\frac{1}{\pi z} + \frac{\pi z}{6} + \frac{7(\pi z)^3}{360} + \dots \right) \right. \\
& \cdot \left(z + \gamma z^2 + \left(\frac{\gamma^2}{2} - \frac{\pi^2}{12} \right) z^3 + \dots \right) \cdot \left[4^z \left(-\frac{1}{n} \right)^{-2z} \cdot \frac{2(z-1/2)!}{2z-1} \right. \\
& + 4n^z (1+n)^{1-z} \sqrt{\pi} \left[1 + \frac{z \cdot (1+1/n)}{2-z} + \frac{z(1+z) \cdot (1+1/n)^2}{(2-z) \cdot (3-z)} \right. \\
& + \left. \left. \frac{z(z+1)(z+2)(1/n+1)^3}{(2-z)(3-z)(4-z)} + \dots \right] \right]^{-1} \cdot 2 \left[\frac{n^{1-2z} + (1+n)^{1-2z}}{2(1-2z)} \right. \\
& + \frac{1}{4} n^{-2z} \left[-4^z \left(-\frac{1}{n} \right)^{-2z} \sqrt{\pi} \cdot \left(\frac{1}{\pi z} + \frac{\pi z}{6} + \frac{7(\pi z)^3}{360} + \dots \right) \cdot \left(z + \gamma z^2 + \left(\frac{\gamma^2}{2} - \frac{\pi^2}{12} \right) z^3 + \dots \right) \right. \\
& \cdot \frac{2(z-1/2)!}{2z-1} + \frac{4}{2z-1} \left(\frac{n}{1+n} \right)^{z-1} \cdot \left(n + z(-z)! \left[1 + \frac{(1+z) \cdot (1+1/n)}{2-z} + \frac{(1+z)(2+z)(1+1/n)^2}{(2-z)(3-z)} \right. \right. \\
& + \left. \left. \frac{(1+z)(2+z)(3+z)(1+1/n)^3}{(2-z)(3-z)(4-z)} + \dots \right] \right) \left. \right] \left. \right] + \frac{3}{2} \\
& = \frac{1}{2}(1 - i\lambda_n),
\end{aligned} \tag{88}$$

where the hypergeometric functions can be approximated using the techniques found in Ref. [9].

n	$\Im(z)$ [10]	$\Im(z)$ Eq. (88)	absolute error
1	14.134725	14.134725	4.420537×10^{-29}
2	21.022039	21.022039	2.974456×10^{-43}
3	25.010857	25.010857	1.839215×10^{-51}
4	30.424876	30.424876	2.28×10^{-62}
5	32.935061	32.935061	1×10^{-65}
6	37.586178	37.586178	1×10^{-65}
7	40.918719	40.918719	1×10^{-65}
\vdots	\vdots	\vdots	\vdots
100	236.524229	236.524229	1×10^{-64}

Table II: Imaginary Nontrivial Zeros of the Riemann Zeta Function

IV. CONCLUSION

In this study, we have discussed the domain and eigenvalues of the BBM Hamiltonian. Moreover, a second quantization procedure was performed for the BBM Schrödinger analogue equation. Finally, a closed-form expression for the nontrivial zeros of the Riemann zeta function was obtained, and a convergence test for the closed-form expression was performed.

-
- [1] Bender, C.M., Brody, D.C. and Mller, M.P., 2016. Hamiltonian for the zeros of the Riemann zeta function. arXiv preprint arXiv:1608.03679.
- [2] Riemann, B., On the Number of Prime Numbers less than a Given Quantity.(Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse.).
- [3] Berry, M.V. and Keating, J.P., 1999. $H = xp$ and the Riemann zeros. In *Supersymmetry and Trace Formulae* (pp. 355-367). Springer US.
- [4] Connes, A., 1999. Trace formula in noncommutative geometry and the zeros of the Riemann zeta function. *Selecta Mathematica, New Series*, 5(1), pp.29-106.
- [5] Odlyzko, A.M., 2001. The 10-nd zero of the Riemann zeta function. *Dynamical, Spectral, and Arithmetic Zeta Functions: AMS Special Session on Dynamical, Spectral, and Arithmetic Zeta Functions, January 15-16, 1999, San Antonio, Texas*, 290, p.139.

- [6] Faris, W.G. and Lavine, R.B., 1974. Commutators and self-adjointness of Hamiltonian operators. Communications in Mathematical Physics, 35(1), pp.39-48.
- [7] Glimm, J. and Jaffe, A., 1972. The $\lambda\varphi$ 24 Quantum Field Theory without Cutoffs. IV. Perturbations of the Hamiltonian. Journal of Mathematical Physics, 13(10), pp.1568-1584.
- [8] Mortici, C., 2010. Improved convergence towards generalized EulerMascheroni constant. Applied Mathematics and Computation, 215(9), pp.3443-3448.
- [9] Pearson, J., 2009. Computation of hypergeometric functions. University of Oxford.
- [10] <http://www.plouffe.fr/simon/constants/zeta100.html>

Appendix A: CONVERGENCE

For brevity, let us examine the convergence of the integral representation of the discrete nontrivial zeros of the Riemann zeta function on the positive half line $x \in \mathbb{R}^+$, $z \in \mathbb{C}$, $\Re(z) > 1$, and $\Re(x+1) > 0$. From Eq. (21), the integral representation of the discrete nontrivial zeros of the Riemann zeta function are given by

$$\begin{aligned} z_n = & -\frac{1}{N} \int_{\mathbb{R}^+} \zeta(z, x+1)xz\zeta(z+1, x+1)dx \\ & -\frac{1}{N} \int_{\mathbb{R}^+} \zeta(z, x)xz\zeta(z+1, x)dx \\ & +\frac{1}{N} \int_{\mathbb{R}^+} \zeta(z, x+1)xz\zeta(z+1, x) + \zeta(z, x)xz\zeta(z+1, x+1)dx + \frac{3}{2}, \end{aligned} \quad (\text{A1})$$

where

$$N = \int_{\mathbb{R}^+} \left[(n+x+1)^{-2z} - 2(n+x)^{-z}(n+x+1)^{-z} + (n+x)^{-2z} \right] dx. \quad (\text{A2})$$

Lemma 3.1. *From the first term on the RHS of Eq. (A1), if*

$$\int_0^t \zeta(z, x+1)xz\zeta(z+1, x+1)dx \quad (\text{A3})$$

exists for every number $t \geq 0$, then

$$\int_0^\infty \zeta(z, x+1)xz\zeta(z+1, x+1)dx = \lim_{t \rightarrow \infty} \int_0^t \zeta(z, x+1)xz\zeta(z+1, x+1)dx, \quad (\text{A4})$$

provided this limit exists as a finite number.

Proof.

$$\int_0^t \zeta(z, x+1)xz\zeta(z+1, x+1)dx = \frac{((n+t+1)^{-2z}((n+1)(\frac{t}{n+1}+1)^{2z} - n - 2tz - 1))}{(2z(2z-1))}. \quad (\text{A5})$$

From L'Hospital's Rule, we have

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{((n+t+1)^{-2z}((n+1)(\frac{t}{n+1}+1)^{2z} - n - 2tz - 1))}{(2z(2z-1))} \\ = & \lim_{t \rightarrow \infty} \frac{((n+t+1)^{-2z}((n+1)(\frac{t}{n+1}+1)^{2z} - n - 2tz - 1))}{(2z(2z-1))} \cdot \frac{(n+t+1)^{-2z}}{(n+t+1)^{-2z}} \\ = & \lim_{t \rightarrow \infty} \frac{-2z(n+t+1)^{-4z-1}(n(\frac{t}{n+1}+1)^{2z} + (\frac{t}{n+1}+1)^{2z} - n - 4tz + t - 1)}{4(1-2z)z^2(n+t+1)^{-2z-1}}. \end{aligned} \quad (\text{A6})$$

Upon evaluating Eq. (A6) with a series expansion at $t = \infty$, we obtain

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{(-1 - n + t + (1+n)(1 + \frac{t}{1+n})^{2z} - 4tz)}{(2(1+n+t)^{2z}z(-1+2z))} \\ = & \frac{((n+t+1)^{-2z}((n+1)(\frac{t}{n+1}+1)^{2z} - n + t(1-4z) - 1))}{(2z(2z-1))}. \end{aligned} \quad (\text{A7})$$

Hence, it can be seen that the first term on the RHS of Eq. (A1) is convergent, given that the limit seen in Eq. (A6) exists as a finite number as seen in Eq. (A7). Here, it should be pointed out that as $t = -n$, and $\Re(z) = 1/2$, Eq. (A7) is of indeterminate form. As such, we apply L'Hopital's rule to obtain

$$\begin{aligned} & \frac{(n+t+1)^{-2z} \left((n+1) \left(\frac{t}{n+1} + 1 \right)^{2z} - n + t(1-4z) - 1 \right)}{(2z(2z-1))} \\ &= \frac{2(n+1) \left(1 - \frac{n}{n+1} \right)^{(2z)} \log \left(1 - \frac{n}{n+1} \right) + 4n}{8z-2}. \end{aligned} \quad (\text{A8})$$

□

Lemma 3.2. *From the second term on the RHS of Eq. (A1), if*

$$\int_0^t \zeta(z, x) x z \zeta(z+1, x) dx \quad (\text{A9})$$

exists for every number $t \geq 0$, then

$$\int_0^\infty \zeta(z, x) x z \zeta(z+1, x) dx = \lim_{t \rightarrow \infty} \int_0^t \zeta(z, x) x z \zeta(z+1, x) dx, \quad (\text{A10})$$

provided this limit exists as a finite number.

Proof.

$$\int_0^t \zeta(z, x) x z \zeta(z+1, x) dx = - \frac{((n+t)^{-2z} \left(-n \left(\frac{n+t}{n} \right)^{2z} + n + 2tz \right))}{(2(2z-1))}. \quad (\text{A11})$$

From L'Hospital's Rule, we have

$$\begin{aligned} & - \lim_{t \rightarrow \infty} \frac{((n+t)^{-2z} \left(-n \left(\frac{n+t}{n} \right)^{2z} + n + 2tz \right))}{(2(2z-1))} \\ &= - \lim_{t \rightarrow \infty} \frac{((n+t)^{-2z} \left(-n \left(\frac{n+t}{n} \right)^{2z} + n + 2tz \right))}{(2(2z-1))} \cdot \frac{(n+t)^{-2z}}{(n+t)^{-2z}} \\ &= - \lim_{t \rightarrow \infty} \frac{(n+t)^{-4z} \left(-n \left(\frac{n+t}{n} \right)^{2z} + n + 2tz \right)}{2(2z-1)(n+t)^{-2z}} \end{aligned} \quad (\text{A12})$$

Upon evaluating Eq. (A12) with a series expansion at $t = \infty$, we obtain

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{((n+t)^{-2z} \left(-n \left(\frac{n+t}{n} \right)^{2z} + n + 2tz \right))}{((n+t)^{-2z} \left(-n \left(\frac{n+t}{n} \right)^{2z} + n + 2tz \right))} \\ &= \frac{(n+t)^{-2z} \left(-n \left(\frac{n+t}{n} \right)^{2z} + n + 2tz \right)}{(2(2z-1))}. \end{aligned} \quad (\text{A13})$$

Hence, it can be seen that the second term on the RHS of Eq. (A1) is convergent, given that the limit seen in Eq. (A12) exists as a finite number as seen in Eq. (A13). Here, it should be pointed out that as $t = -n$, and $\Re(z) = 1/2$, Eq. (A13) is undefined. □

Lemma 3.3. *From the third term on the RHS of Eq. (A1), if*

$$\int_0^t \zeta(z, x+1) x z \zeta(z+1, x) + \zeta(z, x) x z \zeta(z+1, x+1) dx \quad (\text{A14})$$

exists for every number $t \geq 0$, then

$$\begin{aligned} & \int_0^\infty \zeta(z, x+1) x z \zeta(z+1, x) + \zeta(z, x) x z \zeta(z+1, x+1) dx \\ &= \lim_{t \rightarrow \infty} \int_0^t \zeta(z, x+1) x z \zeta(z+1, x) + \zeta(z, x) x z \zeta(z+1, x+1) dx, \end{aligned} \quad (\text{A15})$$

provided this limit exists as a finite number.

Proof. From the RHS of Eq. (26) it can be seen that

$$\begin{aligned} & \int_0^t \zeta(z, x+1)xz\zeta(z+1, x) + \zeta(z, x)xz\zeta(z+1, x+1)dx \\ &= \frac{((n+t)^{-z}(n+t+1)^{-z}((n+t)_2F_1(1, 1-2z, 1-z, n+t+1) - n - 2tz))}{(2z-1)} \\ & - \frac{((n)^{-z}(n+1)^{-z}((n)_2F_1(1, 1-2z, 1-z, n+1) - n))}{(2z-1)}. \end{aligned} \quad (\text{A16})$$

Since the second term on the RHS of Eq. (A16) is independent of t , we are only concerned with the limit of the first term on the RHS of Eq. (A16). As such, we consider the limit

$$\lim_{t \rightarrow \infty} \frac{((n+t)_2F_1(1, 1-2z, 1-z, n+t+1) - n - 2tz)}{(n+t)^z(n+t+1)^z(2z-1)}. \quad (\text{A17})$$

Here, it is useful to employ Gauss' theorem, i.e.,

$${}_2F_1(1, 1-2z, 1-z, n+t+1) = \frac{\Gamma(1-z)\Gamma(z-1)}{\Gamma(-z)\Gamma(z)} \quad (\text{A18})$$

where $\Re(z) > 1$, $n = -t$, and

$$\Gamma(z) = \int_0^\infty x^{z-1}e^{-x}dx \quad (\text{A19})$$

is the gamma function. Therefore, Eq. (A17) can be written

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{((n+t) \frac{\Gamma(1-z)\Gamma(z-1)}{\Gamma(-z)\Gamma(z)} - n - 2tz)}{(n+t)^z(n+t+1)^z(2z-1)} \\ &= - \lim_{t \rightarrow \infty} \frac{(n+t)^{-z}(n+t+1)^{-z}(n+t)}{(z-1)}. \end{aligned} \quad (\text{A20})$$

Upon evaluating Eq. (A20) with a series expansion at $t = \infty$, we obtain

$$\lim_{t \rightarrow \infty} \frac{((n+t)_2F_1(1, 1-2z, 1-z, n+t+1) - n - 2tz)}{(n+t)^z(n+t+1)^z(2z-1)} = (n+t)^{-z}(n+t+1)^{-z} \left[-\frac{n}{(z-1)} - \frac{(tz)}{(z-1)} \right]. \quad (\text{A21})$$

Hence, it can be seen that the third term on the RHS of Eq. (A1) is convergent, given that the limit seen in Eq. (A17) exists as a finite number as seen in Eq. (A21). Here, it should be pointed out that as $t = -n$, and $\Re(z) = 1/2$, Eq. (A21) is undefined. Moreover, the second term on the RHS of Eq. (A16) is indeterminate at $\Re(z)$. \square

Finally, we must consider the convergence of the normalization factor N .

Lemma 3.4. *From the first three terms on the RHS of Eq. (A1), if*

$$\int_0^t \left[(n+x+1)^{-2z} - 2(n+x)^{-z}(n+x+1)^{-z} + (n+x)^{-2z} \right] dx \quad (\text{A22})$$

exists for every number $t \geq 0$, then

$$\begin{aligned} & \int_0^\infty \left[(n+x+1)^{-2z} - 2(n+x)^{-z}(n+x+1)^{-z} + (n+x)^{-2z} \right] dx \\ &= \lim_{t \rightarrow \infty} \int_0^t \left[(n+x+1)^{-2z} - 2(n+x)^{-z}(n+x+1)^{-z} + (n+x)^{-2z} \right] dx \end{aligned} \quad (\text{A23})$$

provided this limit exists as a finite number.

Proof.

$$\begin{aligned}
& \int_0^\infty \left[(n+x+1)^{-2z} - 2(n+x)^{-z}(n+x+1)^{-z} + (n+x)^{-2z} \right] dx \\
&= \lim_{t \rightarrow \infty} \frac{((n+t+1)^{-2z}((n+1)(\frac{t}{n+1}) + 1)^{2z} - n - t - 1)}{(2z-1)} \\
&+ \lim_{t \rightarrow \infty} \frac{((n+t)^{-2z}(n((\frac{n+t}{n})^{2z} - 1) - t))}{(2z-1)} \\
&+ \lim_{t \rightarrow \infty} \frac{(2(-n-t)^z(n+t)^{-z}(n+t+1)^{1-z} {}_2F_1(1-z, z, 2-z, n+t+1))}{(z-1)} \\
&- \frac{(2(-n)^z(n)^{-z}(n+1)^{1-z} {}_2F_1(1-z, z, 2-z, n+1))}{(z-1)}, \tag{A24}
\end{aligned}$$

where the last term on the RHS of Eq. (A24) omits the limit, as it is independent of t . The limits seen on the RHS of Eq. (A24) can be evaluated in a similar manner to those seen in Eqs. (A7), (A13), and (A17), respectively. \square