

# On the Riemann Hypothesis, Complex Scalings and Logarithmic Time Reversal

Carlos Castro

Center for Theoretical Studies of Physical Systems  
Clark Atlanta University, Atlanta, GA. 30314, castro@ctps.cau.edu

May 2017

## Abstract

An approach to solving the Riemann Hypothesis is revisited within the framework of the special properties of  $\Theta$  (theta) functions, and the notion of  $\mathcal{CT}$  invariance. The conjugation operation  $\mathcal{C}$  amounts to complex scaling transformations, and the  $\mathcal{T}$  operation  $t \rightarrow (1/t)$  amounts to the reversal  $\log(t) \rightarrow -\log(t)$ . A judicious scaling-like operator is constructed whose spectrum  $E_s = s(1-s)$  is real-valued, leading to  $s = \frac{1}{2} + i\rho$ , and/or  $s = \text{real}$ . These values are the location of the non-trivial and trivial zeta zeros, respectively. A thorough analysis of the one-to-one correspondence among the zeta zeros, and the orthogonality conditions among pairs of eigenfunctions, reveals that *no* zeros exist off the critical line. The role of the  $\mathcal{C}, \mathcal{T}$  transformations, and the properties of the Mellin transform of  $\Theta$  functions were essential in our construction.

Keywords: Quantum Mechanics, Dirac Operators, Riemann Hypothesis, Hilbert-Polya conjecture, Modularity.

Riemann's outstanding hypothesis (RH) [1] that the non-trivial complex zeros of the zeta-function  $\zeta(s)$  must be of the form  $s_n = 1/2 \pm i\rho_n$ , is one of most important open problems in pure mathematics. The zeta-function has a relation with the number of prime numbers less than a given quantity and the zeros of zeta are deeply connected with the distribution of primes [1]. References [2] are devoted to the mathematical properties of the zeta-function. The RH has also been studied from the point of view of mathematics and physics by [7], [13], [19], [23], [10], [24], [8], [9], [25], [12], [22], [27], [18], [28], among many others. And most recently by [29], [14] and [15].

Let us begin with the one-dimensional differential operators [21], [16]

$$D_1 = -\frac{d}{d \ln t} + \frac{dV}{d \ln t} + k. \quad (1.1)$$

where  $k$  is an arbitrary parameter. The eigenvalues  $s$  can be complex-valued in general, and its eigenfunctions are

$$\psi_s(t) = t^{-s+k} e^{V(t)}. \quad (1.2a)$$

$D_1$  is *not* self-adjoint since it is an operator that does not admit an adjoint extension to the whole real line characterized by the *real* variable  $t$ . The parameter  $k$  is also real-valued. For this reason the eigenvalues of  $D_1$  can be complex valued numbers  $s$ . The conjugation operation  $\mathcal{C}$  acting on the eigenfunctions is defined as

$$\psi_s(t) = t^{-s+k} e^{V(t)} \rightarrow \psi_{s^*}(t) = t^{-s^*+k} e^{V(t)} = t^{-s^*+s} \psi_s(t). \quad (1.2b)$$

one learns that the conjugation operation  $\mathcal{C}$  can also be recast as a *scaling* transformation of  $\psi_s(t)$  by  $t$ -dependent (local) scaling factors

$$t^{-s^*+s} = e^{(-s^*+s) (\ln t)} = e^{2i \operatorname{Im}(s) (\ln t)} = e^{i\theta_s(t)} \quad (1.2c)$$

which amounts to a  $t$ -dependent phase rotation  $\theta_s(t)$  which is proportional to the imaginary part of  $s$  and to  $\ln t$ .

We also define the “mirror” operator to  $D_1$  as follows,

$$D_2 = \frac{d}{d \ln t} - \frac{dV(1/t)}{d \ln t} + k. \quad (1.3)$$

that is related to  $D_1$  by the substitution  $t \rightarrow 1/t$  and by noticing that

$$\frac{dV(1/t)}{d \ln(1/t)} = -\frac{dV(1/t)}{d \ln t}. \quad (1.4)$$

where  $V(1/t)$  is not equal to  $V(t)$  and  $D_2$  is *not* self-adjoint either. The eigenfunctions of the  $D_2$  operator are  $\Psi_s(\frac{1}{t})$ , with the same eigenvalue  $s$

$$D_2 \Psi_s\left(\frac{1}{t}\right) = s \Psi_s\left(\frac{1}{t}\right) \quad (1.5)$$

A “Wick rotation” of variables  $t = iz$  furnishes  $z \rightarrow -(1/z)$  which is a truly modular  $SL(2, Z)$  transformation  $z \rightarrow (az + b/cz + d)$  with unit determinant  $ad - bc = 1$ .

Out of the infinity of possible choices for  $V(t)$ , one may choose  $V(t)$  which is related to the Bernoulli string spectral counting function, and given by the Jacobi theta series as follows

$$e^{2V(t)} = \sum_{n=-\infty}^{\infty} e^{-\pi n^2 t^l} = 2\omega(t^l) + 1. \quad (1.6)$$

where  $l$  is another real parameter introduced corresponding to the scaling exponent  $t^l$  in eq-(1.6).

The related theta function defined by Gauss is given by

$$G(1/x) = \sum_{n=-\infty}^{\infty} e^{-\pi n^2/x} = 2\omega(1/x) + 1. \quad (1.7)$$

where  $\omega(x) = \sum_{n=1}^{\infty} e^{-\pi n^2 x}$ . The Gauss-Jacobi series obeys the relation

$$G\left(\frac{1}{x}\right) = \sqrt{x} G(x). \quad (1.8)$$

resulting from the Poisson re-summation formula.

After setting  $e^{2V(t)} = G(t^l) = G(x)$ , where  $x \equiv t^l$ , by recurring to the properties of the Gauss-Jacobi theta series under the  $x \rightarrow 1/x$  transformations (1.8), and when the parameters  $l, k$  are *constrained* to obey the condition  $l = 4(2k - 1)$ , one can show that the eigenfunctions of the  $D_2$  operator  $\Psi_s(\frac{1}{t})$ , satisfy the key relation  $\Psi_{1-s}(t) = \Psi_s(\frac{1}{t})$  [16], [21].

A little algebra reveals that the pair of mirror ‘‘Hamiltonians’’  $H_A = D_2 D_1$  and  $H_B = D_1 D_2$ , when  $l = 4(2k - 1)$  have for eigenvalues and eigenfunctions the following

$$H_A \Psi_s(t) = s(1 - s)\Psi_s(t). \quad H_B \Psi_s\left(\frac{1}{t}\right) = s(1 - s)\Psi_s\left(\frac{1}{t}\right). \quad (1.9)$$

due to the relation  $\Psi_s(1/t) = \Psi_{1-s}(t)$  based on the modular properties of the Gauss-Jacobi series,  $G(\frac{1}{x}) = \sqrt{x} G(x)$ . Therefore, despite that  $H_A, H_B$  are *not* Hermitian they have the same spectrum  $s(1 - s)$  which is *real*-valued only in the critical line *and* in the real line. Eq-(1.9) is the one-dimensional version of the eigenfunctions of the two-dimensional hyperbolic Laplacian given in terms of the Eisenstein’s series.

The inner product is defined as follows <sup>1</sup>

$$\langle f|g \rangle = \int_0^{\infty} f^* g \frac{dt}{t}.$$

Based on this definition, the inner product of two eigenfunctions of  $D_1$  is

$$\langle \psi_{s_1} | \psi_{s_2} \rangle = \int_0^{\infty} e^{2V} t^{-s_{12}+2k-1} dt; \quad s_{12} \equiv s_1^* + s_2 \quad (1.10a)$$

A regularization of the integral (1.10a) can be attained by removing the zero  $n = 0$  mode of the Gauss-Jacobi series. Upon doing so and performing the change of variables  $x = t^l$ , it leads to

$$\frac{2}{l} \int_0^{\infty} e^{2V} x^{\frac{2(-s_{12}+2k)}{2l}-1} dx = \frac{2}{l} Z \left[ \frac{2}{l}(2k - s_{12}) \right] = \frac{2}{l} Z[s] \quad (1.10b)$$

---

<sup>1</sup>At the moment we are not concerned if one has a Banach or a Hilbert space

where we have denoted  $s_{12} = s_1^* + s_2 = x_1 + x_2 + i(y_2 - y_1)$ , and  $s = \frac{2}{l}(-s_{12} + 2k)$ . The completed zeta function  $Z[s]$  results from the evaluation of the Mellin transform as shown next. It is known that the completed zeta function  $Z[s]$  can be expressed in terms of the Jacobi theta series,  $\omega(x)$  defined by eqs-(1.6, 1.7) as the integral [2]

$$\begin{aligned}
\int_0^\infty \sum_{n=1}^\infty e^{-\pi n^2 x} x^{s/2-1} dx &= \\
&= \int_0^\infty x^{s/2-1} \omega(x) dx \\
&= \frac{1}{s(s-1)} + \int_1^\infty [x^{s/2-1} + x^{(1-s)/2-1}] \omega(x) dx \\
&= Z(s) = Z(1-s),
\end{aligned} \tag{1.11}$$

where the completed zeta function is defined as

$$Z(s) \equiv \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s). \tag{1.12}$$

and which obeys the functional relation  $Z(s) = Z(1-s)$ .

To sum up, a family of scaling-like operators  $D_1, D_2$  in one dimension allows to evaluate the inner product of their eigenfunctions  $\Psi_s(t)$  (after removing the zero mode of the Gauss-Jacobi theta series) giving  $(2/l)Z[\frac{2}{l}(2k - s^* - s)]$ , where  $Z(s)$  is the Riemann completed zeta function and the  $l, k$  parameters are constrained to obey  $l = 4(2k-1)$  in order to have the relation  $\Psi_s(1/t) = \Psi_{1-s}(t)$ . Hence, by using the properties of the Gauss-Jacobi series  $G(\frac{1}{x}) = \sqrt{x} G(x)$  it follows that under the log-time reversal  $\mathcal{T}$  operation  $lnt \rightarrow -lnt$  (equivalent to  $t \rightarrow \frac{1}{t}$ ) the eigenfunctions  $\Psi_s(t)$  behave as

$$\mathcal{T} \Psi_s(t) = \Psi_s\left(\frac{1}{t}\right) = \Psi_{1-s}(t). \tag{1.13}$$

In order to avoid the removal of the zero mode  $n = 0$  of the Gauss-Jacobi theta series and evaluate the integrals appearing in the inner products, in [16] we proposed a family of theta series where *no* regularization is needed in the construction of the inner products. There is a two-parameter family of theta series  $\Theta_{j,m}(t)$  that yield well defined inner products *without* the need to extract the zero mode  $n = 0$  divergent contribution. We found that the two parameter family of theta series related to a different choice for  $V(t)$  is given by

$$e^{2V_{j,m}(t)} = \Theta_{j,m}(t^l) \equiv \sum_{n=-\infty}^{n=\infty} n^{2m} H_{2j}(n\sqrt{2\pi t^l}) e^{-\pi n^2 t^l}, \quad m = 1, 2, \dots; \quad j = 0, 1, 2, \dots \tag{1.14}$$

Due to the *weighted* theta series in eq-(1.14) the zero mode  $n = 0$  does *not* contribute to the sum in eq-(1.14), since  $m$  is a positive integer, and the Mellin transform of  $\Theta_{j,m}(t^l) = \Theta_{j,m}(x)$  ( $x = t^l$ ) after exploiting the symmetry of the even-degree Hermite polynomials, is given by [5], [6]

$$\int_0^\infty \left[ \frac{2}{l} \sum_{n=1}^{n=\infty} n^{2m} H_{2j}(n\sqrt{2\pi x}) e^{-\pi n^2 x} \right] x^{s/2-1} dx = -\frac{2}{l} (8\pi)^j P_j(s) \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s-2m); \quad \text{Re } s > 1+2m, \quad m = 1, 2, \dots \quad (1.15)$$

The polynomial pre-factor in (1.15) is given in terms of a terminating Hypergeometric series [6]

$$P_j(s) = (8\pi)^{-j} (-1)^j \frac{(2j)!}{j!} {}_2F_1\left(-j, \frac{s}{2}; \frac{1}{2}; 2\right). \quad (1.16)$$

The polynomial  $P_j(s)$  has simple zeros on the critical line  $\text{Re } s = \frac{1}{2}$ , obeys the functional relation  $P_j(s) = (-1)^j P_j(1-s)$  and in particular  $P_j(s = \frac{1}{2}) = 0$  when  $j = \text{odd}$  [6]. It is only when  $j = \text{even}$  that  $P_j(s = \frac{1}{2}) \neq 0$ . In order to find the analytical continuation of the Mellin transform (1.15) for *all* values of  $s$  in the complex plane we must use the analytical continuation of  $\zeta(s)$  as found by Riemann in his celebrated paper. A Poisson re-summation formula for  $\Theta_{j,m}(x)$  (1.14) leads to the important relation

$$\frac{(-1)^j}{\sqrt{x}} \Theta_{j,m}\left(\frac{1}{x}\right) = \Theta_{j,m}(x). \quad (1.17)$$

which allows us to show that only when  $j = \text{even}$  one can implement the  $\mathcal{T}$  transformations of the new eigenfunctions  $\Psi_s^{j,m}(t) = t^{-s+k} e^{V_{j,m}(t)}$  of  $D_1$ , and corresponding to the weighted theta series  $\Theta_{j,m}(t^l)$  of eq-(1.14), giving

$$\mathcal{T} \Psi_s^{j,m}(t) = \Psi_s^{j,m}\left(\frac{1}{t}\right) = \Psi_{1-s}^{j,m}(t) \quad (1.18)$$

this relationship requires that one must have

$$j = \text{even}, \quad l = 4(2k-1) \quad (1.19)$$

Therefore, the eigenfunctions and eigenvalues of the pair of ‘‘Hamiltonians’’ is

$$H_A \Psi_s^{j,m}(t) = s(1-s) \Psi_s^{j,m}(t), \quad H_B \Psi_s^{j,m}\left(\frac{1}{t}\right) = s(1-s) \Psi_s^{j,m}\left(\frac{1}{t}\right) \quad (1.20)$$

subjected to the conditions in eq-(1.18).

We explicitly inserted the superscripts  $j, m$  in  $\Psi_s^{j,m}(t) = t^{-s+k} e^{V_{j,m}(t)}$  to denote the  $j, m$  dependence in the definition of  $V(t)$  given by eq-(1.14).

In what follows we shall omit the superscripts  $j, m$  for convenience. Defining

$$\Psi_s^{\mathcal{CT}}(t) \equiv \mathcal{C} \mathcal{T} \Psi_s(t) = \mathcal{C} \Psi_s\left(\frac{1}{t}\right) = \mathcal{C} \Psi_{1-s}(t) = \Psi_{1-s^*}(t) \quad (1.21)$$

one finds that it is also an eigenfunction of  $H_A$  with an eigenvalue  $s^*(1-s^*)$  :

$$\begin{aligned} H_A | \Psi_s^{\mathcal{CT}}(t) \rangle &= H_A \mathcal{CT} | \Psi_s(t) \rangle = H_A | \Psi_{1-s^*}(t) \rangle = \\ s^*(1-s^*) | \Psi_{1-s^*}(t) \rangle &= s^*(1-s^*) \mathcal{CT} | \Psi_s(t) \rangle = (E_s)^* | \Psi_s^{\mathcal{CT}}(t) \rangle . \end{aligned} \quad (1.22)$$

where we have defined  $(E_s)^* = s^*(1-s^*)$ .

If the  $\mathcal{CT}$  action on  $s(1-s) \Psi_s$  is *linear* :  $\mathcal{CT} s(1-s) \Psi_s = s(1-s) \mathcal{CT} \Psi_s$ , instead of *antilinear* :  $\mathcal{CT} s(1-s) \Psi_s = s^*(1-s^*) \mathcal{CT} \Psi_s$ , and if

$$\begin{aligned} \langle \Psi_s | [H_A, \mathcal{CT}] | \Psi_s \rangle &= 0 \Rightarrow \\ \langle \Psi_s | H_A \mathcal{CT} | \Psi_s \rangle - \langle \Psi_s | \mathcal{CT} H_A | \Psi_s \rangle &= \\ (E_s)^* \langle \Psi_s | \mathcal{CT} | \Psi_s \rangle - E_s \langle \Psi_s | \mathcal{CT} | \Psi_s \rangle &= \\ (E_s^* - E_s) \langle \Psi_s | \mathcal{CT} | \Psi_s \rangle &= 0. \end{aligned} \quad (1.23)$$

Similar results follow for the  $H_B$  operator. From eq-(1.23) one has two cases to consider.

- Case A : If the pseudo-norm is null

$$\langle \Psi_s | \mathcal{CT} | \Psi_s \rangle = 0 \Rightarrow (E_s - E_s^*) \neq 0 \quad (1.24)$$

then the *complex* eigenvalues  $E_s = s(1-s)$  and  $E_s^* = s^*(1-s^*)$  are *complex* conjugates of each other. In this case one cannot prove the RH, and there exists the possibility that there are quartets of non-trivial Riemann zeta zeros (off the critical line) given by  $s_n, 1-s_n, s_n^*, 1-s_n^*$ .

- Case B : If the pseudo-norm is *not* null :

$$\langle \Psi_s | \mathcal{CT} | \Psi_s \rangle \neq 0 \Rightarrow (E_s - E_s^*) = 0 \quad (1.25)$$

then the eigenvalues are *real* given by  $E_s = s(1-s) = E_s^* = s^*(1-s^*)$  and which implies that  $s = \text{real}$  (location of the trivial zeta zeros) and/or  $s = \frac{1}{2} + i\rho$  (location of the non-trivial zeta zeros). In this case the RH would be true and the non-trivial Riemann zeta zeros are given by  $s_n = \frac{1}{2} + i\rho_n$  and  $1-s_n = s_n^* = \frac{1}{2} - i\rho_n$ . We are going to prove next why Case **A** does and cannot occur, therefore the RH is true because we are left with case **B**.<sup>2</sup>

Therefore one has now at our disposal a well defined inner product of the states  $\Psi_s^{j,m}(t)$  (*without* the need to regularize the integrals by extracting out the zero  $n=0$  mode of the theta series). In particular, from eq-(1.15) one learns

<sup>2</sup>The authors [4] were the first to my knowledge who explored the possibility that *PT* symmetry might be relevant to the RH. It was their work which inspired us.

that the inner product of the states  $\Psi_s^{j,m}(t)$  with the states  $\Psi_{\frac{1}{2}+2m}(t)$ , where  $m = 1, 2, \dots$ , is given by

$$\langle \Psi_{\frac{1}{2}+2m}^{j,m}(t) | \Psi_s^{j,m}(t) \rangle = -\frac{2}{l} (8\pi)^j P_j(s+2m) \pi^{-(s+2m)/2} \Gamma\left(\frac{s+2m}{2}\right) \zeta(s). \quad (1.26)$$

to arrive at this result above requires performing the change of variables  $t^l = x$ , and *fixing* uniquely the values  $l = -2; k = \frac{1}{4}$  obeying the required constraint  $l = 4(2k - 1)$  in eq-(1.18). Therefore, the non-trivial zeta zeros  $s_n = \frac{1}{2} \pm \rho_n$  are in a one-to-one correspondence with the states  $\Psi_{s_n}^{j,m}(t)$  *orthogonal* to the states  $\Psi_{\frac{1}{2}+2m}^{j,m}(t)$  in eq-(1.26) :

$$\begin{aligned} & \langle \Psi_{\frac{1}{2}+2m}^{j,m}(t) | \Psi_{s_n}^{j,m}(t) \rangle = \\ & -\frac{2}{l} (8\pi)^j P_j(s_n + 2m) \pi^{-(s_n+2m)/2} \Gamma\left(\frac{s_n + 2m}{2}\right) \zeta(s_n) = 0; \quad m = 1, 2, 3, \dots \end{aligned} \quad (1.27)$$

It remains to prove, when  $l = -2, k = \frac{1}{4}, t^l = x$ , and  $s_{12} = s_1^* + s_2 = s_1^* + (1 - s_1^*) = 1$ , that

$$\begin{aligned} & \langle \Psi_s^{j,m}(t) | \mathcal{CT} | \Psi_s^{j,m}(t) \rangle = \langle \Psi_s^{j,m}(t) | | \Psi_{1-s^*}^{j,m}(t) \rangle = \\ & \int_0^\infty \left[ \frac{2}{l} \sum_{n=1}^{n=\infty} n^{2m} H_{2j}(n\sqrt{2\pi x}) e^{-\pi n^2 x} \right] x^{\frac{2(-s_{12}+2k)}{2l}-1} dx = \\ & -\frac{2}{l} (8\pi)^j P_j\left(s = \frac{1}{2}\right) \pi^{-1/4} \Gamma\left(\frac{1}{4}\right) \zeta\left(\frac{1}{2} - 2m\right) \neq 0; \quad j = \text{even}, m = 1, 2, 3, \dots \end{aligned} \quad (1.28)$$

Hence, one arrives at a definite solid conclusion in eq-(1.28). Because  $\zeta\left(\frac{1}{2} - 2m\right) \neq 0$  when  $m = 1, 2, \dots$ , and  $P_j\left(\frac{1}{2}\right) \neq 0$  when  $j = \text{even}$  in eq-(1.28), then  $\langle \Psi_s | \mathcal{CT} | \Psi_s \rangle \neq 0$ , and this rules out case **A** in eq-(1.24), and singles out case **B** in eq-(1.25) leading to the conclusion that  $E_s = s(1-s) = \text{real} \Rightarrow s = \frac{1}{2} + i\rho$  ( and/or  $s = \text{real}$  ), which is the *location* of the non-trivial zeta zeros (if the RH is true) and trivial zeta zeros, respectively.

Armed with these findings that the eigenvalues  $s$  which define the eigenfunctions  $\Psi_s(t)$  must be real and/or reside in the critical line, we can proceed further than we did back in [16] and gain more information about the location of the zeta zeros. Let us analyze the scenario in case the RH were *not* true. Given any real number  $s' = \frac{1}{2} + \xi \in \mathcal{R}$ , such that  $\xi \neq 2m$ , and  $s = \frac{1}{2} + i\lambda \in \mathcal{L}$  residing in the critical line, let us imagine that the inner product

$$\begin{aligned} & \langle \Psi_{s'}^{j,m}(t) | \Psi_s^{j,m}(t) \rangle = \langle \Psi_{\frac{1}{2}+\xi}^{j,m}(t) | \Psi_s^{j,m}(t) \rangle = \\ & -\frac{2}{l} (8\pi)^j P_j(s + \xi) \pi^{-(s+\xi)/2} \Gamma\left(\frac{s+\xi}{2}\right) \zeta(s + \xi - 2m) = 0 \end{aligned} \quad (1.29)$$

and its complex conjugate

$$\begin{aligned}
\langle \Psi_{s'}^{j,m}(t) | \Psi_{s^*}^{j,m}(t) \rangle &= \langle \Psi_{\frac{1}{2}+\xi}^{j,m}(t) | \Psi_{s^*}^{j,m}(t) \rangle = \\
-\frac{2}{l} (8\pi)^j P_j(s^* + \xi) \pi^{-(s^*+\xi)/2} \Gamma\left(\frac{s^* + \xi}{2}\right) \zeta(s^* + \xi - 2m) &= 0 \quad (1.30)
\end{aligned}$$

generate other nontrivial zeros *off* the critical line given by  $z \equiv s + \xi - 2m$ ;  $z^* \equiv s^* + \xi - 2m$ , respectively. Then, by symmetry,  $1 - z^* = s - \xi + 2m$ , and  $1 - z = s^* - \xi + 2m$  should also be another pair of complex conjugate (putative) zeros *off* the critical line since the number of zeros *off* the critical line must appear in *quartets* resulting from the symmetry property of the completed zeta function  $Z(s) = Z(1 - s)$ .

However, due to the fact that the numbers  $m$  are *positive* integers, and from inspection of the fundamental integral in eq-(1.15), one can infer that this latter pair of complex conjugate zeros  $s - \xi + 2m$ , and  $s^* - \xi + 2m$ , *cannot* be obtained from an orthogonality condition among the  $\Psi_s^{j,m}(t)$  and  $\Psi_{s'}^{j,m}(t)$ , for any  $s$  lying in the critical line, and any  $s'$  lying in the real line. Consequently, if there were zeros off the critical line, only *half* of those could be obtained from imposing the orthogonality conditions. The only way one could generate all the (non-trivial) zeros from the orthogonality conditions is when *all* of them reside in the critical line and which is consistent with the RH.

It is true that one could have the inner products

$$\langle \Psi_{\frac{1}{2}+\xi'}^{j,m'}(t) | \Psi_s^{j,m'}(t) \rangle = -\frac{2}{l} (8\pi)^j P_j(s+\xi') \pi^{-(s+\xi')/2} \Gamma\left(\frac{s+\xi'}{2}\right) \zeta(s+\xi'-2m') \quad (1.31)$$

corresponding to states associated to *different* values of  $m' \neq m$ , and  $\xi' \neq \xi$ , such that

$$\zeta(s + \xi' - 2m') = \zeta(s - \xi + 2m) = 0 \quad (1.32)$$

namely, one could perform the identification

$$s + \xi' - 2m' = s - \xi + 2m \Rightarrow \xi + \xi' = 2m + 2m', \quad \xi \neq 2m; \quad \xi' \neq 2m' \quad (1.33)$$

and claim that one has found the sought-after pair of orthogonal states  $\Psi_{\frac{1}{2}+\xi'}^{jm'}(t)$ ,  $\Psi_s^{jm'}(t)$  which generates the putative zero off the critical line given by  $s - \xi + 2m$ .

However, in this case one would be choosing eigenfunctions  $\Psi_s^{jm'}(t)$ ,  $\Psi_{\frac{1}{2}+\xi'}^{jm'}(t)$  of another *different* operator  $D'_1$  resulting from the different value of  $m' \neq m$ , and which leads to a different weighted theta series in eq-(1.14). Therefore, in order to generate the *quartets* of putative zeros off the critical line one would be forced to look at the orthogonality conditions involving eigenfunctions of many *different* operators  $D_1, D'_1, D''_1, \dots$  associated with many *different* functions  $V_{j,m}(t), V_{j,m'}(t), V_{j,m''}(t), \dots$ , instead of focusing on the orthogonality conditions involving eigenfunctions  $\Psi_s^{j,m}(t)$  of only *one* operator  $D_1$ , associated to only *one* function  $V_{j,m}(t)$ .



The above arguments are reminiscent of our prior physical interpretation of the location of the nontrivial Riemann zeta zeros. These locations corresponded to the presence of tachyonic-resonances/tachyonic-condensates in bosonic string theory [11]. We found that if there were zeros *off* the critical line violating the RH these zeros do *not* correspond to any poles of the string scattering amplitude. We believe that complex scalings and logarithmic time reversal transformations hold important clues as to why the Riemann hypothesis is true. The role of the  $\mathcal{C}, \mathcal{T}$  transformations, and the properties of the Mellin transform [6] were essential in our construction.

### Acknowledgements

We thank to M. Bowers for her hospitality and assistance.

### References

- [1] B. Riemann, On the number of prime numbers less than a given quantity, Monatsberichte der Berliner Akademie, November, 1859. (Translated by D. R. Wilkins, 1998).
- [2] A. A. Karatsuba, S. M. Voronin, *The Riemann zeta function*. [Translated from the Russian by Neal Koblitz ] (Berlin-New York, Walter de Gruyter Pub., 1992); S. J. Patterson, *An introduction to the theory of the Riemann zeta function* (Cambridge, Univ. Press, 1988); H. M. Edwards, *Riemann's Zeta Function* ( New York, Dover Pub., 2001); E. C. Titchmarsh, *The theory of the Riemann zeta-function* (Oxford, Clarendon Press, 1986).
- [3] J. V. Armitage, The Riemann hypothesis and the Hamiltonian of a quantum mechanical system, in *Number theory and dynamical systems*, (London Math Soc Lecture Notes, series 134, Cambridge University Press). V. Armitage, Zeta Functions with a Zero at  $s = \frac{1}{2}$ , *Inven. Mathematicae* **15** (1972) 199.
- [4] Z. Ahmed and S. Jain, "A pseudo-unitary ensemble of random matrices, PT symmetry and the Riemann hypothesis" *Mod. Phys. Lett.* **A 21** (2005) 331.
- [5] S. Keating, An infinite family of zeta functions indexed by Hermite polynomials, *J. Math. Analysis Appls.* **140** (1989) 218.
- [6] M. Coffey, Theta and Riemann  $\xi$  function representations from Harmonic Oscillator eigensolutions, *Phys. Letts* **A 362** (2007) 352; Special functions and the Mellin transforms of Laguerre and Hermite functions, [arXiv : math-ph/0612085].

- [7] L. de Branges, *J. Funct. Anal.* **107** (1992) 122; J. Alcántara-Bode, *Int. Eqs. Op. Theory* **17** (1993) 151; L. de Branges, *Funct. Anal.* **121** (1994) 117; X.-J. Li, *Math. Nachr.* **166** (1994) 229; N. Nikolski, *Ann. Inst. Fourier (Grenoble)* **45** (1995) 143; V.I. Vasyunin, *Algebra i Analiz* **7** (1995) 118 [Trans. St. Petersburg Math. J. **7** (1996) 405].
- [8] D. Schumayer and D. A. W. Hutchinson, “Physics of the Riemann Hypothesis” , *Rev. Mod. Phys* **83** (2011) 307 [arXiv: 1101.3116].
- [9] G. Mussardo, “The quantum mechanical potential for the prime numbers”, the Abdus Salam International Centre For Theoretical Physics, ISAS/EP/97/153.
- [10] M. Pitkanen, “A further step in the proof of Riemann hypothesis” arXiv : genmath/ 0109072. See also <http://www.physics.helsinki.fi/~matpitka/>.
- [11] C. Castro, *Int. J. Geom Methods in Mod Phys* **3**, no.2 (2006) 187.
- [12] H. Montgomery, *Proc. Int. Congress of Mathematics* (Vancouver 1974, vol. 1, 379).
- [13] N. Katz and P. Sarnak, *Bull. Am. Math. Soc.*, **36** (1999) 1. J. Main, *Phys. Rep* **316** (1999) 233.
- [14] C. Bender, D. Brody and M. Muller, “Hamiltonians for the zeros of the Riemann Zeta Function”, *Phys. Rev. Lett*, **118** (2017) 130201.
- [15] J. Bellissard, “Comment on Hamiltonians for the zeros of the Riemann Zeta Function” arXiv : 1704.02644.
- [16] C. Castro, *Int. J. Geom Methods in Mod Phys* **5**, no.1 (2008) 187.  
C.Castro, “On the Riemann Hypothesis, Area Quantization, Dirac Operators, Modularity and Renormalization Group” *International Journal of Geometric Methods in Modern Physics* **7**, No. 1 (2010) 1.
- [17] R. Gorenflo and F. Mainardi, ”Fractional Calculus : Integral and Differential Equations of Fractional Order” arXiv : 0805.3823.
- [18] H. Wu and D. W. L. Sprung, *Phys. Rev. E* **48** (1993) 2595.
- [19] M. Berry and J. Keating, *SIAM Review*, **41**, no. 2 (1999) 236. M. V. Berry and Z. V. Lewis, *Proc. Roy. Soc. Lond. A* **370** (1980) 459. M. Berry, ”  $H = xp$  and the Riemann zeros” in *Supersymmetry and Trace Formulae ” Chaos and Disorder* , ed. J. Keating, D. Khmelnitskii and I. Lerner, Kluwer 1999.
- [20] B. P. van Zyl and D. A. W. Hutchinson, *Phys. Rev. E* **67** (2003) 066211.

- [21] C. Castro and J. Mahecha, Fractal SUSY QM, Geometric Probability and the Riemann Hypothesis, *Int. J. Geom Methods in Mod Phys* **1** no.6 (2004) 751. C. Castro, "On strategies towards the Riemann Hypothesis : Fractal SUSY-QM and a Trace formula", *Int. J. Geom Methods in Mod Phys* **4**, no. 5 (2007) 881. C. Castro and J. Mahecha, Final steps towards a proof of the Riemann hypothesis, [arXiv:hep-th/0208221].
- [22] P. Slater, A Numerical Examination of the Castro-Mahecha Supersymmetric Model of the Riemann Zeros, [arXiv: math.NT/0511188]; " Fractal Fits to Riemann zeros", *Canadian J. Physics* **85** (2007), 345 [arXiv: math-ph/0606005].
- [23] A. Connes, *Selecta Mathematica (New Series)* **5** (1999) 29.
- [24] G. Sierra, "The Riemann zeros as spectrum and the Riemann hypothesis" arXiv : 1601.01797.  
 G. Sierra and P. Townsend, "Landau Levels and Riemann Zeros" arXiv : 0805.4079 [ math-ph] .  
 G. Sierra,  $xp$  with interactions and the Riemann zeros, [arXiv: math-ph/0702034].  
 G. Sierra, *New Journal of Physics* **10** (2008) 033016.
- [25] A. LeClair, "Riemann Hypothesis and Random Walks: the Zeta case", arXiv : 1601.00914.  
 G. Franca and A. LeClair, "A theory for the zeros of Riemann Zeta and other L-functions" areXiv : 1407.4358.
- [26] A. Karatsuba and M. Korolev, *Russian Math. Surveys* **60**, no. 3 (2005) 433.
- [27] N. Katz and P. Sarnka,, "Random Matrices, Frobenius Eigenvalues and monodromy" *Amer. Math. Soc. Colq, Publ.* **45**, AMS, Providence, RI. 1999. M. Mehta, *Random Matrices*, second edition, Acad Press, Boston, 1991.
- [28] A. Odlyzko, <http://www.dtc.umn.edu/odlyzko>.
- [29] R. Aschhheim, C. Castro and K. Irwin, "The search for a Hamiltonian whose Energy Spectrum coincides with the Riemann Zeta Zeros" to appear in the IJGMMP 2017.