

New Principles of Differential Equations I

Hong Lai Zhu *

School of Physics and Electronic Information, Huaibei Normal University, Anhui 235000, China

Abstract

This is the first part of the total paper. Since the theory of partial differential equations (PDEs) has been established nearly 300 years, there are many important problems have not been resolved, such as what are the general solutions of Laplace equation, acoustic wave equation, Helmholtz equation, heat conduction equation, Schrodinger equation and other important equations? How to solve the problems of definite solutions which have universal significance for these equations? What are the laws of general solution of the m th-order linear PDEs with n variables ($n, m \geq 2$)? Is there any general rule for the solution of a PDE in arbitrary orthogonal coordinate systems? Can we obtain the general solution of vector PDEs? Are there very simple methods to quickly and efficiently solve the exact solutions of nonlinear PDEs? And even general solution? Etc. These problems are all effectively solved in this paper. Substituting the results into the original equations, we have verified that they are all correct.

keywords: concise general solution; series general solution; exact solutions; transformational equations; problems of definite solutions.

Abbreviations: concise general solution (CGS); series general solution (SGS); independent variable transformational equations (IVTEs), dependent variable transformational equations (DVTEs); symmetric vector partial differential equations (SVPDEs); corresponding scalar equation (CSE); independent variable transformation vector equation (IVTVE).

Introduction	2
1. General solutions and exact solutions of Cauchy problem of mathematical physics equations	3
1.1. New principles and methods 1	3
1.2. 1D wave equation	8
1.3. 2D wave equation	11
1.4. Acoustic wave equation	13
1.5. Laplace equation	12
1.6. New principles and methods 2	13
1.7. Poisson equation	15
1.8. Helmholtz equation	16
1.9. Heat conduction equation and diffusion equation	20
1.10. Schrödinger equation	24
1.11. Singular general solution of the Helmholtz equation	29
2. Principles of transformational equations	32
2.1. The principle of independent variable transformational equations	32
2.2. The principle of dependent variable transformational equations	34

*E-mail address: honglaizhu@gmail.com, honglaizhu@qq.com

2.3. General Solutions of Vector Equation in Various Orthogonal Coordinate Systems	36
3. General solutions laws of linear partial differential equations	42
3.1. General solutions laws of linear partial differential equations with variable coefficients	43
3.2. General solutions laws of linear partial differential equations with constant coefficients	49
4. General solutions and particular solutions of nonlinear partial differential equations	51
5. Extention and conclusion	56
5.1. Two axioms and a conjecture	56
5.2. General Equations and Restricted Equations	58
5.3. Conclusion	59
Appendix	63
Appendix A	63
Appendix B	63
Appendix C	65
References	67

1 Introduction

In the establishment period of PDE theory, because of the general solution of the one-dimensional wave equation solved by d'Alembert, the mathematicians at that time believed that the general solutions of PDEs existed universally. Since there was no substantive breakthrough, with Cauchy's advice, they had to turn their attention to a variety of problems of definite solutions.

Using the new defines, laws and methods presented in this paper, the general solutions of many important PDEs had been solved for the first time, such as the Laplace equation, the wave equation, the Helmholtz equation, heat equation and so on, the exact solutions of relevant Cauchy problems have been solved too. In some cases, the general solutions and the exact solutions of the Cauchy problem for the Poisson equation and the Schrödinger equation have been solved. The types and numbers of the PDEs whose general solution could be solved by the new analytic method system are far more than the sum of the other methods can solve, and the solving process is very clear and concise.

The new theory also further reveals two major flaws and errors in the existing theory:

1. The new theory found that the general solutions of many PDEs have two forms: concise general solutions and series general solutions, in theory, infinite series solutions of a PDE can be obtained by its series general solution. The general solution of the one-dimensional wave equation introduced by mainstream textbooks and professional books is not a general solution in fact. It is only a special case of real general solution which can deduce the Fourier series solution. According to the obtained general solutions, we find why the state and the change of many natural phenomena that can be described by PDEs are both infinite. In this paper we find the relationship between the general solution and the series solution of PDEs, and point out that the alleged general solution of 1D wave equation obtained in current textbooks and professional books is not the general solution in fact. We find that any series solutions of PDEs can be obtained by its series general solution theoretically and show the root cause why the states and changes of some natural phenomena described by PDEs are all infinity.

2. In the theory of PDEs, almost all of the textbooks and professional books directly or indirectly declare that the number of the arbitrary functions in the general solution of m th-order PDEs is m , but no related rigorous proof up to now. In this paper we find a singularity of general solutions of Helmholtz equation the first time, namely the number of arbitrary functions in the general solutions is more than 2.

In Chapter 2 of this paper we present three types transformational equations: independent variable transformational equations (IVTEs), dependent variable transformational equations (DVTEs), independent variable transformation vector equation (IVTVE), and get a law of partial differential equations solution in various orthogonal coordinate system. The general solution of vector wave equation in Cartesian, cylindrical and spherical coordinate systems have been solved for the first time. We point out that the general solutions or particular solutions of various symmetric vector partial differential equations can be obtained similarly in any orthogonal coordinate system, such as vector Helmholtz equation, the magnetic vector potential equation and so on.

The laws of the general solution of m th-order linear partial differential equations with n variables have been studied deeply in Chapter 3 ($n, m \geq 2$). We have solved some typical nonlinear partial differential equations general solutions, particular solutions or solitary wave solutions in Chapter 4 and other relevant chapter, such as Emden-Fowler equation, Klein-Gordon equation, sine-Gordon equation, Burgers equation, KdV equation, etc. Based on the large number of results obtained in this book, we find that the general solutions of some PDEs have similarities, and find the roots of these similarities by the concepts of general equations and restricted equations.

1. General solutions and exact solutions of Cauchy problem of mathematical physics equations

In recent decades, for solving partial differential equations (PDEs) many analytic methods [1-3] and numerical methods [4-6] have been developed rapidly, the solitary wave solutions [7-9] plays an important role in nonlinear PDEs (NLPDEs), the existence [10, 11], uniqueness [12, 13], and stability [14, 15] of the PDEs solution have been well studied. The formulas of differential equations general solution have the same important value and significance as the algebraic equations Root Formulas, although the exact or numerical solutions of many PDEs have been found, species of PDEs which have the general solution are extremely rare yet.

Since mathematical physics equations (MPEs) are very important in PDEs, their progress is always been noticed especially [16, 17]. In the professional books of MPEs only one dimensional wave equation [18, 19] and some linear PDEs (LPDEs) with two variables [20] have general solution. Using the new analytic methods proposed in this chapter, the general solutions of many important PDEs had been solved for the first time, the exact solutions of relevant problems of definite solutions have been solved too.

1.1. New principles and methods 1

We first study the laws of one multivariate function is a composite function of another. In \mathbb{R}^n space ($n \geq 2$), assuming $u(x_1, \dots, x_n), v(x_1, \dots, x_n)$ are both smooth functions, and u is a composite function of v

$$u(x_1, \dots, x_n) = f(v), (f'_v \neq 0), \quad (1)$$

where f is an unary smooth function, according to the laws of differential and total differential

$$\begin{aligned} du &= f'_v dv = f'_v v_{x_1} dx_1 + f'_v v_{x_2} dx_2 + \dots + f'_v v_{x_n} dx_n \\ &= u_{x_1} dx_1 + u_{x_2} dx_2 + \dots + u_{x_n} dx_n. \end{aligned}$$

So

$$u_{x_i} = f'_v v_{x_i}. \quad (2)$$

By (2), we obtain further

$$u_{x_i x_i} = f''_v v_{x_i x_i} + f''_v v_{x_i}^2, u_{x_i x_j} = f'_v v_{x_i x_j} + f''_v v_{x_i} v_{x_j}. \quad (3)$$

Higher order laws may be deduced analogously. We set

$$v(x_1, \dots, x_n) = k_1x_1 + k_2x_2 + \dots + k_nx_n + k_{n+1}, \quad (4)$$

where k_i are all arbitrary constants ($i = 1, 2, \dots, n + 1$), then

$$v_{x_i} = k_i, v_{x_ix_i} = v_{x_ix_j} = 0, (i \neq j), (i, j \in \{1, 2, \dots, n\}). \quad (5)$$

By (3) and under the condition of (4), we have

$$u_{x_i} = k_i f'_v, u_{x_ix_i} = k_i^2 f''_v, u_{x_ix_j} = k_i k_j f''_v \quad (6)$$

Using mathematical induction we can get

$$u_{x_i}^{(m)} = k_i^m f_v^{(m)}, u_{x_ix_j}^{(pq)} = k_i^p k_j^q f_v^{(p+q)}, (0 \leq m < \infty, 0 \leq p, q < \infty), \quad (7)$$

$$u_{x_1x_2 \dots x_n}^{(i_1 i_2 \dots i_n)} = k_1^{i_1} k_2^{i_2} \dots k_n^{i_n} f_v^{(m)}, (i_1 + i_2 + \dots + i_n = m), \quad (8)$$

where

$$\begin{aligned} f_v^{(m)} &\equiv \frac{d^m f}{dv^m}, f_v^{(p+q)} \equiv \frac{d^{p+q} f}{dv^{p+q}}, u_{x_i}^{(m)} \equiv \frac{\partial^m u}{\partial x_i^m}, \\ u_{x_ix_j}^{(pq)} &\equiv \frac{\partial^{p+q} u}{\partial x_i^p \partial x_j^q}, u_{x_1x_2 \dots x_n}^{(i_1 i_2 \dots i_n)} \equiv \frac{\partial^m u}{\partial x_1^{i_1} \partial x_2^{i_2} \dots \partial x_n^{i_n}}. \end{aligned} \quad (9)$$

By the above laws, we present a new transformational method to solve the general solutions or exact solutions of some PDEs.

Transformational Method 1. In the domain D , ($D \subset \mathbb{R}^n$), any established m th-order PDE with n space variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$, set $v = v(x_1, \dots, x_n)$ and $u = f(v)$ are both undetermined m th-differentiable functions ($u, v \in C^m(D)$), then substitute $u = f(v)$ and its partial derivatives into $F = 0$

1. In case of working out $v(x_1, \dots, x_n)$ and $f(v)$, then $u = f(v(x_1, \dots, x_n))$ is the solution of $F = 0$,
2. In case of dividing out $f(v)$ and its partial derivative, also working out $v(x_1, \dots, x_n)$, then $u = f(v(x_1, \dots, x_n))$ is the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function,
3. In case of dividing out $f(v)$ and its partial derivative, also getting $k = 0$, but in fact $k \neq 0$, then $u = f(v(x_1, \dots, x_n))$ is not the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function.

In Transformational Method 1 $v = v(x_1, \dots, x_n)$ may be an unknown function completely or has a determinate form with unknown parameters, the solution of f may be an arbitrary or a certain unary m th-differentiable function, the solution of v and f may not be single and so on. Here we use Transformational Method 1 to solve three important PDEs.

Example 1.1.

$$u_{xx} + u_{yy} + u_{zz} = 0. \quad (10)$$

Eq. (10) is Laplace equation in Cartesian coordinate system. According to Transformational Method 1, set

$$u(x, y, z) = f(v) = f(k_1x + k_2y + k_3z + k_4), \quad (11)$$

where $v = k_1x + k_2y + k_3z + k_4$, $k_1 - k_4$ are unascertained parameters and f is an undetermined 2th-differentiable function, by (7)

$$u_{xx} + u_{yy} + u_{zz} = (k_1^2 + k_2^2 + k_3^2)f_v^{(2)} = 0.$$

The first case is

$$f_v^{(2)} = 0, \quad (12)$$

according to Transformational Method 1 the solution of Eq. (10) is

$$u = f(v) = k_1x + k_2y + k_3z + k_4, \quad (13)$$

where $k_1 - k_4$ are all arbitrary constants. The second case is

$$k_1^2 + k_2^2 + k_3^2 = 0 \Rightarrow k_1 = \pm\sqrt{-k_2^2 - k_3^2}, \quad (14)$$

where k_2 and k_3 are all arbitrary constants. By Transformational Method 1 the solution of Eq. (10) is

$$u = f_1(x\sqrt{-k_2^2 - k_3^2} + k_2y + k_3z + k_4) + f_2(-x\sqrt{-k_2^2 - k_3^2} + k_2y + k_3z + k_4) \quad (15)$$

where k_2, k_3 and k_4 are all arbitrary constants. Note the k_2, k_3 and k_4 in $f_1(x\sqrt{-k_2^2 - k_3^2} + k_2y + k_3z + k_4)$ may be different with them in $f_2(-x\sqrt{-k_2^2 - k_3^2} + k_2y + k_3z + k_4)$, the $k_1 - k_4$ in Eq. (13) may be different with them in Eq. (15), and Eq. (10) is a linear equation, in order to facilitate writing, the general solution of Laplace equation may be written as

$$u = f_1(x\sqrt{-k_2^2 - k_3^2} + k_1y + k_2z + k_3) + f_2(-x\sqrt{-k_4^2 - k_5^2} + k_4y + k_5z + k_6) + k_7x + k_8y + k_9z + k_{10}, \quad (16)$$

where f_1 and f_2 are arbitrary unary second differentiable functions, $k_1 - k_{10}$ are arbitrary parameters.

Since $k_1 - k_{10}$ are arbitrary constants and Eq. (10) is a linear equation, (16) can also be written as the form of a function series

$$u = \sum_{i=1}^s (f_{1_i}(x\sqrt{-k_{1_i}^2 - k_{2_i}^2} + k_{1_i}y + k_{2_i}z + k_{3_i}) + f_{2_i}(-x\sqrt{-k_{4_i}^2 - k_{5_i}^2} + k_{4_i}y + k_{5_i}z + k_{6_i}) + k_{7_i}x + k_{8_i}y + k_{9_i}z + k_{10_i}), \quad (17)$$

where f_{1_i} and f_{2_i} are arbitrary unary second differentiable functions, $(1 \leq s \leq \infty)$, $k_{1_i} - k_{10_i}$ are arbitrary determined parameters.

We call (16) the **concise general solution** (CGS) which has the most simple form and call (17) the **series general solution** (SGS) which could have infinite arbitrary functions in it. Theoretically all specific series solutions of Eq. (10) can be obtained by SGS (17).

Example 1.2.

$$(u_x^{(m)})^r + \frac{a(x,y,z)(k_1y+k_2z+k_4)^{mr}}{(k_1x+k_3z+k_5)^{mr}}(u_y^{(m)})^r - \frac{(1+a(x,y,z))(k_1y+k_2z+k_4)^{mr}}{(k_2x+k_3y+k_6)^{mr}}(u_z^{(m)})^r = 0, \quad (18)$$

where $k_1 - k_6$ are arbitrary known constant, $a(x, y, z)$ is an arbitrary known function with 3 variables, by Transformational Method 1, set

$$u(x, y, z) = f(v) = f(k_1xy + k_2xz + k_3yz + k_4x + k_5y + k_6z + k_7), \quad (19)$$

where $v = k_1xy + k_2xz + k_3yz + k_4x + k_5y + k_6z + k_7$, k_7 is an undetermined parameter and f is an unary m th-differentiable function to be determined, then

$$u_x = f'_v v_x = (k_1y + k_2z + k_4)f'_v, \quad (20)$$

$$u_y = f'_v v_y = (k_1 x + k_3 z + k_5) f'_v, \quad (21)$$

$$u_z = f'_v v_z = (k_2 x + k_3 y + k_6) f'_v. \quad (22)$$

According to (20)-(22) and mathematical induction we get

$$u_x^{(m)} = (k_1 y + k_2 z + k_4)^m f_v^{(m)}, \quad (23)$$

$$u_y^{(m)} = (k_1 x + k_3 z + k_5)^m f_v^{(m)}, \quad (24)$$

$$u_z^{(m)} = (k_2 x + k_3 y + k_6)^m f_v^{(m)}. \quad (25)$$

Then

$$\begin{aligned} & (u_x^{(m)})^r + \frac{a(x,y,z)(k_1 y + k_2 z + k_4)^{mr}}{(k_1 x + k_3 z + k_5)^{mr}} (u_y^{(m)})^r - \frac{(1+a(x,y,z))(k_1 y + k_2 z + k_4)^{mr}}{(k_2 x + k_3 y + k_6)^{mr}} (u_z^{(m)})^r \\ &= (k_1 y + k_2 z + k_4)^{mr} (f_v^{(m)})^r + \frac{a(x,y,z)(k_1 y + k_2 z + k_4)^{mr}}{(k_1 x + k_3 z + k_5)^{mr}} (k_1 x + k_3 z + k_5)^{mr} (f_v^{(m)})^r \\ & - \frac{(1+a(x,y,z))(k_1 y + k_2 z + k_4)^{mr}}{(k_2 x + k_3 y + k_6)^{mr}} (k_2 x + k_3 y + k_6)^{mr} (f_v^{(m)})^r = 0 \\ & \Rightarrow (k_1 y + k_2 z + k_4)^{mr} (f_v^{(m)})^r - (k_1 y + k_2 z + k_4)^{mr} (f_v^{(m)})^r = 0. \end{aligned}$$

According to Transformational Method 1, the solution of Eq. (18) is

$$u = f(k_1 x y + k_2 x z + k_3 y z + k_4 x + k_5 y + k_6 z + k_7), \quad (26)$$

where $f(v)$ is an arbitrary unary m th-differentiable function and k_7 is an arbitrary constants. If $m = 1$, (26) is the general solution of Eq. (18).

Example 1.3.

$$a_1 (u_{x_1}^{(m)})^r + a_2 (u_{x_2}^{(m)})^r + \dots + a_n (u_{x_n}^{(m)})^r + a_{n+1} (u_{x_2 x_3}^{(pq)})^r = 0, \quad (27)$$

where a_i , ($i = 1, 2, \dots, n+1$) are arbitrary known constants, $r \geq 1$, $1 \leq p+q = m$, the left of Eq. (27) could be added any number and types of $(u_{x_1 i_2 \dots i_n}^{(i_1 i_2 \dots i_n)})^r$, ($i_1 + i_2 + \dots + i_n = m$) with any constant coefficient, since the similar calculation method, for facilitating writing there is only the $a_{n+1} (u_{x_2 x_3}^{(pq)})^r$ in Eq. (27).

By Transformational Method 1, set $u(x_1, \dots, x_n) = f(v)$, $v(x_1, \dots, x_n) = k_1 x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1}$, where k_1, k_2, \dots, k_{n+1} are unascertained parameters and f is an undetermined unary m th-differentiable function, by (7)

$$\begin{aligned} & a_1 (u_{x_1}^{(m)})^r + a_2 (u_{x_2}^{(m)})^r + \dots + a_n (u_{x_n}^{(m)})^r + a_{n+1} (u_{x_2 x_3}^{(pq)})^r \\ &= (a_1 k_1^{mr} + a_2 k_2^{mr} + \dots + a_n k_n^{mr} + a_{n+1} k_2^{pr} k_3^{qr}) (f_v^{(m)})^r = 0. \end{aligned}$$

The first case is

$$(f_v^{(m)})^r = 0, \quad (28)$$

according to Transformational Method 1 the solution of Eq. (27) is

$$u = f(v) = c_{m-1} v^{m-1} + c_{m-2} v^{m-2} + \dots + c_1 v, \quad (29)$$

where $v(x_1, \dots, x_n) = k_1 x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1}$, $k_1 - k_{n+1}$ and $c_1 - c_{m-1}$ are all arbitrary constants.

Since v contains arbitrary constants k_{n+1} , so there is no arbitrary constants c_0 in (29).

The second case is

$$a_1 k_1^{mr} + a_2 k_2^{mr} + \dots + a_n k_n^{mr} + a_{n+1} k_2^{pr} k_3^{qr} = 0, \quad (30)$$

if m, r are both odd, then

$$k_1 = \left(-\frac{a_2 k_2^{mr} + a_3 k_3^{mr} + \dots + a_n k_n^{mr} + a_{n+1} k_2^{pr} k_3^{qr}}{a_1} \right)^{\frac{1}{mr}}, \quad (31)$$

where $k_2 - k_{n+1}$ are all arbitrary constants. By Transformational Method 1 the solution of Eq. (27) is

$$u = f\left(\left(-\frac{a_2 k_2^{mr} + \dots + a_n k_n^{mr} + a_{n+1} k_2^{pr} k_3^{qr}}{a_1} \right)^{\frac{1}{mr}} x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1} \right), \quad (32)$$

where f is an arbitrary unary m th-differentiable function.

If there is at least one even number among m and r in Eq. (27), then

$$k_1 = \pm \left(-\frac{a_2 k_2^{mr} + a_3 k_3^{mr} + \dots + a_n k_n^{mr} + a_{n+1} k_2^{pr} k_3^{qr}}{a_1} \right)^{\frac{1}{mr}}. \quad (33)$$

By Transformational Method 1, except (29) and (32) another solution of Eq. (27) is

$$u = f\left(-\left(-\frac{a_2 k_2^{mr} + \dots + a_n k_n^{mr} + a_{n+1} k_2^{pr} k_3^{qr}}{a_1} \right)^{\frac{1}{mr}} x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1} \right). \quad (34)$$

In the case of $r = 1$, Eq. (27) becomes linear equation

$$a_1 u_{x_1}^{(m)} + a_2 u_{x_2}^{(m)} + \dots + a_n u_{x_n}^{(m)} + a_{n+1} u_{x_2 x_3}^{(pq)} = 0. \quad (35)$$

If m is odd, by (29) and (32) the solution of Eq. (35) is

$$u = f\left(\left(-\frac{a_2 k_2^m + \dots + a_n k_n^m + a_{n+1} k_2^p k_3^q}{a_1} \right)^{\frac{1}{m}} x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1} \right) + c_{m-1} v^{m-1} + c_{m-2} v^{m-2} + \dots + c_1 v, \quad (36)$$

where $v = C_1 x_1 + C_2 x_2 + \dots + C_n x_n + C_{n+1}$, $C_1 - C_{n+1}$ are arbitrary constants. If m is even, by (29), (32) and (34) the solution of Eq. (35) is

$$u = f_1\left(\left(-\frac{a_2 k_2^m + \dots + a_n k_n^m + a_{n+1} k_2^p k_3^q}{a_1} \right)^{\frac{1}{m}} x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1} \right) + f_2\left(-\left(-\frac{a_2 l_2^m + \dots + a_n l_n^m + a_{n+1} l_2^p l_3^q}{a_1} \right)^{\frac{1}{m}} x_1 + l_2 x_2 + \dots + l_n x_n + l_{n+1} \right) + c_{m-1} v^{m-1} + c_{m-2} v^{m-2} + \dots + c_1 v \quad (37)$$

where f_1 and f_2 are arbitrary unary m th-differentiable functions, $k_2 - k_{n+1}$ and $l_2 - l_{n+1}$ are arbitrary parameters. In Appendix A we proved that if $k_1, l_1 \neq 0$ and $k_1, l_1 \rightarrow 0$ in (37), $c_1 v$ can be described by f_1 and f_2 . If $m = 2, r = p = q = 1$, Eq. (27) becomes

$$a_1 u_{x_1}^{(2)} + a_2 u_{x_2}^{(2)} + \dots + a_n u_{x_n}^{(2)} + a_{n+1} u_{x_2 x_3} = 0. \quad (38)$$

According to (37) and (16), the CGS and SGS of Eq. (38) can be get respectively

$$u = f_1\left(\left(-\frac{a_2 k_2^2 + \dots + a_n k_n^2 + a_{n+1} k_2 k_3}{a_1} \right)^{\frac{1}{2}} x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1} \right) + f_2\left(-\left(-\frac{a_2 l_2^2 + \dots + a_n l_n^2 + a_{n+1} l_2 l_3}{a_1} \right)^{\frac{1}{2}} x_1 + l_2 x_2 + \dots + l_n x_n + l_{n+1} \right) + c_1 v. \quad (39)$$

$$\begin{aligned}
u = & \sum_{i=1}^s (f_{1_i} \left(\left(-\frac{a_2 k_{i_2}^2 + \dots + a_n k_{i_n}^2 + a_{n+1} k_{i_2} k_{i_3}}{a_1} \right)^{\frac{1}{2}} x_1 + k_{i_2} x_2 + \dots + k_{i_n} x_n + k_{i_{n+1}} \right) \\
& + f_{2_i} \left(-\left(-\frac{a_2 l_{i_2}^2 + \dots + a_n l_{i_n}^2 + a_{n+1} l_{i_2} l_{i_3}}{a_1} \right)^{\frac{1}{2}} x_1 + l_{i_2} x_2 + \dots + l_{i_n} x_n + l_{i_{n+1}} \right) + c_1 v
\end{aligned} \tag{40}$$

Consider the following Cauchy problem of Eq. (38)

$$u(0, x_2, \dots, x_n) = \sum_{i=1}^s \varphi_i (k_{i_2} x_2 + k_{i_3} x_3 + \dots + k_{i_n} x_n + k_{i_{n+1}}), \tag{41}$$

$$u_{x_1}(0, x_2, \dots, x_n) = \sum_{i=1}^s \psi_i (k_{i_2} x_2 + k_{i_3} x_3 + \dots + k_{i_n} x_n + k_{i_{n+1}}), \tag{42}$$

where $1 \leq s \leq \infty$, x_1 sometimes equal to time t . In (40), set $c_1 = 0$, $k_{i_j} = l_{i_j}$, ($i = 1, 2, \dots, s, j = 2, 3, \dots, n+1$), by further calculation which is in Appendix B, the exact solution of Eq. (38) in the conditions of (41) and (42) is

$$\begin{aligned}
u = & \frac{1}{2} \sum_{i=1}^s (\varphi_i (k_{i_1} x_1 + k_{i_2} x_2 + \dots + k_{i_n} x_n + k_{i_{n+1}}) \\
& + \varphi_i (-k_{i_1} x_1 + k_{i_2} x_2 + \dots + k_{i_n} x_n + k_{i_{n+1}}) \\
& + \frac{1}{k_{i_1}} \int_{-k_{i_1} x_1 + k_{i_2} x_2 + \dots + k_{i_n} x_n + k_{i_{n+1}}}^{k_{i_1} x_1 + k_{i_2} x_2 + \dots + k_{i_n} x_n + k_{i_{n+1}}} \psi(\xi_i) d\xi_i)
\end{aligned} \tag{43}$$

where

$$k_{i_1} = \left(-\left(a_2 k_{i_2}^2 + \dots + a_n k_{i_n}^2 + a_{n+1} k_{i_2} k_{i_3} \right) / a_1 \right)^{\frac{1}{2}}. \tag{44}$$

According to the above three typical cases, we know that the solutions of some linear or nonlinear PDEs can be obtained by using Transformational Method 1.

1.2. 1D wave equation

1D wave equation

$$u_{tt} - a^2 u_{xx} = 0, \tag{45}$$

is the first PDE studied deeply. Almost all current textbooks and professional books have pointed out that the general solution of Eq. (45) is

$$u = f_1(x + at) + f_2(x - at). \tag{46}$$

Fourier series solution of Eq. (45) is

$$u = \sum_{i=1}^s \left(A_n \cos\left(\frac{n\pi at}{l}\right) + B_n \sin\left(\frac{n\pi at}{l}\right) \sin\left(\frac{n\pi x}{l}\right) \right). \tag{47}$$

By (46) we cannot get (47) obviously, there is no answer why the particular solution (47) cannot be got by the general solution (46).

Eq. (45) is a special case of Eq. (38), according to (39) and (40), its CGS and SGS can be get respectively

$$u = f_1(k_1 x + k_1 at + k_2) + f_2(k_3 x - k_3 at + k_4) + k_5 x + k_6 t + k_7, \tag{48}$$

$$\begin{aligned}
u = & \sum_{i=1}^s (f_{1_i}(k_{1_i} x + k_{1_i} at + k_{2_i}) + f_{2_i}(k_{3_i} x - k_{3_i} at + k_{4_i})) \\
& + k_5 x + k_6 t + k_7,
\end{aligned} \tag{49}$$

where f_1, f_{1_i}, f_2 and f_{2_i} are arbitrary unary second differentiable functions, $k_1 - k_7$ are arbitrary parameters, $k_{1_i} - k_{4_i}$ are arbitrary determined parameters, ($1 \leq s \leq \infty$). Of course, the general solution of Eq. (45) can also be written as

$$u = f_1(k_1x + k_1at + k_2) + \sum_{i=1}^s f_{2_i}(k_{3_i}x - k_{3_i}at + k_{4_i}) + k_5x + k_6t + k_7 \quad (50)$$

and so on, (50) is also a SGS, but in this paper we will not discuss the general solutions in special forms.

By the above results we can see that (46) is a special case of (48) and (49), and is not a general solution of Eq. (45) in fact, so using (46) we cannot get the Fourier series solution.

Theoretically every specific series solution of Eq. (45) can be obtained by the SGS (49), as a case, we will obtain the Fourier series solution (47).

Set

$$f_{1_n}(k_{1_n}x + k_{1_n}at + k_{2_n}) = C_n \sin(k_{1_n}x + k_{1_n}at + k_{2_n}), \quad (51)$$

$$f_{2_n}(k_{3_n}x - k_{3_n}at + k_{4_n}) = D_n \cos(k_{3_n}x - k_{3_n}at + k_{4_n}), \quad (52)$$

So

$$\begin{aligned} u &= \sum_{i=1}^s (f_{1_n}(k_{1_n}x + k_{1_n}at + k_{2_n}) + f_{2_n}(k_{3_n}x - k_{3_n}at + k_{4_n})) \\ &= \sum_{i=1}^s (C_n \sin(k_{1_n}x) \cos(k_{1_n}at) \cos k_{2_n} + \cos(k_{1_n}x) \sin(k_{1_n}at) \cos k_{2_n} \\ &\quad + \cos(k_{1_n}x) \cos(k_{1_n}at) \sin k_{2_n} - \sin(k_{1_n}x) \sin(k_{1_n}at) \sin k_{2_n}) \\ &\quad + D_n (\cos(k_{3_n}x) \cos(k_{3_n}at) \cos k_{4_n} + \sin(k_{3_n}x) \sin(k_{3_n}at) \cos k_{4_n} \\ &\quad - \sin(k_{3_n}x) \cos(k_{3_n}at) \sin k_{4_n} + \cos(k_{3_n}x) \sin(k_{3_n}at) \sin k_{4_n}). \end{aligned}$$

Set $k_{1_n} = k_{3_n} = k_n$, then

$$\begin{aligned} u &= \sum_{i=1}^s ((C_n \cos k_{2_n} - D_n \sin k_{4_n}) \sin(k_nx) \cos(k_nat) \\ &\quad + (C_n \cos k_{2_n} + D_n \sin k_{4_n}) \cos(k_nx) \sin(k_nat) \\ &\quad + (C_n \sin k_{2_n} + D_n \cos k_{4_n}) \cos(k_nx) \cos(k_nat) \\ &\quad + (-C_n \sin k_{2_n} + D_n \cos k_{4_n}) \sin(k_nx) \sin(k_nat)) \end{aligned} \quad (53)$$

Set

$$C_n \cos k_{2_n} + D_n \sin k_{4_n} = C_n \sin k_{2_n} + D_n \cos k_{4_n} = 0.$$

We have

$$k_{4_n} = \frac{(2m+1)\pi}{2} - k_{2_n}.$$

Set $k_{4_n} = \pi/2 - k_{2_n}$, we can get $C_n = -D_n$. Substituting the above results into (53)

$$\begin{aligned} u &= \sum_{i=1}^s ((C_n \cos k_{2_n} - D_n \sin k_{4_n}) \sin(k_nx) \cos(k_nat) \\ &\quad + (-C_n \sin k_{2_n} + D_n \cos k_{4_n}) \sin(k_nx) \sin(k_nat)) \\ &= \sum_{i=1}^s (2C_n \cos k_{2_n} \sin(k_nx) \cos(k_nat) - 2C_n \sin k_{2_n} \sin(k_nx) \sin(k_nat)). \end{aligned}$$

Namely

$$u = \sum_{i=1}^s 2C_n (\cos k_{2_n} \cos(k_nat) - \sin k_{2_n} \sin(k_nat)) \sin(k_nx). \quad (54)$$

Since C_n, k_n and k_{2n} are all arbitrary parameters, set

$$k_n = \frac{n\pi}{l}, 2C_n \cos k_{2n} = A_n, -2C_n \sin k_{2n} = B_n. \quad (55)$$

Then (54) may be translated into (47). (46) was first discovered by d' Alembert, then Daniel Bernoulli discovered an infinite series solution

$$u = \sum_{i=1}^{\infty} a_n \sin\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi at}{l}\right). \quad (56)$$

The relationship between (46) and (56) led to a well-known controversy in the history of mathematics [21], many famous mathematicians have been involved in this drastic and lengthy debate, even after discovering the Fourier series solution (47), the relationship between (46) and (47) is still unclear, now the problem is finally solved successfully, (46) and (47) are two different closed solutions, not the general solution, only the form of (46) is relatively close the general solution.

1.3. 2D wave equation

The form of 2D wave equation in Cartesian coordinate system is

$$u_{tt} - a^2 u_{xx} - a^2 u_{yy} = 0. \quad (57)$$

Eq. (57) is an especial case of Eq. (38), by (39) its CGS can be obtained

$$\begin{aligned} u &= f_1 \left(k_1 x + k_2 y + at \sqrt{k_1^2 + k_2^2} + k_3 \right) \\ &+ f_2 \left(k_4 x + k_5 y - at \sqrt{k_4^2 + k_5^2} + k_6 \right) + k_7 x + k_8 y + k_9 t + k_{10} \\ &= g \left(\frac{k_1 x}{\sqrt{k_1^2 + k_2^2}} + \frac{k_2 y}{\sqrt{k_1^2 + k_2^2}} + at + g_0 \right) \\ &+ h \left(\frac{k_4 x}{\sqrt{k_4^2 + k_5^2}} + \frac{k_5 y}{\sqrt{k_4^2 + k_5^2}} - at + h_0 \right) + k_7 x + k_8 y + k_9 t + k_{10} \\ &= g(x \cos \theta + y \sin \theta + at + g_0) \\ &+ h(x \cos \varphi + y \sin \varphi - at + h_0) + k_7 x + k_8 y + k_9 t + k_{10}, \end{aligned} \quad (58)$$

where f_1, f_2, g and h are arbitrary unary second differentiable functions, $k_1 - k_{10}, \theta, \varphi, g_0$ and h_0 are arbitrary parameters. $g(x \cos \theta + y \sin \theta + at + g_0)$ is a parallel wave with the speed a , the angle between x axis and spread direction of g is θ which is arbitrary. The SGS of Eq. (57) is

$$u(x, y, t) = \sum_i (g_i(x \cos \theta_i + y \sin \theta_i + at + g_{i0}) + h_i(x \cos \varphi_i + y \sin \varphi_i - at + h_{i0})) + k_7 x + k_8 y + k_9 t + k_{10}, \quad (59)$$

where g_i and h_i are arbitrary unary second differentiable functions, $\theta_i, \varphi_i, g_{i0}$ and h_{i0} are arbitrary determined parameters.

A research hotspot is using numerical methods to study the 2D wave equation [21]. Consider the following initial value problem of Eq. (57)

$$\begin{aligned} u(x, y, 0) &= \sum_i \varphi_i(k_{i1}x + k_{i2}y + k_{i3}), \\ u_t(x, y, 0) &= \sum_i \psi_i(k_{i1}x + k_{i2}y + k_{i3}). \end{aligned} \quad (60)$$

Similar to the solving method of (43), the exact solution of Eq. (57) on the conditions of (60) can be got

$$\begin{aligned}
u = & \frac{1}{2} \sum_i (\varphi_i(k_{i_1}x + k_{i_2}y - at\sqrt{k_{i_1}^2 + k_{i_2}^2} + k_{i_3}) \\
& + \varphi_i(k_{i_1}x + k_{i_2}y + at\sqrt{k_{i_1}^2 + k_{i_2}^2} + k_{i_3})) \\
& + \frac{1}{a\sqrt{k_{i_1}^2 + k_{i_2}^2}} \int_{k_{i_1}x + k_{i_2}y - at\sqrt{k_{i_1}^2 + k_{i_2}^2} + k_{i_3}}^{k_{i_1}x + k_{i_2}y + at\sqrt{k_{i_1}^2 + k_{i_2}^2} + k_{i_3}} \psi(\xi_i) d\xi_i.
\end{aligned} \tag{61}$$

1.4. Acoustic wave equation

The form of acoustic wave equation is

$$p_{tt} - c_0^2 p = 0 \tag{62}$$

where Δ is the Laplace operator, p is the sound pressure, and c_0 is the sound speed. Eq. (62) is a special case of Eq. (38), according to (39) its CGS in Cartesian coordinate system is

$$\begin{aligned}
p = & f_1(k_1x + k_2y + k_3z + c_0t\sqrt{k_1^2 + k_2^2 + k_3^2} + k_4) \\
& + f_2(k_5x + k_6y + k_7z - c_0t\sqrt{k_5^2 + k_6^2 + k_7^2} + k_8) + k_9x + k_{10}y + k_{11}z + k_{12}t + k_{13} \\
= & g\left(\frac{k_1x}{\sqrt{k_1^2 + k_2^2 + k_3^2}} + \frac{k_2y}{\sqrt{k_1^2 + k_2^2 + k_3^2}} + \frac{k_3z}{\sqrt{k_1^2 + k_2^2 + k_3^2}} + c_0t + \frac{k_4}{\sqrt{k_1^2 + k_2^2 + k_3^2}}\right) \\
& + h\left(\frac{k_5x}{\sqrt{k_5^2 + k_6^2 + k_7^2}} + \frac{k_6y}{\sqrt{k_5^2 + k_6^2 + k_7^2}} + \frac{k_7z}{\sqrt{k_5^2 + k_6^2 + k_7^2}} - c_0t + \frac{k_8}{\sqrt{k_5^2 + k_6^2 + k_7^2}}\right) \\
& + k_9x + k_{10}y + k_{11}z + k_{12}t + k_{13} \\
= & g(x \sin \theta \cos \varphi + y \sin \theta \sin \varphi + z \cos \theta + c_0t + g_0) \\
& + h(x \sin \phi \cos \psi + y \sin \phi \sin \psi + z \cos \phi + c_0t + h_0) \\
& + k_9x + k_{10}y + k_{11}z + k_{12}t + k_{13},
\end{aligned} \tag{63}$$

where f_1, f_2, g and h are arbitrary unary second differentiable functions, $k_1 - k_{13}, \theta, \varphi, \phi, \psi, g_0$ and h_0 are arbitrary parameters.

$g(x \sin \theta \cos \varphi + y \sin \theta \sin \varphi + z \cos \theta + c_0t + g_0)$ is a parallel wave with the speed c_0 , θ is the angle between z axis and spread direction of g , φ is the angle between x axis and the projection in xy plane of spread direction of g . The SGS of Eq. (62) is

$$\begin{aligned}
p = & \sum_i g_i(x \sin \theta_i \cos \varphi_i + y \sin \theta_i \sin \varphi_i + z \cos \theta_i + c_0t + g_{i0}) \\
& + \sum_i h_i(x \sin \phi_i \cos \psi_i + y \sin \phi_i \sin \psi_i + z \cos \phi_i - c_0t + h_{i0}) \\
& + k_9x + k_{10}y + k_{11}z + k_{12}t + k_{13},
\end{aligned} \tag{64}$$

where g_i and h_i are arbitrary unary second differentiable functions, $\theta_i, \varphi_i, \phi_i, \psi_i, g_{i0}, h_{i0}$ are arbitrary determined parameters.

Consider the following initial value problem of Eq. (62)

$$p(x, y, z, 0) = \sum_i \varphi_i(k_{i_1}x + k_{i_2}y + k_{i_3}z + k_{i_4}), \tag{65}$$

$$p_t(x, y, z, 0) = \sum_i \psi_i(k_{i_1}x + k_{i_2}y + k_{i_3}z + k_{i_4}). \tag{66}$$

Similar to the solving method of (43), the exact solution of Eq. (62) on the conditions of (65) and (66) is

$$\begin{aligned}
p = & \frac{1}{2} \sum_i (\varphi_i(k_{i_1}x + k_{i_2}y + k_{i_3}z + c_0t\sqrt{k_{i_1}^2 + k_{i_2}^2 + k_{i_3}^2} + k_{i_4}) \\
& + \varphi_i(k_{i_1}x + k_{i_2}y + k_{i_3}z - c_0t\sqrt{k_{i_1}^2 + k_{i_2}^2 + k_{i_3}^2} + k_{i_4}) \\
& + \frac{1}{c_0\sqrt{k_{i_1}^2 + k_{i_2}^2 + k_{i_3}^2}} \int_{k_{i_1}x + k_{i_2}y + k_{i_3}z - c_0t\sqrt{k_{i_1}^2 + k_{i_2}^2 + k_{i_3}^2} + k_{i_4}}^{k_{i_1}x + k_{i_2}y + k_{i_3}z + c_0t\sqrt{k_{i_1}^2 + k_{i_2}^2 + k_{i_3}^2} + k_{i_4}} \psi_i(\xi) d\xi)
\end{aligned} \tag{67}$$

Nonlinear acoustic wave equation is a hot area of current research [22, 23], the solving method of nonlinear PDEs will be studied in our other papers.

1.5. Laplace equation

Laplace equation is importantly used not only in classical electrodynamics, thermodynamics and fluid dynamics etc., but also in the modern theory of the invisible [25, 26]. In recent decades a research hotspot is using many numerical methods for solving Laplace's equation in various geometries and boundary conditions, such as the moment methods [27], quasi-reversibility methods [28, 29], finite difference methods [30] and so on.

According to the previous calculation results, the CGS and SGS of $u_{xx} + u_{yy} + u_{zz} = 0$ are (16) and (17) respectively. Assuming Eq. (10) satisfies the following boundary conditions

$$\begin{aligned}
u(0, y, z) &= \sum_{i=1}^s \varphi_i(k_{i_1}y + k_{i_2}z + k_{i_3}) \\
u_x(0, y, z) &= \sum_{i=1}^s \psi_i(k_{i_1}y + k_{i_2}z + k_{i_3}).
\end{aligned} \tag{68}$$

According to (43) and (44), the exact solution of Eq. (10) on the conditions of (68) is

$$\begin{aligned}
u = & \frac{1}{2} \sum_{i=1}^s (\varphi_i(x\sqrt{-k_{i_1}^2 - k_{i_2}^2} + k_{i_1}y + k_{i_2}z + k_{i_3}) \\
& + \varphi_i(-x\sqrt{-k_{i_1}^2 - k_{i_2}^2} + k_{i_1}y + k_{i_2}z + k_{i_3}) \\
& + \frac{1}{\sqrt{-k_{i_1}^2 - k_{i_2}^2}} \int_{-x\sqrt{-k_{i_1}^2 - k_{i_2}^2} + k_{i_1}y + k_{i_2}z + k_{i_3}}^{x\sqrt{-k_{i_1}^2 - k_{i_2}^2} + k_{i_1}y + k_{i_2}z + k_{i_3}} \psi(\xi_i) d\xi_i)
\end{aligned} \tag{69}$$

1.6. Basic principles and methods 2

$v(x_1, \dots, x_n)$ and f are both undetermined in Transformational Method 1, to solve some PDEs we may be required to set f pending and $v(x_1, \dots, x_n)$ known, so put forward Transformational Method 2.

Transformational Method 2. In the domain D , ($D \subset \mathbb{R}^n$), any established m th-order PDE with n space variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$, set $v = v(x_1, \dots, x_n)$ known and $u = f(v)$ undetermined ($u, v \in C^m(D)$), then substitute $u = f(v)$ and its partial derivatives into $F = 0$

1. In case of working out f , then $u = f(v)$ is the solution of $F = 0$,
2. In case of dividing out f and its partial derivative, also getting $0 = 0$, then $u = f(v)$ is the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function,
3. In case of dividing out f and its partial derivative, also getting $k = 0$, but in fact $k \neq 0$, then $u = f(v)$ is not the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function.

We will research the application of Transformational Method 2 later in this book. Through the comparison, we can find that the traveling wave method and the solitary wave method are the concrete applications of Transformation Method 1, 2.

Now we study another important compound law of multivariate functions. In \mathbb{R}^n space ($n \geq 2$), assuming u , v and g are smooth functions, set

$$u(x_1, \dots, x_n) = g(x_1, \dots, x_n)f(v), (f'_v \neq 0), \quad (70)$$

where $v = v(x_1, \dots, x_n)$, f is an unary smooth function, then

$$\begin{aligned} du &= u_{x_1} dx_1 + u_{x_2} dx_2 + \dots + u_{x_n} dx_n = f dg + g df = f dg + g f'_v dv \\ &= \left(f g_{x_1} + g f'_v v_{x_1} \right) dx_1 + \left(f g_{x_2} + g f'_v v_{x_2} \right) dx_2 + \dots + \left(f g_{x_n} + g f'_v v_{x_n} \right) dx_n. \end{aligned}$$

So

$$u_{x_i} = f g_{x_i} + g f'_v v_{x_i}. \quad (71)$$

By (71) we could obtain

$$u_{x_i x_i} = f g_{x_i x_i} + 2g_{x_i} f'_v v_{x_i} + g f''_v v_{x_i}^2 + g f'_v v_{x_i x_i}, \quad (72)$$

$$u_{x_i x_j} = f g_{x_i x_j} + g_{x_i} f'_v v_{x_j} + g_{x_j} f'_v v_{x_i} + g f''_v v_{x_i} v_{x_j} + g f'_v v_{x_i x_j}. \quad (73)$$

Higher order law may be deduced analogously.

According to the above laws we present Transformational Method 3.

Transformational Method 3. In the domain D , ($D \subset \mathbb{R}^n$), any established m th-order PDE with n space variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$, setting $f(v), g(x_1, \dots, x_n)$ and $v(x_1, \dots, x_n)$ are all undetermined function, $g, v \in C^m(D)$, then substitute $u = gf(v)$ and its partial derivatives into $F = 0$

1. In case of working out f, g and v , then $u = gf(v)$ is the solution of $F = 0$,
2. In case of dividing out f and its partial derivative, also working out g and v , then $u = gf(v)$ is the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function,
3. In case of getting $k = 0$, but in fact $k \neq 0$, then $u = gf(v)$ is not the solution of $F = 0$.

In Transformational Method 3 $v(x_1, \dots, x_n)$ and $g(x_1, \dots, x_n)$ may be unknown completely or have definite forms with unknown parameters, the solution of f, v and g may not be single. To solve some PDEs we may be required to set $f(v), v(x_1, \dots, x_n)$ pending and $g(x_1, \dots, x_n)$ known, so put forward Transformational Method 4.

Transformational Method 4. In the domain D , ($D \subset \mathbb{R}^n$), any established m th-order PDE with n space variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$, setting $g(x_1, \dots, x_n)$ is known and $f(v), v(x_1, \dots, x_n)$ are undetermined, $g, v \in C^m(D)$, then substitute $u = gf(v)$ and its partial derivatives into $F = 0$

1. In case of working out f and v , then $u = gf(v)$ is the solution of $F = 0$,
2. In case of dividing out f and its partial derivative, also working out $v(x_1, \dots, x_n)$, then $u = gf(v)$ is the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function,

3. In case of getting $k = 0$, but in fact $k \neq 0$, then $u = gf(v)$ is not the solution of $F = 0$.

To solve some PDEs we may be required to set $g(x_1, \dots, x_n)$, $f(v)$ undetermined and $v(x_1, \dots, x_n)$ known and so on. The forms of these laws are similar to Transformational Method 3-4, we will not present here.

1.7. Poisson equation

Consider the following Poisson equation

$$\Delta u = c(x, y, z). \quad (74)$$

Supposing

$$c(x, y, z) = r(v), v(x, y, z) = k_1x + k_2y + k_3z + k_4, \quad (75)$$

$c(x, y, z)$ is known in practical problems, so $v(x, y, z) = k_1x + k_2y + k_3z + k_4$ is known too. According to Transformational Method 2, set $u = f(v)$, f is an undetermined unary function, by (7), then

$$\begin{aligned} \Delta u &= c(x, y, z) \\ \implies (k_1^2 + k_2^2 + k_3^2) f_v'' &= r(v) \\ \implies u(x, y, z) = f(v) &= \frac{\iint r(v) dv dv}{k_1^2 + k_2^2 + k_3^2} + C_1v + C_2, \end{aligned}$$

where C_1 and C_2 are arbitrary constants. So particular solution of Eq. (74) is

$$u(x, y, z) = \frac{\iint r(v) dv dv}{k_1^2 + k_2^2 + k_3^2} + C_1v + C_2. \quad (76)$$

According to the general solution of Laplace equation, the CGS of Eq. (74) may be get

$$\begin{aligned} u(x, y, z) &= f_1 \left(x\sqrt{-l_1^2 - l_2^2} + l_1y + l_2z + l_3 \right) + f_2 \left(-x\sqrt{-l_4^2 - l_5^2} + l_4y + l_5z + l_6 \right) \\ &+ \frac{\iint r(v) dv dv}{k_1^2 + k_2^2 + k_3^2} + l_7x + l_8y + l_9z + l_{10}, \end{aligned} \quad (77)$$

where f_1 and f_2 are arbitrary unary second differentiable functions, since $l_1 - l_{10}$ are arbitrary constants, (77) can be written as

$$\begin{aligned} u(x, y, z) &= \sum_{i=1}^s \left(f_{i_1} \left(x\sqrt{-l_{i_1}^2 - l_{i_2}^2} + l_{i_1}y + l_{i_2}z + l_{i_3} \right) + f_{i_2} \left(-x\sqrt{-l_{i_4}^2 - l_{i_5}^2} + l_{i_4}y + l_{i_5}z + l_{i_6} \right) \right) \\ &+ \frac{\iint r(v) dv dv}{k_1^2 + k_2^2 + k_3^2} + l_7x + l_8y + l_9z + l_{10}, \quad (1 \leq s < \infty), \end{aligned} \quad (78)$$

where f_{i_1} and f_{i_2} are arbitrary unary second differentiable functions, $l_{i_1} - l_{i_6}$ are arbitrary determined constants.

Currently, using numerical methods to analyse Poisson equation is a hot research area [31], under the condition of (75) we set

$$u(0, y, z) = p(k_2y + k_3z + k_4) + \sum_{i=1}^s \varphi_i(l_{i_1}y + l_{i_2}z + l_{i_3}), \quad (79)$$

$$u_x(0, y, z) = k_1 p'(k_2 y + k_3 z + k_4) + \sum_{i=1}^s \psi_i(l_{i_1} y + l_{i_2} z + l_{i_3}), \quad (80)$$

where φ_i, ψ_i and p are known functions, and

$$p(k_1 x + k_2 y + k_3 z + k_4) = \frac{\iint r(v) dv dv}{k_1^2 + k_2^2 + k_3^2} + C_1 v + C_2. \quad (81)$$

In (78) set $l_{i_1} = l_{i_4}, l_{i_2} = l_{i_5}, l_{i_3} = l_{i_6}$ and $l_7 x + l_8 y + l_9 z + l_{10} = C_1 v + C_2$, similar to the calculation of (43) we get

$$\begin{aligned} & f_{i_1}(l_{i_1} y + l_{i_2} z + l_{i_3}) + f_{i_2}(l_{i_1} y + l_{i_2} z + l_{i_3}) = \varphi_i(l_{i_1} y + l_{i_2} z + l_{i_3}), \\ & f_{i_1}(l_{i_1} y + l_{i_2} z + l_{i_3}) - f_{i_2}(l_{i_1} y + l_{i_2} z + l_{i_3}) \\ &= \frac{1}{\sqrt{-l_{i_1}^2 - l_{i_2}^2}} \int_{l_{i_1} y_0 + l_{i_2} z_0 + l_{i_3}}^{l_{i_1} y + l_{i_2} z + l_{i_3}} \psi_i(\xi_i) d\xi_i + f_{i_1}(l_{i_1} y_0 + l_{i_2} z_0 + l_{i_3}) - f_{i_2}(l_{i_1} y_0 + l_{i_2} z_0 + l_{i_3}). \end{aligned}$$

By the further calculation, the exact solution of Eq. (74) on the conditions of (75), (79) and (80) is

$$\begin{aligned} u(x, y, z) &= p(k_1 x + k_2 y + k_3 z + k_4) \\ &+ \sum_{i=1}^s \left(\frac{1}{2} \varphi_i \left(x \sqrt{-l_{i_1}^2 - l_{i_2}^2} + l_{i_1} y + l_{i_2} z + l_{i_3} \right) + \frac{1}{2} \varphi_i \left(-x \sqrt{-l_{i_1}^2 - l_{i_2}^2} + l_{i_1} y + l_{i_2} z + l_{i_3} \right) \right) \\ &+ \frac{1}{2\sqrt{-l_{i_1}^2 - l_{i_2}^2}} \int_{-x\sqrt{-l_{i_1}^2 - l_{i_2}^2} + l_{i_1} y + l_{i_2} z + l_{i_3}}^{x\sqrt{-l_{i_1}^2 - l_{i_2}^2} + l_{i_1} y + l_{i_2} z + l_{i_3}} \psi_i(\xi_i) d\xi_i \end{aligned} \quad (82)$$

2D wave equation and acoustic wave equation which we studied previously are homogeneous equations, the general solution and the exact solution of the Cauchy problem for their non-homogeneous equation can be obtained similar to the Poisson equation.

1.8. Helmholtz equation

Before research Helmholtz equation, we first consider a PDE as follows

$$a_1 u_{xx} + a_2 u_{yy} + a_3 u_{zz} + a_4 u_{xy} + a_5 u_{yz} + a_6 u_{zx} = a_7, \quad (83)$$

where $a_i = a_i(x, y, z, u)$, ($i = 1, 2, \dots, 7$), according to Transformational Method 1, set

$$u(x, y, z) = f(v) = f(k_1 x + k_2 y + k_3 z + k_4), \quad (84)$$

where $k_1 - k_4$ are parameters to be determined, f is an undetermined unary function, then

$$\begin{aligned} & a_1 u_{xx} + a_2 u_{yy} + a_3 u_{zz} + a_4 u_{xy} + a_5 u_{yz} + a_6 u_{zx} \\ &= k_1^2 a_1 f_v'' + k_2^2 a_2 f_v'' + k_3^2 a_3 f_v'' + k_1 k_2 a_4 f_v'' + k_2 k_3 a_5 f_v'' + k_1 k_3 a_6 f_v'' \\ &= a_7. \end{aligned}$$

Namely

$$f_v'' = \frac{a_7}{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3 + k_1 k_2 a_4 + k_2 k_3 a_5 + k_1 k_3 a_6}. \quad (85)$$

If $\frac{a_7}{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3 + k_1 k_2 a_4 + k_2 k_3 a_5 + k_1 k_3 a_6}$ can be converted into $g(v)$ or $h(f)$, it can be further computed, set

$$a_i(x, y, z, u) = a_i(v), (i = 1, 2, \dots, 7), \quad (86)$$

So the particular solution of Eq. (83) on the condition of (86) is

$$u(x, y, z) = \iint \frac{a_7 dv dv}{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3 + k_1 k_2 a_4 + k_2 k_3 a_5 + k_1 k_3 a_6} + C_1 v + C_2, \quad (87)$$

where C_1 and C_2 are arbitrary constant, $k_1 - k_4$ are determinate parameters. For instance

$$u_{xx} + (k_1 x + k_2 y + k_3 z + k_4)^m u_{yy} + (k_1 x + k_2 y + k_3 z + k_4)^n u_{zz} = \sin(k_1 x + k_2 y + k_3 z + k_4). \quad (88)$$

According to (87) its particular solution is

$$u(x, y, z) = \iint \frac{\sin v dv dv}{k_1^2 + k_2^2 v^m + k_3^2 v^n} + C_1 v + C_2,$$

where $v(x, y, z) = k_1 x + k_2 y + k_3 z + k_4$. Set

$$a_i(x, y, z, u) = a_i(u), (i = 1, 2, \dots, 7). \quad (89)$$

From (84)-(85) we have

$$\begin{aligned} f_v'' &= \frac{a_7}{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3 + k_1 k_2 a_4 + k_2 k_3 a_5 + k_1 k_3 a_6} \\ \implies v &= C_1 \pm \int \left(C_2 + 2 \int \frac{a_7 df}{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3 + k_1 k_2 a_4 + k_2 k_3 a_5 + k_1 k_3 a_6} \right)^{-\frac{1}{2}} df, \end{aligned}$$

where $k_1 - k_4, C_1$ and C_2 are arbitrary constant. Namely

$$a_1(u) u_{xx} + a_2(u) u_{yy} + a_3(u) u_{zz} + a_4(u) u_{xy} + a_5(u) u_{yz} + a_6(u) u_{zx} = a_7(u). \quad (90)$$

The particular solution of Eq. (90) is

$$v = C_1 \pm \int \left(C_2 + 2 \int \frac{a_7 du}{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3 + k_1 k_2 a_4 + k_2 k_3 a_5 + k_1 k_3 a_6} \right)^{-\frac{1}{2}} du. \quad (91)$$

The solving method of Eq. (83) can be extended to any similar PDEs with n space variables. Emden-Fowler equation [32, 33], Klein-Gordon equation [34, 35] and sine-Gordon equation [36] are special cases of Eq. (90), which are the hotspots of current research.

Consider the following PDE

$$a_1 u_{xx} + a_2 u_{yy} + a_3 u_{zz} + k^2 u = 0 \quad (92)$$

It's a special case of Eq. (90), according to (91)

$$\begin{aligned} v &= C_1 \pm \int \left(C_2 - 2 \int \frac{k^2 u du}{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3} \right)^{-\frac{1}{2}} du \\ &= C_1 \pm \frac{\sqrt{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3} \arcsin(C_3 u)}{k} \\ \implies u &= \frac{1}{C_3} \sin \left(\frac{\pm k (v - C_1)}{\sqrt{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3}} \right) \\ &= \pm C_4 \sin \left(\frac{C_5 + k (k_1 x + k_2 y + k_3 z)}{\sqrt{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3}} \right). \end{aligned}$$

Since C_4 is an arbitrary constant, so the particular solution of Eq. (92) can be written as

$$u(x, y, z) = C_4 \sin \left(\frac{C_5 + k(k_1x + k_2y + k_3z)}{\sqrt{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3}} \right), \quad (93)$$

where $k_1 - k_3, C_4$ and C_5 are arbitrary constant. We use Transformational Method 3 to obtain the general solution of Eq. (92), set

$$u(x, y, z) = g(x, y, z) h(w) = g(x, y, z) h(l_1x + l_2y + l_3z + l_4), \quad (94)$$

where $w(x, y, z) = l_1x + l_2y + l_3z + l_4, l_1 - l_4$ are undetermined parameters, $h(w)$ and $g(x, y, z)$ are undetermined second differentiable functions, according to (72) and (94) we get

$$\begin{aligned} u_{xx} &= hg_{xx} + 2l_1g_xh'_w + l_1^2gh''_w, \\ u_{yy} &= hg_{yy} + 2l_2g_yh'_w + l_2^2gh''_w, \\ u_{zz} &= hg_{zz} + 2l_3g_zh'_w + l_3^2gh''_w. \end{aligned}$$

So

$$\begin{aligned} &a_1u_{xx} + a_2u_{yy} + a_3u_{zz} + k^2u \\ &= a_1hg_{xx} + 2a_1l_1g_xh'_w + a_1l_1^2gh''_w + a_2hg_{yy} + 2a_2l_2g_yh'_w \\ &+ a_2l_2^2gh''_w + a_3hg_{zz} + 2a_3l_3g_zh'_w + a_3l_3^2gh''_w + k^2gh. \end{aligned}$$

Namely

$$(a_1l_1^2 + a_2l_2^2 + a_3l_3^2)gh''_w + 2(a_1l_1g_x + a_2l_2g_y + a_3l_3g_z)h'_w + (a_1g_{xx} + a_2g_{yy} + a_3g_{zz} + k^2g)h = 0. \quad (95)$$

Set $h(w)$ an arbitrary unary second differentiable function, according to (95) we obtain

$$a_1l_1^2 + a_2l_2^2 + a_3l_3^2 = 0 \Rightarrow l_1 = \pm \sqrt{\frac{-a_2l_2^2 - a_3l_3^2}{a_1}}, \quad (96)$$

$$a_1l_1g_x + a_2l_2g_y + a_3l_3g_z = 0, \quad (97)$$

$$a_1g_{xx} + a_2g_{yy} + a_3g_{zz} + k^2g = 0. \quad (98)$$

By (93) the particular solution of Eq. (98) is

$$g(x, y, z) = C_4 \sin \left(\frac{C_5 + k(k_1x + k_2y + k_3z)}{\sqrt{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3}} \right) \quad (99)$$

Substituting from (99) into (97) we get

$$\begin{aligned} &a_1l_1g_x + a_2l_2g_y + a_3l_3g_z \\ &= \frac{a_1l_1C_4kk_1 + a_2l_2C_4kk_2 + a_3l_3C_4kk_3}{\sqrt{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3}} \cos \left(\frac{C_5 + k(k_1x + k_2y + k_3z)}{\sqrt{k_1^2 a_1 + k_2^2 a_2 + k_3^2 a_3}} \right) = 0 \\ &\Rightarrow a_1l_1C_4kk_1 + a_2l_2C_4kk_2 + a_3l_3C_4kk_3 = 0. \end{aligned}$$

Namely

$$k_1 = \frac{-a_2k_2l_2 - a_3k_3l_3}{a_1l_1}. \quad (100)$$

Then

$$\begin{aligned}
u(x, y, z) &= g(x, y, z) h(w) \\
&= \sin\left(\frac{C_5 + k(k_1x + k_2y + k_3z)}{\sqrt{k_1^2a_1 + k_2^2a_2 + k_3^2a_3}}\right) h(l_1x + l_2y + l_3z + l_4) \\
&= \sin\left(\frac{C_5a_1l_1 - k(a_2k_2l_2 + a_3k_3l_3)x + ka_1l_1(k_2y + k_3z)}{\sqrt{(a_2k_2l_2 + a_3k_3l_3)^2 + (a_2k_2^2 + a_3k_3^2)a_1^2l_1^2}}\right) h(l_1x + l_2y + l_3z + l_4).
\end{aligned}$$

So the general solution of Eq. (92) is

$$\begin{aligned}
u &= \sin\left(\frac{l_5 - k(a_2k_2l_2 + a_3k_3l_3)x + k\sqrt{-a_1a_2l_2^2 - a_1a_3l_3^2}(k_2y + k_3z)}{\sqrt{(a_2k_2l_2 + a_3k_3l_3)^2 - a_1(a_2k_2^2 + a_3k_3^2)(a_2l_2^2 + a_3l_3^2)}}\right) \\
&h_1\left(\sqrt{\frac{-a_2l_2^2 - a_3l_3^2}{a_1}}x + l_2y + l_3z + l_4\right) \\
&+ \sin\left(\frac{l_{15} - k(a_2k_{12}l_{12} + a_3k_{13}l_{13})x - k\sqrt{-a_1a_2l_{12}^2 - a_1a_3l_{13}^2}(k_{12}y + k_{13}z)}{\sqrt{(a_2k_{12}l_{12} + a_3k_{13}l_{13})^2 - a_1(a_2k_{12}^2 + a_3k_{13}^2)(a_2l_{12}^2 + a_3l_{13}^2)}}\right) \\
&h_2\left(-\sqrt{\frac{-a_2l_{12}^2 - a_3l_{13}^2}{a_1}}x + l_{12}y + l_{13}z + l_{14}\right),
\end{aligned} \tag{101}$$

where h_1 and h_2 are arbitrary unary second differentiable functions, $k_2, k_3, k_{12}, k_{13}, l_2 - l_5$ and $l_{12} - l_{15}$ are arbitrary constants.

Consider the following 3D Helmholtz equation

$$u_{xx} + u_{yy} + u_{zz} + k^2u = 0. \tag{102}$$

According to (101) we can get the general solution of Eq. (102) is

$$\begin{aligned}
u &= \sin\left(\frac{l_5 - k(k_2l_2 + k_3l_3)x + k\sqrt{-l_2^2 - l_3^2}(k_2y + k_3z)}{\sqrt{(k_2l_2 + k_3l_3)^2 - (k_2^2 + k_3^2)(l_2^2 + l_3^2)}}\right) h_1\left(\sqrt{-l_2^2 - l_3^2}x + l_2y + l_3z + l_4\right) \\
&+ \sin\left(\frac{l_{15} - k(k_{12}l_{12} + k_{13}l_{13})x - k\sqrt{-l_{12}^2 - l_{13}^2}(k_{12}y + k_{13}z)}{\sqrt{(k_{12}l_{12} + k_{13}l_{13})^2 - (k_{12}^2 + k_{13}^2)(l_{12}^2 + l_{13}^2)}}\right) \\
&h_2\left(-\sqrt{-l_{12}^2 - l_{13}^2}x + l_{12}y + l_{13}z + l_{14}\right),
\end{aligned} \tag{103}$$

Consider the following 2D Helmholtz equation

$$u_{xx} + u_{yy} + k^2u = 0. \tag{104}$$

By (101) the general solution of Eq. (102) could be got

$$\begin{aligned}
u &= \sin \left(\frac{C_6 - k(a_2 k_2 l_2 + a_3 k_3 l_3)x + k\sqrt{-a_1 a_2 l_2^2 - a_1 a_3 l_3^2}(k_2 y + k_3 z)}{\sqrt{(a_2 k_2 l_2)^2 - (a_2 k_2^2)(a_2 l_2^2)}} \right) \\
&h_1 \left(\sqrt{\frac{-a_2 l_2^2}{a_1}} x + l_2 y + l_4 \right) \\
&+ \sin \left(\frac{C_8 - k(a_2 k_{12} l_{12} + a_3 k_{13} l_{13})x - k\sqrt{-a_1 a_2 l_{12}^2 - a_1 a_3 l_{13}^2}(k_{12} y + k_{13} z)}{\sqrt{(a_2 k_{12} l_{12})^2 - (a_2 k_{12}^2)(a_2 l_{12}^2)}} \right) \\
&h_2 \left(-\sqrt{\frac{-a_2 l_{12}^2 - a_3 l_{13}^2}{a_1}} x + l_{12} y + l_{13} z + l_{14} \right).
\end{aligned}$$

The denominator of the above equation is equal to zero, so we can preliminarily judge that Eq. (104) has no general solution.

For the 1D Helmholtz equation $u_x x + k^2 u = 0$, according to (101) we can get that the denominator is equal to zero, so it can be judged preliminarily that 1D Helmholtz equation does not have any general solution.

Currently analysing the Helmholtz equation is mainly used numerical methods [37-40]. Here we consider the following boundary value problem of Eq. (102)

$$u(0, y, z) = \sin(\sqrt{2}k(y + 2z)) \varphi(y + z), \quad (105)$$

$$u_x(0, y, z) = \sqrt{-2} \sin(\sqrt{2}k(y + 2z)) \phi'(x + y) + 3k \cos(\sqrt{2}k(y + 2z)) \phi(x + y), \quad (106)$$

where φ, ϕ are known function, comparing (103) with (105) we obtain

$$k_2 = k_{12} = l_2 = l_3 = l_{12} = l_{13} = 1, k_3 = k_{13} = 2, l_4 = l_{14} = C_6 = C_8 = 0.$$

Namely

$$\begin{aligned}
u &= \sin(3kix + \sqrt{2}k(y + 2z)) h_1(\sqrt{-2}x + y + z) \\
&+ \sin(3kix - \sqrt{2}k(y + 2z)) h_2(-\sqrt{-2}x + y + z).
\end{aligned} \quad (107)$$

Then

$$\begin{aligned}
u(0, y, z) &= \sin(\sqrt{2}k(y + 2z)) h_1(y + z) - \sin(\sqrt{2}k(y + 2z)) h_2(y + z) \\
&= \sin(\sqrt{2}k(y + 2z)) \varphi(y + z) \Rightarrow h_1(y + z) - h_2(y + z) = \varphi(y + z), \\
u_x(0, y, z) &= \sqrt{-2} \sin(\sqrt{2}k(y + 2z)) (h_1'(y + z) + h_2'(y + z)) \\
&+ 3k \cos(\sqrt{2}k(y + 2z)) (h_1(y + z) + h_2(y + z)) \\
&= \sqrt{-2} \sin(\sqrt{2}k(y + 2z)) \phi'(x + y) + 3k \cos(\sqrt{2}k(y + 2z)) \phi(x + y) \\
&\Rightarrow h_1(y + z) + h_2(y + z) = \phi(x + y).
\end{aligned}$$

Namely

$$h_1(y + z) - h_2(y + z) = \varphi(y + z) \quad (108)$$

$$h_1(y+z) + h_2(y+z) = \phi(x+y) \quad (109)$$

Then

$$\begin{aligned} h_1(y+z) &= \frac{1}{2} (\phi(y+z) + \varphi(y+z)) \\ \Rightarrow h_1(\sqrt{-2}x + y + z) &= \frac{1}{2} (\phi(\sqrt{-2}x + y + z) + \varphi(\sqrt{-2}x + y + z)), \\ h_2(y+z) &= \frac{1}{2} (\phi(y+z) - \varphi(y+z)) \\ \Rightarrow h_2(-\sqrt{-2}x + y + z) &= \frac{1}{2} (\phi(-\sqrt{-2}x + y + z) - \varphi(-\sqrt{-2}x + y + z)). \end{aligned}$$

So the exact solution of Eq. (102) on the conditions of (105) and (106) can be get

$$\begin{aligned} u &= \frac{1}{2} \sin(3k_1x + \sqrt{2}k(y+2z)) (\phi(\sqrt{-2}x + y + z) + \varphi(\sqrt{-2}x + y + z)) \\ &+ \frac{1}{2} \sin(3k_1x - \sqrt{2}k(y+2z)) (\phi(\sqrt{-2}x + y + z) - \varphi(\sqrt{-2}x + y + z)). \end{aligned} \quad (110)$$

1.9. heat equation and diffusion equation

Consider the following PDE

$$a_0u_t + a_1u_{xx} + a_2u_{yy} + a_3u_{zz} = 0, \quad (111)$$

where a_i are known constants. For solving its particular solution, by Transformational Method 1 we set

$$u(t, x, y, z) = f(v) = f(k_0t + k_1x + k_2y + k_3z + k_4), \quad (112)$$

where $v(t, x, y, z) = k_0t + k_1x + k_2y + k_3z + k_4$, $k_0 - k_4$ are parameters to be determined, f is an undetermined unary second differentiable function. By (7)

$$a_0u_t + a_1u_{xx} + a_2u_{yy} + a_3u_{zz} = a_0k_0f'_v + (a_1k_1^2 + a_2k_2^2 + a_3k_3^2) f''_v = 0.$$

Set $w = f'_v$, then

$$\begin{aligned} a_0k_0f'_v + (a_1k_1^2 + a_2k_2^2 + a_3k_3^2) f''_v = 0 &\Rightarrow (a_1k_1^2 + a_2k_2^2 + a_3k_3^2) w'_v = -a_0k_0w \\ \Rightarrow w = k_7e^{\frac{-a_0k_0v}{a_1k_1^2 + a_2k_2^2 + a_3k_3^2}} &\Rightarrow f(v) = -k_7 \frac{a_1k_1^2 + a_2k_2^2 + a_3k_3^2}{a_0k_0} e^{\frac{-a_0k_0v}{a_1k_1^2 + a_2k_2^2 + a_3k_3^2}} + k_6. \end{aligned}$$

So the particular solution of Eq. (111) is

$$u(t, x, y, z) = k_5e^{\frac{-a_0k_0(k_0t + k_1x + k_2y + k_3z)}{a_1k_1^2 + a_2k_2^2 + a_3k_3^2}} + k_6, \quad (113)$$

where $k_0 - k_6$ are arbitrary constants.

In order to obtain the general solution of Eq. (111), according to Transformational Method 3 we set

$$u(t, x, y, z) = gh(w) = g(t, x, y, z) h(l_0t + l_1x + l_2y + l_3z + l_4), \quad (114)$$

where $w(t, x, y, z) = l_0t + l_1x + l_2y + l_3z + l_4$, $l_0 - l_4$ are parameters to be determined, h and g are undetermined second differentiable functions. By (71) and (72) we get

$$u_t = l_0gh'_w + hg_t,$$

$$\begin{aligned}
u_{xx} &= l_1^2 g h_w'' + 2l_1 g_x h_w' + h g_{xx}, \\
u_{yy} &= l_2^2 g h_w'' + 2l_2 g_y h_w' + h g_{yy}, \\
u_{zz} &= l_3^2 g h_w'' + 2l_3 g_z h_w' + h g_{zz}.
\end{aligned}$$

Then

$$\begin{aligned}
&a_0 u_t + a_1 u_{xx} + a_2 u_{yy} + a_3 u_{zz} \\
&= a_0 l_0 g h_w' + a_0 h g_t + a_1 l_1^2 g h_w'' + 2a_1 l_1 g_x h_w' + a_1 h g_{xx} + a_2 l_2^2 g h_w'' + 2a_2 l_2 g_y h_w' + a_2 h g_{yy} \\
&+ a_3 l_3^2 g h_w'' + 2a_3 l_3 g_z h_w' + a_3 h g_{zz}.
\end{aligned}$$

Namely

$$\begin{aligned}
&(a_1 l_1^2 + a_2 l_2^2 + a_3 l_3^2) g h_w'' + (a_0 l_0 g + 2a_1 l_1 g_x + 2a_2 l_2 g_y + 2a_3 l_3 g_z) h_w' \\
&+ (a_0 g_t + a_1 g_{xx} + a_2 g_{yy} + a_3 g_{zz}) h = 0.
\end{aligned} \tag{115}$$

Set $h(w)$ an arbitrary unary second differentiable function, according to (115) we get

$$a_1 l_1^2 + a_2 l_2^2 + a_3 l_3^2 = 0 \Rightarrow l_1 = \pm \sqrt{\frac{-a_2 l_2^2 - a_3 l_3^2}{a_1}}, \tag{116}$$

$$a_0 l_0 g + 2a_1 l_1 g_x + 2a_2 l_2 g_y + 2a_3 l_3 g_z = 0, \tag{117}$$

$$a_0 g_t + a_1 g_{xx} + a_2 g_{yy} + a_3 g_{zz} = 0. \tag{118}$$

By (113) the particular solution of Eq. (118) is

$$g(t, x, y, z) = k_5 e^{\frac{-a_0 k_0 (k_0 t + k_1 x + k_2 y + k_3 z)}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2}} + k_6, \tag{119}$$

Set $k_6 = 0$, and substituting from (119) into (117), then

$$\begin{aligned}
&a_0 l_0 g + 2a_1 l_1 g_x + 2a_2 l_2 g_y + 2a_3 l_3 g_z \\
&= a_0 l_0 k_5 e^{\frac{-a_0 k_0 (k_0 t + k_1 x + k_2 y + k_3 z)}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2}} - 2a_0 k_0 k_5 e^{\frac{-a_0 k_0 (k_0 t + k_1 x + k_2 y + k_3 z)}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2}} \frac{a_1 l_1 k_1 + a_2 l_2 k_2 + a_3 l_3 k_3}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2} = 0 \\
&\Rightarrow l_0 = 2k_0 \frac{a_1 l_1 k_1 + a_2 l_2 k_2 + a_3 l_3 k_3}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2}
\end{aligned}$$

We have

$$l_0 = 2k_0 \frac{a_1 l_1 k_1 + a_2 l_2 k_2 + a_3 l_3 k_3}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2}. \tag{120}$$

Therefore

$$\begin{aligned}
u(x, y, z, t) &= g(x, y, z, t) h(w) = k_5 e^{\frac{-a_0 k_0 (k_0 t + k_1 x + k_2 y + k_3 z)}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2}} h(l_0 t + l_1 x + l_2 y + l_3 z + l_4) \\
&= e^{\frac{-a_0 k_0 (k_0 t + k_1 x + k_2 y + k_3 z)}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2}} h\left(\frac{2k_0 (a_1 l_1 k_1 + a_2 l_2 k_2 + a_3 l_3 k_3) t}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2} + l_1 x + l_2 y + l_3 z + l_4\right).
\end{aligned}$$

So the general solution of Eq. (111) is

$$\begin{aligned}
u = e^{\frac{-a_0 k_0 (k_0 t + k_1 x + k_2 y + k_3 z)}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2}} & \\
h_1 \left(\frac{2k_0 \left(k_1 \sqrt{-a_1 (a_2 l_2^2 + a_3 l_3^2)} + a_2 l_2 k_2 + a_3 l_3 k_3 \right) t}{a_1 k_1^2 + a_2 k_2^2 + a_3 k_3^2} + \sqrt{\frac{-a_2 l_2^2 - a_3 l_3^2}{a_1} x + l_2 y + l_3 z + l_4} \right) & \\
+ e^{\frac{-a_0 k_{10} (k_{10} t + k_{11} x + k_{12} y + k_{13} z)}{a_1 k_{11}^2 + a_2 k_{12}^2 + a_3 k_{13}^2}} h_2 \left(\frac{2k_{10} (-k_{11} \sqrt{-a_1 (a_2 l_{12}^2 + a_3 l_{13}^2)} + a_2 l_{12} k_{12} + a_3 l_{13} k_{13}) t}{a_1 k_{11}^2 + a_2 k_{12}^2 + a_3 k_{13}^2} \right. & \\
\left. - \sqrt{\frac{-a_2 l_{12}^2 - a_3 l_{13}^2}{a_1} x + l_{12} y + l_{13} z + l_{14}} \right) & \tag{121}
\end{aligned}$$

where h_1 and h_2 are arbitrary unary second differentiable functions, $k_0 - k_3, k_{10} - k_{13}, l_2 - l_4$ and $l_{12} - l_{14}$ are arbitrary constants.

The form of 3D heat equation and diffusion equation is

$$u_t - a^2 (u_{xx} + u_{yy} + u_{zz}) = 0, \tag{122}$$

According to (24) we can get the general solution of Eq. (122) is

$$\begin{aligned}
u = e^{\frac{k_0 (k_0 t + k_1 x + k_2 y + k_3 z)}{(k_1^2 + k_2^2 + k_3^2) a^2}} h_1 \left(\frac{2k_0 \left(\sqrt{-l_2^2 - l_3^2} k_1 + l_2 k_2 + l_3 k_3 \right) t + \sqrt{-l_2^2 - l_3^2} x + l_2 y + l_3 z + l_4}{k_1^2 + k_2^2 + k_3^2} \right) & \\
+ e^{\frac{k_{10} (k_{10} t + k_{11} x + k_{12} y + k_{13} z)}{(k_{11}^2 + k_{12}^2 + k_{13}^2) a^2}} h_2 \left(\frac{2k_{10} \left(-\sqrt{-l_{12}^2 - l_{13}^2} k_{11} + l_{12} k_{12} + l_{13} k_{13} \right) t - \sqrt{-l_{12}^2 - l_{13}^2} x + l_{12} y + l_{13} z + l_{14}}{k_{11}^2 + k_{12}^2 + k_{13}^2} \right) & \tag{123}
\end{aligned}$$

The form of 2D heat equation and diffusion equation is

$$u_t - a^2 (u_{xx} + u_{yy}) = 0 \tag{124}$$

By (121) the general solution of Eq. (124) could be got

$$\begin{aligned}
u = e^{\frac{k_0 (k_0 t + k_1 x + k_2 y)}{(k_1^2 + k_2^2) a^2}} h_1 \left(\frac{2k_0 (il_2 k_1 + l_2 k_2) t + il_2 x + l_2 y + l_4}{k_1^2 + k_2^2} \right) & \\
+ e^{\frac{k_{10} (k_{10} t + k_{11} x + k_{12} y)}{(k_{11}^2 + k_{12}^2) a^2}} h_2 \left(\frac{2k_{10} (-il_{12} k_{11} + l_{12} k_{12}) t - il_{12} x + l_{12} y + l_{14}}{k_{11}^2 + k_{12}^2} \right) & \tag{125}
\end{aligned}$$

The form of 1D heat equation and diffusion equation is

$$u_t - a^2 u_{xx} = 0 \tag{126}$$

According to (121) we have

$$u = C e^{\frac{k_0 (k_0 t + k_1 x)}{a^2 k_1^2}} \tag{127}$$

Therefore, it can be preliminarily determined that Eq. (126) has no general solution.

Nonlinear problem [41-44] and numerical methods [45-47] are the research hotspots of the heat equation, here we consider the following initial value problem of Eq. (122)

$$u(x, y, z, 0) = e^{\frac{x+y+z}{a^2}} (\varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z)) \quad (128)$$

Comparing (123) with (128) we get

$$k_1 = k_2 = k_3 = \frac{k_4}{3}, k_{11} = k_{12} = k_{13} = \frac{k_{14}}{3}, l_2 = l_3 = l_{12} = l_{13} = 1, l_5 = l_{15} = 0$$

So

$$u(x, y, z, t) = e^{\frac{x+y+z+3t}{a^2}} (h_1(\sqrt{-2}x + y + z + (4 + 2\sqrt{-2})t) + h_2(-\sqrt{-2}x + y + z + (4 - 2\sqrt{-2})t)) \quad (129)$$

Then

$$\begin{aligned} u(x, y, z, 0) &= e^{\frac{x+y+z}{a^2}} (\varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z)) \\ &= e^{\frac{x+y+z}{a^2}} (h_1(\sqrt{-2}x + y + z) + h_2(-\sqrt{-2}x + y + z)) \\ &\Rightarrow \varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z) = h_1(\sqrt{-2}x + y + z) + h_2(-\sqrt{-2}x + y + z) \\ &\Rightarrow \varphi_1(\sqrt{-2}x + y + z + (4 + 2\sqrt{-2})t) = h_1(\sqrt{-2}x + y + z + (4 + 2\sqrt{-2})t) \\ &\varphi_2(-\sqrt{-2}x + y + z + (4 - 2\sqrt{-2})t) = h_2(-\sqrt{-2}x + y + z + (4 - 2\sqrt{-2})t) \end{aligned}$$

Namely

$$\begin{aligned} h_1(\sqrt{-2}x + y + z + (4 + 2\sqrt{-2})t) &= \varphi_1(\sqrt{-2}x + y + z + (4 + 2\sqrt{-2})t) \\ h_2(-\sqrt{-2}x + y + z + (4 - 2\sqrt{-2})t) &= \varphi_2(-\sqrt{-2}x + y + z + (4 - 2\sqrt{-2})t) \end{aligned}$$

So the exact solution of Eq. (122) on the conditions of (128) can be get

$$u(x, y, z, t) = e^{\frac{x+y+z+3t}{a^2}} (\varphi_1(\sqrt{-2}x + y + z + (4 + 2\sqrt{-2})t) + \varphi_2(-\sqrt{-2}x + y + z + (4 - 2\sqrt{-2})t)) \quad (130)$$

1.10. Schrödinger Equation

Linear [48-50] and nonlinear [51, 52] stationary state Schrödinger equation are the focus of current research, Consider the following linear equation

$$\frac{\hbar^2}{2m} \Delta u - (V(x, y, z) - E)u = 0, \quad (131)$$

where m is the mass of the described particle and \hbar is the reduced Plank constant, by Transformational Method 2, set

$$u(x, y, z) = f(v) = f(k_1x + k_2y + k_3z + k_4), \quad (132)$$

$$V(x, y, z) - E = a(v) = a(k_1x + k_2y + k_3z + k_4), \quad (133)$$

where $v(x, y, z) = k_1x + k_2y + k_3z + k_4$, $k_1 - k_4$ are known parameters, $V(x, y, z) - E = a(v)$ is a known function, f is an undetermined unary second differentiable function, then

$$\begin{aligned} \frac{\hbar^2}{2m} \Delta u - (V(x, y, z) - E)u &= \frac{\hbar^2}{2m} (k_1^2 + k_2^2 + k_3^2) f_v'' - a(v) f = 0 \\ \Rightarrow f_v'' + b(v) f &= 0. \end{aligned}$$

Namely

$$f_v'' + b(v) f = 0. \quad (134)$$

where

$$b(v) = \frac{-2ma(v)}{\hbar^2 (k_1^2 + k_2^2 + k_3^2)} = \frac{-2m(V(x, y, z) - E)}{\hbar^2 (k_1^2 + k_2^2 + k_3^2)}. \quad (135)$$

If $b(v)$ is some special function [53], Eq. (134) has a particular solution and its general solution may be obtained by the law of second-order linear ODEs (LODEs), such as

$$b(v) = -c(v^{2n} + nv^{n-1}), \quad (136)$$

$$V(x, y, z) = a(v) + E = \frac{c\hbar^2 (k_1^2 + k_2^2 + k_3^2)}{2m} (cv^{2n} + nv^{n-1}) + E, \quad (137)$$

where c is an arbitrary constant, the particular solution of Eq. (134) under the condition of (137) is

$$f(v) = \exp\left(\frac{cv^{n+1}}{n+1}\right) = \exp\left(\frac{c(k_1x + k_2y + k_3z + k_4)^{n+1}}{n+1}\right).$$

So the particular solution of Eq. (131) under the condition of (137) is

$$u(x, y, z) = \exp\left(\frac{c(k_1x + k_2y + k_3z + k_4)^{n+1}}{n+1}\right). \quad (138)$$

For getting the general solution of Eq. (131) under the condition of (137), according to Transformational Method 3, we set

$$u(x, y, z) = g(x, y, z) h(w) = g(x, y, z) h(l_1x + l_2y + l_3z + l_4), \quad (139)$$

where $w(x, y, z) = l_1x + l_2y + l_3z + l_4$, $l_1 - l_4$ are parameters to be determined, $h(w)$ and $g(x, y, z)$ are undetermined second differentiable function, by (129) and (72) we obtain

$$u_{xx} = hg_{xx} + 2l_1g_x h_w' + l_1^2 g h_w'', \quad (140)$$

$$u_{yy} = hg_{yy} + 2l_2g_y h_w' + l_2^2 g h_w'', \quad (141)$$

$$u_{zz} = hg_{zz} + 2l_3g_z h_w' + l_3^2 g h_w''. \quad (142)$$

Then

$$\begin{aligned}
& \frac{\hbar^2}{2m} \Delta u - V((x, y, z) - E)u \\
&= \frac{\hbar^2}{2m} h g_{xx} + \frac{\hbar^2}{m} l_1 g_x h'_w + \frac{\hbar^2}{2m} l_1^2 g h''_w + \frac{\hbar^2}{2m} h g_{yy} + \frac{\hbar^2}{m} l_2 g_y h'_w + \frac{\hbar^2}{2m} l_2^2 g h''_w + \frac{\hbar^2}{2m} h g_{zz} \\
&+ \frac{\hbar^2}{m} l_3 g_z h'_w + \frac{\hbar^2}{2m} l_3^2 g h''_w + (V(x, y, z) - E)gh = 0.
\end{aligned}$$

We have

$$\begin{aligned}
& \frac{\hbar^2}{2m} (l_1^2 + l_2^2 + l_3^2) g h''_w + \frac{\hbar^2}{m} (l_1 g_x + l_2 g_y + l_3 g_z) h'_w \\
&+ \left(\frac{\hbar^2}{2m} g_{xx} + \frac{\hbar^2}{2m} g_{yy} + \frac{\hbar^2}{2m} g_{zz} + (V(x, y, z) - E)g \right) h = 0.
\end{aligned} \tag{143}$$

Set $h(w)$ an arbitrary second differentiable function, by (143) we get

$$l_1^2 + l_2^2 + l_3^2 = 0 \implies l_1 = \pm \sqrt{-l_2^2 - l_3^2}, \tag{144}$$

$$l_1 g_x + l_2 g_y + l_3 g_z = 0, \tag{145}$$

$$\frac{\hbar^2}{2m} g_{xx} + \frac{\hbar^2}{2m} g_{yy} + \frac{\hbar^2}{2m} g_{zz} + (V(x, y, z) - E)g = 0. \tag{146}$$

By (138), the particular solution of Eq. (146) on the condition of (137) is

$$g(x, y, z) = \exp\left(\frac{c(k_1 x + k_2 y + k_3 z + k_4)^{n+1}}{n+1}\right). \tag{147}$$

Substituting from (147) into (145) we get

$$\begin{aligned}
& l_1 c k_1 (k_1 x + k_2 y + k_3 z + k_4)^n \exp\left(\frac{c(k_1 x + k_2 y + k_3 z + k_4)^{n+1}}{n+1}\right) \\
&+ l_2 c k_2 (k_1 x + k_2 y + k_3 z + k_4)^n \exp\left(\frac{c(k_1 x + k_2 y + k_3 z + k_4)^{n+1}}{n+1}\right) \\
&+ l_3 c k_3 (k_1 x + k_2 y + k_3 z + k_4)^n \exp\left(\frac{c(k_1 x + k_2 y + k_3 z + k_4)^{n+1}}{n+1}\right) = 0 \\
\implies l_1 &= \frac{-k_2 l_2 - k_3 l_3}{k_1} = \pm \sqrt{-l_2^2 - l_3^2} \\
\implies l_2 &= \frac{-k_1 k_2 l_3 \pm \sqrt{-k_1^4 l_3^2 - k_1^2 k_3^2 l_3^2 - k_2^2 k_3^2 l_3^2}}{k_1^2 + k_2^2}.
\end{aligned}$$

Namely

$$l_2 = \frac{-k_1 k_2 l_3 \pm \sqrt{-k_1^4 l_3^2 - k_1^2 k_3^2 l_3^2 - k_2^2 k_3^2 l_3^2}}{k_1^2 + k_2^2}. \tag{148}$$

Then

$$u(x, y, z) = g(x, y, z) h(w) = \exp\left(\frac{c(k_1 x + k_2 y + k_3 z + k_4)^{n+1}}{n+1}\right) h(l_1 x + l_2 y + l_3 z + l_4).$$

So the general solution of Eq. (131) on the condition of (137) is

$$u = \exp\left(\frac{c(k_1x + k_2y + k_3z + k_4)^{n+1}}{n+1}\right) \left(h_1\left(\sqrt{-l_2^2 - l_3^2}x + l_2y + l_3z + l_4\right) + h_2\left(-\sqrt{-l_{12}^2 - l_{13}^2}x + l_{12}y + l_{13}z + l_{14}\right) \right), \quad (149)$$

where h_1 and h_2 are arbitrary second differentiable unary function, $k_1 - k_4$ and c are determinate parameters, l_3, l_4, l_{13} and l_{14} are arbitrary constants, $l_{12} = \frac{-k_1k_2l_{13} \pm l_{13}\sqrt{-k_1^4 - k_1^2k_3^2 - k_2^2k_3^2}}{k_1^2 + k_2^2}$.

Time dependent Schrödinger equation is always the focus of research [54-59], in addition, the related nonlinear equation [60, 61] and the time fractional Schrödinger equations (TFSEs) [62, 63] are the deeply researched field. Consider the following linear equation

$$i\hbar u_t + \frac{\hbar^2}{2m} \Delta u - V(x, y, z, t) u = 0. \quad (150)$$

According to Method 2, set

$$u(x, y, z, t) = f(v) = f(k_1x + k_2y + k_3z + k_4t + k_5), \quad (151)$$

$$V(x, y, z, t) = a(v) = a(k_1x + k_2y + k_3z + k_4t + k_5), \quad (152)$$

where $v = k_1x + k_2y + k_3z + k_4t + k_5$, $k_1 - k_5$ are known parameters, $V(x, y, z, t) = a(v)$ is a known function, f is an undetermined unary second differentiable function, then

$$i\hbar u_t + \frac{\hbar^2}{2m} \Delta u - V(x, y, z, t) u = i\hbar k_4 f'_v + \frac{\hbar^2}{2m} (k_1^2 + k_2^2 + k_3^2) f''_v - a(v) f = 0.$$

Namely

$$f''_v + k f'_v + b(v) f = 0, \quad (153)$$

where

$$k = \frac{i2mk_4}{\hbar(k_1^2 + k_2^2 + k_3^2)}, \quad b(v) = \frac{-2ma(v)}{\hbar^2(k_1^2 + k_2^2 + k_3^2)}. \quad (154)$$

If $b(v)$ is some special function [53], Eq. (153) has a particular solution and its general solution may be obtained by the law of second-order LODEs, such as

$$b(v) = c(-cv^{2n} + kv^n + nv^{n-1}).$$

The particular solution of Eq. (153) is

$$f(v) = \exp\left(-\frac{cv^{n+1}}{n+1}\right).$$

Namely

$$V(x, y, z, t) = a(v) = \frac{-c\hbar^2(k_1^2 + k_2^2 + k_3^2)}{2m} (-cv^{2n} + kv^n + nv^{n-1}). \quad (155)$$

The particular solution of Eq. (150) on the condition of (155) is

$$u(x, y, z, t) = \exp\left(-\frac{c(k_1x + k_2y + k_3z + k_4t + k_5)^{n+1}}{n+1}\right). \quad (156)$$

By

$$f_v'' + (g(v) + h(v)) f_v' + (g(v)h(v) + g_v') f = 0. \quad (157)$$

The particular solution of Eq. (157) is

$$f(v) = \exp\left(-\int g(v) dv\right). \quad (158)$$

Set $h(v) = -g(v) + k$, where $g(v)$ is an arbitrary unary first differentiable function, then

$$b(v) = -g^2(v) + kg(v) + g_v'. \quad (159)$$

Namely

$$V(x, y, z, t) = a(v) = \frac{\hbar^2 (k_1^2 + k_2^2 + k_3^2)}{2m} (g^2(v) - kg(v) - g_v'). \quad (160)$$

The particular solution of Eq. (150) on the condition of (160) is

$$u(x, y, z, t) = f(v) = \exp\left(-\int g(v) dv\right). \quad (161)$$

For getting the general solution of Eq. (150) on the condition of (155), according to Transformational Method 3, we set

$$u(x, y, z, t) = g(x, y, z, t) h(w) = g(x, y, z, t) h(l_1x + l_2y + l_3z + l_4t + l_5), \quad (162)$$

where $w = l_1x + l_2y + l_3z + l_4t + l_5$, $l_1 - l_5$ are parameters to be determined, h and g are undetermined second differentiable functions, by (71)-(72) and (162)

$$\begin{aligned} & i\hbar u_t + \frac{\hbar^2}{2m} \Delta u - V(x, y, z, t) u \\ &= i\hbar l_4 g h_w' + i\hbar h g_t + \frac{\hbar^2}{2m} l_1^2 g h_w'' + \frac{\hbar^2}{m} l_1 g_x h_w' + \frac{\hbar^2}{2m} h g_{xx} + \frac{\hbar^2}{2m} l_2^2 g h_w'' + \frac{\hbar^2}{m} l_2 g_y h_w' \\ &+ \frac{\hbar^2}{2m} h g_{yy} + \frac{\hbar^2}{2m} l_3^2 g h_w'' + \frac{\hbar^2}{m} l_3 g_z h_w' + \frac{\hbar^2}{2m} h g_{zz} - V g h \frac{\hbar^2}{2m} (l_1^2 + l_2^2 + l_3^2) g h_w'' \\ &+ \hbar \left(i l_4 g + \frac{\hbar}{m} l_1 g_x + \frac{\hbar}{m} l_2 g_y + \frac{\hbar}{m} l_3 g_z \right) h_w' + \left(i\hbar g_t + \frac{\hbar^2}{2m} g_{xx} + \frac{\hbar^2}{2m} g_{yy} + \frac{\hbar^2}{2m} g_{zz} - V g \right) h = 0. \end{aligned}$$

Namely

$$\begin{aligned} & \frac{\hbar^2}{2m} (l_1^2 + l_2^2 + l_3^2) g h_w'' + \hbar \left(i l_4 g + \frac{\hbar}{m} l_1 g_x + \frac{\hbar}{m} l_2 g_y + \frac{\hbar}{m} l_3 g_z \right) h_w' \\ &+ \left(i\hbar g_t + \frac{\hbar^2}{2m} g_{xx} + \frac{\hbar^2}{2m} g_{yy} + \frac{\hbar^2}{2m} g_{zz} - V g \right) h = 0. \end{aligned} \quad (163)$$

Set $h(w)$ an arbitrary second differentiable function, by (163) we get

$$l_1^2 + l_2^2 + l_3^2 = 0 \implies l_1 = \pm \sqrt{-l_2^2 - l_3^2}, \quad (164)$$

$$i l_4 g + \frac{\hbar}{m} l_1 g_x + \frac{\hbar}{m} l_2 g_y + \frac{\hbar}{m} l_3 g_z = 0, \quad (165)$$

$$i\hbar g_t + \frac{\hbar^2}{2m} g_{xx} + \frac{\hbar^2}{2m} g_{yy} + \frac{\hbar^2}{2m} g_{zz} - V g = 0. \quad (166)$$

By (156) the particular solution of Eq. (166) on the condition of (155) is

$$g(x, y, z, t) = \exp\left(-\frac{c(k_1x + k_2y + k_3z + k_4t + k_5)^{n+1}}{n+1}\right). \quad (167)$$

Substituting from (167) into (165) we get

$$\begin{aligned} & il_4 \exp\left(-\frac{c(k_1x + k_2y + k_3z + k_4t + k_5)^{n+1}}{n+1}\right) \\ & - \frac{\hbar}{m} l_1 c k_1 (k_1x + k_2y + k_3z + k_4t + k_5)^n \exp\left(-\frac{c(k_1x + k_2y + k_3z + k_4t + k_5)^{n+1}}{n+1}\right) \\ & - \frac{\hbar}{m} l_2 c k_2 (k_1x + k_2y + k_3z + k_4t + k_5)^n \exp\left(-\frac{c(k_1x + k_2y + k_3z + k_4t + k_5)^{n+1}}{n+1}\right) \\ & - \frac{\hbar}{m} l_3 c k_3 (k_1x + k_2y + k_3z + k_4t + k_5)^n \exp\left(-\frac{c(k_1x + k_2y + k_3z + k_4t + k_5)^{n+1}}{n+1}\right) = 0 \\ \implies l_4 & = -\frac{\hbar c}{m} (l_1 k_1 + l_2 k_2 + l_3 k_3) (k_1x + k_2y + k_3z + k_4t + k_5)^n. \end{aligned}$$

Namely

$$l_4 = -\frac{\hbar c}{m} (l_1 k_1 + l_2 k_2 + l_3 k_3) (k_1x + k_2y + k_3z + k_4t + k_5)^n. \quad (168)$$

Since l_4 is a constant and is not a function of x, y, z and t , if (167) is the particular solution of Eq. (166), by (168) n must equal 0, then

$$V = \frac{-c\hbar^2 (k_1^2 + k_2^2 + k_3^2)}{2m} (-cv^{2n} + kv^n + nv^{n-1}) = \frac{-c\hbar^2 (k_1^2 + k_2^2 + k_3^2)}{2m} (-c + k). \quad (169)$$

Since $k_1 - k_5, k$ and c are determinate constants, so $V(x, y, z, t)$ is an determinate constants too, namely

$$l_4 = -\frac{\hbar c}{m} (l_1 k_1 + l_2 k_2 + l_3 k_3). \quad (170)$$

Then

$$u = gh = \exp\left(-\frac{c(k_1x + k_2y + k_3z + k_4t + k_5)^{n+1}}{n+1}\right) h(l_1x + l_2y + l_3z + l_4t + l_5).$$

So the general solution of Eq. (150) in the condition of (169) is

$$u = e^{-c(k_1x + k_2y + k_3z + k_4t + k_5)} \left(h_1 \left(\sqrt{-l_2^2 - l_3^2} x + l_2 y + l_3 z + l_4 t + l_5 \right) + h_2 \left(-\sqrt{-l_{12}^2 - l_{13}^2} x + l_{12} y + l_{13} z + l_{14} t + l_{15} \right) \right) \quad (171)$$

where h_1 and h_2 are arbitrary unary second differentiable functions, $l_{14} = -\frac{\hbar c}{m} (l_{11} k_1 + l_{12} k_2 + l_{13} k_3)$, $l_2, l_3, l_5, l_{12}, l_{13}$ and l_{15} are arbitrary parameters.

Consider the following initial value problem of Eq. (150) on the condition of (169)

$$u(x, y, z, 0) = e^{x+y+z} (\varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z)), \quad (172)$$

$$\begin{aligned}
u_t(x, y, z, 0) &= e^{x+y+z} (\varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z)) \\
&+ \frac{\hbar}{m} e^{x+y+z} \left((2 + \sqrt{-2}) \varphi_1'(\sqrt{-2}x + y + z) + (2 - \sqrt{-2}) \varphi_2'(-\sqrt{-2}x + y + z) \right).
\end{aligned} \tag{173}$$

Comparing (171) with (172) we have

$$k_1 = k_2 = k_3 = -\frac{1}{c}, l_2 = l_3 = l_{12} = l_{13} = 1, k_5 = l_5 = l_{15} = 0.$$

By further calculation which is in Appendix C, the exact solutions of the initial value problem is

$$\begin{aligned}
u &= e^{x+y+z+ct} \left(\varphi_1(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t) \right. \\
&\left. + \varphi_2(-\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 - \sqrt{-2})t) \right).
\end{aligned} \tag{174}$$

When $V(x, y, z, t)$ is a constant, Eq. (150) is also an important case of the diffusion equation with a source [64], its general solution and the exact solutions of the Cauchy problem are applicable to the diffusion equation.

1.11. Singular general solution of the Helmholtz equation

For the 3D Helmholtz equation

$$\Delta u + k^2 u = 0. \tag{175}$$

By (93), its particular solution is

$$u(x, y, z) = C_4 \sin \left(\frac{C_5 + k(k_1 x + k_2 y + k_3 z)}{\sqrt{k_1^2 + k_2^2 + k_3^2}} \right), \tag{176}$$

where $k_1 - k_3, C_4$ and C_5 are arbitrary constant. Similar to the solving method of (101), by (176) and Transformational Method 3 the SGS of Eq. (175) is

$$\begin{aligned}
u &= \sin \left(\frac{C_6 - k(k_2 l_2 + k_3 l_3)x + k\sqrt{-l_2^2 - l_3^2}(k_2 y + k_3 z)}{\sqrt{(k_2 l_2 + k_3 l_3)^2 - (k_2^2 + k_3^2)(l_2^2 + l_3^2)}} \right) h_1 \left(\sqrt{-l_2^2 - l_3^2}x + l_2 y + l_3 z + l_4 \right) \\
&+ \sin \left(\frac{C_8 - k(k_{12} l_{12} + k_{13} l_{13})x - k\sqrt{-l_{12}^2 - l_{13}^2}(k_{12} y + k_{13} z)}{\sqrt{(k_{12} l_{12} + k_{13} l_{13})^2 - (k_{12}^2 + k_{13}^2)(l_{12}^2 + l_{13}^2)}} \right) \\
&h_2 \left(-\sqrt{-l_{12}^2 - l_{13}^2}x + l_{12} y + l_{13} z + l_{14} \right),
\end{aligned} \tag{177}$$

where h_1 and h_2 are arbitrary unary second differentiable functions, $k_2, k_3, k_{12}, k_{13}, l_2 - l_4, l_{12} - l_{14}, C_6$ and C_8 are arbitrary constants.

According to [65]

$$u = (k_1 \cosh \alpha x + k_2 \sinh \alpha x) (k_3 \cosh \beta y + k_4 \sinh \beta y) (k_5 \cosh \gamma z + k_6 \sinh \gamma z), \quad k^2 = -\alpha^2 - \beta^2 - \gamma^2, \tag{178}$$

where $k_1 - k_6$ are arbitrary constants, (178) is a particular solution of Eq. (175). Other particular solutions of the Helmholtz equation can be referred to [65, 66].

Here we shall use (178) and Transformational Method 3 to obtain a new solution of Eq. (175), and shall analyze the relationship between it and (177).

According to Transformational Method 3, we set

$$u(x, y, z) = g(x, y, z) h(w) = g(x, y, z) h(l_1x + l_2y + l_3z + l_4), \quad (179)$$

where $h(w)$ and $g(x, y, z)$ are undetermined second differentiable functions, $w(x, y, z) = l_1x + l_2y + l_3z + l_4$, $l_1 - l_4$ are undetermined parameters, by (179) we have

$$u_{xx} = hg_{xx} + 2l_1g_xh'_w + l_1^2gh''_w, \quad (180)$$

$$u_{yy} = hg_{yy} + 2l_2g_yh'_w + l_2^2gh''_w, \quad (181)$$

$$u_{zz} = hg_{zz} + 2l_3g_zh'_w + l_3^2gh''_w. \quad (182)$$

So

$$\begin{aligned} \Delta u + k^2u \\ = hg_{xx} + 2l_1g_xh'_w + l_1^2gh''_w + hg_{yy} + 2l_2g_yh'_w + l_2^2gh''_w + hg_{zz} + 2l_3g_zh'_w + l_3^2gh''_w + k^2gh = 0. \end{aligned}$$

Namely

$$(l_1^2 + l_2^2 + l_3^2)gh''_w + 2(l_1g_x + l_2g_y + l_3g_z)h'_w + (g_{xx} + g_{yy} + g_{zz} + k^2g)h = 0. \quad (183)$$

Set $h(w)$ an arbitrary unary second differentiable function, according to (183) we obtain

$$l_1^2 + l_2^2 + l_3^2 = 0 \implies l_1 = \pm\sqrt{-l_2^2 - l_3^2}, \quad (184)$$

$$l_1g_x + l_2g_y + l_3g_z = 0, \quad (185)$$

$$g_{xx} + g_{yy} + g_{zz} + k^2g = 0. \quad (186)$$

By (178), a particular solution of Eq. (186) is

$$g = (k_1\cosh\alpha x + k_2\sinh\alpha x)(k_3\cosh\beta y + k_4\sinh\beta y)(k_5\cosh\gamma z + k_6\sinh\gamma z), \quad k^2 = -\alpha^2 - \beta^2 - \gamma^2, \quad (187)$$

Substituting (187) into (185)

$$\begin{aligned} l_1g_x + l_2g_y + l_3g_z \\ = \alpha l_1(k_2\cosh\alpha x + k_1\sinh\alpha x)(k_3\cosh\beta y + k_4\sinh\beta y)(k_5\cosh\gamma z + k_6\sinh\gamma z) \\ + \beta l_2(k_1\cosh\alpha x + k_2\sinh\alpha x)(k_4\cosh\beta y + k_3\sinh\beta y)(k_5\cosh\gamma z + k_6\sinh\gamma z) \\ + \gamma l_3(k_1\cosh\alpha x + k_2\sinh\alpha x)(k_3\cosh\beta y + k_4\sinh\beta y)(k_6\cosh\gamma z + k_5\sinh\gamma z) = 0, \end{aligned}$$

we get

$$k_1 = k_2, k_3 = k_4, k_5 = k_6, \quad l_1 = \frac{-\beta l_2 - \gamma l_3}{\alpha}. \quad (188)$$

Since

$$\begin{aligned} l_1 &= \pm\sqrt{-l_2^2 - l_3^2} = \frac{-\beta l_2 - \gamma l_3}{\alpha} \\ \implies \left(\sqrt{k^2 + \gamma^2}l_2 - \frac{\beta\gamma l_3}{\sqrt{k^2 + \gamma^2}} \right)^2 + \left(k^2 + \beta^2 - \frac{\beta^2\gamma^2}{k^2 + \gamma^2} \right) l_3^2 &= 0 \\ \implies \sqrt{k^2 + \gamma^2}l_2 - \frac{\beta\gamma l_3}{\sqrt{k^2 + \gamma^2}} = 0, \quad k^2 + \beta^2 - \frac{\beta^2\gamma^2}{k^2 + \gamma^2} &= 0 \\ \implies l_2 = \frac{\beta\gamma l_3}{k^2 + \gamma^2}, \quad \beta = \pm\sqrt{-k^2 - \gamma^2}, \quad \alpha &= 0. \end{aligned}$$

For $\alpha = 0$, we have

$$k_3 = k_4, k_5 = k_6, l_3 = \frac{-\beta l_2}{\gamma}, k^2 = -\beta^2 - \gamma^2, \quad (189)$$

and

$$k^2 = -\beta^2 - \gamma^2 \implies \beta = \pm\sqrt{-k^2 - \gamma^2} \implies l_1 = \pm\sqrt{-l_2^2 - l_3^2} = \pm\sqrt{-l_2^2 - \frac{\beta^2 l_2^2}{\gamma^2}} = \pm\frac{kl_2}{\gamma}$$

Namely

$$\beta = \pm\sqrt{-k^2 - \gamma^2}, \alpha = 0, l_1 = \pm\frac{kl_2}{\gamma} \quad (190)$$

We set $\omega = \sqrt{-k^2 - \gamma^2}$, then

$$\begin{aligned} u(x, y, z) &= g(x, y, z) h(w) \\ &= (k_2 \cosh \alpha x + k_1 \sinh \alpha x) (k_3 \cosh \beta y + k_4 \sinh \beta y) (k_5 \cosh \gamma z + k_6 \sinh \gamma z) h(l_1 x + l_2 y + l_3 z + l_4) \\ &= (\cosh \beta y + \sinh \beta y) (\cosh \gamma z + \sinh \gamma z) h(l_1 x + l_2 y + l_3 z + l_4) = e^{\pm \omega y + \gamma z} h(l_1 x + l_2 y + l_3 z + l_4). \end{aligned}$$

So

$$\begin{aligned} u(x, y, z) &= e^{\omega y + \gamma z} (h_1(kx + \gamma y + \omega z + C_1) + h_2(-kx + \gamma y + \omega z + C_2) + h_3(kx + \gamma y - \omega z + C_3) \\ &+ h_4(-kx + \gamma y - \omega z + C_4)) + e^{-\omega y + \gamma z} (h_5(kx + \gamma y + \omega z + C_5) + h_6(-kx + \gamma y + \omega z + C_6) \\ &+ h_7(kx + \gamma y - \omega z + C_7) + h_8(-kx + \gamma y - \omega z + C_8)) \end{aligned} \quad (191)$$

where $h_1 - h_8$ are arbitrary unary second differentiable functions, $C_1 - C_8$ are arbitrary constants. Substituting (191) into Eq. (175), we find the necessary to set $h_1 + h_2 = C_9$ and $h_7 + h_8 = C_{10}$, where C_9 and C_{10} are arbitrary constants, so the verified new general solution of Eq. (175) is:

$$\begin{aligned} u(x, y, z) &= e^{\omega y + \gamma z} (h_3(kx + \gamma y - \omega z + C_3) + h_4(-kx + \gamma y - \omega z + C_4) + C_9) \\ &+ e^{-\omega y + \gamma z} (h_5(kx + \gamma y + \omega z + C_5) + h_6(-kx + \gamma y + \omega z + C_6) + C_{10}). \end{aligned}$$

In order to facilitate the writing and application, we rewrite the above equation

$$\begin{aligned} u(x, y, z) &= e^{\omega y + \gamma z} (h_1(kx + \gamma y - \omega z + C_1) + h_2(-kx + \gamma y - \omega z + C_2) + C_5) \\ &+ e^{-\omega y + \gamma z} (h_3(kx + \gamma y + \omega z + C_3) + h_4(-kx + \gamma y + \omega z + C_4) + C_6). \end{aligned} \quad (192)$$

where $h_1 - h_4$ are arbitrary unary second differentiable functions, $C_1 - C_6$ are arbitrary constants.

Note that the number of arbitrary functions in (192) is four! Namely the number of arbitrary functions in the SGS of 2th-order linear PDE (LPDE) is more than 2!

For the modified Helmholtz equation

$$\Delta u - k^2 u = 0. \quad (193)$$

Similar to the calculation of (101), the SGS of Eq. (193) is

$$\begin{aligned}
u = & \sin \left(\frac{C_1 - ik(k_2l_2 + k_3l_3)x + ik\sqrt{-l_2^2 - l_3^2}(k_2y + k_3z)}{\sqrt{(k_2l_2 + k_3l_3)^2 - (k_2^2 + k_3^2)(l_2^2 + l_3^2)}} \right) \\
& h_1 \left(\sqrt{-l_2^2 - l_3^2}x + l_2y + l_3z + l_4 \right) \\
& + \sin \left(\frac{C_2 + ik(k_{12}l_{12} + k_{13}l_{13})x - ik\sqrt{-l_{12}^2 - l_{13}^2}(k_{12}y + k_{13}z)}{\sqrt{(k_{12}l_{12} + k_{13}l_{13})^2 - (k_{12}^2 + k_{13}^2)(l_{12}^2 + l_{13}^2)}} \right) \\
& h_2 \left(\sqrt{-l_{12}^2 - l_{13}^2}x + l_{12}y + l_{13}z + l_{14} \right) \\
& + \sin \left(\frac{C_3 - ik(k_{22}l_{22} + k_{23}l_{23})x - ik\sqrt{-l_{22}^2 - l_{23}^2}(k_{22}y + k_{23}z)}{\sqrt{(k_{22}l_{22} + k_{23}l_{23})^2 - (k_{22}^2 + k_{23}^2)(l_{22}^2 + l_{23}^2)}} \right) \\
& h_3 \left(-\sqrt{-l_{22}^2 - l_{23}^2}x + l_{22}y + l_{23}z + l_{24} \right) \\
& + \sin \left(\frac{C_4 + ik(k_{32}l_{32} + k_{33}l_{33})x + ik\sqrt{-l_{32}^2 - l_{33}^2}(k_{32}y + k_{33}z)}{\sqrt{(k_{32}l_{32} + k_{33}l_{33})^2 - (k_{32}^2 + k_{33}^2)(l_{32}^2 + l_{33}^2)}} \right) \\
& h_4 \left(-\sqrt{-l_{32}^2 - l_{33}^2}x + l_{32}y + l_{33}z + l_{34} \right),
\end{aligned} \tag{193}$$

where h_1-h_4 are arbitrary unary second differentiable functions, $k_2, k_3, k_{12}, k_{13}, k_{22}, k_{23}, k_{32}, k_{33}, l_2-l_4, l_{12}-l_{14}, l_{22}-l_{24}, l_{32}-l_{34}$ and C_1-C_4 are arbitrary constants. Note that the number of arbitrary functions in (194) is four, the SGS of Eq. (193) which similar to (192) is more complex.

According to the above calculation, we propose a new problem: Comparing (192) with (177), whether they are independent of each other? How to prove?

In the theory of ordinary differential equations (ODEs), it has been proven that the number of the arbitrary constants in the general solution of m th-order ODEs is m [67, 68]. In the theory of PDEs, almost all of the textbooks and professional books directly or indirectly declare that the number of the arbitrary functions in the general solution of m th-order PDEs is m [20, 64, 69, 70], but no related rigorous proof up to now, so the above problem is very important, it relates to how many arbitrary functions in the SGS of m th-order PDEs. By the law of superposition, if (192) and (177) are independent, the number of arbitrary functions in the SGS of Eq. (175) is six.

Because the concise general solution of a PDE may not be the only, we can use symbols G_n^m express these different solution, where n is the number of arbitrary functions and m is the number of discretionary parameters in the corresponding concise general solution.

2. Principles of transformational equations

We presents five new concepts and five new theorems in this chapter, using Theorem 1 if the solution of a PDE is known, the solutions of its various independent variable transformational equations (IVTEs) can be obtained directly. According to Theorem 2 we can use two solvable equations to obtain a new solvable PDE. By Theorem 4 and 5, the general solutions of the vector wave equation in cartesian, cylindrical and spherical coordinate systems have been solved for the first time. We point out that the general solutions or particular solutions of various symmetric vector partial differential equations can be obtained similarly in any orthogonal coordinate

system, such as vector Helmholtz equation, vector Laplace equation and the magnetic vector potential equation and so on.

2.1. The principle of independent variable transformational equations

Many MPEs' solutions under non-Cartesian coordinate system have been investigated deeply [71-73]. One PDE in different coordinate systems has different forms, such as the Laplace equation in spherical coordinates and cylindrical coordinates are written as

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial u}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial u}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 u}{\partial \varphi^2} = 0, \quad (195)$$

$$\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial u}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2} = 0. \quad (196)$$

Here we begin to study the relationship law of the PDEs' solutions in different coordinate systems. First, we define independent variable transformational equations (IVTEs).

Definition. In the domain $D, (D \subset \mathbb{R}^n), (n \geq 2)$, set x_i are independent and y_i are independent too, ($i = 1, 2, \dots, n$), by $x_i = x_i(y_1, y_2, \dots, y_n), y_i = y_i(x_1, x_2, \dots, x_n)$, which are known, m th-order PDE $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$ is converted into another m th-order PDE $G(y_1, \dots, y_n, u, u_{y_1}, \dots, u_{y_n}, u_{y_1 y_2}, \dots) = 0$, then $G = 0$ is called a IVTE of $F = 0$.

Since the concrete forms of $x_i = x_i(y_1, y_2, \dots, y_n), y_i = y_i(x_1, x_2, \dots, x_n)$ are infinite, so every PDE has infinite IVTEs. In the orthogonal coordinate system theory the forms of IVTEs are always obtained by the metric and exterior calculus method, in fact, they could also be get by known $x_i = x_i(y_1, \dots, y_n)$ and $y_i = y_i(x_1, \dots, x_n)$, such as the Laplace equation in cylindrical coordinates, using

$$x = r \cos \theta, y = r \sin \theta, z = z. \quad (197)$$

We obtain

$$r = \sqrt{x^2 + y^2}, \theta = \arctan \frac{y}{x}, z = z, \quad (198)$$

and further

$$\begin{aligned} r_x &= \frac{x}{\sqrt{x^2 + y^2}} = \cos \theta, r_y = \frac{y}{\sqrt{x^2 + y^2}} = \sin \theta, r_z = 0 \\ \theta_x &= \frac{-y}{x^2 + y^2} = \frac{-\sin \theta}{r}, \theta_y = \frac{x}{x^2 + y^2} = \frac{\cos \theta}{r}, \theta_z = 0, z_x = z_y = 0 \\ u_x &= u_r r_x + u_\theta \theta_x + u_z z_x = u_r \cos \theta - u_\theta \frac{\sin \theta}{r} \\ u_{xx} &= u_{rr} r_x \cos \theta - u_r \sin \theta \theta_x - u_{\theta\theta} \theta_x \frac{\sin \theta}{r} - u_\theta \frac{\cos \theta}{r} \theta_x + u_\theta \frac{\sin \theta}{r^2} r_x \\ &= u_{rr} \cos^2 \theta + u_r \frac{\sin^2 \theta}{r} + u_{\theta\theta} \frac{\sin^2 \theta}{r^2} + 2u_\theta \frac{\sin \theta \cos \theta}{r^2} \\ u_y &= u_r r_y + u_\theta \theta_y + u_z z_y = u_r \sin \theta + u_\theta \frac{\cos \theta}{r} \\ u_{yy} &= u_{rr} r_y \sin \theta + u_r \cos \theta \theta_y + u_{\theta\theta} \theta_y \frac{\cos \theta}{r} - u_\theta \frac{\sin \theta}{r} \theta_y - u_\theta \frac{\cos \theta}{r^2} r_y \\ &= u_{rr} \sin^2 \theta + u_r \frac{\cos^2 \theta}{r} + u_{\theta\theta} \frac{\cos^2 \theta}{r^2} - 2u_\theta \frac{\sin \theta \cos \theta}{r^2}. \end{aligned}$$

Then

$$\begin{aligned}
& u_{xx} + u_{yy} + u_{zz} \\
&= u_{rr} \cos^2 \theta + u_r \frac{\sin^2 \theta}{r} + u_{\theta\theta} \frac{\sin^2 \theta}{r^2} + 2u_\theta \frac{\sin \theta \cos \theta}{r^2} + u_{rr} \sin^2 \theta + u_r \frac{\cos^2 \theta}{r} + u_{\theta\theta} \frac{\cos^2 \theta}{r^2} \\
&- 2u_\theta \frac{\sin \theta \cos \theta}{r^2} + u_{zz} = u_{rr} + \frac{1}{r} u_r + \frac{1}{r^2} u_{\theta\theta} + u_{zz} = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial u}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2}.
\end{aligned}$$

Namely

$$u_{xx} + u_{yy} + u_{zz} = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial u}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2}.$$

Using the definition of IVTEs we present Theorem 1.

Theorem 1. In the domain $D, (D \subset \mathbb{R}^n)$, if the solution $u = f(x_1, \dots, x_n)$ of a m th-order PDE $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$ is known, then the solution of its IVTE $G(y_1, \dots, y_n, u, u_{y_1}, \dots, u_{y_n}, u_{y_1 y_2}, \dots) = 0$ is $u = f(x_1, \dots, x_n) = g(y_1, \dots, y_n)$.

Proof. Using $x_i = x_i(y_1, \dots, y_n)$ and $y_i = y_i(x_1, \dots, x_n)$, $G(y_1, \dots, y_n, u, u_{y_1}, \dots, u_{y_n}, u_{y_1 y_2}, \dots) = 0$ could be converted into $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$, because the solution $u = f(x_1, \dots, x_n)$ of $F = 0$ is known, so $u = f(x_1, \dots, x_n)$ is the solution of $G = 0$ too, by $x_i = x_i(y_1, \dots, y_n)$, $u = f(x_1, \dots, x_n)$ may be varied into $u = g(y_1, \dots, y_n)$.

According to Theorem 1, if the solution of a PDE is known, the solutions of its various IVTEs can be obtained directly, as in spherical coordinates

$$x = r \sin \theta \cos \varphi, y = r \sin \theta \sin \varphi, z = r \cos \theta. \quad (199)$$

By the general solution (16) of Laplace equation in the Cartesian coordinate system, its general solution in the spherical coordinate system can be got directly

$$\begin{aligned}
u(r, \theta, \varphi) &= f_1 \left(r \sin \theta \cos \varphi \sqrt{-k_1^2 - k_2^2} + k_1 r \sin \theta \sin \varphi + k_2 r \cos \theta + k_3 \right) \\
&+ f_2 \left(-r \sin \theta \cos \varphi \sqrt{-k_4^2 - k_5^2} + k_4 r \sin \theta \sin \varphi + k_5 r \cos \theta + k_6 \right) \\
&+ k_7 r \sin \theta \cos \varphi + k_8 r \sin \theta \sin \varphi + k_9 r \cos \theta + k_{10}.
\end{aligned} \quad (200)$$

Using (197) the general solution of Laplace equation in the cylindrical coordinate system is

$$\begin{aligned}
u(r, \theta, z) &= f_1 \left(r \cos \theta \sqrt{-k_1^2 - k_2^2} + k_1 r \sin \theta + k_2 z + k_3 \right) \\
&+ f_2 \left(-r \cos \theta \sqrt{-k_4^2 - k_5^2} + k_4 r \sin \theta + k_5 z + k_6 \right) \\
&+ k_7 r \cos \theta + k_8 r \sin \theta + k_9 z + k_{10}.
\end{aligned} \quad (201)$$

All the solutions of PDEs obtained in this paper, we can use them to obtain the solutions in all orthogonal coordinate system by Theorem 1, exact solutions of Cauchy problems can be similarly analysed yet.

2.2. The principle of dependent variable transformational equations

First, we define the dependent variable transformational equations (DVTEs).

Definition. In the domain $D, (D \subset \mathbb{R}^n)$, set $v = h(x_1, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots)$, so a PDE

$F(x_1, \dots, x_n, v, v_{x_1}, \dots, v_{x_n}, v_{x_1 x_2}, \dots) = 0$ may be converted into $G(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$, then $G = 0$ is called a DVTE of $F = 0$.

Using the definition of DVTE we present Theorem 2.

Theorem 2. In the domain $D, (D \subset \mathbb{R}^n)$, if the solution $v = f(x_1, \dots, x_n)$ of a PDE $F(x_1, \dots, x_n, v, v_{x_1}, \dots, v_{x_n}, v_{x_1 x_2}, \dots) = 0$ is known, set $v = h(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots)$, then the solution of its DVTE $G(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$ is the solution of $h(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = f(x_1, \dots, x_n)$.

Proof. Set $v = h(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots)$, then $G(y_1, \dots, y_n, u, u_{y_1}, \dots, u_{y_n}, u_{y_1 y_2}, \dots) = 0$ could be converted into $F(x_1, \dots, x_n, v, v_{x_1}, \dots, v_{x_n}, v_{x_1 x_2}, \dots) = 0$, because the solution $v = f(x_1, \dots, x_n)$ of $F = 0$ is known, so if the solution $u = g(x_1, \dots, x_n)$ of $h(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = f(x_1, \dots, x_n)$ can be solved, then $u = g(x_1, \dots, x_n)$ is the solution of $G = 0$ yet.

Essentially, Theorem 2 uses two solvable equations to obtain a new solvable PDE, the two solvable equations are

$$F(x_1, \dots, x_n, v, v_{x_1}, \dots, v_{x_n}, v_{x_1 x_2}, \dots) = 0, \quad (202)$$

$$h(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = f(x_1, x_2, \dots, x_n) = v. \quad (203)$$

The new solvable PDE which is the DVTE is

$$G(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0. \quad (204)$$

Eq. (202) may be a solvable PDE or ODE, Eq. (203) may be a solvable PDE, ODE or function equation, so the concretely using methods of Theorem 2 may be infinite. Actually, Transformational Method 1-4 are specific applications of Theorem 2. Here we use Theorem 2 to get the general solution of two linear PDEs first, in Chapter 4, we'll use it to solve the general solution of two nonlinear PDEs.

Example 2.1

$$u_{xy} + a(x, y)u_x + b(x, y)u_y + (a_x(x, y) + b(x, y)a(x, y))u = c(x, y), \quad (205)$$

where a, b, c are any known binary functions, Eq. (205) is a second order linear hyperbolic PDE, due to

$$v_x + b(x, y)v = c(x, y), \quad (206)$$

the general solution of Eq. (206) is [74]:

$$v(x, y) = e^{-\int b(x, y)dx} \left(\varphi(y) + \int c(x, y) e^{\int b(x, y)dx} dx \right), \quad (207)$$

where $\varphi(y)$ is an arbitrary unary function, by Theorem 2 we set

$$v(x, y) = u_y + a(x, y)u. \quad (208)$$

According to (207) we get

$$\begin{aligned} u(x, y) &= e^{-\int a(x, y)dy} \left(\phi(x) + \int v(x, y) e^{\int a(x, y)dy} dy \right) \\ &= e^{-\int a(x, y)dy} \left(\phi(x) + \int \left(e^{-\int b(x, y)dx} \left(\varphi(y) + \int c(x, y) e^{\int b(x, y)dx} dx \right) \right) e^{\int a(x, y)dy} dy \right), \end{aligned}$$

where $\phi(x)$ is an arbitrary unary function, by (208) we could get that Eq. (205) is a DVTE of Eq. (206), according to Theorem 2, the general solution of Eq. (205) is

$$u = e^{-\int a(x,y)dy} \left(\phi(x) + \int \left(e^{-\int b(x,y)dx} \left(\varphi(y) + \int c(x,y) e^{\int b(x,y)dx} dx \right) \right) e^{\int a(x,y)dy} dy \right). \quad (209)$$

Example 2.2

$$a_y u_{xy} + a_x u_{yy} + a_y b u_x + a_x b u_y + (a_y b_x + a_x b_y) u = 0, \quad (210)$$

where a,b are any known binary functions, due to

$$a_y v_x + a_x v_y = 0, \quad (211)$$

the general solution of Eq. (211) is [74]:

$$v = g(a(x,y)), \quad (212)$$

where g is an arbitrary unary first differentiable function, by Theorem 2 we set

$$v(x,y) = u_y + b(x,y)u. \quad (213)$$

According to (207) and (212) we get

$$\begin{aligned} u(x,y) &= e^{-\int b(x,y)dy} \left(\phi(x) + \int v(x,y) e^{\int a(x,y)dy} dy \right) \\ &= e^{-\int b(x,y)dy} \left(\phi(x) + \int g(a(x,y)) e^{\int b(x,y)dy} dy \right), \end{aligned}$$

where $\phi(x)$ is an arbitrary unary function, by (213) we could get that Eq. (210) is a DVTE of Eq. (211), according to Theorem 2, the general solution of Eq. (210) is

$$u(x,y) = e^{-\int b(x,y)dy} \left(\phi(x) + \int g(a(x,y)) e^{\int b(x,y)dy} dy \right). \quad (214)$$

The definition and rule of DVTEs can be extended to ODEs.

Definition. In the domain D , ($D \subset \mathbb{R}^1$), set $w = h(x, y, y', y'', \dots, y^{(m)})$, so an n th-order ODE $F(x, w, w', w'', \dots, w^{(n)}) = 0$ may be converted into an $n+m$ th-order ODE $G(x, y, y', y'', \dots, y^{(m+n)}) = 0$ then $G = 0$ is called a DVTE of $F = 0$.

Theorem 3. In the domain D , ($D \subset \mathbb{R}^1$), if the solution $w = f(x)$ of an ODE $F(x, w, w', w'', \dots, w^{(n)}) = 0$ is known, set $w = h(x, y, y', y'', \dots, y^{(m)})$, then the solution of its DVTE $G(x, y, y', y'', \dots, y^{(m+n)}) = 0$, is the solution of $h(x, y, y', y'', \dots, y^{(m)}) = f(x)$.

For the theoretical system' completeness we present Theorem 3, the proof method of Theorem 3 is similar to Theorem 2. In fact, some ODEs have been solved by Theorem 3 [53], so here we will not study and example.

2.3. General Solutions of Vector Equation in Various Orthogonal Coordinate Systems

In \mathbb{R}^n space ($n \geq 2$), the expression of a vector function \mathbf{u} is

$$u = u_1 e_1 + u_2 e_2 + \dots + u_n e_n, \quad (215)$$

where $u_i = u_i(x_1, x_2, \dots, x_n)$, $i = (1, 2, \dots, n)$, Suppose \mathbf{u} satisfies the PDE

$$F \left(x_1, \dots, x_n, u, \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}, \frac{\partial^2 u}{\partial x_1 \partial x_2}, \dots \right) = 0. \quad (216)$$

If Eq. (216) satisfies

$$\begin{aligned} & F \left(x_1, \dots, x_n, u, \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}, \frac{\partial^2 u}{\partial x_1 \partial x_2}, \dots \right) \\ &= \sum_{i=1}^n F_i \left(x_1, \dots, x_n, u_i, \frac{\partial u_i}{\partial x_1}, \dots, \frac{\partial u_i}{\partial x_n}, \frac{\partial^2 u_i}{\partial x_1 \partial x_2}, \dots \right) e_i = 0. \end{aligned} \quad (217)$$

That is, only the dependent variable u_i in the i th component equation, and the form of each component $F_i \left(x_1, \dots, x_n, u_i, \frac{\partial u_i}{\partial x_1}, \dots, \frac{\partial u_i}{\partial x_n}, \frac{\partial^2 u_i}{\partial x_1 \partial x_2}, \dots \right)$ in Eq. (217) are the same, we call the vector PDEs as the **symmetric vector partial differential equations** (SVPDEs).

By Maxwell equations we can get the vector wave equation which represents the electromagnetic wave spreading in the vacuum [75]

$$\Delta u - \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = 0, \quad (218)$$

where c is the speed of light, \mathbf{u} is the electric-field strength \mathbf{E} or magnetic induction \mathbf{B} , in 3-dimensional space

$$u = u_x e_x + u_y e_y + u_z e_z, \quad (219)$$

$$\Delta u - \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = \left(\Delta u_x - \frac{1}{c^2} \frac{\partial^2 u_x}{\partial t^2} \right) e_x + \left(\Delta u_y - \frac{1}{c^2} \frac{\partial^2 u_y}{\partial t^2} \right) e_y + \left(\Delta u_z - \frac{1}{c^2} \frac{\partial^2 u_z}{\partial t^2} \right) e_z = 0, \quad (220)$$

the form of each component $F_i \left(x, y, z, u_i, \frac{\partial u_i}{\partial x}, \dots, \frac{\partial u_i}{\partial z}, \frac{\partial^2 u_i}{\partial x \partial x}, \dots \right)$ in Eq. (220) are the same, so Eq. (218) is a SVPDEs. Another example is that time harmonic electromagnetic waves distribution in space satisfies the vector Helmholtz equation

$$\Delta u + k^2 u = 0, \quad (221)$$

where k is a constant related to the medium and the frequency of electromagnetic wave, \mathbf{u} is the electric-field strength \mathbf{E} or magnetic induction \mathbf{B} , for

$$\Delta u + k^2 u = (\Delta u_x + k^2 u_x) e_x + (\Delta u_y + k^2 u_y) e_y + (\Delta u_z + k^2 u_z) e_z = 0. \quad (222)$$

the form of each component F_i in Eq. (222) are the same, so it is a SVPDEs too. A vector PDE may be a SVPDE in a certain orthogonal coordinate system and may not in other orthogonal coordinate system, such as the form of vector Helmholtz equation in the cylindrical coordinate system is [20]

$$\left(\Delta u_r - \frac{u_r}{r^2} - \frac{2}{r^2} \frac{\partial u_\theta}{\partial \theta} + k^2 u_r \right) e_r + \left(\Delta u_\theta - \frac{u_\theta}{r^2} + \frac{2}{r^2} \frac{\partial u_r}{\partial \theta} + k^2 u_\theta \right) e_\theta + (\Delta u_z + k^2 u_z) e_z = 0. \quad (223)$$

Obviously Eq. (223) is not a SVPDE.

Any a n -dimensional SVPDE $F = 0$ can be decomposed into n independent equations $F_i = 0$, ($i = 1, 2, \dots, n$), we call $F_i = 0$ the **corresponding scalar equation** (CSE) of $F = 0$,




Figure 1.1:




Figure 1.2:

due to each F_i in $F = 0$ has the same form, so every SVPDE has a unique CSE. The CSE of Eq. (218) is:

$$\Delta u_i - \frac{1}{c^2} \frac{\partial^2 u_i}{\partial t^2} = 0 \quad (224)$$

The CSE of Eq. (221) is

$$\Delta u_i + k^2 u_i = 0 \quad (225)$$

Here we propose a new theorem based on the above two new definitions.

Theorem 4. If there are m arbitrary functions in the general solution of the CSE $F_i = 0$, then the number of arbitrary functions in the general solution of the n -dimensional SVPDE $F = 0$ is mn

Proof. Because a n -dimensional SVPDE $F = 0$ can be decomposed into n independent equations $F_i = 0$, ($i = 1, 2, \dots, n$), and each $F_i = 0$ has the same form, so the numbers of arbitrary functions in the general solution of every $F_i = 0$ are all m , according to Eq. (215) we can obtain the number of arbitrary functions in the general solution of $F = 0$ is mn .

Now we begin to analyze the solution law of vector PDEs in various orthogonal coordinate system. By exterior differential and the form in Cartesian coordinate system, the form of a vector PDE in various orthogonal coordinate systems can be calculated [20]. In this paper, we propose a new simple method which can not only easily calculate the form of a vector PDE in any orthogonal coordinate systems, and can directly get its solutions in all kinds of orthogonal coordinate systems by the solution of any an orthogonal coordinate system, therefore it has great advantages.

According to the definition of different orthogonal coordinate system and the magnitude of every unit vector is equal to 1, we can write out the mathematical relationship between the unit vectors of different orthogonal coordinate system, such as the relation between the unit vector e_r, e_θ, e_z of cylinder coordinate system and the unit vectors e_x, e_y, e_z of Cartesian coordinate system, for

$$e_r = \cos\theta e_x + \sin\theta e_y, e_\theta = -\sin\theta e_x + \cos\theta e_y, e_z = e_z. \quad (226)$$

By Eq. (226) we can get

$$e_x = \cos\theta e_r - \sin\theta e_\theta, e_y = \sin\theta e_r + \cos\theta e_\theta, e_z = e_z. \quad (227)$$

For the spherical coordinate system

From Figure 1.1 we get

$$e_r = \sin\theta \cos\varphi e_x + \sin\theta \sin\varphi e_y + \cos\theta e_z \quad (228)$$

$$e_\varphi = -\sin\varphi e_x + \cos\varphi e_y \quad (229)$$

According to Figure 1.2

$$e_\theta = \cos\theta \cos\varphi e_x + \cos\theta \sin\varphi e_y - \sin\theta e_z. \quad (230)$$

By further calculation we have

$$e_x = \cos\varphi \sin\theta e_r + \cos\varphi \cos\theta e_\theta - \sin\varphi e_\varphi \quad (231)$$

$$e_y = \sin\varphi \sin\theta e_r + \sin\varphi \cos\theta e_\theta + \cos\varphi e_\varphi \quad (232)$$

$$e_z = \cos\theta e_r - \sin\theta e_\theta \quad (233)$$

By the functional relationship between the unit vector in different coordinate systems, the function relation of dependent variable components in diversified coordinate systems can be deduced, and the form of a vector PDE in various coordinate systems can be got too, such as the cylindrical coordinate system

$$\begin{aligned} u &= u_x e_x + u_y e_y + u_z e_z = u_x (\cos\theta e_r - \sin\theta e_\theta) + u_y (\sin\theta e_r + \cos\theta e_\theta) + u_z e_z \\ &= (u_x \cos\theta + u_y \sin\theta) e_r + (-u_x \sin\theta + u_y \cos\theta) e_\theta + u_z e_z = u_r e_r + u_\theta e_\theta + u_z e_z \end{aligned} \quad (234)$$

From Eq. (234) we get

$$u_r = u_x \cos\theta + u_y \sin\theta, u_\theta = -u_x \sin\theta + u_y \cos\theta \quad (235)$$

By Eq. (235) we can obtain

$$u_x = \cos\theta u_r - \sin\theta u_\theta, u_y = \sin\theta u_r + \cos\theta u_\theta \quad (236)$$

So

$$\begin{aligned} \Delta u &= (\Delta u_x) e_x + (\Delta u_y) e_y + (\Delta u_z) e_z \\ &= \left(\left(\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} + \frac{\partial^2}{\partial z^2} \right) (\cos\theta u_r - \sin\theta u_\theta) \right) (\cos\theta e_r - \sin\theta e_\theta) \\ &+ \left(\left(\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} + \frac{\partial^2}{\partial z^2} \right) (\sin\theta u_r + \cos\theta u_\theta) \right) (\sin\theta e_r + \cos\theta e_\theta) \\ &+ (\Delta u_z) e_z \\ &= (\cos^2\theta \Delta u_r - \frac{2\sin\theta \cos\theta}{r^2} \frac{\partial u_r}{\partial \theta} - \frac{\cos^2\theta u_r}{r^2} - \sin\theta \cos\theta \Delta u_\theta - \frac{2\cos^2\theta}{r^2} \frac{\partial u_\theta}{\partial \theta} + \frac{\sin\theta \cos\theta u_\theta}{r^2} \\ &+ \sin^2\theta \Delta u_r + \frac{2\sin\theta \cos\theta}{r^2} \frac{\partial u_r}{\partial \theta} - \frac{\sin^2\theta u_r}{r^2} + \sin\theta \cos\theta \Delta u_\theta - \frac{2\sin^2\theta}{r^2} \frac{\partial u_\theta}{\partial \theta} - \frac{\sin\theta \cos\theta u_\theta}{r^2}) e_r \\ &+ (-\sin\theta \cos\theta \Delta u_r + \frac{2\sin^2\theta}{r^2} \frac{\partial u_r}{\partial \theta} + \frac{\sin\theta \cos\theta u_r}{r^2} + \sin^2\theta \Delta u_\theta + \frac{2\sin\theta \cos\theta}{r^2} \frac{\partial u_\theta}{\partial \theta} - \frac{\sin^2\theta u_\theta}{r^2} \\ &+ \sin\theta \cos\theta \Delta u_r + \frac{2\cos^2\theta}{r^2} \frac{\partial u_r}{\partial \theta} - \frac{\sin\theta \cos\theta u_r}{r^2} + \cos^2\theta \Delta u_\theta - \frac{2\sin\theta \cos\theta}{r^2} \frac{\partial u_\theta}{\partial \theta} - \frac{\cos^2\theta u_\theta}{r^2}) e_\theta \\ &+ (\Delta u_z) e_z \end{aligned}$$

Namely

$$u = \left(\Delta u_r - \frac{\Delta u_r}{r^2} - \frac{2}{r^2} \frac{\partial u_\theta}{\partial \theta} \right) e_r + \left(\Delta u_\theta - \frac{u_\theta}{r^2} + \frac{2}{r^2} \frac{\partial u_r}{\partial \theta} \right) e_\theta + (\Delta u_z) e_z \quad (237)$$

Similar to the independent variable transformational equations (IVTE) presented in [10], essentially the vector PDE in different coordinate systems are the IVTE each other. Therefore,

we propose the concept of an independent variable transformation vector equation (IVTVE).

Definition. In the domain $D, (D \subset \mathbb{R}^n)(n \geq 2)$, set x_i are independent and y_i are independent too, $(i = 1, 2, \dots, n)$, by $x_i = x_i(y_1, y_2, \dots, y_n), y_i = y_i(x_1, x_2, \dots, x_n), e_{x_i} = f_i(y_1, y_2, \dots, y_n, e_{y_1}, e_{y_2}, \dots, e_{y_n})$ and $e_{y_i} = g_i(x_1, x_2, \dots, x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n})$ which are known, m th-order vector PDE $F(x_1, \dots, x_n, u, \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}, \frac{\partial^2 u}{\partial x_1 \partial x_2}, \dots) = 0$ can be converted into another m th-order vector PDE $G(y_1, \dots, y_n, u, \frac{\partial u}{\partial y_1}, \dots, \frac{\partial u}{\partial y_n}, \frac{\partial^2 u}{\partial y_1 \partial y_2}, \dots) = 0$, then $G = 0$ is called a IVTVE of $F = 0$.

According to the define of IVTVE we propose theorem 5.

Theorem 5. In the domain $D, (D \subset \mathbb{R}^n)$, if the solution $u = f(x_1, x_2, \dots, x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n})$ of a m th-order vector PDE $F(x_1, \dots, x_n, u, \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}, \frac{\partial^2 u}{\partial x_1 \partial x_2}, \dots) = 0$ is known, then the solution of its IVTVE $G(y_1, \dots, y_n, u, \frac{\partial u}{\partial y_1}, \dots, \frac{\partial u}{\partial y_n}, \frac{\partial^2 u}{\partial y_1 \partial y_2}, \dots) = 0$, is $u = f(x_1, x_2, \dots, x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n}) = g(y_1, y_2, \dots, y_n, e_{y_1}, e_{y_2}, \dots, e_{y_n})$.

Proof. Using $x_i = x_i(y_1, y_2, \dots, y_n), y_i = y_i(x_1, x_2, \dots, x_n), e_{x_i} = f_i(y_1, y_2, \dots, y_n, e_{y_1}, e_{y_2}, \dots, e_{y_n})$ and $e_{y_i} = g_i(x_1, x_2, \dots, x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n}), G(y_1, \dots, y_n, u, \frac{\partial u}{\partial y_1}, \dots, \frac{\partial u}{\partial y_n}, \frac{\partial^2 u}{\partial y_1 \partial y_2}, \dots) = 0$, could be converted into $F(x_1, \dots, x_n, u, \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}, \frac{\partial^2 u}{\partial x_1 \partial x_2}, \dots) = 0$ because the solution $u = f(x_1, x_2, \dots, x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n})$ of $F = 0$ is known, so $u = f(x_1, x_2, \dots, x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n})$ is the solution of $G = 0$ too, by $x_i = x_i(y_1, y_2, \dots, y_n)$, and $e_{x_i} = f_i(y_1, y_2, \dots, y_n, e_{y_1}, e_{y_2}, \dots, e_{y_n})$, $u = f(x_1, x_2, \dots, x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n})$ may be varied into $u = g(y_1, y_2, \dots, y_n, e_{y_1}, e_{y_2}, \dots, e_{y_n})$.

Using theorem 5, we can solve the general solutions of vector wave equation and Helmholtz equation in Cartesian coordinate system, the CSE Eq. (224) of Eq. (218) is the acoustic equation, we had solved its general solution in Section 1.4

$$u_i = f_{1i} \left(k_{1i}x + k_{2i}y + k_{3i}z + ct\sqrt{k_{1i}^2 + k_{2i}^2 + k_{3i}^2 + k_{4i}} \right) + f_{2i} \left(k_{5i}x + k_{6i}y + k_{7i}z - ct\sqrt{k_{5i}^2 + k_{6i}^2 + k_{7i}^2 + k_{8i}} \right) + k_{9i}x + k_{10i}y + k_{11i}z + k_{12i}t + k_{13i} \quad (238)$$

where f_{1i} and f_{2i} are arbitrary unary second differentiable functions, $k_{1i} - k_{13i}$ are arbitrary parameters, $(i = x, y, z)$. Using Eq. (219) and (238), we can obtain the general solution of Eq. (218) with six arbitrary second differentiable functions

$$u = \sum_{i=x,y,z} \left(f_{1i} \left(k_{1i}x + k_{2i}y + k_{3i}z + ct\sqrt{k_{1i}^2 + k_{2i}^2 + k_{3i}^2 + k_{4i}} \right) + f_{2i} \left(k_{5i}x + k_{6i}y + k_{7i}z - ct\sqrt{k_{5i}^2 + k_{6i}^2 + k_{7i}^2 + k_{8i}} \right) + k_{9i}x + k_{10i}y + k_{11i}z + k_{12i}t + k_{13i} \right) e_i \quad (239)$$

It is necessary to pay an attention to that the wave velocity u_i are all the speed of light velocity c .

The general solution of the Helmholtz equation Eq. (225) is

$$\begin{aligned}
u_i = & \sin \left(\frac{k_{3i} - k(k_{1i}l_{1i} + k_{2i}l_{2i})x + k\sqrt{-l_{1i}^2 - l_{2i}^2}(k_{1i}y + k_{2i}z)}{\sqrt{(k_{1i}l_{1i} + k_{2i}l_{2i})^2 - (k_{1i}^2 + k_{2i}^2)(l_{1i}^2 + l_{2i}^2)}} \right) \\
& h_{1i} \left(\sqrt{-l_{1i}^2 - l_{2i}^2}x + l_{1i}y + l_{2i}z + l_{3i} \right) \\
& + \sin \left(\frac{k_{6i} - k(k_{4i}l_{4i} + k_{5i}l_{5i})x - k\sqrt{-l_{4i}^2 - l_{5i}^2}(k_{4i}y + k_{5i}z)}{\sqrt{(k_{4i}l_{4i} + k_{5i}l_{5i})^2 - (k_{4i}^2 + k_{5i}^2)(l_{4i}^2 + l_{5i}^2)}} \right) \\
& h_{2i} \left(-\sqrt{-l_{4i}^2 - l_{5i}^2}x + l_{4i}y + l_{5i}z + l_{6i} \right),
\end{aligned} \tag{240}$$

where h_{1i} and h_{2i} are arbitrary unary second differentiable functions, $k_{1i} - k_{6i}$ and $l_{1i} - l_{6i}$ are arbitrary constants. Using Eq. (219) and (240), we can obtain the general solution of the vector Helmholtz equation Eq. (221) with six arbitrary second differentiable functions

$$\begin{aligned}
u = & \sum_{i=x,y,z} \left(\sin \left(\frac{k_{3i} - k(k_{1i}l_{1i} + k_{2i}l_{2i})x + k\sqrt{-l_{1i}^2 - l_{2i}^2}(k_{1i}y + k_{2i}z)}{\sqrt{(k_{1i}l_{1i} + k_{2i}l_{2i})^2 - (k_{1i}^2 + k_{2i}^2)(l_{1i}^2 + l_{2i}^2)}} \right) \right. \\
& h_{1i} \left(\sqrt{-l_{1i}^2 - l_{2i}^2}x + l_{1i}y + l_{2i}z + l_{3i} \right) \\
& + \sin \left(\frac{k_{6i} - k(k_{4i}l_{4i} + k_{5i}l_{5i})x - k\sqrt{-l_{4i}^2 - l_{5i}^2}(k_{4i}y + k_{5i}z)}{\sqrt{(k_{4i}l_{4i} + k_{5i}l_{5i})^2 - (k_{4i}^2 + k_{5i}^2)(l_{4i}^2 + l_{5i}^2)}} \right) \\
& \left. h_{2i} \left(-\sqrt{-l_{4i}^2 - l_{5i}^2}x + l_{4i}y + l_{5i}z + l_{6i} \right) \right) e_i
\end{aligned} \tag{241}$$

The solutions of PDEs in different orthogonal coordinate systems have been the focus of research [71-73]. According to Theorem 5, if a solution of a vector PDE is known, the solutions of its various IVTVEs can be obtained directly, as in the cylindrical coordinate system

$$x = r\cos\theta, y = r\sin\theta, z = z. \tag{242}$$

Using Eq. (227), (239), (242) and Theorem 5, the general solution of vector wave equation in cylindrical coordinate system can be obtained directly

$$\begin{aligned}
u = & (f_{1x} + f_{2x} + g_x)(\cos\theta e_r - \sin\theta e_\theta) + (f_{1y} + f_{2y} + g_y)(\sin\theta e_r + \cos\theta e_\theta) + (f_{1z} + f_{2z} + g_z)e_z \\
= & (\cos\theta f_{1x} + \cos\theta f_{2x} + \cos\theta g_x + \sin\theta f_{1y} + \sin\theta f_{2y} + \sin\theta g_y)e_r \\
& + (-\sin\theta f_{1x} - \sin\theta f_{2x} - \sin\theta g_x + \cos\theta f_{1y} + \cos\theta f_{2y} + \cos\theta g_y)e_\theta + (f_{1z} + f_{2z} + g_z)e_z
\end{aligned}$$

Namely

$$\begin{aligned}
u = & (\cos\theta(f_{1x} + f_{2x} + g_x) + \sin\theta(f_{1y} + f_{2y} + g_y))e_r \\
& + (-\sin\theta(f_{1x} + f_{2x} + g_x) + \cos\theta(f_{1y} + f_{2y} + g_y))e_\theta + (f_{1z} + f_{2z} + g_z)e_z,
\end{aligned} \tag{243}$$

where

$$f_{1i} = f_{1i} \left(k_{1i}r\cos\theta + k_{2i}r\sin\theta + k_{3i}z + ct\sqrt{k_{1i}^2 + k_{2i}^2 + k_{3i}^2} + k_{4i} \right) \tag{244}$$

$$f_{2i} = f_{2i} \left(k_{5i}r\cos\theta + k_{6i}r\sin\theta + k_{7i}z - ct\sqrt{k_{5i}^2 + k_{6i}^2 + k_{7i}^2} + k_{8i} \right) \tag{245}$$

$$g_i = k_{9i}r\cos\theta + k_{10i}r\sin\theta + k_{11i}z + k_{12i}t + k_{13i} \quad (246)$$

For the spherical coordinate system

$$x = r\sin\theta\cos\varphi, y = r\sin\theta\sin\varphi, z = r\cos\theta. \quad (247)$$

By Eq. (227), (231-233), (247) and Theorem 5, the general solution of vector wave equation in spherical coordinate system can be obtained straightway

$$\begin{aligned} u = & (f_{1x} + f_{2x} + g_x)(\cos\varphi\sin\theta e_r + \cos\varphi\cos\theta e_\theta - \sin\varphi e_\varphi) \\ & + (f_{1y} + f_{2y} + g_y)(\sin\varphi\sin\theta e_r + \sin\varphi\cos\theta e_\theta + \cos\varphi e_\varphi) \\ & + (f_{1z} + f_{2z} + g_z)(\cos\theta e_r - \sin\theta e_\theta) \end{aligned}$$

Namely

$$\begin{aligned} u = & (f_{1x} + f_{2x} + g_x)(\cos\varphi\sin\theta e_r + \cos\varphi\cos\theta e_\theta - \sin\varphi e_\varphi) \\ & + (f_{1y} + f_{2y} + g_y)(\sin\varphi\sin\theta e_r + \sin\varphi\cos\theta e_\theta + \cos\varphi e_\varphi) \\ & + (f_{1z} + f_{2z} + g_z)(\cos\theta e_r - \sin\theta e_\theta) \end{aligned}$$

Namely

$$\begin{aligned} u = & (\cos\varphi\sin\theta(f_{1x} + f_{2x} + g_x) + \sin\varphi\sin\theta(f_{1y} + f_{2y} + g_y) + \cos\theta(f_{1z} + f_{2z} + g_z))e_r \\ & + \cos\varphi\cos\theta(f_{1x} + f_{2x} + g_x) + \sin\varphi\cos\theta(f_{1y} + f_{2y} + g_y) - \sin\theta(f_{1z} + f_{2z} + g_z)e_\theta \\ & + (-\sin\varphi(f_{1x} + f_{2x} + g_x) + \cos\varphi(f_{1y} + f_{2y} + g_y))e_\varphi, \end{aligned} \quad (243)$$

where

$$f_{1i} = f_{1i} \left(k_{1i}r\sin\theta\cos\varphi + k_{2i}r\sin\theta\sin\varphi + k_{3i}r\cos\theta + ct\sqrt{k_{1i}^2 + k_{2i}^2 + k_{3i}^2 + k_{4i}} \right), \quad (249)$$

$$f_{2i} = f_{2i} \left(k_{5i}r\sin\theta\cos\varphi + k_{6i}r\sin\theta\sin\varphi + k_{7i}r\cos\theta + ct\sqrt{k_{5i}^2 + k_{6i}^2 + k_{7i}^2 + k_{8i}} \right), \quad (250)$$

$$g_i = k_{9i}r\sin\theta\cos\varphi + k_{10i}r\sin\theta\sin\varphi + k_{11i}r\cos\theta + k_{12i}t + k_{13i}, (i = x, y, z). \quad (251)$$

The general solutions of the vector Laplace equation and Helmholtz equation and so on in cylindrical coordinate system and spherical coordinate system can be obtained similarly.

3. General solutions laws of linear partial differential equations

In recent years, many numerical methods have been developed to solve LPDEs, such as Finite integration method [76, 77], Bernoulli matrix method [78], Chebyshev matrix method [79] and so on, the existence [80], uniqueness [81, 82] and stability [83] of the solution are also the focus of research. Using the principle of transformational equations, if the solution of a LPDE is known, the solutions of its infinite IVTEs and DVTEs can be obtained, which may be LPDEs or NLPDEs. In this chapter, we will research new general solutions laws of LPDEs.

3.1. General solutions laws of linear partial differential equations with variable coefficients

In this section, if there is no special interpretation, $a_i = a_i(x_1, \dots, x_n)$, $a_{j_i} = a_{j_i}(x_1, \dots, x_n)$, $a_{i_1 i_2 \dots i_n} = a_{i_1 i_2 \dots i_n}(x_1, \dots, x_n)$, $b_i = b_i(x_1, \dots, x_n)$, $b_{j_i} = b_{j_i}(x_1, \dots, x_n)$, l_i and l_{j_i} are arbitrary constants, f and f_i are arbitrary unary smooth functions ($i, j = 1, 2, \dots$).

Proposition 1. If $v(x_1, \dots, x_n)$ is the particular solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = 0$, then its

general solution is $u = f(v(x_1, \dots, x_n))$.

Prove. By Transformational Method 1 set $u(x_1, \dots, x_n) = f(v) = f(v(x_1, \dots, x_n))$ then

$$a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = a_1 v_{x_1} f' + a_2 v_{x_2} f' + \dots + a_n v_{x_n} f' = 0.$$

Namely $a_1 v_{x_1} + a_2 v_{x_2} + \dots + a_n v_{x_n} = 0$, if its particular solution is known, the general solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = 0$ is $u(x_1, \dots, x_n) = f(v(x_1, \dots, x_n))$.

Proposition 2. If $a_1 = \frac{-a_2 a_{j x_2} - a_3 a_{j x_3} - \dots - a_n a_{j x_n}}{a_{j x_1}}$ then the general solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = 0$, is $u = f(a_j(x_1, \dots, x_n))$.

Prove. According to Transformational Method 2, set $u(x_1, \dots, x_n) = f(a_j(x_1, \dots, x_n))$, ($j = 2, 3, \dots, n$), then

$$a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = a_1 a_{j x_1} f' + a_2 a_{j x_2} f' + \dots + a_n a_{j x_n} f' = 0.$$

Namely

$$a_1 = \frac{-a_2 a_{j x_2} - a_3 a_{j x_3} - \dots - a_n a_{j x_n}}{a_{j x_1}}.$$

So the general solution of

$a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = 0$ under the condition of $a_1 = \frac{-a_2 a_{j x_2} - a_3 a_{j x_3} - \dots - a_n a_{j x_n}}{a_{j x_1}}$ is $u = f(a_j(x_1, \dots, x_n))$.

Proposition 3.

$$a_1 v_{x_1} + a_2 v_{x_2} + \dots + a_n v_{x_n} = 0, \quad (252)$$

$$a_1 g_{x_1} + a_2 g_{x_2} + \dots + a_n g_{x_n} + a_{n+1} g = 0. \quad (253)$$

If the particular solutions of Eq. (252) and Eq. (253) are known, the general solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} + a_{n+1} u = 0$ is $u = g(x_1, \dots, x_n) f(v(x_1, \dots, x_n))$.

Prove. According to Transformational Method 3, set $u(x_1, \dots, x_n) = g(x_1, \dots, x_n) f(v(x_1, \dots, x_n))$, the

$$\begin{aligned} & a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} + a_{n+1} u \\ &= (a_1 v_{x_1} + a_2 v_{x_2} + \dots + a_n v_{x_n}) g f'_v + (a_1 g_{x_1} + a_2 g_{x_2} + \dots + a_n g_{x_n} + a_{n+1} g) f = 0. \end{aligned}$$

Setting f an arbitrary unary first differentiable function, we obtain

$$a_1 v_{x_1} + a_2 v_{x_2} + \dots + a_n v_{x_n} = 0, \quad (252)$$

$$a_1 g_{x_1} + a_2 g_{x_2} + \dots + a_n g_{x_n} + a_{n+1} g = 0. \quad (253)$$

So if the particular solutions of Eq. (252) and Eq. (253) are known, the general solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} + a_{n+1} u = 0$ is

$$u(x_1, \dots, x_n) = g(x_1, \dots, x_n) f(v(x_1, \dots, x_n)). \quad (254)$$

Proposition 4. If the general solution $u = f(v(x_1, \dots, x_n))$ of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = 0$ is known, then the general solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} + a_{n+1} u = 0$ is $u = f(v(x_1, \dots, x_n)) + g(x_1, \dots, x_n)$, where $u = g(x_1, \dots, x_n)$ is the particular solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} + a_{n+1} u = 0$.

Proposition 4 is obvious, and almost needs no proof.

Proposition 5. If the general solution $u = f(v(x, y, z))$ of $(b_1D_x + b_2D_y + b_3D_z)u = 0$ is known, where $D_{x_j} \equiv \frac{\partial}{\partial x_j}$, ($x_j = x, y, z$) then the general solution of $(b_1D_x + b_2D_y + b_3D_z)^2u = 0$ is $u = f_1(v(x, y, z)) + (l_1x + l_2y + l_3z + l_4) f_2(v(x, y, z))$.

Prove. Because the general solution $u = f(v(x, y, z))$ of $(b_1D_x + b_2D_y + b_3D_z)u = 0$ is known, apparently $u = f(v(x, y, z))$ is also the solution of $(b_1D_x + b_2D_y + b_3D_z)^2u = 0$, setting $u = g(x, y, z)f(v(x, y, z))$ is the solution of is the solution of $(b_1D_x + b_2D_y + b_3D_z)^2u = 0$ too, then

$$\begin{aligned}
& (b_1D_x + b_2D_y + b_3D_z)^2u \\
&= (b_1^2D_x^2 + b_2^2D_y^2 + b_3^2D_z^2 + 2b_1b_2D_xD_y + 2b_1b_3D_xD_z + 2b_2b_3D_yD_z) gf \\
&= b_1^2(gf_{xx} + 2f_xg_x + g_{xx}f) + b_2^2(gf_{yy} + 2f_yg_y + g_{yy}f) + b_3^2(gf_{zz} + 2f_zg_z + g_{zz}f) \\
&\quad + 2b_1b_2(g_xf_y + gf_{xy} + g_{xy}f + g_yf_x) + 2b_1b_3(g_xf_z + gf_{xz} + g_{xz}f + g_zf_x) \\
&\quad + 2b_2b_3(g_yf_z + gf_{yz} + g_{yz}f + g_zf_y) \\
&= (b_1^2f_{xx} + b_2^2f_{yy} + b_3^2f_{zz} + 2b_1b_2f_{xy} + 2b_1b_3f_{xz} + 2b_2b_3f_{yz}) g \\
&\quad + (b_1^2g_{xx} + b_2^2g_{yy} + b_3^2g_{zz} + 2b_1b_2g_{xy} + 2b_1b_3g_{xz} + 2b_2b_3g_{yz}) f \\
&\quad + (2b_1^2g_x + 2b_1b_2g_y + 2b_1b_3g_z) f_x + (2b_2^2g_y + 2b_1b_2g_x + 2b_2b_3g_z) f_y \\
&\quad + (2b_3^2g_z + 2b_1b_3g_x + 2b_2b_3g_y) f_z \\
&= (b_1^2g_{xx} + 2b_1b_2g_{xy} + b_2^2g_{yy} + 2b_1b_3g_{xz} + 2b_2b_3g_{yz} + b_3^2g_{zz}) f \\
&\quad + 2b_1g_x(b_1f_x + b_2f_y + b_3f_z) + 2b_2g_y(b_1f_x + b_2f_y + b_3f_z) + 2b_3g_z(b_1f_x + b_2f_y + b_3f_z) \\
&= (b_1^2g_{xx} + 2b_1b_2g_{xy} + b_2^2g_{yy} + 2b_1b_3g_{xz} + 2b_2b_3g_{yz} + b_3^2g_{zz}) f = 0 \\
&\implies b_1^2g_{xx} + 2b_1b_2g_{xy} + b_2^2g_{yy} + 2b_1b_3g_{xz} + 2b_2b_3g_{yz} + b_3^2g_{zz} = 0.
\end{aligned}$$

Namely

$$b_1^2g_{xx} + 2b_1b_2g_{xy} + b_2^2g_{yy} + 2b_1b_3g_{xz} + 2b_2b_3g_{yz} + b_3^2g_{zz} = 0, \quad (255)$$

$g(x, y, z)$ has to be a particular solution of Eq. (255), due to

$$g(x, y, z) = l_1x + l_2y + l_3z + l_4, \quad (256)$$

(256) is a particular solution of Eq. (255), so the general solution of $(b_1D_x + b_2D_y + b_3D_z)^2u = 0$ is $u = f_1(v(x, y, z)) + (l_1x + l_2y + l_3z + l_4) f_2(v(x, y, z))$.

Proposition 6. If the general solution $u = f(v(x_1, \dots, x_n))$ of $(b_1D_{x_1} + b_2D_{x_2} + \dots + b_nD_{x_n})u = 0$ is known, then the general solution of $(b_1D_{x_1} + b_2D_{x_2} + \dots + b_nD_{x_n})^2u = 0$ is $u = f_1(v(x_1, \dots, x_n)) + (l_1x_1 + l_2x_2 + \dots + l_nx_n) f_2(v(x_1, \dots, x_n))$.

Prove. Because the general solution $u = f(v(x_1, \dots, x_n))$ of $(b_1D_{x_1} + b_2D_{x_2} + \dots + b_nD_{x_n})u = 0$ is known, apparently $u = f(v(x_1, \dots, x_n))$ is the solution of $(b_1D_{x_1} + b_2D_{x_2} + \dots + b_nD_{x_n})^2u = 0$ yet, according to Proposition 5, we set that $u = g(x_1, \dots, x_n)f = l_sx_s f$ is the solution of $(b_1D_{x_1} + b_2D_{x_2} + \dots + b_nD_{x_n})^2u = 0$ too, namely $g(x_1, \dots, x_n) = l_sx_s$, l_s is an arbitrary

constant, ($s = 1, 2, \dots, n$), then

$$\begin{aligned}
& (b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n})^2 u = \left(\sum_{i=1}^n b_i^2 D_{x_i}^2 + 2 \sum_{i<j} b_i b_j D_{x_i} D_{x_j} \right) (gf) \\
& = \sum_{i=1}^n b_i^2 (g_{x_i x_i} f + 2g_{x_i} f_{x_i} + g f_{x_i x_i}) + 2 \sum_{i \neq j} b_i b_j (g_{x_i x_j} f + g_{x_i} f_{x_j} + g_{x_j} f_{x_i} + g f_{x_i x_j}) \\
& = g \left(\sum_{i=1}^n b_i^2 f_{x_i x_i} + 2 \sum_{i<j} b_i b_j f_{x_i x_j} \right) + f \left(\sum_{i=1}^n b_i^2 g_{x_i x_i} + 2 \sum_{i<j} b_i b_j g_{x_i x_j} \right) + 2 \sum_{i=1}^n b_i^2 g_{x_i} f_{x_i} \\
& + 2 \sum_{i \neq j} b_i b_j (g_{x_i} f_{x_j} + g_{x_j} f_{x_i}) = 2 \sum_{i=1}^n b_i^2 g_{x_i} f_{x_i} + 2 \sum_{i<j} b_i b_j g_{x_i} f_{x_j} + 2 \sum_{i<j} b_i b_j g_{x_j} f_{x_i} \\
& = 2b_s^2 l_s^2 f_{x_s} + 2l_s b_s \sum_{s<j} b_j f_{x_j} + 2l_s b_s \sum_{i<s} b_i f_{x_i} = 2l_s b_s \sum_{i=1}^n b_i f_{x_i} = 0.
\end{aligned}$$

That $u = l_s x_s f(v(x_1, \dots, x_n))$ is the solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n})^2 u = 0$ is proved. So its general solution is

$$u = f_1(v(x_1, \dots, x_n)) + (l_1 x_1 + l_2 x_2 + \dots + l_n x_n) f_2(v(x_1, \dots, x_n)). \quad (257)$$

Note (257) may be written as

$$u = f_1(v(x_1, \dots, x_n)) + (l_1 x_1 + l_2 x_2 + \dots + l_n x_n + l_{n+1}) f_2(v(x_1, \dots, x_n)).$$

According to Proposition 6, we present a conjecture

Conjecture 1. If the general solution $u = f(v(x_1, \dots, x_n))$ of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n})u = 0$ is known, then the general solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n})^m u = 0$ is

$$u = \sum_{j=1}^m (l_{j1} x_1 + l_{j2} x_2 + \dots + l_{jn} x_n)^{j-1} f_j(v(x_1, \dots, x_n)).$$

Theoretically Conjecture 1 can be proved by mathematical induction, we shall not analyse it further.

For the m th-order linear PDE with variable coefficients

$$\sum_{i_1+i_2+\dots+i_n=m} a_{i_1 i_2 \dots i_n} u_{x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}} = 0, \quad (258)$$

where i_j are positive integers, $1 \leq j \leq n$, If Eq. (258) can be translated into

$$\begin{aligned}
& (b_{11} D_{x_1} + b_{12} D_{x_2} + \dots + b_{1n} D_{x_n}) (b_{21} D_{x_1} + b_{22} D_{x_2} + \dots + b_{2n} D_{x_n}) \dots \\
& (b_{m1} D_{x_1} + b_{m2} D_{x_2} + \dots + b_{mn} D_{x_n}) u = 0.
\end{aligned} \quad (259)$$

For

$$(b_{j1} D_{x_1} + b_{j2} D_{x_2} + \dots + b_{jn} D_{x_n}) u = b_{j1} u_{x_1} + b_{j2} u_{x_2} + \dots + b_{jn} u_{x_n} = 0, \quad (j = 1, 2, \dots, m). \quad (260)$$

If the particular solutions $u = v_j(x_1, \dots, x_n)$, ($j = 1, 2, \dots, m$) of Eq. (260) are all known, by Proposition 1 the general solution of Eq. (258) is

$$u = \sum_{j=1}^m f_j(v_j(x_1, x_2, \dots, x_n)). \quad (261)$$

If Eq. (258) can be translated into:

$$\prod_{j=1}^q (b_{j_1} D_{x_1} + b_{j_2} D_{x_2} + \dots + b_{j_n} D_{x_n})^{p_j} u = 0, \quad (262)$$

where $\sum_{j=1}^q p_j = m$, its general solution of conjecture may be written by Conjecture 1.

Proposition 7. If the general solution $u = g(x_1, \dots, x_n) f(v(x_1, \dots, x_n))$ of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})u = 0$ is known, then the general solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^2 u = 0$ is $u = g(x_1, \dots, x_n) (f_1(v(x_1, \dots, x_n)) + (l_1 x_1 + l_2 x_2 + \dots + l_n x_n) f_2(v(x_1, \dots, x_n)))$.

Prove. If the general solution $u = g(x_1, \dots, x_n) f(v(x_1, \dots, x_n))$ of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})u = 0$ is known, is known, apparently $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^2 u = 0$ is $u = g(x_1, \dots, x_n) f(v(x_1, \dots, x_n))$ is the solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^2 u = 0$ yet, set

$$u = ht = l_s x_s g(x_1, x_2, \dots, x_n) f(v(x_1, x_2, \dots, x_n)), \quad (263)$$

where $h = l_s x_s, t = g(x_1, x_2, \dots, x_n) f(v(x_1, \dots, x_n))$ and l_s is an arbitrary constant ($s = 1, 2, \dots, n$). Assuming (263) is a particular solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^2 u = 0$, then

$$\begin{aligned} & (b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^2 u \\ &= \sum_{i=1}^n b_i^2 (h_{x_i x_i} t + 2h_{x_i} t_{x_i} + h t_{x_i x_i}) + b_{n+1}^2 h t \\ & \quad + 2 \sum_{1 \leq i < j \leq n} b_i b_j (h_{x_i x_j} t + h_{x_i} t_{x_j} + h_{x_j} t_{x_i} + h t_{x_i x_j}) + 2b_{n+1} \sum_{i=1}^n b_i (h_{x_i} t + h t_{x_i}) \\ &= h \left(\sum_{i=1}^n b_i^2 t_{x_i x_i} + b_{n+1}^2 t + 2 \sum_{1 \leq i < j \leq n} b_i b_j t_{x_i x_j} + 2b_{n+1} \sum_{i=1}^n b_i t_{x_i} \right) + t \sum_{i=1}^n b_i^2 h_{x_i x_i} \\ & \quad + 2 \sum_{i=1}^n b_i^2 h_{x_i} t_{x_i} + 2t \sum_{1 \leq i < j \leq n} b_i b_j h_{x_i x_j} + 2 \sum_{1 \leq i < j \leq n} b_i b_j h_{x_i} t_{x_j} + 2 \sum_{1 \leq i < j \leq n} b_i b_j h_{x_j} t_{x_i} \\ & \quad + 2b_{n+1} t \sum_{i=1}^n b_i h_{x_i} \\ &= 2 \sum_{i=1}^n b_i^2 h_{x_i} t_{x_i} + 2 \sum_{1 \leq i < j \leq n} b_i b_j h_{x_i} t_{x_j} + 2 \sum_{1 \leq i < j \leq n} b_i b_j h_{x_j} t_{x_i} + 2b_{n+1} t \sum_{i=1}^n b_i h_{x_i} \\ &= 2b_s^2 h_{x_s} t_{x_s} + 2b_s h_{x_s} \sum_{1 \leq s < j \leq n} b_j t_{x_j} + 2b_s h_{x_s} \sum_{1 \leq i < s \leq n} b_i t_{x_i} + 2b_{n+1} b_s h_{x_s} t \end{aligned}$$

That (263) is a solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^2 u = 0$ is proved, so its general solution is $u = g(x_1, \dots, x_n) (f_1(v(x_1, \dots, x_n)) + (l_1 x_1 + l_2 x_2 + \dots + l_n x_n) f_2(v(x_1, \dots, x_n)))$.

According to Proposition 7, we present Conjecture 2.

Conjecture 2. If the general solution $u = g(x_1, \dots, x_n) f(v(x_1, \dots, x_n))$ of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n})u = 0$ is known, the general solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^m u = 0$ is $u = g(x_1, \dots, x_n) \sum_{j=1}^m (l_{j_1} x_1 + l_{j_2} x_2 + \dots + l_{j_n} x_n)^{j-1} f_j(v(x_1, \dots, x_n)), (m \geq 2)$.

For the m th-order linear PDE with variable coefficients

$$\sum_{0 \leq i_1 + i_2 + \dots + i_n \leq m} a_{i_1 i_2 \dots i_n} u_{x_1 x_2 \dots x_n}^{(i_1 i_2 \dots i_n)} = 0, \quad (264)$$

where i_j are positive integers, $1 \leq j \leq n$. Suppose Eq. (264) can be translated into:

$$(b_{1_1} D_{x_1} + b_{1_2} D_{x_2} + \dots + b_{1_n} D_{x_n} + b_{1_{n+1}}) (b_{2_1} D_{x_1} + b_{2_2} D_{x_2} + \dots + b_{2_n} D_{x_n} + b_{2_{n+1}}) \dots (b_{m_1} D_{x_1} + b_{m_2} D_{x_2} + \dots + b_{m_n} D_{x_n} + b_{m_{n+1}}) u = 0. \quad (265)$$

If the particular solutions $g_j(x_1, \dots, x_n)$ of $(b_{j_1} D_{x_1} + b_{j_2} D_{x_2} + \dots + b_{j_n} D_{x_n} + b_{j_{n+1}})u = 0$ and the particular solutions $v_j(x_1, \dots, x_n)$ of $(b_{j_1} D_{x_1} + b_{j_2} D_{x_2} + \dots + b_{j_n} D_{x_n})u = 0$ are all known ($j = 1, 2, \dots, n$), by Proposition 3 the general solution of Eq. (265) is

$$u(x_1, \dots, x_n) = \sum_{j=1}^m (g_j(x_1, \dots, x_n) f_j(v_j(x_1, \dots, x_n))). \quad (266)$$

Suppose Eq. (264) can be translated into

$$\prod_{j=1}^q (b_{j_1} D_{x_1} + b_{j_2} D_{x_2} + \dots + b_{j_n} D_{x_n} + b_{j_{n+1}})^{p_j} u = 0, \quad (267)$$

where $\sum_{j=1}^q p_j = m$, its general solution of conjecture may be written by Conjecture 2.

For the m th-order linear PDE with variable coefficients

$$\sum_{0 \leq i_1 + i_2 + \dots + i_n \leq m} a_{i_1 i_2 \dots i_n} u_{x_1 x_2 \dots x_n}^{(i_1 i_2 \dots i_n)} = h(x_1, x_2, \dots, x_n), \quad (268)$$

where $h(x_1, x_2, \dots, x_n)$ is an arbitrary known function, we need first solve the particular solution of Eq. (268), by the general solution of its homogeneous equation, the general solution of Eq. (268) could be got.

3.2. General solutions laws of linear partial differential equations with constant coefficients

Here we will research the general solutions laws of LPDEs with constant coefficients, which are the special cases of LPDEs with variable coefficients. In this section, if there is no special interpretation $a_i, a_{j_i}, b_i, b_{j_i}, c_i, k_i, k_{j_i}, l_i, l_{j_i}$ and $a_{i_1 i_2 \dots i_n}$ are arbitrary constants, f and f_i are arbitrary unary smooth functions ($i, j = 1, 2, \dots$).

Proposition 8. The general solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = 0$ is $u = f(\frac{-1}{a_1}(a_2 k_2 + a_3 k_3 + \dots + a_n k_n)x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1})$.

Prove. According to Transformational Method 1, set $u(x_1, \dots, x_n) = f(v)$, $v(x_1, \dots, x_n) = k_1 x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1}$, then

$$\begin{aligned} a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} &= a_1 k_1 f'_v + a_2 k_2 f'_v + \dots + a_n k_n f'_v = 0 \\ \implies k_1 &= \frac{-1}{a_1} (a_2 k_2 + a_3 k_3 + \dots + a_n k_n). \end{aligned}$$

So the general solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} = 0$ is $u(x_1, x_2, \dots, x_n) = f(\frac{-1}{a_1}(a_2 k_2 + a_3 k_3 + \dots + a_n k_n)x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1})$.

Proposition 9. The general solution of $(b_1D_{x_1} + b_2D_{x_2} + \dots + b_nD_{x_n})^2u = 0$ is $u = f_1\left(\frac{-1}{b_1}(b_2k_{1_2} + b_3k_{1_3} + \dots + b_nk_{1_n})x_1 + k_{1_2}x_2 + \dots + k_{1_n}x_n + k_{1_{n+1}}\right) + (l_1x_1 + l_2x_2 + \dots + l_nx_n)f_2\left(\frac{-1}{b_1}(b_2k_{2_2} + b_3k_{2_3} + \dots + b_nk_{2_n})x_1 + k_{2_2}x_2 + \dots + k_{2_n}x_n + k_{2_{n+1}}\right)$.

Prove. According to Proposition 6 and Proposition 8, we can get Proposition 9 directly.

By Proposition 9 and Conjecture 1, we present Conjecture 3.

Conjecture 3. The general solution of $(b_1D_{x_1} + b_2D_{x_2} + \dots + b_nD_{x_n})^m u = 0$ is $u = \sum_{j=1}^m (l_{j_1}x_1 + \dots + l_{j_n}x_n)^{j-1} f_j \left(\frac{-1}{b_1} (b_2k_{j_2} + \dots + b_nk_{j_n}) x_1 + k_{j_2}x_2 + \dots + k_{j_n}x_n + k_{j_{n+1}} \right)$.

For the m th-order LPDE with constant coefficients

$$\sum_{i_1+i_2+\dots+i_n=m} a_{i_1i_2\dots i_n} u_{x_1^{i_1}x_2^{i_2}\dots x_n^{i_n}} = 0, \quad (269)$$

where i_j are positive integers, $1 \leq j \leq n$. If Eq. (269) can be translated into

$$(b_{1_1}D_{x_1} + b_{1_2}D_{x_2} + \dots + b_{1_n}D_{x_n})(b_{2_1}D_{x_1} + b_{2_2}D_{x_2} + \dots + b_{2_n}D_{x_n}) \dots (b_{m_1}D_{x_1} + b_{m_2}D_{x_2} + \dots + b_{m_n}D_{x_n})u = 0. \quad (270)$$

By (261) and Proposition 8, the general solution of Eq. (270) is

$$u = \sum_{j=1}^m f_j \left(\frac{-1}{b_{j_1}} (b_{j_2}k_{j_2} + b_{j_3}k_{j_3} + \dots + b_{j_n}k_{j_n}) x_1 + k_{j_2}x_2 + \dots + k_{j_n}x_n + k_{j_{n+1}} \right). \quad (271)$$

If Eq. (269) can be converted into

$$\prod_{j=1}^q (b_{j_1}D_{x_1} + b_{j_2}D_{x_2} + \dots + b_{j_n}D_{x_n})^{p_j} u = 0, \quad (272)$$

where $\sum_{j=1}^q p_j = m$, its general solution of conjecture may be written by Conjecture 3.

Proposition 10. The general solution of $a_1u_{x_1} + a_2u_{x_2} + \dots + a_nu_{x_n} + a_{n+1}u = 0$ is $u = f\left(\frac{-1}{a_1}(a_2k_2 + a_3k_3 + \dots + a_nk_n)x_1 + k_2x_2 + \dots + k_nx_n + k_{n+1}\right) \sum_{i=1}^n c_i e^{\frac{-a_{n+1}x_i}{a_i}}$.

Prove. We set $u = g(x_1, \dots, x_n)f(v)$, where $v(x_1, \dots, x_n) = k_1x_1 + \dots + k_nx_n + k_{n+1}$, then

$$\begin{aligned} & a_1u_{x_1} + a_2u_{x_2} + \dots + a_nu_{x_n} + a_{n+1}u \\ &= a_1g_{x_1}f + a_1k_1gf'_v + a_2g_{x_2}f + a_2k_2gf'_v + \dots + a_ng_{x_n}f + a_nk_ngf'_v + a_{n+1}gf \\ &= (a_1k_1 + a_2k_2 + \dots + a_nk_n)gf'_v + (a_1g_{x_1} + a_2g_{x_2} + \dots + a_ng_{x_n} + a_{n+1}g)f = 0. \end{aligned}$$

Setting f an arbitrary unary first differentiable function, then we get

$$a_1k_1 + a_2k_2 + \dots + a_nk_n = 0 \implies k_1 = \frac{-1}{a_1}(a_2k_2 + a_3k_3 + \dots + a_nk_n), \quad (273)$$

$$a_1g_{x_1} + a_2g_{x_2} + \dots + a_ng_{x_n} + a_{n+1}g = 0. \quad (274)$$

Set $g(x_1, \dots, x_n) = h(x_i), (i = 1, 2, \dots, n)$, then

$$a_1 g_{x_1} + \dots + a_n g_{x_n} + a_{n+1} g = a_i h_{x_i} + a_{n+1} h = 0 \implies h(x_i) = g(x_1, \dots, x_n) = c_i e^{\frac{-a_{n+1} x_i}{a_i}},$$

so $u = \sum_{i=1}^n c_i e^{\frac{-a_{n+1} x_i}{a_i}}$ is the particular solution of $a_1 u_{x_1} + a_2 u_{x_2} + \dots + a_n u_{x_n} + a_{n+1} u = 0$, thus its general solution is

$$u = f \left(\frac{-1}{a_1} (a_2 k_2 + a_3 k_3 + \dots + a_n k_n) x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1} \right) \sum_{i=1}^n c_i e^{\frac{-a_{n+1} x_i}{a_i}}. \quad (275)$$

Proposition 11. The general solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^2 u = 0$ is $u = \left(\sum_{j=1}^2 (l_{j_1} x_1 + l_{j_2} x_2 + \dots + l_{j_n} x_n)^{j-1} f_j \left(\frac{-1}{b_1} (b_2 k_{j_2} + b_3 k_{j_3} + \dots + b_n k_{j_n}) x_1 + k_{j_2} x_2 + \dots + k_{j_n} x_n + k_{j_{n+1}} \right) \right) \sum_{i=1}^n c_i e^{\frac{-b_{n+1} x_i}{b_i}}$.

Prove. According to Proposition 7 and Proposition 10, we can get Proposition 11 directly.

By Proposition 11, we present Conjecture 4.

Conjecture 4. The general solution of $(b_1 D_{x_1} + b_2 D_{x_2} + \dots + b_n D_{x_n} + b_{n+1})^m u = 0$ is $u = \left(\sum_{j=1}^m (l_{j_1} x_1 + l_{j_2} x_2 + \dots + l_{j_n} x_n)^{j-1} f_j \left(\frac{-1}{b_1} (b_2 k_{j_2} + b_3 k_{j_3} + \dots + b_n k_{j_n}) x_1 + k_{j_2} x_2 + \dots + k_{j_n} x_n + k_{j_{n+1}} \right) \right) \sum_{i=1}^n c_i e^{\frac{-b_{n+1} x_i}{b_i}}, (m \geq 2)$.

For the m th-order linear PDE with constant coefficients

$$\sum_{0 \leq i_1 + i_2 + \dots + i_n \leq m} a_{i_1 i_2 \dots i_n} u_{x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}} = 0, \quad (276)$$

where i_j are positive integers, $1 \leq j \leq n$. Suppose Eq. (276) can be translated into

$$(b_{1_1} D_{x_1} + b_{1_2} D_{x_2} + \dots + b_{1_n} D_{x_n} + b_{1_{n+1}}) (b_{2_1} D_{x_1} + b_{2_2} D_{x_2} + \dots + b_{2_n} D_{x_n} + b_{2_{n+1}}) \dots (b_{m_1} D_{x_1} + b_{m_2} D_{x_2} + \dots + b_{m_n} D_{x_n} + b_{m_{n+1}}) u = 0. \quad (277)$$

According to Proposition 10 the general solution of Eq. (277) is

$$u = \sum_{j=1}^m \left(f_j \left(\frac{-1}{b_{j_1}} (b_{j_2} k_{j_2} + \dots + b_{j_n} k_{j_n}) x_1 + k_{j_2} x_2 + \dots + k_{j_n} x_n + k_{j_{n+1}} \right) \sum_{i=1}^n c_{j_i} e^{\frac{-b_{j_{n+1}} x_i}{b_{j_i}}} \right). \quad (278)$$

Suppose Eq. (276) can be translated into

$$\prod_{j=1}^q (b_{j_1} D_{x_1} + b_{j_2} D_{x_2} + \dots + b_{j_n} D_{x_n} + b_{j_{n+1}})^{p_j} u = 0, \quad (279)$$

where $\sum_{j=1}^q p_j = m$, its general solution of conjecture may be written by Conjecture 4.

For the m th-order linear PDE with constant coefficients

$$\sum_{0 \leq i_1 + i_2 + \dots + i_n \leq m} a_{i_1 i_2 \dots i_n} u_{x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}} = g(x_1, x_2, \dots, x_n). \quad (280)$$

If $g(x_1, \dots, x_n) = h(v), v = k_1x_1 + \dots + k_nx_n + k_{n+1}$, h is a known unary function, k_1, k_2, \dots, k_{n+1} are known parameters, according to Transformational Method 2 set $u(x_1, x_2, \dots, x_n) = f(v)$, Eq. (280) could be converted into a m th-order nonhomogeneous LODE with constant coefficients, and its particular solution $f(v)$ could be obtained. Using (278) and so on, the general solution of the homogeneous equation of Eq. (280) can be had, thus the general solution of Eq. (280) may be gained further.

4. General solutions and particular solutions of nonlinear partial differential equations

We have studied Eq. (18), (27) and Eq. (83), which are NLPDEs, and have obtained their exact solutions or general solutions. Since the importance of NLPDEs, many effective methods have been proposed to obtain their exact solutions, such as F-expansion method [84], tanh-sech method [85C87], extended tanh method [88C90], multiple exp-function method [91, 92], hyperbolic function method [93], Jacobi elliptic function expansion method [94], homogeneous balance method [95C97], sineCcosine method [98C100] and so on. In this section we will solve some typical examples by the new laws and methods proposed in this paper.

According to (1), (2) we present Theorem 6 first.

Theorem 6. In the domain $D, (D \subset \mathbb{R}^n)$, if $u(x_1, x_2, \dots, x_n)$ is a first differentiable function, then

$$\frac{\partial \int f(u) du}{\partial x_i} = f(u) u_{x_i}. \quad (281)$$

Proof. Set

$$v(x_1, x_2, \dots, x_n) = g(u) = \int f(u) du. \quad (282)$$

By (2) and (282)

$$v_{x_i} = \frac{\partial \int f(u) du}{\partial x_i} = g'(u) u_{x_i} = f(u) u_{x_i}.$$

Thus the theorem is proved.

Example 4.1.

$$b(u) u_x = a(x, y), \quad (283)$$

where $a(x, y)$ is an any known binary function, $b(u)$ is an arbitrary known unary function, by Theorem 6

$$b(u) u_x = \frac{\partial \int b(u) du}{\partial x} = a(x, y) \int b(u) du = \phi(y) + \int a(x, y) dx,$$

where $\phi(y)$ is an arbitrary unary function, so the general solution of Eq. (283) is:

$$\int b(u) du = \phi(y) + \int a(x, y) dx. \quad (284)$$

Example 4.2.

$$a_y u_{xx} + a_x u_{xy} + a_y b(u) u_x^2 + a_x b(u) u_x u_y = 0, \quad (285)$$

where a is an any known binary function, $b(u)$ is an arbitrary known unary function, due to

$$a_y v_x + a_x v_y = 0, \quad (211)$$

the general solution of Eq. (211) is

$$v = g(a(x, y)), \quad (212)$$

where g is an arbitrary unary first differentiable function, by Theorem 2 we set

$$v(x, y) = f(u) u_x, \quad (286)$$

where f is an undetermined unary function, according to (284) and (212)

$$\int f(u) du = \phi(y) + \int v(x, y) dx = \phi(y) + \int g(a(x, y)) dx,$$

where $\phi(y)$ is an arbitrary unary function, by (286)

$$\begin{aligned} a_y v_x + a_x v_y &= a_y f(u) u_{xx} + a_y f'(u) u_x^2 + a_x f(u) u_{xy} + a_x f'(u) u_x u_y = 0 \\ \implies a_y u_{xx} + a_x u_{xy} + a_y \frac{f'(u)}{f(u)} u_x^2 + a_x \frac{f'(u)}{f(u)} u_x u_y \\ &= a_y u_{xx} + a_x u_{xy} + a_y b(u) u_x^2 + a_x b(u) u_x u_y = 0 \\ \implies b(u) &= \frac{f'(u)}{f(u)} \implies f(u) = e^{\int b(u) du}. \end{aligned}$$

So Eq. (285) is a DVTE of Eq. (211), according to Theorem 2, the general solution of Eq. (285) is

$$\int e^{\int b(u) du} du = \phi(y) + \int g(a(x, y)) dx. \quad (287)$$

Example 4.3.

$$u_{xy} + a(u) u_x u_y + b(x, y) u_y = 0, \quad (288)$$

where $a(u)$ is an any known unary function, $b(x, y)$ is an arbitrary known binary function, due to

$$v_x + b(x, y) v = 0, \quad (289)$$

the general solution of Eq. (289) is [74]

$$v = \varphi(y) e^{-\int b(x, y) dx}, \quad (290)$$

where $\varphi(y)$ is an arbitrary unary function, by Theorem 2 we set

$$v = f(u) u_y, \quad (291)$$

where f is an undetermined unary first differentiable function, according to (284) and (290)

$$\int f(u) du = \int v dy = \phi(x) + \int \varphi(y) e^{-\int b(x, y) dx} dy,$$

where $\phi(x)$ is an arbitrary unary function, by (291)

$$\begin{aligned} v_x + b(x, y) v &= f(u) u_{xy} + f'(u) u_x u_y + b(x, y) f(u) u_y = 0 \\ \implies u_{xy} + \frac{f'(u)}{f(u)} u_x u_y + b(x, y) u_y &= u_{xy} + a(u) u_x u_y + b(x, y) u_y = 0 \\ \implies \frac{f'(u)}{f(u)} &= a(u) \implies f(u) = e^{\int a(u) du}. \end{aligned}$$

So Eq. (288) is a DVTE of Eq. (289), according to Theorem 2, the general solution of Eq. (288) is

$$\int e^{\int a(u)du} du = \phi(x) + \int \varphi(y) e^{-\int b(x,y)dx} dy. \quad (292)$$

Example 4.4.

$$a_1 u_{x_1} + a_2 u_{x_2} + \dots a_n u_{x_n} + a_{n+1} h(u) = 0, \quad (293)$$

where $a_i = a_i(x_1, \dots, x_n)$, ($i = 1, 2, \dots, n+1$), and $h(u)$ is an arbitrary known unary function. By Proposition 3, supposing the general solution $w = g(x_1, \dots, x_n) f(v(x_1, \dots, x_n))$ of $a_1 w_{x_1} + a_2 w_{x_2} + \dots a_n w_{x_n} + a_{n+1} w = 0$ is known, set

$$w = p(u). \quad (294)$$

Then

$$w_{x_i} = p'_u u_{x_i}, \quad (295)$$

$$a_1 w_{x_1} + a_2 w_{x_2} + \dots a_n w_{x_n} + a_{n+1} w = a_1 p'_u u_{x_1} + a_2 p'_u u_{x_2} + \dots a_n p'_u u_{x_n} + a_{n+1} p(u) = 0$$

$$\implies a_1 u_{x_1} + a_2 u_{x_2} + \dots a_n u_{x_n} + a_{n+1} \frac{p(u)}{p'_u} = a_1 u_{x_1} + a_2 u_{x_2} + \dots a_n u_{x_n} + a_{n+1} h(u) = 0$$

$$\implies \frac{p(u)}{p'_u} = h(u)$$

$$\implies w = p(u) = e^{\int \frac{du}{h(u)}} = g(x_1, x_2, \dots, x_n) f(v(x_1, x_2, \dots, x_n)).$$

So Eq. (293) is a DVTE of $a_1 w_{x_1} + a_2 w_{x_2} + \dots a_n w_{x_n} + a_{n+1} w = 0$, according to Theorem 2, the general solution of Eq. (293) is

$$e^{\int \frac{du}{h(u)}} = g(x_1, x_2, \dots, x_n) f(v(x_1, x_2, \dots, x_n)). \quad (296)$$

If a_i are constants, by Proposition 10 and (296), the general solution of Eq. (293) can be get

$$e^{\int \frac{du}{h(u)}} = f\left(\frac{-1}{a_1} (a_2 k_2 + a_3 k_3 + \dots a_n k_n) x_1 + k_2 x_2 + \dots + k_n x_n + k_{n+1}\right) \sum_{i=1}^n c_i e^{\frac{-a_{n+1} x_i}{a_i}}, \quad (297)$$

where $k_2 - k_{n+1}$ and $c_1 - c_n$ are arbitrary constants.

In the theory of PDEs, converting a PDE to a relatively simple ODE is a classical method, such as Laplace transform, Fourier transform, and so on integral transformation belong to this type of way. According to Eq. (7) and (8), two kinds of PDEs can be transformed into ODEs, and now we use two new theorems to express this idea:

Theorem 7. In the domain D , ($D \subset \mathbb{R}^n$), any established m th-order PDE with n space variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$, If all the known functions satisfy $a_i(x_1, \dots, x_n) = a_i(k_1 x_1 + \dots + k_n x_n + k_{n+1})$, where k_1, k_2, \dots, k_{n+1} are known parameters, set $u(x_1, \dots, x_n) = f(k_1 x_1 + \dots + k_n x_n + k_{n+1})$, then substitute $u = f(k_1 x_1 + \dots + k_n x_n + k_{n+1})$ and its partial derivatives into $F = 0$

1. If $F = 0$ is a linear PDE, then it can be converted to a linear ODE,

2. If $F = 0$ is a non-linear PDE, then it can be converted to a non-linear ODE.

Proof. Set $v(x_1, \dots, x_n) = k_1 x_1 + \dots + k_n x_n + k_{n+1}$, then $a_i(x_1, \dots, x_n) = a_i v$, $u = f(v)$, by (7) and (8) $F = 0$ can be converted to an ODE whose dependent variable is f and independent variable is v , since each linear term in $F = 0$ is transformed into a new linear term, each non-linear term is transformed into a new nonlinear term, so the theorem is proved.

Theorem 8. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n space variables $F(u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$, namely in the equation there is no known function $a_i(x_1, \dots, x_n)$ set $u(x_1, \dots, x_n) = f(k_1 x_1 + \dots + k_n x_n + k_{n+1})$, where k_1, k_2, \dots, k_{n+1} are unascertained parameters, then substitute $u = f(k_1 x_1 + \dots + k_n x_n + k_{n+1})$ and its partial derivatives into $F = 0$

1. If $F = 0$ is a linear PDE, then it can be converted to a linear ODE,

2. If $F = 0$ is a non-linear PDE, then it can be converted to a non-linear ODE.

Proof. Set $v(x_1, \dots, x_n) = k_1 x_1 + \dots + k_n x_n + k_{n+1}$, then $u = f(v)$, by (2.7) and (2.8) $F = 0$ can be converted to an ODE whose dependent variable is f and independent variable is v , since each linear term in $F = 0$ is transformed into a new linear term, each non-linear term is transformed into a new nonlinear term, so the theorem is proved.

Theorem 7 and 8 is a further application of Transformational Method 1 and 2. In the previous, the method of solving the particular solution of Poisson equation is in fact using theorem 7, the method of solving Eq. (27) is actually the application of theorem 8. Now we use theorem 8 to solve two typical nonlinear PDEs.

Example 6.4.

$$a(u) u_t + b(u) u_x + c(u) u_{xx} = 0. \quad (298)$$

According to Theorem 8, set $u(x, t) = f(v) = f(k_1 x + k_2 t + k_3)$, $k_1 - k_3$ are parameters to be determined, f is an undetermined unary second differentiable function, then

$$a(u) u_t + b(u) u_x + c(u) u_{xx} = k_2 a(f) f'_v + k_1 b(f) f'_v + k_1^2 c(f) f''_v = 0.$$

Namely

$$f''_v + \frac{k_2 a(f) + k_1 b(f)}{k_1^2 c(f)} f'_v = 0 \quad (299)$$

Because

$$y'' + b(y) (y')^2 + c(y) (y')^m = 0. \quad (300)$$

The general solution of Eq. (300) is [53]

$$x = C_2 + \int \left(\frac{-e^{(2-m) \int b(y) dy}}{C_1 + (2-m) \int c(y) e^{(2-m) \int b(y) dy} dy} \right)^{\frac{1}{2-m}} dy, \quad (301)$$

where C_1 and C_2 are arbitrary constants, so the exact solution of Eq. (298) is

$$v = k_1 x + k_2 t + k_3 = \int \frac{du}{C_1 - \int \frac{k_2 a(u) + k_1 b(u)}{k_1^2 c(u)} du}. \quad (302)$$

where $k_1 - k_3$ are arbitrary constants, Burgers equation

$$u_t + uu_x + \alpha u_{xx} = 0, \quad (303)$$

is a special case of Eq. (298), according to (302) its exact solution is

$$k_1 x + k_2 t + k_3 = -2k_1^2 \alpha \int \frac{du}{C_1 + 2k_2 u + k_1 u^2}. \quad (304)$$

Example 6.5.

$$a(u)u_t + b(u)u_x + c(u)u_{xxx} = 0. \quad (305)$$

According to Theorem 8, set $u(x, t) = f(v) = f(k_1x + k_2t + k_3)$, $k_1 - k_3$ are parameters to be determined, f is an undetermined unary third differentiable function, then

$$a(u)u_t + b(u)u_x + c(u)u_{xxx} = k_2a(u)f'_v + k_1b(u)f'_v + k_1^3c(u)f'''_v = 0.$$

Namely

$$f'''_v + \frac{k_2a(u) + k_1b(u)}{k_1^3c(u)}f'_v = 0. \quad (306)$$

Because

$$y''' + b(y)y'(y'')^m = 0. \quad (307)$$

The general solution of Eq. (307) is [53]

$$x = C_1 + \int \left(C_2 - 2 \int \left(C_3 + (1-m) \int b(y) dy \right)^{\frac{1}{1-m}} dy \right)^{\frac{-1}{2}} dy, \quad (308)$$

where $C_1 - C_3$ are arbitrary constants, so the exact solution of Eq. (305) is

$$k_1x + k_2t + k_3 = \int \left(C_1 - C_2u - 2 \int \int \frac{k_2a(u) + k_1b(u)}{k_1^3c(u)} du du \right)^{\frac{-1}{2}} du. \quad (309)$$

where $k_1 - k_3$ are arbitrary constants, KdV equation, mKdV equation and KdV-mKdV equation

$$u_t + uu_x + \beta u_{xxx} = 0, \quad (310)$$

$$u_t + \alpha u^2 u_x + \beta u_{xxx} = 0, \quad (311)$$

$$u_t + \gamma uu_x + \alpha u^2 u_x + \beta u_{xxx} = 0, \quad (312)$$

are special cases of Eq. (305), according to (309) their exact solutions are

$$k_1x + k_2t + k_3 = \int \left(C_1 - C_2u - \frac{k_2}{k_1^3\beta}u^2 - \frac{k_1}{3k_1^3\beta}u^3 \right)^{\frac{-1}{2}} du, \quad (313)$$

$$k_1x + k_2t + k_3 = \int \left(C_1 - C_2u - \frac{k_2}{k_1^3\beta}u^2 - \frac{k_1\alpha}{6k_1^3\beta}u^4 \right)^{\frac{-1}{2}} du, \quad (314)$$

$$k_1x + k_2t + k_3 = \int \left(C_1 - C_2u - \frac{k_2}{k_1^3\beta}u^2 - \frac{k_1\gamma}{3k_1^3\beta}u^3 - \frac{k_1\alpha}{6k_1^3\beta}u^4 \right)^{\frac{-1}{2}} du, \quad (315)$$

respectively. Since $k_1 - k_3$ are arbitrary constants, (304), (313)-(315) are all solitary wave solutions.

5. Extention and conclusion

5.1. Two axioms and a conjecture

For revealing the relationship between an arbitrary m th-order partial differential equation (PDE) of n variables and an arbitrary m th-differentiable function of n variables, we present Axiom 1.

Axiom 1. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$, set $f(x_1, \dots, x_n)$ an arbitrary known function, $f \in C^m(D)$, then substitute from $u = f(x_1, \dots, x_n)$ and its partial derivatives into $F = 0$

1. In case of getting $0 = 0$, then $u = f(x_1, \dots, x_n)$ is the solution of $F = 0$,
2. In case of getting $k = 0$, but in fact $k \neq 0$, then $u = f(x_1, \dots, x_n)$ is not the solution of $F = 0$,
3. In case of getting $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = f(x_1, \dots, x_n)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$.

Among the three conclusions of Axiom 1, the first two are obvious and the relevant cases could be illustrated easily, we mainly analyse the third conclusion, such as

Example 5.1.

$$z_x^3 + z_y^3 + z^3 = x^3y^3. \quad (316)$$

We may choose an arbitrary binary first differentiable function, for instance substitute $z = xy$ into Eq. (316)

$$z_x^3 + z_y^3 + z^3 = y^3 + x^3 + x^3y^3 = x^3y^3 \implies y = -x.$$

So $z = xy$ is a solution of Eq. (316) under the condition of $y = -x$. Note $z = -x^2$ is not a particular solution of Eq. (316), in 3-dimensional space, $z = -x^2$ is a three-dimensional surface perpendicular to the y -axis, $z = xy$ under the condition of $y = -x$ is a three-dimensional curve:

$$\begin{cases} x = x \\ y = -x \\ z = xy = -x^2 \end{cases}$$

or set $x = t$, then:

$$\begin{cases} x = t \\ y = -t \\ z = -t^2 \end{cases}$$

So the geometric meaning and the mathematical meaning of these two cases are completely different.

Substituting from any function of n variables into an arbitrary PDE of n variables may obtain other special circumstances, such as denominator equal zero and so on, the probabilities of these circumstances are very low like getting $0 = 0$ or $k = 0$, so Axiom 1 reveals that almost any m th-differentiable function of n variables is a conditional solution of an arbitrary m th-order PDE of n variables.

Axiom 1 may be extended to ordinary differential equations (ODEs), so we present Axiom 2.

Axiom 2. In the domain $D, (D \subset \mathbb{R}^1)$, any established m th-order ordinary differential equation (ODE) $F(x, y, y^{(1)}, y^{(2)} \dots, y^{(m)}) = 0$, set $f(x)$ known and $f \in C^m(D)$, then substitute from $y = f(x)$ and its derivatives into $F = 0$

1. In case of getting $0 = 0$, then $y = f(x)$ is the solution of $F = 0$,
2. In case of getting $k = 0$, but in fact $k \neq 0$, then $y = f(x)$ is not the solution of $F = 0$,
3. In case of getting $g(x) = 0$, then $y = f(x)$ is the solution of $F = 0$ under the condition of $g(x) = 0$,
4. In case of getting $x = k_1, k_2 \cdots k_l (l \geq 1)$, then $y = f(x)$ is the discrete solution of $F = 0$ under the condition of $x = k_1, k_2 \cdots k_l$.

The probability of the above first and second results is very little, so Axiom 2 reveals that almost every unary m th-differentiable function is a conditional solution of an arbitrary m th-order ODE. Such as Abel equation.

Example 5.2.

$$y' + y^3 + a(x) = 0, \quad (317)$$

where $a(x)$ is an arbitrary unary functions, discretionarily set $y = cx$, then

$$y' + y^3 + a(x) = c + c^3x^3 + a(x) = 0 \implies a(x) = -c - c^3x^3.$$

Therefore, under the condition of $a(x) = -c - c^3x^3$, the particular solution of the Eq. (317) is $y = cx$.

Example 5.3.

$$y' + y^3 + x^3 = 0, \quad (318)$$

set $y = cx$, then

$$y' + y^3 + x^3 = c + (c^3 + 1)x^3 = 0 \implies x = \left(\frac{-c}{c^3 + 1} \right)^{\frac{1}{3}}.$$

Therefore, on the point $x = \left(\frac{-c}{c^3 + 1} \right)^{\frac{1}{3}}$, the particular solution of the Eq. (318) is $y = cx$. In some specific case, the general solution of the Abel equation may be referred to [101, 102].

We know that a univariate function satisfying certain conditions can be expanded into a Taylor series or a Fourier series. When we consider the Cauchy problem of Eq. (38), we assume that the conditions are (41) and (42). Now we propose a **conjecture** about the n-ary function:

Conjecture 5. A n-ary function satisfying certain conditions can be expanded into a series:

$$f(x_1, x_2, \dots, x_n) = \sum_{i=1}^s \varphi_i(k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}), (1 \leq s \leq \infty) \quad (319)$$

Where φ_i are arbitrarily determined unary functions and k_{i_j} are arbitrarily determined parameters.

For the unary function, Conjecture 5 is obviously correct, because the Taylor series and Fourier series are all special cases of $\sum_{i=1}^s \varphi_i(k_{i_1}x + k_{i_2})$ for the n-ary function ($n \geq 2$), how to strictly prove Conjecture 5 is a new mathematical problem. Even if some $f(x_1, x_2, \dots, x_n)$ cannot be strictly expanded to (318), since φ_i, k_{i_j} can be arbitrarily chosen, it can be further envisioned

that $f(x_1, x_2, \dots, x_n)$ should be approximatively replaced to $\sum_{i=1}^s \varphi_i (k_{i_1} x_1 + k_{i_2} x_2 + \dots + k_{i_n} x_n + k_{i_{n+1}})$ in a restricted domain.

5.2. General Equations and Restricted Equations

In this book, we have solved general solutions of many important PDEs, such as the concise general solution of acoustic equation in Cartesian coordinate system is

$$p = f_1 \left(k_1 x + k_2 y + k_3 z + c_0 t \sqrt{k_1^2 + k_2^2 + k_3^2 + k_4} \right) + f_2 \left(k_5 x + k_6 y + k_7 z - c_0 t \sqrt{k_5^2 + k_6^2 + k_7^2 + k_8} \right) + k_9 x + k_{10} y + k_{11} z + k_{12} t + k_{13}. \quad (63?)$$

If we set $k_3 = k_7 = k_{11} = 0$, then

$$p = f_1 \left(k_1 x + k_2 y + c_0 t \sqrt{k_1^2 + k_2^2 + k_4} \right) + f_2 \left(k_5 x + k_6 y - c_0 t \sqrt{k_5^2 + k_6^2 + k_8} \right) + k_9 x + k_{10} y + k_{12} t + k_{13} \quad (320)$$

(320) is essentially the general solution (58) of 2D wave equation

$$u = f_1 \left(k_1 x + k_2 y + at \sqrt{k_1^2 + k_2^2 + k_3} \right) + f_2 \left(k_4 x + k_5 y - at \sqrt{k_4^2 + k_5^2 + k_6} \right) + k_7 x + k_8 y + k_9 t + k_{10}. \quad (58?)$$

If we set $k_2 = k_3 = k_6 = k_7 = k_{11} = 0$, then

$$p = f_1 (k_1 x + c_0 k_1 t + k_4) + f_2 (k_5 x - c_0 k_5 t + k_8) + k_9 x + k_{12} t + k_{13} \quad (321)$$

(321) is essentially the general solution (48) of the 1D wave equation

$$u = f_1 (k_1 x + k_1 at + k_2) + f_2 (k_3 x - k_3 at + k_4) + k_5 x + k_6 t + k_7, \quad (48)$$

Such as the concise general solution of 3D heat equation in Cartesian coordinate system is

$$u = e^{\frac{k_0(k_0 t + k_1 x + k_2 y + k_3 z)}{(k_1^2 + k_2^2 + k_3^2) a^2}} h_1 \left(\frac{2k_0 \left(\sqrt{-l_2^2 - l_3^2} k_1 + l_2 k_2 + l_3 k_3 \right)}{k_1^2 + k_2^2 + k_3^2} t + \sqrt{-l_2^2 - l_3^2} x + l_2 y + l_3 z + l_4 \right) + e^{\frac{k_{10}(k_{10} t + k_{11} x + k_{12} y + k_{13} z)}{(k_{11}^2 + k_{12}^2 + k_{13}^2) a^2}} h_2 \left(\frac{2k_{10} \left(-\sqrt{-l_{12}^2 - l_{13}^2} k_{11} + l_{12} k_{12} + l_{13} k_{13} \right)}{k_{11}^2 + k_{12}^2 + k_{13}^2} t - \sqrt{-l_{12}^2 - l_{13}^2} x + l_{12} y + l_{13} z + l_{14} \right) \quad (123)$$

If we set $k_0 = k_{10} = 0$, then

$$u = h_1 \left(\sqrt{-l_2^2 - l_3^2} x + l_2 y + l_3 z + l_4 \right) + h_2 \left(-\sqrt{-l_{12}^2 - l_{13}^2} x + l_{12} y + l_{13} z + l_{14} \right), \quad (322)$$

Contrast (322) with (16)

$$u = f_1(x \sqrt{-k_2^2 - k_3^2} + k_1 y + k_2 z + k_3) + f_2(-x \sqrt{-k_4^2 - k_5^2} + k_4 y + k_5 z + k_6) + k_7 x + k_8 y + k_9 z + k_{10}, \quad (16)$$

(322) is a part of the general solution of Laplace equation.

Now let us ask a question: Why there is such a relation in the general solutions of these different equations?

In order to solve this problem, we first propose three new defines: **general equations, restricted equations and homologous restricted equations.**

If the equation $F_i = 0$ is a special case of the equation $F = 0, (i = 1, 2, \dots)$, then $F_i = 0$ is the restricted equation of $F = 0, F = 0$ is the general equation of $F_i = 0, F_i = 0$ between each other known as the homologous restricted equations.

In theory, any equation can have infinite general equations; if an equation contains arbitrary known functions or parameters, the equation can have infinite restricted equations.

Using the above defines, we propose **Axiom 3:**

Axiom 3: If there is no meaningless case, the solution of a general equation is known, then the solutions of all its restricted equations are known; if the solution of a restricted equation is unknown, then the solutions of all its general equations are unknown.

Axiom 3 is not hard to be understood. Since restricted equations are special cases of their general equations, the solutions of restricted equations are also special cases of the solutions of their general equations. Unless nonsensical cases occur, the solutions of all the restricted equations can be directly obtained by the known solutions of their general equations. On the other hand, if the solution of a restricted equation is not solved, it is impossible to solve the solutions of its general equations which are more complex.

According to the above defines and laws, we can explain why the solutions of some PDEs are similar, or even there are some definite relationships within them. Because they are homologous restricted equations, such as one-dimensional, two-dimensional and three-dimensional wave equation are all the restricted equations of Eq. (38); the heat equation and the Laplace equation are both the restricted equations of Eq. (111).

Since there is $k_7x + k_8y + k_9z + k_{10}$ in the general solution of the Laplace equation, we can make a preliminary judgment that these terms may be absent in the general solutions of (111) and (122).

5.3. Conclusion

In this paper, we have proposed ten new concepts, three new axioms and eight new theorems, they are:

concise general solution (CGS); series general solution (SGS); independent variable transformational equations (IVTEs); dependent variable transformational equations (DVTEs); symmetric vector partial differential equations (SVPDEs); corresponding scalar equation (CSE); independent variable transformation vector equation (IVTVE); general equations; restricted equations; homologous restricted equations.

Axiom 1. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$, set $f(x_1, \dots, x_n)$ an arbitrary known function, $f \in C^m(D)$, then substitute from $u = f(x_1, \dots, x_n)$ and its partial derivatives into $F = 0$

1. In case of getting $0 = 0$, then $u = f(x_1, \dots, x_n)$ is the solution of $F = 0$,

2. In case of getting $k = 0$, but in fact $k \neq 0$, then $u = f(x_1, \dots, x_n)$ is not the solution of $F = 0$,

3. In case of getting $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = f(x_1, \dots, x_n)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$.

Axiom 2. In the domain $D, (D \subset \mathbb{R}^1)$, any established m th-order ordinary differential equation (ODE) $F(x, y, y^{(1)}, y^{(2)} \dots y^{(m)}) = 0$, set $f(x)$ known and $f \in C^m(D)$, then substitute from $y = f(x)$ and its derivatives into $F = 0$

1. In case of getting $0 = 0$, then $y = f(x)$ is the solution of $F = 0$,

2. In case of getting $k = 0$, but in fact $k \neq 0$, then $y = f(x)$ is not the solution of $F = 0$,

3. In case of getting $g(x) = 0$, then $y = f(x)$ is the solution of $F = 0$ under the condition of $g(x) = 0$,

4. In case of getting $x = k_1, k_2 \dots k_l (l \geq 1)$, then $y = f(x)$ is the discrete solution of $F = 0$ under the condition of $x = k_1, k_2 \dots k_l$.

Above two new axioms reveal that almost any m th-differentiable function with n variables is a conditional solution of an arbitrary m th-order PDE with n variables and almost any unary m th-differentiable function is a conditional solution of an arbitrary m th-order ODE.

Axiom 3. If there is no meaningless case, the solution of a general equation is known, then the solutions of all its restricted equations are known; if the solution of a restricted equation is unknown, then the solutions of all its general equations are unknown.

Theorem 1. In the domain $D, (D \subset \mathbb{R}^n)$, if the solution $u = f(x_1, \dots, x_n)$ of a m th-order PDE $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$ is known, then the solution of its IVTE $G(y_1, \dots, y_n, u, u_{y_1}, \dots, u_{y_n}, u_{y_1y_2}, \dots) = 0$ is $u = f(x_1, \dots, x_n) = g(y_1, \dots, y_n)$.

Theorem 1 reveals the law of partial differential equations solution in various orthogonal coordinate system.

Theorem 2. In the domain $D, (D \subset \mathbb{R}^n)$, if the solution $v = f(x_1, \dots, x_n)$ of a PDE $F(x_1, \dots, x_n, v, v_{x_1}, \dots, v_{x_n}, v_{x_1x_2}, \dots) = 0$ is known, set $v = h(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots)$ then the solution of its DVTE $G(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$ is the solution of $h(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = f(x_1, \dots, x_n)$.

Theorem 3. In the domain $D, (D \subset \mathbb{R}^1)$, if the solution $w = f(x)$ of an ODE $F(x, w, w', w'', \dots, w^{(n)}) = 0$ is known, set $w = h(x, y, y', y'', \dots, y^{(m)})$, then the solution of its DVTE $G(x, y, y', y'', \dots, y^{(m+n)}) = 0$ is the solution of $h(x, y, y', y'', \dots, y^{(m)}) = f(x)$.

Theorem 4. If there are m arbitrary functions in the general solution of the CSE $F_i = 0$, then the number of arbitrary functions in the general solution of the n -dimensional SVPDE $F = 0$ is mn .

Theorem 5. In the domain $D, (D \subset \mathbb{R}^n)$, if the solution $u = f(x_1, x_2 \dots x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n})$ of a m th-order vector PDE $F(x_1, \dots, x_n, u, \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}, \frac{\partial^2 u}{\partial x_1 \partial x_2}, \dots) = 0$ is known, then the solution of its IVTVE $G(y_1, \dots, y_n, u, \frac{\partial u}{\partial y_1}, \dots, \frac{\partial u}{\partial y_n}, \frac{\partial^2 u}{\partial y_1 \partial y_2}, \dots) = 0$ is $u = f(x_1, x_2 \dots x_n, e_{x_1}, e_{x_2}, \dots, e_{x_n}) = g(y_1, y_2, \dots, y_n, e_{y_1}, e_{y_2}, \dots, e_{y_n})$

Theorem 6. In the domain $D, (D \subset \mathbb{R}^n)$, if $u(x_1, x_2, \dots, x_n)$ is a first differentiable function, then

$$\frac{\partial \int f(u) du}{\partial x_i} = f(u) u_{x_i}. \quad (281)$$

Theorem 7. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n space variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$, If all the known functions satisfy $a_i(x_1, \dots, x_n) = a_i(k_1x_1 + \dots + k_nx_n + k_{n+1})$ where k_1, k_2, \dots, k_{n+1} are known parameters, set $u(x_1, \dots, x_n) = f(k_1x_1 + \dots + k_nx_n + k_{n+1})$, then substitute $u = f(k_1x_1 + \dots + k_nx_n + k_{n+1})$ and its partial derivatives into $F = 0$

1. If $F = 0$ is a linear PDE, then it can be converted to a linear ODE,
2. If $F = 0$ is a non-linear PDE, then it can be converted to a non-linear ODE.

Theorem 8. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n space variables $F(u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$, namely in the equation there is no known function $a_i(x_1, \dots, x_n)$ set $u(x_1, \dots, x_n) = f(k_1x_1 + \dots + k_nx_n + k_{n+1})$, where k_1, k_2, \dots, k_{n+1} are unascertained parameters, then substitute $u = f(k_1x_1 + \dots + k_nx_n + k_{n+1})$ and its partial derivatives into $F = 0$

1. If $F = 0$ is a linear PDE, then it can be converted to a linear ODE,
2. If $F = 0$ is a non-linear PDE, then it can be converted to a non-linear ODE.

In Chapter 2, we indicate that Transformational Method 1-4 are specific applications of Theorem 2. In fact, Transformational Method 1-4 are also concrete applications of Axiom 1. According to Axiom 1, the complete representation of the four transformation methods should be:

Transformational Method 1. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$, set $v = v(x_1, \dots, x_n)$ and $u = f(v)$ are both undetermined m th-differentiable functions ($u, v \in C^m(D)$), then substitute $u = f(v)$ and its partial derivatives into $F = 0$

1. In case of working out $v(x_1, \dots, x_n)$ and $f(v)$, then $u = f(v)$ is the solution of $F = 0$,
2. In case of dividing out $f(v)$ and its partial derivative, also working out $v(x_1, \dots, x_n)$, then $u = f(v)$ is the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function,
3. In case of dividing out $f(v)$ and its partial derivative, also getting $k = 0$, but in fact $k \neq 0$, then $u = f(v)$ is not the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function.
4. In case of working out $v(x_1, \dots, x_n)$ and $f(v)$ under the condition of $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = f(v)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$,
5. In case of dividing out $f(v)$ and its partial derivative, also working out $v(x_1, \dots, x_n)$ under the condition of $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = f(v)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$, and f is an arbitrary unary m th-differentiable function,

Transformational Method 2. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1x_2}, \dots) = 0$, set $v = v(x_1, \dots, x_n)$ known and $u = f(v)$ undetermined ($u, v \in C^m(D)$), then substitute $u = f(v)$ and its partial derivatives into $F = 0$

1. In case of working out f , then $u = f(v)$ is the solution of $F = 0$,
2. In case of dividing out f and its partial derivative, also getting $0 = 0$, then $u = f(v)$ is the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function,
3. In case of dividing out f and its partial derivative, also getting $k = 0$, but in fact $k \neq 0$, then $u = f(v)$ is not the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function

4. In case of working out f under the condition of $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = f(v)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$,

5. In case of dividing out f and its partial derivative, also getting $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = f(v)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$, and f is an arbitrary unary m th-differentiable function.

Transformational Method 3. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$, setting $f(v), g(x_1, \dots, x_n)$ and $v(x_1, \dots, x_n)$ are all undetermined function, $g, v \in C^m(D)$, then substitute $u = gf(v)$ and its partial derivatives into $F = 0$

1. In case of working out f, g and v , then $u = gf(v)$ is the solution of $F = 0$,

2. In case of dividing out f and its partial derivative, also working out g and v , then $u = gf(v)$ is the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function,

3. In case of getting $k = 0$, but in fact $k \neq 0$, then $u = gf(v)$ is not the solution of $F = 0$,

4. In case of working out f, g and v under the condition of $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = gf(v)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$,

5. In case of dividing out f and its partial derivative, also working out g and v under the condition of $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = gf(v)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$, and f is an arbitrary unary m th-differentiable function.

Transformational Method 4. In the domain $D, (D \subset \mathbb{R}^n)$, any established m th-order PDE with n variables $F(x_1, \dots, x_n, u, u_{x_1}, \dots, u_{x_n}, u_{x_1 x_2}, \dots) = 0$, setting $g(x_1, \dots, x_n)$ is known and $f(v), v(x_1, \dots, x_n)$ are undetermined, $g, v \in C^m(D)$, then substitute $u = gf(v)$ and its partial derivatives into $F = 0$

1. In case of working out f and v , then $u = gf(v)$ is the solution of $F = 0$,

2. In case of dividing out f and its partial derivative, also working out $v(x_1, \dots, x_n)$, then $u = gf(v)$ is the solution of $F = 0$, and f is an arbitrary unary m th-differentiable function,

3. In case of getting $k = 0$, but in fact $k \neq 0$, then $u = gf(v)$ is not the solution of $F = 0$,

4. In case of working out f and v under the condition of $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = gf(v)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$,

5. In case of dividing out f and its partial derivative, also working out $v(x_1, \dots, x_n)$ under the condition of $h(x_1, \dots, x_l) = 0, (l \leq n)$, then $u = gf(v)$ is the solution of $F = 0$ under the condition of $h(x_1, \dots, x_l) = 0$, and f is an arbitrary unary m th-differentiable function.

Using above four new transformational methods, the general solutions and the exact solutions of the Cauchy problem for the Laplace equation, 2D wave equation, the acoustic wave equation, Helmholtz equation, heat equation and the diffusion equation have been solved. In some cases, the general solutions and the exact solutions of the Cauchy problem for the Poisson equation and Schrödinger equation have been solved too. We also find a singularity of general solutions of Helmholtz equation for the first time, namely the number of arbitrary functions in the general solutions is more than 2.

For research the laws of the general solution of m th-order LPDEs with n variables, we also

present 11 Propositions and 4 conjectures in Chapter 3.

Appendix

Appendix A

In (37) it can be proved that if $k_1, l_1 \neq 0$ and $k_1, l_1 \rightarrow 0$, $c_1 v$ can be described by f_1 and f_2 , set

$$k_i = l_i = C_i, (i = 2, 3, \dots, n+1).$$

Then

$$\begin{aligned} f_1 &= f_1(kx_1 + C_2x_2 + \dots + C_nx_n + C_{n+1}), \\ f_2 &= f_2(-kx_1 + C_2x_2 + \dots + C_nx_n + C_{n+1}), \end{aligned}$$

where

$$k = \left(-\frac{a_2C_2^2 + a_3C_3^2 + \dots + a_nC_n^2 + a_{n+1}C_2C_3}{a_1} \right)^{\frac{1}{2}}.$$

Set

$$A c_1(kx_1 + C_2x_2 + \dots + C_nx_n + C_{n+1}) + B c_1(-kx_1 + C_2x_2 + \dots + C_nx_n + C_{n+1}) = (A+B)c_1(C_1x_1 + C_2x_2 + \dots + C_nx_n + C_{n+1}) \Rightarrow C_1 = \frac{A-B}{A+B}k.$$

If $A = B \neq 0$, then $\frac{A-B}{A+B} = 0$. If $B = -A + 1$, then

$$\lim_{A \rightarrow \infty} \frac{A-B}{A+B} = \lim_{A \rightarrow \infty} (2A-1) \rightarrow \infty, \quad \lim_{A \rightarrow \infty} \frac{A-B}{A+B} = \lim_{A \rightarrow \infty} (2A-1) \rightarrow -\infty.$$

Namely $\frac{A-B}{A+B} \in (-\infty, \infty)$, if $k \neq 0$ and $k \rightarrow 0$, selecting A, B felicitously, C_1 may equal to arbitrary real number, so $c_1 v$ can be described by f_1, f_2 , and

$$\begin{aligned} c_1 v &= c_1(C_1x_1 + C_2x_2 + \dots + C_nx_n + C_{n+1}) \\ &= \frac{A c_1}{A+B}(kx_1 + C_2x_2 + \dots + C_nx_n + C_{n+1}) \\ &\quad + \frac{B c_1}{A+B}(-kx_1 + C_2x_2 + \dots + C_nx_n + C_{n+1}), \end{aligned}$$

where $C_1 = \frac{A-B}{A+B}k$.

Appendix B

The calculation of (43) as follows. In (40), set $c_1 = 0, k_{ij} = l_{ij}, (i = 1, 2, \dots, s, j = 2, 3, \dots, n+1)$

$$k_{i1} = \left(-(a_2k_{i2}^2 + \dots + a_nk_{in}^2 + a_{n+1}k_{i2}k_{i3}) / a_1 \right)^{\frac{1}{2}}. \quad (44)$$

According to (40)-(42)

$$\begin{aligned} u(0, x_2, \dots, x_n) &= \sum_{i=1}^s (f_{1i}(k_{i2}x_2 + \dots + k_{in}x_n + k_{i_{n+1}}) + f_{2i}(k_{i2}x_2 + \dots + k_{in}x_n + k_{i_{n+1}})) \\ &= \sum_{i=1}^s \varphi_i(k_{i2}x_2 + \dots + k_{in}x_n + k_{i_{n+1}}), \end{aligned}$$

$$\begin{aligned} u_{x_1}(0, x_2, \dots, x_n) &= \sum_{i=1}^s (k_{i1} f'_{1i}(k_{i2}x_2 + \dots + k_{in}x_n + k_{i_{n+1}}) - k_{i1} f'_{2i}(k_{i2}x_2 + \dots + k_{in}x_n + k_{i_{n+1}})) \\ &= \sum_{i=1}^s (k_{i2}x_2 + \dots + k_{in}x_n + k_{i_{n+1}}). \end{aligned}$$

We have

$$\begin{aligned} & f_{1_i} (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) + f_{2_i} (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \\ &= \varphi_i (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}), \end{aligned} \quad (323)$$

$$\begin{aligned} & k_{i_1}f'_{1_i} (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) - k_{i_1}f'_{2_i} (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \\ &= \psi_i (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \end{aligned} \quad (324)$$

According to (324) we get

$$\begin{aligned} & f_{1_i}(k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) - f_{2_i}(k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \\ &= \frac{1}{k_{i_1}} \int_{k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}}^{k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}} \psi(\xi_i) d\xi_i + \\ & f_{1_i}(k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) - f_{2_i}(k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) \end{aligned} \quad (325)$$

Combining (323) and (325), then

$$\begin{aligned} & f_{1_i} (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \\ &= \frac{1}{2}\varphi_i (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) + \frac{1}{2k_{i_1}} \int_{k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}}^{k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}} \psi(\xi_i) d\xi_i \\ &+ \frac{1}{2}f_{1_i} (k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) - \frac{1}{2}f_{2_i} (k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) \\ &\Rightarrow f_{1_i} (k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \\ &= \frac{1}{2}\varphi_i (k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) + \frac{1}{2k_{i_1}} \int_{k_{i_2}x_{2_0} + k_{i_3}x_{3_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}}^{k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}} \psi(\xi_i) d\xi_i \\ &+ \frac{1}{2}f_{1_i} (k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) - \frac{1}{2}f_{2_i} (k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) \end{aligned}$$

$$\begin{aligned} & f_{2_i} (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \\ &= \frac{1}{2}\varphi_i (k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) - \frac{1}{2k_{i_1}} \int_{k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}}^{k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}} \psi(\xi_i) d\xi_i - \\ & \frac{1}{2}f_{1_i} (k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) + \frac{1}{2}f_{2_i} (k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) \\ &\Rightarrow f_{2_i} (-k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \\ &= \frac{1}{2}\varphi_i (-k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) - \frac{1}{2k_{i_1}} \int_{k_{i_2}x_{2_0} + k_{i_3}x_{3_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}}^{-k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}} \psi(\xi_i) d\xi_i \\ &- \frac{1}{2}f_{1_i} (k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) + \frac{1}{2}f_{2_i} (k_{i_2}x_{2_0} + \dots + k_{i_n}x_{n_0} + k_{i_{n+1}}) \end{aligned}$$

In the conditions of (41) and (42), the exact solution of Eq. (38) is

$$\begin{aligned} u &= \frac{1}{2} \sum_{i=1}^s (\varphi_i(k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}) \\ &+ \varphi_i(-k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}})) \\ &+ \frac{1}{k_{i_1}} \int_{-k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}}^{k_{i_1}x_1 + k_{i_2}x_2 + \dots + k_{i_n}x_n + k_{i_{n+1}}} \psi(\xi_i) d\xi_i \end{aligned} \quad (43)$$

Appendix C

Consider the following initial value problem of Eq. (150) on the condition of (169)

$$u(x, y, z, 0) = e^{x+y+z} (\varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z)), \quad (172)$$

$$\begin{aligned} & u_t(x, y, z, 0) \\ &= e^{x+y+z} (\varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z)) \\ &+ \frac{\hbar}{m} e^{x+y+z} \left((2 + \sqrt{-2}) \varphi'_1(\sqrt{-2}x + y + z) + (2 - \sqrt{-2}) \varphi'_2(-\sqrt{-2}x + y + z) \right). \end{aligned} \quad (173)$$

Comparing (171) with (172) we have

$$k_1 = k_2 = k_3 = -\frac{1}{c}, l_2 = l_3 = l_{12} = l_{13} = 1, k_5 = l_5 = l_{15} = 0.$$

Then

$$\begin{aligned} u(x, y, z, t) &= e^{x+y+z-ck_4t} \left(h_1(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t) \right. \\ &\quad \left. + h_2(-\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 - \sqrt{-2})t) \right), \\ (x, y, z, t) &= -ck_4 e^{x+y+z-ck_4t} \left(h_1(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t) \right. \\ &\quad \left. + h_2(-\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 - \sqrt{-2})t) \right) \\ &\quad + e^{x+y+z-ck_4t} \left(\frac{\hbar}{m}(2 + \sqrt{-2})h_1'(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t) \right. \\ &\quad \left. + \frac{\hbar}{m}(2 - \sqrt{-2})h_2'(-\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 - \sqrt{-2})t) \right). \end{aligned}$$

Therefore

$$\begin{aligned} u(x, y, z, 0) &= e^{x+y+z} \left(\varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z) \right) \\ &= e^{x+y+z} \left(h_1(\sqrt{-2}x + y + z) + h_2(-\sqrt{-2}x + y + z) \right) \\ &\implies h_1(\sqrt{-2}x + y + z) = \varphi_1(\sqrt{-2}x + y + z) \\ &\implies h_1\left(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t\right) = \varphi_1\left(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t\right). \end{aligned}$$

Namely

$$h_1\left(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t\right) = \varphi_1\left(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t\right). \quad (326)$$

$$h_2\left(-\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 - \sqrt{-2})t\right) = \varphi_2\left(-\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 - \sqrt{-2})t\right). \quad (327)$$

Thus

$$\begin{aligned} u_t(x, y, z, 0) &= e^{x+y+z} \left(\varphi_1(\sqrt{-2}x + y + z) + \varphi_2(-\sqrt{-2}x + y + z) \right) \\ &\quad + \frac{\hbar}{m} e^{x+y+z} \left((2 + \sqrt{-2})\varphi_1'(\sqrt{-2}x + y + z) + (2 - \sqrt{-2})\varphi_2'(-\sqrt{-2}x + y + z) \right) \\ &= -ck_4 e^{x+y+z} \left(h_1(\sqrt{-2}x + y + z) + h_2(-\sqrt{-2}x + y + z) \right) \\ &\quad + e^{x+y+z} \left(\frac{\hbar}{m}(2 + \sqrt{-2})h_1'(\sqrt{-2}x + y + z) + \frac{\hbar}{m}(2 - \sqrt{-2})h_2'(-\sqrt{-2}x + y + z) \right) \\ &\implies k_4 = -\frac{1}{c}. \end{aligned}$$

So the exact solutions of the initial value problem is

$$\begin{aligned} u &= e^{x+y+z+ct} \left(\varphi_1(\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 + \sqrt{-2})t) \right. \\ &\quad \left. + \varphi_2(-\sqrt{-2}x + y + z + \frac{\hbar}{m}(2 - \sqrt{-2})t) \right). \end{aligned} \quad (174)$$

References

- [1] H. Triki, A.-M. Wazwaz, Traveling wave solutions for fifth-order KdV type equations with time-dependent coefficients, *Commun. Nonlinear Sci. Numer. Simulat.* 19 (2014) 404-408.
- [2] Y. Sun, New exact traveling wave solutions for double SineCGordon equation, *Appl. Math. Comput.* 258 (2015) 100-104.
- [3] K. Mallory, R.A. Van Gorder, K. Vajravelu, Several classes of exact solutions to the 1 + 1 BornCGInfeld Equation, *Commun. Nonlinear Sci. Numer. Simulat.* 19 (2014) 1669-1674.
- [4] T. Ellis, L. Demkowicz, J. Chan, Locally conservative discontinuous PetrovCGalerkin finite element methods for fluid problems, *Comput. Math. Appl.* 68 (2014) 1530-1549.
- [5] Z. Yosibash, Computing singular solutions of elliptic boundary value problems in polyhedral domains using p-FEM, *Appl. Numer. Math.* 33 (2000) 71.
- [6] D. Boffi, On the finite element method on quadrilateral meshes, *Appl. Numer. Math.* 56 (2006) 1271-1282.
- [7] M. Tabor, Modern dynamics and classical analysis, *Nature* 310 (1984) 277-282.
- [8] J. Liu, G. Mu, Z. Dai, X. Liu, Analytic multi-soliton solutions of the generalized Burgers equation, *Comput. Math. Appl.* 61 (2011) 1995-1999.
- [9] M. Song, X.J. Hou, J. Cao, Solitary wave solutions and kink wave solutions for a generalized KDVCmKDV equation, *Appl. Math. Comput.* 217 (2011) 5942-5948.
- [10] P.J. Rabier, C.A. Stuart, Application of elliptic regularity to bifurcation in stationary nonlinear Schrödinger equations, *Nonlinear Anal.* 52 (2003) 869-890..
- [11] R. Kajikiya, Least energy solutions of the Emden-Fowler equation in hollow thin symmetric domains, *J. Math. Anal. Appl.* 406 (2013) 277-286.
- [12] S.-T. YAU, Y. GAO, Obstacle Problem for von Kármán Equations, *Adv. Appl. Math.* 13 (1992) 123-141.
- [13] C.M. Guo, C.B. Zhai, R.P. Song, An existence and uniqueness result for the singular Lane-Emden-Fowler equation, *Nonlinear Anal. TMA* 72 (2010) 1275-1279.
- [14] F. Finster, J. Smoller, S.-T. Yau, Particle-like solutions of the Einstein-Dirac-Maxwell equations, *Phys. Lett. A* 259 (1999) 431-436.
- [15] L. Damascelli, A. Farina, B. Sciunzi, E. Valdinoci, Liouville results for m-Laplace equations of Lane-Emden-Fowler type, *Ann. I. H. Poincaré - AN.* 26 (4) (2009) 1099-1119.
- [16] L.C.L. Botelho, *Lecture Notes in Applied Differential Equations of Mathematical Physics*, World Scientific Pub Co, Hackensack, (2008).

- [17] G. Buttazzo, G.P. Galdi, L. Zanghirati, *Developments in Partial Differential Equations and Applications to Mathematical Physics*, Springer Science + Business Media, New York, (1992).
- [18] I. Rubinstein, L. Rubinstein, *Partial Differential Equations in Classical Mathematical Physics*, Cambridge University Press, Cambridge, (1998).
- [19] J.R. Kirkwood, *Mathematical Physics with Partial Differential Equations*, Academic Press, Waltham, (2013).
- [20] C.S. Wu, *Mathematical and Physical Methods*, Peking University Press, Beijing, China, 2003.
- [21] M. Kline, *Mathematical Thought from Ancient to Modern Times*, Oxford University Press, New York, (1972).
- [22] Y. Mirbagheri, H. Nahvi, J. Parvzian, A. Düster, Reducing spurious oscillations in discontinuous wave propagation simulation using high-order finite elements, *Comput. Math. Appl.* 70 (2015) 1640-1658.
- [23] M. Chen , M. Torres, T. Walsh, Existence of traveling wave solutions of a high-order nonlinear acoustic wave equation, *Phys. Lett. A.* 373 (2009) 1037-1043.
- [24] C.L. Frota, Some Nonlinear Wave Equations with Acoustic Boundary Conditions, *J. Differ. Equations* 164 (2000) 92-109.
- [25] F.G. Vasquez , G. W. Milton , D. Onofrei , Active Exterior Cloaking for the 2D Laplace and Helmholtz Equations, *Phys. Rev. Lett.* 103(7), 073901 (2009).
- [26] Q. Ma, Z.L. Mei, S.K. Zhu, T. Y. Jin, T. J. Cui, Experiments on Active Cloaking and Illusion for Laplace Equation, *Phys. Rev. Lett.* 111, 173901 (2013).
- [27] Y.C. Hon, T. Wei, Backus-Gilbert algorithm for the Cauchy problem of the Laplace equation, *Inverse Probl.* 17 (2) (2001) 261-271.
- [28] L. Bourgeois, A mixed formulation of quasi-reversibility to solve the Cauchy problem for Laplaces equation, *Inverse Probl.* 21 (3) (2005) 1087-1104.
- [29] M.V. Klibanov, F. Santosa, A computational quasi-reversibility method for Cauchy problems for Laplaces equation, *SIAM J. Appl. Math.* 51 (6) (1991) 1653-1675.
- [30] F. Berntsson, L. Eldén, Numerical solution of a Cauchy problem for the Laplace equation, *Inverse Probl.* 17 (4) (2001) 839-853.
- [31] Z. Chai, B. Shi, A novel lattice Boltzmann model for the Poisson equation, *Appl. Math. Model.* 32 (2008) 2050-2058.
- [32] J. Wei, D. Ye, F. Zhou, Analysis of boundary bubbling solutions for an anisotropic Emden-Fowler equation, *Ann. I. H. Poincaré - AN.* 25 (2008) 425-447.

- [33] S.M. Rybicki, Global Bifurcations of Solutions of Emden-Fowler-Type Equation $-\Delta u(x) = \lambda f(u(x))$ on an Annulus in $R^n, n \geq 3$, J. Differ. Equations, 183 (2002) 208-223.
- [34] C. Ye, W. Zhang, New explicit solutions for the Klein-Gordon equation with cubic nonlinearity, Appl. Math. Comput. 217 (2010) 716-724.
- [35] M.G. Hafez, M.N. Alam, M.A. Akbar, Exact traveling wave solutions to the Klein-Gordon equation using the novel (G'/G) -expansion method, Results Phys. 4 (2014) 177-184
- [36] A.V. Porubov, A.L. Fradkov, B.R. Andrievsky, Feedback control for some solutions of the sine-Gordon equation, Appl. Math. Comput. 269 (2015) 17-22.
- [37] T. Wei, Y.C. Hon, L. Ling, Method of fundamental solutions with regularization techniques for Cauchy problems of elliptic operators, Eng. Anal. Bound. Elem. 31 (2007) 373-385.
- [38] B.T. Jin, Y. Zheng, A meshless method for some inverse problems associated with the inhomogeneous Helmholtz equation, Comput. Methods Appl. Mech. Eng. 195 (2006) 2270-2288.
- [39] L. Marin, L. Elliott, P.J. Heggs, D.B. Ingham, D. Lesnic, X. Wen, An alternating iterative algorithm for the Cauchy problem associated to the Helmholtz equation, Comput. Methods Appl. Mech. Eng. 192 (2003) 709-722..
- [40] D. Zhang, F. Ma, E. Zheng, A Herglotz wavefunction method for solving the inverse Cauchy problem connected with the Helmholtz equation, J. Comput. Appl. Math. 237 (2013) 215-222.
- [41] B. Choi, D. Jeong, M.Y. Choi, General method to solve the heat equation, Phys. A 444 (2016) 530-537.
- [42] L.F. Barannyk, B. Kloskowska, On symmetry reduction and invariant solutions to some nonlinear multidimensional heat equations, Reports on Math. Phys. 45 (2000) 1-22.
- [43] A. Ahmad, A.H. Bokhari, A.H. Kara, F.D. Zamana, Symmetry classifications and reductions of some classes of (2+1)-nonlinear heat equation, J. Math. Anal. Appl. 339 (2008) 175-181.
- [44] M.K. Alaoui, S.A. Messaoudi, H.B. Khenous, A blow-up result for nonlinear generalized heat equation, Comput. Math. Appl. 68 (2014) 1723-1732.
- [45] N. Bellomo, L.M.D. Socio, R. Monaco, Random heat equation: solutions by the stochastic adaptive interpolation method, Comput. Math. Applic 16 (1988) 759-766.
- [46] J. Biazar, A.R. Amirtaimoori, An analytic approximation to the solution of heat equation by Adomian decomposition method and restrictions of the method, Appl. Math. Comput. 171 (2005) 738-745.
- [47] W.T. Ang, A.B. Gumel, A boundary integral method for the three-dimensional heat equation subject to specification of energy, J. Comput. Appl. Math. 135 (2001) 303-311.

- [48] T. Graen, H. Grubmüller, NuSol Numerical solver for the 3D stationary nuclear Schrödinger equation, *Comput. Phys. Commun.* 198 (2016) 169-178.
- [49] Y. Fang, X. You, Q. Ming, A new phase-fitted modified RungeCKutta pair for the numerical solution of the radial Schrödinger equation, *Appl. Math. Comput.* 224 (2013) 432-441.
- [50] R.M. Singh, S.B. Bhardwaj, S.C. Mishra, Closed-form solutions of the Schrödinger equation for a coupled harmonic potential in three dimensions, *Comput. Math. Appl.* 66 (2013) 537-541.
- [51] P. Bégout, J.I. Díaz, Localizing estimates of the support of solutions of some nonlinear Schrödinger equations C The stationary case, *Ann. I. H. Poincaré C AN.* 29 (2012) 35-58.
- [52] S. Cingolani, Semiclassical stationary states of Nonlinear Schrödinger equations with an external magnetic field, *J. Differ. Equations* 188 (2003) 52-79.
- [53] A.D. Polyanin and V.F. Zaitsev, *Handbook of Exact Solutions for Ordinary Differential Equations*, CRC Press, Florida (2003).
- [54] D.F. Gordon, B. Hafizi, A. S. Landsman, Amplitude flux, probability flux, and gauge invariance in the finite volume scheme for the Schrödinger equation, *J. Comput. Phys.* 280 (2015) 457-464.
- [55] S. Blanes, F. Casas, A. Murua, An efficient algorithm based on splitting for the time integration of the Schrödinger equation, *J. Comput. Phys.* 303 (2015) 396-412.
- [56] Z. Huang, J. Xu, B. Sun, B. Wu, X. Wu, A new solution of Schrödinger equation based on symplectic Algorithm, *Comput. Math. Appl.* 69 (2015) 1303-1312.
- [57] Z.A. Anastassi, T.E. Simos, A parametric symmetric linear four-step method for the efficient integration of the Schrödinger equation and related oscillatory problems, *J. Comput. Appl. Math.* 236 (2012) 3880-3889.
- [58] I.K. Gainullin, M.A. Sonkin, High-performance parallel solver for 3D time-dependent Schrodinger equation for large-scale nanosystems, *Comput. Phys. Commun.* 188 (2015) 68-75.
- [59] Diwaker, B. Panda, A. Chakraborty, Exact solution of Schrodinger equation for two state problem with time dependent coupling, *Phys. A.* 442 (2016) 380-387.
- [60] J. Lenells, Admissible boundary values for the defocusing nonlinear Schrödinger equation with asymptotically time-periodic data, *J. Differ. Equations* 259 (2015) 5617-5639.
- [61] X. -Y. Xie, B. Tian, W. -R. Sun, Y. Sun, Bright solitons for the (2+1)-dimensional coupled nonlinear Schrödinger equations in a graded-index waveguide, *Commun. Nonlinear Sci. Numer. Simulat.* 29 (2015) 300-306.
- [62] P. Wang, C. Huang, An energy conservative difference scheme for the nonlinear fractional Schrödinger equations, *J. Comput. Phys.* 293 (2015) 238-251.

- [63] S.Z. Rida, H.M. El-Sherbiny, A.A.M. Arafa, On the solution of the fractional nonlinear Schrödinger equation, *Phys. Lett. A* 372 (2008) 553-558.
- [64] K.M. Liang, *Mathematical and Physical Methods*, Higher Education Press, Beijing, China, 1978, pp. 162-163.
- [65] A.D. Polyanin, *Handbook of Linear Partial Differential Equations for Engineers and Scientists*, CRC Press, Florida, (2001).
- [66] S. V. Ershkov, Exact solution of Helmholtz equation for the case of non-paraxial Gaussian beams, *J. King Saud Univ. (Sci.)* 27 (2015) 198-203.
- [67] T.R. Ding, C.Z. Li, *Tutorials of Ordinary Differential Equations*, Second ed., Higher Education Press, Beijing, China, (2004).
- [68] W.A. Adkins, M.G. Davidson, *Ordinary Differential Equations*, Springer, New York, (2012).
- [69] L. C. Evans, *Partial Differential Equations*, American Mathematical Society, Providence, (1998).
- [70] J. Jost, *Partial differential equations*, Third ed., Springer, New York, (2013)
- [71] L. Hu, J. Zou, X. Fu, H. Y. Yang, X.D. Ruan, C.Y. Wang, Divisionally analytical solutions of Laplace's equations for dry calibration of electromagnetic velocity probes, *Appl. Math. Model.* 33 (2009) 3130-3150.
- [72] W.F. Ames, Invariant solutions of the underwater acoustic wave equation, *Comput. Math. Appl.* 11 (1985) 681-685.
- [73] T.F. Chan, L. Shen, A stable explicit scheme for the ocean acoustic wave equation, *Comput. Math. Appl.* 11 (1985) 929-936.
- [74] A.D. Polyanin, V.F. Zaitsev, A. Moussiaux, *Handbook of First Order Partial Differential Equations*, CRC Press, Florida, (2001), pp.4, pp.62.
- [75] S.H. Guo, *Electrodynamics*, Higher Education Press, Beijing, China, 1995.
- [76] M. Li, C.S. Chen, Y.C. Hon, P.H. Wen, Finite integration method for solving multi-dimensional partial differential equations, *Appl. Math. Model.* 39 (2015) 4979-4994.
- [77] P.H. Wen, Y.C. Hon, M. Li, T. Korakianitis, Finite integration method for partial differential equations, *Appl. Math. Model.* 37 (2013) 10092-10106.
- [78] F. Toutounian, E. Tohidi, A new Bernoulli matrix method for solving second order linear partial differential equations with the convergence analysis, *Appl. Math. Comput.* 223 (2013) 298-310.
- [79] C. Kesan, Chebyshev polynomial solutions of second-order linear partial differen-

- tial equations, *Appl. Math. Comput.* 134 (2003) 109-124.
- [80] L. Ehrenpreis, *Fourier Analysis in Several Complex Variables*, Wiley-Interscience, New York, 1970.
- [81] P.-C. Hu, B.Q. Li, On meromorphic solutions of linear partial differential equations of second order, *J. Math. Anal. Appl.* 393 (2012) 200-211.
- [82] D.G. Aronson, Isolated singularities of solutions of second order parabolic equations, *Arch. Rat. Mech. Anal.* 19 (1965) 231-238
- [83] N. Lungu, D. Popa, Hyers-Ulam stability of a first order partial differential equation, *J. Math. Anal. Appl.* 385 (2012) 86-91.
- [84] Z. Sheng, The periodic wave solutions for the (2 + 1) dimensional Konopelchenko-Dubrovsky equations, *Chaos Solitons Fract.* 30 (2006) 1213-1220.
- [85] W.X. Ma, Travelling wave solutions to a seventh order generalized KdV equation, *Phys. Lett. A.* 180 (1993) 221-224.
- [86] W. Malfliet, Solitary wave solutions of nonlinear wave equations, *Amer. J. Phys.* 60 (7) (1992) 650-654.
- [87] A.M. Wazwaz, Two reliable methods for solving variants of the KdV equation with compact and noncompact structures, *Chaos Solitons Fract.* 28 (2) (2006) 454-462.
- [88] S.A. El-Wakil, M.A. Abdou, New exact travelling wave solutions using modified extended tanh-function method, *Chaos Solitons Fract.* 31 (4) (2007) 840-852.
- [89] E. Fan, Extended tanh-function method and its applications to nonlinear equations, *Phys. Lett. A* 277 (4-5) (2000) 212-218.
- [90] A.M. Wazwaz, The tanh-function method: Solitons and periodic solutions for the Dodd-Bullough-Mikhailov and the Tzitzeica-Dodd-Bullough equations, *Chaos Solitons Fract.* 25 (1) (2005) 55-63.
- [91] W.X. Ma, T.W. Huang, Y. Zhang, A multiple exp-function method for nonlinear differential equations and its application, *Phys. Scr.* 82 (2010) 065003.
- [92] I. E. Inan, Y. Ugurlu, Exp-function method for the exact solutions of fifth order KdV equation and modified Burgers equation, *Appl. Math. Comput.* 217 (2010) 1294-1299.
- [93] T.C. Xia, B. Li, H.Q. Zhang, New explicit and exact solutions for the Nizhnik-Novikov-Vesselov equation, *Appl. Math. E-Notes* 1 (2001) 139-142.
- [94] M. Inc, M. Ergut, Periodic wave solutions for the generalized shallow water wave equation by the improved Jacobi elliptic function method, *Appl. Math. E-Notes* 5 (2005) 89-96.
- [95] M. L. Wang, Solitary wave solution for variant Boussinesq equation, *Phys. Lett.*

A 199 (1995) 169-172.

[96] M. Khalfallah, New exact traveling wave solutions of the $(3 + 1)$ dimensional Kadomtsev-Petviashvili (KP) equation, *Commun. Nonlinear Sci. Numer. Simul.* 14 (2009) 1169-1175.

[97] E.G. Fan, Two new applications of the homogeneous balance method, *Phys. Lett. A* 265 (2000) 353-357.

[98] A.M. Wazwaz, A sine-cosine method for handling nonlinear wave equations, *Math. Comput. Model.* 40 (5-6) (2004) 499-508.

[99] E. Yusufoglu, A. Bekir, Solitons and periodic solutions of coupled nonlinear evolution equations by using sine-cosine method, *Int. J. Comput. Math.* 83 (12) (2006) 915-924.

[100] A.M. Wazwaz, The sine-cosine method for obtaining solutions with compact and noncompact structures, *Appl. Math. Comput.* 159 (2) (2004) 559-576.

[101] M.K. Mak, T. Harko, New method for generating general solution of the Abel differential equation, *Comput. Math. Appl.* 43 (2002) 91-94.

[102] M.K. Mak, H.W. Chan, T. Harko, Solutions generating technique for Abel-type nonlinear ordinary differential equations, *Comput. Math. Appl.* 41 (10/11), (2001), 1395-1401.