

SERIES REPRESENTATION OF POWER FUNCTION

KOLOSOV PETRO

ABSTRACT. In this paper described numerical expansion of natural-valued power function x^n , in point $x = x_0$ where n, x_0 - natural numbers. Applying numerical methods, that is calculus of finite differences, namely, discrete case of Binomial expansion is reached. Received results were compared with solutions according to Newton's Binomial theorem and MacMillan Double Binomial sum. Additionally, in section 4 exponential function's e^x representation is shown.

Keywords. Power function, Binomial coefficient, Binomial Theorem, Finite difference, Perfect cube, Exponential function, Pascal's triangle, Series representation, Binomial Sum, Multinomial theorem, Multinomial coefficient, Binomial distribution

2010 Math. Subject Class. 30BXX

e-mail: kolosov_94@mail.ua

ORCID: 0000-0002-6544-8880

CONTENTS

1. Introduction and main results	1
1.1. Binomial Theorem based on definition (1.20)	9
1.2. Multinomial case for $n = 4$	9
1.3. Generalized Multinomial Theorem based on definition (1.20)	9
1.4. Generalized Binomial Series by means of properties (1.24), (1.25)	10
1.5. Comparison of (1.31) to Binomial Theorem and MacMillan Double Binomial sum	10
2. e^x Representation	11
3. Conclusion and future research	11
References	12
4. Application 1. Wolfram Mathematica 11 codes of some expressions	12
5. Application 2. An extended version of triangle (3)	15

1. INTRODUCTION AND MAIN RESULTS

In this paper coefficient $\left\| \begin{matrix} n \\ k \end{matrix} \right\|$ is introduced and its properties are shown, one of which is power's expansion. We will generalize the partial case of power $n = 3$, received using $\left\| \begin{matrix} n \\ k \end{matrix} \right\|$ to high power $u > 3$. First, let basically describe Newton's

Date: November 23, 2017.

Binomial Theorem, since our coefficient $\left\lfloor \begin{matrix} n \\ k \end{matrix} \right\rfloor$ is derived from finite difference of perfect cubes, which is taken regarding Binomial expansion. In elementary algebra, the Binomial theorem describes the algebraic expansion of powers of a binomial. The theorem describes expanding of the power of $(x + y)^n$ into a sum involving terms of the form ax^by^c where the exponents b and c are nonnegative integers with $b+c = n$, and the coefficient a of each term is a specific positive integer depending on n and b . The coefficient a in the term of ax^by^c is known as the Binomial coefficient. The main properties of the Binomial Theorem are next

Properties 1.1. *Binomial Theorem properties*

- (1) *The powers of x go down until it reaches $x_0 = 1$ starting value is n (the n in $(x + y)^n$)*
- (2) *The powers of y go up from 0 ($y^0 = 1$) until it reaches n (also n in $(x + y)^n$)*
- (3) *The n -th row of the Pascal's Triangle (see [1], [13]) will be the coefficients of the expanded binomial.*
- (4) *For each line, the number of products (i.e. the sum of the coefficients) is equal to $x + 1$*
- (5) *For each line, the number of product groups is equal to 2^n*

According to the Binomial theorem, it is possible to expand any power of $x + y$ into a sum of the form (see [2], [4])

$$(1.2) \quad (x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$$

Let expand monomial x^n such that $(x, n) \in \mathbb{N}$ applying finite differences, that are reached by means of Binomial theorem (1.2)

Lemma 1.3. *Power function could be represented as discrete integral of its first order finite difference*

$$(1.4) \quad \begin{aligned} x^n &= \int_0^{x-1} \Delta_h[x^n]h \\ &= \sum_{k=0}^{x-1} \underbrace{nk^{n-1}h + \binom{n}{2}k^{n-2}h^2 + \dots + \binom{n}{n-1}kh^{n-1} + h^n}_{\Delta_h[x^n] = (x+h)^n - x^n} \\ &= \sum_{j=0}^{x-1} \sum_{k=1}^n \binom{n}{k} j^{n-k} h^k, \quad h \in \mathbb{R} \end{aligned}$$

We can reach the same result using Fundamental Theorem of Calculus

$$(1.5) \quad x^n = \int_0^x nt^{n-1} dt = \sum_{k=0}^{x-1} \int_k^{k+1} nt^{n-1} dt = \sum_{k=0}^{x-1} (k+1)^n - k^n$$

Lemma 1.6. *From lemma (1.4) follows that finite difference of power x^n , $n \in \mathbb{N}$ could be reached by Binomial expansion of the form*

$$(1.7) \quad \Delta_h[x^n] = (x + h)^n - x^n = \sum_{k=1}^n \binom{n}{k} x^{n-k} h^k$$

Otherwise, let be a difference table of perfect cubes (see also [6], eq. 7) $\Delta^k[x^3]$, $x \in \mathbb{N}$, $k \in [1, 3] \subseteq \mathbb{N}$ over x from $[0, 7] \subseteq \mathbb{N}$

x	x^3	$\Delta[x^3]$	$\Delta^2[x^3]$	$\Delta^3[x^3]$
0	0	1	6	6
1	1	7	12	6
2	8	19	18	6
3	27	37	24	6
4	64	61	30	6
5	125	91	36	
6	216	127		
7	343			

Figure 1: Difference table of x^3 , $x \in \mathbb{N}$ up to 3rd order, [6], eq. 7

Note that increment h is set to be $h = 1$ and $k > 2$ -order difference is taken regarding to [10]. From Figure 1 the regularity concerning sequence A008458, [8] in OEIS is observed, [9], that is the first order finite difference of consequent perfect cubes equals to $1 + a(n)$, where $a(n)$ is generating function of sequence A008458. Let show, that

$$(1.9) \quad \Delta_{h=1}[x^3] = (x+1)^3 - x^3 = 1 + 3! \cdot 0 + 3! \cdot 1 + \dots + 3! \cdot x$$

Applying compact sigma notation on (1.9), by means of identities,

$$(1.10) \quad \begin{aligned} \Delta_{h=1}[x^3] &= 1 + 3! \cdot 1 + 3! \cdot 2 + \dots + 3! \cdot x \\ \nabla_{h=1}[x^3] &= 1 + 3! \cdot 0 + 3! \cdot 1 + \dots + 3! \cdot (x-1) \end{aligned}$$

the first order forward, as well as backward finite difference of consequent perfect cubes could be represented as

$$(1.11) \quad \begin{aligned} \Delta_{h=1}[x^3] &= \sum_{k=1}^x 3! \cdot k + \frac{1}{x} = 3! \binom{x+1}{2} + 1 \\ &= \sum_{k=1}^x 3! \binom{k}{k-1} + \frac{1}{x}, \quad x = x_0 \in \mathbb{N}, x \neq 0 \end{aligned}$$

$$(1.12) \quad \begin{aligned} \nabla_{h=1}[x^3] &= \sum_{k=0}^{x-1} 3! \cdot k + \frac{1}{x} \\ &= \sum_{k=0}^{x-1} 3! \binom{k}{k-1} + \frac{1}{x}, \quad x = x_0 \in \mathbb{N}, x \neq 0 \end{aligned}$$

By lemma (1.3) discrete integral of first difference is used to reach expansion of monomial x^n , $h \in \mathbb{R}$ and we have an equality between (1.7) and (1.11)

$$(1.13) \quad \Delta_{h=1}[x^3] = \sum_{k=1}^3 \binom{3}{k} x^{3-k} = \sum_{k=1}^x 3! \cdot k + \frac{1}{x}, \quad x = x_0 \in \mathbb{N}, x \neq 0$$

Then we have right to substitute (1.11) into (1.4) instead binomial expansion and represent perfect cube x^3 as summation over k from 0 to $x-1$. Let be extended

example of expansion of x^3 , $x \in \mathbb{N}$

$$\begin{aligned}
 (1.14) \quad x^3 &= (1 + 3! \cdot 0) + (1 + 3! \cdot 0 + 3! \cdot 1) + (1 + 3! \cdot 0 + 3! \cdot 1 + 3! \cdot 2) \\
 &\quad \cdots + (1 + 3! \cdot 0 + 3! \cdot 1 + 3! \cdot 2 + \cdots + 3! \cdot (x-1)) \\
 &= x + (x-0) \cdot 3! \cdot 0 + (x-1) \cdot 3! \cdot 1 + (x-2) \cdot 3! \cdot 2 \\
 &\quad \cdots + (x-(x-1)) \cdot 3! \cdot (x-1)
 \end{aligned}$$

Provided that x is natural. Note that increment h in equation (1.14) is set to be $h = 1$ and not displayed. Using compact sigma notation on (1.14), we received

$$\begin{aligned}
 (1.15) \quad x^3 &= x + 3! \cdot \sum_{m=0}^{x-1} mx - m^2 = x + 3! \cdot \sum_{m=0}^x mx - m^2 \\
 &= x + 3! \cdot \sum_{m=1}^x mx - m^2 = \sum_{m=0}^{x-1} 3! \cdot mx - 3! \cdot m^2 + 1 \\
 &= \sum_{m=1}^x 3! \cdot mx - 3! \cdot m^2 + 1, \quad x \in \mathbb{N}, n \in \mathbb{N}
 \end{aligned}$$

Property 1.16. Let be a sets $\mathfrak{S}(x) := \{1, 2, \dots, x\} \subseteq \mathbb{N}$, $\mathfrak{C}(x) := \{0, 1, \dots, x\} \subseteq \mathbb{N}$, $\mathfrak{U}(x) := \{0, 1, \dots, x-1\} \subseteq \mathbb{N}$, let be right part of (1.15) written as

$$T(x, \mathfrak{U}(x)) := \sum_{m \in \mathfrak{U}(x)} 3! \cdot mx - 3! \cdot m^2 + 1$$

where $x \in \mathbb{N}$ is variable and $\mathfrak{U}(x)$ is iteration set of (1.15), then we have equality

$$(1.17) \quad T(x, \mathfrak{U}(x)) = T(x, \mathfrak{S}(x)), \quad x \in \mathbb{N}$$

Let be central part of (1.15) denoted as

$$U(x, \mathfrak{C}(x)) := x + 3! \cdot \sum_{m \in \mathfrak{C}(x)} mx - m^2,$$

then

$$(1.18) \quad U(x, \mathfrak{C}(x)) = U(x, \mathfrak{S}(x)) = U(x, \mathfrak{U}(x))$$

Other words, change of iteration sets of central and right parts of (1.15) by $\mathfrak{C}(x)$, $\mathfrak{S}(x)$, $\mathfrak{U}(x)$ and $\mathfrak{U}(x)$, $\mathfrak{S}(x)$ respectively, doesn't change resulting value of (1.15) given $x \in \mathbb{N}$.

Proof. Let be a plot of $y = 3! \cdot kx - 3! \cdot k^2 + 1$ by k over $\mathbb{R}_{\leq 10}^+$, given $x = 10$




Figure 2. Plot of $y = 3! \cdot kx - 3! \cdot k^2 + 1$ by k over $\mathbb{R}_{\leq 10}^+$, given $x = 10$

Obviously, being a parabolic function, it's symmetrical over $\frac{x}{2}$, hence equivalent $T(x, \mathfrak{U}(x)) = T(x, \mathfrak{S}(x))$, $x \in \mathbb{N}$ follows. Reviewing central part of (1.15) and denote $u(t) = tx^{n-2} - t^2x^{n-3}$, we can conclude, that $u(0) = u(x)$, then equality of $U(x, \mathfrak{C}(x)) = U(x, \mathfrak{S}(x)) = U(x, \mathfrak{U}(x))$ immediately follows. This completes the proof. \square

Note that for right part of (1.15) property (1.16) holds only in case (1.17). Let analyse property (1.16). Let be an example of triangle¹ built using right part of expression (1.15) over x from 0 up to $x = 4$. We can see that for each row corresponded to variable $x = 0, 1, 2, 3, 4, \dots$, we have Binomial distribution of row items, more detailed




Figure 3. Triangle generated by $3! \cdot kx - 3! \cdot k^2 + 1$, that is right part of (1.15) from 0 up $x = 4$ over k from 0 to x , sequence A287326 in OEIS, [9], [12].

One could compare Figure (1.19) with Pascal's triangle [1], [13]

¹Note that each $x = 0, 1, 2, 3, \dots, \mathbb{N}$ row's item of triangle is generated by $3! \cdot kx - 3! \cdot k^2 + 1$ over $k \in [0, x] \subseteq \mathbb{N}$

$$\begin{array}{cccccc}
& & & & & 1 \\
& & & & & 1 & 1 \\
& & & & 1 & 2 & 1 \\
& & 1 & 3 & 3 & 1 \\
1 & 4 & 6 & 4 & 1
\end{array}$$

Figure 4. Pascal's triangle built up given $n = 4$ over k from 0 to n , sequence A007318 in OEIS, [9], [1], [13].

Review the triangle (1.19), define the k -th, $0 \leq k \leq n$, item of n -th row of triangle as

Definition 1.20.

$$(1.21) \quad \left\| \begin{array}{c} n \\ k \end{array} \right\| := 3! \cdot nk - 3! \cdot k^2 + 1, \quad 0 \leq k \leq n$$

Note that definition (1.20) also could be rewritten as

$$(1.22) \quad \left\| \begin{array}{c} n \\ k \end{array} \right\| = 3! \cdot nk - 3! \cdot n^0 k^2 + n^0, \quad 0 \leq k \leq n$$

Let us approach to show a few properties of triangle (1.19)

Properties 1.23. *Properties of triangle (1.19). For each $(n, m) \in \mathbb{N}$ holds*

- (1) *Summation of items $\left\| \begin{array}{c} n \\ k \end{array} \right\|$ of n -th row of triangle (1.19) over k from 0 to $n - 1$ returns perfect cube n^3 as binomial of the form*

$$(1.24) \quad \sum_{k=0}^{n-1} \left\| \begin{array}{c} n \\ k \end{array} \right\| = A_{0,n}n - B_{0,n} = n^3,$$

Since the property (1.17) holds, (1.24) could be rewritten as

$$(1.25) \quad \sum_{k=1}^n \left\| \begin{array}{c} n \\ k \end{array} \right\| = A_{1,n}n - B_{1,n} = n^3,$$

*where $A_{\overline{0},1,n}$ and $B_{\overline{0},1,n}$ - integers **depending** on variable $n \in \mathbb{N}$ and on sets $\mathfrak{A}(n)$, $\mathfrak{B}(n)$, respectively.*

- (2) *Recurrence relation between $A_{0,n}$ and $A_{1,n}$*

$$A_{0,n+1} = A_{1,n}, \quad n \geq 1$$

- (3) *Induction by power. Summation of items of n -th row of triangle (1.19), multiplied by n^{m-3} , from 0 to $n - 1$ returns n^m*

- (4) *Summation of items $\left\| \begin{array}{c} n \\ k \end{array} \right\|$ of n -th row of triangle (1.19) over k from 0 to n returns $n^3 + n^0$*

$$(1.26) \quad \sum_{k=0}^n \left\| \begin{array}{c} n \\ k \end{array} \right\| = n^3 + 1$$

- (5) *Induction by power. Summation of each n -th row of triangle (1.19) multiplied by n^{m-3} from 0 to n returns $n^m + n^{m-3}$*

- (6) First item of each row's number corresponding to central polygonal numbers sequence $a(n)$ (sequence A000124 in OEIS, [9], [14]) returns finite difference $\Delta[n^3]$ of consequent perfect cubes. For example, let be the k -th row of triangle (1.19), such that k is central polygonal number, i.e $k = \frac{n^2+n+2}{2}$, $n = 0, 1, 2, \dots, \mathbb{N}$, then item

$$(1.27) \quad \left\| \left\| \frac{n^2+n+2}{2} \right\| \right\| = \Delta_h[n^3], \quad h = 1$$

- (7) Items of (1.19) have Binomial distribution over rows
 (8) Triangle (1.19) being arranged as right triangle has the k -th column generating formula $x^k \cdot (1 + (6 \cdot k - 1) \cdot x)/(1 - x)^2$, [12].
 (9) The linear recurrence, for any k and $n > 0$

$$(1.28) \quad 2 \left\| \left\| n \right\| \right\| = \left\| \left\| n+1 \right\| \right\| + \left\| \left\| n-1 \right\| \right\|$$

- (10) Linear recurrence, for each $n > k$

$$(1.29) \quad 2 \left\| \left\| n \right\| \right\| = \left\| \left\| 2n-k \right\| \right\| + \left\| \left\| 2n-k \right\| \right\|$$

- (11) From (1.27) follows that

$$(1.30) \quad \sum_{k=0}^{n-1} \left\| \left\| n \right\| \right\| = \sum_{k=0}^{n-1} \left\| \left\| \frac{k^2+k+2}{2} \right\| \right\|$$

In (1.24) is noticed, that summation of each n -th row of Triangle (1.19) from 0 to $n-1$ returns perfect cube n^3 , then, by properties (1.27), (1.28), (1.29), for each given number $x \in \mathbb{N}$ the x^n could be easy found via multiplication of each term of (1.24) by x^{n-3}

$$(1.31) \quad \begin{aligned} x^n &= \sum_{k=0}^{x-1} \left\| \left\| x \right\| \right\| x^{n-3} = \frac{1}{2} \sum_{k=0}^{x-1} \left[\left\| \left\| x+1 \right\| \right\| + \left\| \left\| x-1 \right\| \right\| \right] x^{n-3} \\ &= \sum_{k=0}^{x-1} \frac{1}{2} \left[\left\| \left\| 2x-k \right\| \right\| + \left\| \left\| 2x-k \right\| \right\| \right] x^{n-3} \\ &= \sum_{k=0}^{x-1} \frac{1}{2} \left\| \left\| \frac{x^2+x+2}{2} \right\| \right\| x^{n-3} = \sum_{k=0}^{x-1} \frac{1}{2} \left[\left\| \left\| \frac{x^2+x}{2} \right\| \right\| + \left\| \left\| \frac{x^2+x+4}{2} \right\| \right\| \right] x^{n-3} \\ &= \sum_{k=0}^{x-1} \frac{1}{2} \left[\left\| \left\| \binom{n+1}{2} \right\| \right\| + \left\| \left\| \binom{n+1}{2} + \binom{2}{1} \right\| \right\| \right] x^{n-3} \end{aligned}$$

To show other way of representation of power, let move the x from (1.15), $x + 3! \sum mx - m^2$, under the sum operator and change iteration set from $\{0, x-1\}$ to $\{1, x-1\}$, then we get

$$(1.32) \quad x^3 = \sum_{m=1}^{x-1} 3! \cdot mx - 3! \cdot m^2 + \frac{x}{(x-1)}, \quad x \neq 1, \quad x \in \mathbb{N}$$

Review right part of (1.32), let be item $\frac{x}{x-1}$ written as $\frac{x}{x-1} = \frac{x+1-1}{x-1} = 1 + \frac{1}{x-1}$, given the power $n > 3$, multiplying each term of (1.32) by x^{n-3} we can observe

that

$$(1.33) \quad x^n - 1 = \sum_{m=1}^{x-1} \left\| \begin{matrix} x \\ m \end{matrix} \right\| x^{n-3} + x^{n-4} + x^{n-5} + \dots + x + 1$$

Applying properties (1.27), (1.29), (1.30), we can rewrite (1.22) as

$$(1.34) \quad \begin{aligned} x^n - 1 &= \sum_{k=1}^{x-1} \frac{1}{2} \left[\left\| \begin{matrix} 2x-k \\ k \end{matrix} \right\| + \left\| \begin{matrix} 2x-k \\ 0 \end{matrix} \right\| \right] x^{n-3} + x^{n-4} + \dots + x + 1 \\ &= \sum_{k=1}^{x-1} \frac{1}{2} \left[\left\| \begin{matrix} x+1 \\ k \end{matrix} \right\| + \left\| \begin{matrix} x-1 \\ k \end{matrix} \right\| \right] x^{n-3} + x^{n-4} + \dots + x + 1 \\ &= \sum_{m=0}^{x-1} \frac{1}{2} \left[\left\| \begin{matrix} m^2+m \\ 1 \end{matrix} \right\| + \left\| \begin{matrix} m^2+m+4 \\ 1 \end{matrix} \right\| \right] x^{n-3} + x^{n-4} + \dots + x + 1 \\ &= \sum_{k=1}^{x-1} \left\| \begin{matrix} k^2+k+2 \\ 1 \end{matrix} \right\| x^{n-3} + x^{n-4} + \dots + x + 1 \end{aligned}$$

Moving 1 from left part of (1.33) under sum operator, we add a term $\frac{1}{x-1}$ to initial function $\left\| \begin{matrix} x \\ m \end{matrix} \right\| x^{n-3} + x^{n-4} + x^{n-5} + \dots + x + 1$. By means of expansion $\frac{1}{1-x} = -\frac{1}{x-1} = 1 + x + x^2 + x^3 + \dots$, the (1.33) could be rewritten accordingly

$$(1.35) \quad x^n = \sum_{m=1}^{x-1} \left\| \begin{matrix} x \\ m \end{matrix} \right\| x^{n-3} + x^{n-4} + \dots + x + 1 - 1 - x^2 - x^3 - \dots$$

Generalizing (1.35) we have

$$(1.36) \quad \begin{aligned} x^n &= \sum_{m=1}^{x-1} \left[\left\| \begin{matrix} x \\ m \end{matrix} \right\| - 1 \right] x^{n-3} - x^{n-2} - x^{n-1} - \dots \\ &= \sum_{m=1}^{x-1} \left[\left\| \begin{matrix} m^2+m+2 \\ 1 \end{matrix} \right\| - 1 \right] x^{n-3} - x^{n-2} - x^{n-1} - \dots \end{aligned}$$

A product generalization of even and odd power for each $x \in \mathbb{N}$, in case we define $\left\| \begin{matrix} n \\ k \end{matrix} \right\|$ as (1.22)

$$(1.37) \quad x^n = \begin{cases} \prod_{i=0}^j \sum_{k=0}^{x-1} \left\| \begin{matrix} x^0 \\ k \end{matrix} \right\|, & n = 2j, j > 0, n \in \mathbb{N} \\ \prod_{i=0}^j \sum_{k=0}^{x-1} \left\| \begin{matrix} x^0 \\ k \end{matrix} \right\| x^{\frac{1}{j}}, & n = 2j + 1, j > 0, n \in \mathbb{N} \end{cases}$$

1.1. Binomial Theorem based on definition (1.20). Let be two positive integers x, y raised to power $n \in \mathbb{N}$, then, using definition 1.20, we receive

$$\begin{aligned}
 (1.38) \quad (x+y)^n &= \sum_{k=0}^{x+y-1} \sum_{j=0}^{n-3} \left\| \begin{matrix} x+y \\ k \end{matrix} \right\| \binom{n-3}{j} x^{n-3-j} y^j \\
 &= \frac{1}{2} \sum_{k=0}^{x+y-1} \sum_{j=0}^{n-3} \left[\left\| \begin{matrix} x+y+1 \\ k \end{matrix} \right\| + \left\| \begin{matrix} x+y-1 \\ k \end{matrix} \right\| \right] \binom{n-3}{j} x^{n-3-j} y^j \\
 &= \frac{1}{2} \sum_{k=0}^{x+y-1} \sum_{j=0}^{n-3} \left[\left\| \begin{matrix} 2(x+y)-k \\ k \end{matrix} \right\| + \left\| \begin{matrix} 2(x+y)-k \\ 0 \end{matrix} \right\| \right] \binom{n-3}{j} x^{n-3-j} y^j \\
 &= \sum_{k=0}^{x+y-1} \sum_{j=0}^{n-3} \left\| \begin{matrix} k^2+k+2 \\ 1 \end{matrix} \right\| \binom{n-3}{j} x^{n-3-j} y^j \\
 &= \sum_{k=0}^{x+y-1} \frac{1}{2} \sum_{j=0}^{n-3} \left[\left\| \begin{matrix} k^2+k \\ 1 \end{matrix} \right\| + \left\| \begin{matrix} k^2+k+4 \\ 1 \end{matrix} \right\| \right] \binom{n-3}{j} x^{n-3-j} y^j
 \end{aligned}$$

Mathematica code of above expression is attached in Application 1.

1.2. Multinomial case for $n = 4$. . For each $u, j, \dots, m \in \mathbb{N}$ and $n = 4$ holds

$$\begin{aligned}
 (u+j+\dots+m)^4 &= \sum_{k=0}^{u+j+\dots+m-1} \left\| \begin{matrix} u+j+\dots+m \\ k \end{matrix} \right\| u + \left\| \begin{matrix} u+j+\dots+m \\ k \end{matrix} \right\| j + \dots \\
 &\quad \dots + \left\| \begin{matrix} u+j+\dots+m \\ k \end{matrix} \right\| m
 \end{aligned}$$

1.3. Generalized Multinomial Theorem based on definition (1.20). Expression (1.21) could be generalized accordingly

$$\begin{aligned}
 (1.39) \quad &(u+j+\dots+m)^n \\
 = &\sum_{v=0}^{u+j+\dots+m-1} \sum_{k_u+k_j+\dots+k_m=n-4} \binom{n-4}{k_u, k_j, \dots, k_m} \left\| \begin{matrix} u+j+\dots+m \\ v \end{matrix} \right\| u^{k_u+1} j^{k_j} \dots m^{k_m} \\
 + &\sum_{k_u+k_j+\dots+k_m=n-4} \binom{n-4}{k_u, k_j, \dots, k_m} \left\| \begin{matrix} u+j+\dots+m \\ v \end{matrix} \right\| u^{k_u} j^{k_j+1} \dots m^{k_m} + \dots \\
 \dots &\sum_{k_u+k_j+\dots+k_m=n-4} \binom{n-4}{k_u, k_j, \dots, k_m} \left\| \begin{matrix} u+j+\dots+m \\ v \end{matrix} \right\| u^{k_u} j^{k_j} \dots m^{k_m+1}
 \end{aligned}$$

where

$$(1.40) \quad \binom{n}{k_u, k_j, \dots, k_m} = \frac{n!}{k_u! k_j! \dots k_m!}$$

is multinomial coefficient.

1.4. Generalized Binomial Series by means of properties (1.24), (1.25).

Reviewing properties (1.24), (1.25) we can say that

$$(1.41) \quad x^n = A_{0,1,x} x^{n-2} - B_{0,1,x} x^{n-3}, \quad x = x_0 \in \mathbb{N}$$

Rewrite the right part of (1.42) regarding to itself as recursion

$$(1.42) \quad \begin{aligned} x^n &= A_{0,1,x} (A_{0,1,x} x^{n-4} - B_{0,1,x} x^{n-5}) - B_{0,1,x} (A_{0,1,x} x^{n-5} - B_{0,1,x} x^{n-6}) \\ &= A_{0,1,x}^2 x^{n-4} - 2A_{0,1,x} B_{0,1,x} x^{n-5} + B_{0,1,x}^2 x^{n-6} \end{aligned}$$

Reviewing (1.43) we can observe binomial coefficients before each $A_{0,1,x} \cdot B_{0,1,x}$. Continuous j -times recursion of right part of (1.42) gives us

$$(1.43) \quad x^n = \sum_{k=0}^j (-1)^k \binom{j}{k} A_{0,1,x}^{j-k} B_{0,1,x}^k x^{n-2j-k}$$

Suppose that we want to repeat action (1.43) infinite-many times, then

$$\begin{aligned} x^n &= \binom{j}{0} A_{0,1,x}^j B_{0,1,x}^0 x^{n-2j} - \binom{j}{1} A_{0,1,x}^{j-1} B_{0,1,x}^1 x^{n-2j-1} + \dots \\ &+ \binom{j}{2} A_{0,1,x}^{j-2} B_{0,1,x}^2 x^{n-2j-2} - \dots + \sum_k^{\infty} (-1)^k \binom{j}{k} A_{0,1,x}^{j-k} B_{0,1,x}^k x^{n-2j-k} + \dots \end{aligned}$$

We know the solutions of above equation (1.42) for all $A_{0,1,x}$, $B_{0,1,x}$ that present in follow table. The table arranged next way

$x =$	$A_{0,x} =$	$B_{0,x} =$	$A_{1,x} =$	$B_{1,x} =$
1	1	0	6	5
2	6	4	18	28
3	18	25	36	81
4	36	80	60	176
5	60	175	90	325
6	90	324	126	540
7	126	539	168	833
8	168	832	216	1216
9	216	1215	270	1701
10	270	1700	330	2300

Table 1. Array of coefficients $A_{0,1,x}$, $B_{0,1,x}$ over x from 1 to 10.

Sequence $B_{1,x}$ is generated by $2x^3 + 3x^2$, $x \in \mathbb{N}$, sequence A275709 in OEIS, [9]. Sequence $A_{1,x}$ is generated by $3x^2 + 3x$, $x \in \mathbb{N}$, sequence A028896 in OEIS, [9].

1.5. Comparison of (1.31) to Binomial Theorem and MacMillan Double Binomial sum. In this subsection let compare (1.31) to Binomial Theorem and MacMillan Double Binomial sum² (see [3], eq. 12) hereby follow changes hold

²MacMillan Double Binomial sum is the other one numerical expansion of the form

$$\sum_{k=1}^n \sum_{j=1}^k (-1)^{k-j} j^n \binom{k}{j} \binom{x}{k} = x^n$$

- (1) While expansion is taken by Binomial theorem or MacMillan Double Binomial sum, the summation (1.2) is done over $k = 0, 1, 2, \dots, \mathbb{N}$ from 0 and 1 to n , respectively, where $n \in \mathbb{N}$ - power. Reviewing (1.31) we can observe that summation is taken over k from 0 to x , where x - variable, provided that natural-valued.
- (2) According to property (1.16) the iteration sets of right and central parts (1.15) could be changed to (1.17), (1.18) without changes in result.
- (3) We observed that expansion (1.31) returns the power of each particular number $x \in \mathbb{N}$ as binomial of the form

$$(1.44) \quad A_{0,1,x} x^{n-2} - B_{0,1,x} x^{n-3},$$

which holds in particular case $x = x_0 \in \mathbb{N}$.

2. e^x REPRESENTATION

Since the exponential function $f(x) = e^x$, $x \in \mathbb{N}$ is defined as infinite summation of $\frac{x^n}{n!}$, $n = 0, 1, 2, \dots, \infty$ over n (see [5]). Then (1.31) could be applied, hereby

$$(2.1) \quad \begin{aligned} e^x &= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{m=1}^{x-1} \left[\left\| \begin{matrix} x \\ m \end{matrix} \right\| - 1 \right] x^{n-3} - x^{n-2} - x^{n-1} - \dots \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{m=1}^{x-1} \left[\left\| \begin{matrix} \frac{m^2+m+2}{2} \\ 1 \end{matrix} \right\| - 1 \right] x^{n-3} - x^{n-2} - x^{n-1} - \dots \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{k=1}^{x-1} \left[\frac{1}{2} \left[\left\| \begin{matrix} x+1 \\ k \end{matrix} \right\| + \left\| \begin{matrix} x-1 \\ k \end{matrix} \right\| \right] - 1 \right] x^{n-3} - x^{n-2} - x^{n-1} - \dots \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{k=1}^{x-1} \left[\frac{1}{2} \left[\left\| \begin{matrix} 2x-k \\ k \end{matrix} \right\| + \left\| \begin{matrix} 2x-k \\ 0 \end{matrix} \right\| \right] - 1 \right] x^{n-3} - x^{n-2} - x^{n-1} - \dots \end{aligned}$$

From (2.1) we get

$$(2.2) \quad \begin{aligned} e^x - e &= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{m=1}^{x-1} \left\| \begin{matrix} x \\ m \end{matrix} \right\| x^{n-3} + x^{n-4} + \dots + x + 1 \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{m=1}^{x-1} \frac{1}{2} \left[\left\| \begin{matrix} x+1 \\ m \end{matrix} \right\| + \left\| \begin{matrix} x-1 \\ m \end{matrix} \right\| \right] x^{n-3} + x^{n-4} + \dots + x + 1 \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{m=1}^{x-1} \left\| \begin{matrix} \frac{m^2+m+2}{2} \\ 1 \end{matrix} \right\| x^{n-3} + x^{n-4} + \dots + x + 1 \\ &= \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{m=1}^{x-1} \frac{1}{2} \left[\left\| \begin{matrix} \frac{m^2+m}{2} \\ 1 \end{matrix} \right\| + \left\| \begin{matrix} \frac{m^2+m+4}{2} \\ 1 \end{matrix} \right\| \right] x^{n-3} + x^{n-4} + \dots + x + 1 \end{aligned}$$

3. CONCLUSION AND FUTURE RESEARCH

In this paper the coefficient $\left\| \begin{matrix} n \\ k \end{matrix} \right\|$ was introduced in definition (1.20), its properties are shown in (1.23). Power function's x^n , $x, n \in \mathbb{N}$ expansion firstly shown in (1.20) and other versions are shown below. Going from (1.31) Power's expansion applying (5) from properties (1.23) is shown below the subsection (1.3), "Generalized Multinomial Theorem based definition (1.20)". In subsection (1.1) Binomial

Theorem based on definition (1.20) is shown. In section 2 we show a various representations of exponential function e^x , $x \in \mathbb{N}$. In subsection (1.2) we put example of Multinomial theorem for $n = 4$, based on $\left\| \begin{matrix} n \\ k \end{matrix} \right\|$, and generalized it in subsection (1.3). We attach a Wolfram Mathematica codes of most important equations in Application 1. Extended version of Triangle (1.19) is attached in Application 2. Future research should be done in $\left\| \begin{matrix} n^m \\ k^j \end{matrix} \right\|$, $m, j \in \mathbb{N}$ to verify its properties. A research on finding difference and consequently derivative using (1.31) and extend it over real functions applying Taylor's theorem also could be done.

REFERENCES

- [1] Conway, J. H. and Guy, R. K. "Pascal's Triangle." In The Book of Numbers. New York: Springer-Verlag, pp. 68-70, 1996.
- [2] Abramowitz, M. and Stegun, I. A. (Eds.). Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing. New York: Dover, pp. 10, 1972.
- [3] Weisstein, Eric W. "Power", From Mathworld.
- [4] Arfken, G. Mathematical Methods for Physicists, 3rd ed. Orlando, FL: Academic Press, pp. 307-308, 1985.
- [5] Rudin, Walter (1987). Real and complex analysis (3rd ed.). New York: McGraw-Hill. p. 1. ISBN 978-0-07-054234-1.
- [6] Weisstein, Eric W. "Finite Difference." From Mathworld.
- [7] Richardson, C. H. An Introduction to the Calculus of Finite Differences. p. 5, 1954.
- [8] N. J. A. Sloane et al., Entry A008458 in [9], 2002-present.
- [9] The OEIS Foundation Inc.,The On-Line Encyclopedia of Integer Sequences, 1964-present <https://oeis.org/>
- [10] Bakhvalov N. S. Numerical Methods: Analysis, Algebra, Ordinary Differential Equations p. 59, 1977. (In russian)
- [11] N. J. A. Sloane., "Triangle of Eulerian numbers", Entry A008292 in [9], 1996-present.
- [12] Kolosov Petro, et al., Entry A287326 in [9], 2017.
- [13] N. J. A. Sloane and Mira Bernstein et al., Entry A007318 in [9], 1994-present.
- [14] N. J. A. Sloane. "Central polygonal numbers" Entry A000124 in [9], 1994-present.
- [15] Early, Nick. Combinatorics and Representation Theory for Generalized Permutohedra I: Simplicial Plates, p. 3, 2016. Available online at arXiv:1611.06640.
- [16] T. K. Petersen. "Eulerian numbers." Springer New York, 2015. 3-18.
- [17] Joshua Giambalvo. Entry A275709 in [9], 2016.
- [18] Joe Keane. Entry A028896 in [9], 2016.

4. APPLICATION 1. WOLFRAM MATHEMATICA 11 CODES OF SOME EXPRESSIONS

In this section Wolfram Mathematica codes of most expressions are shown. Note that Mathematica `.cdf`-file of all mentioned expressions is available for download at this link. The `.txt`-file reader could find here.

Define coefficient $\left\| \begin{matrix} n \\ k \end{matrix} \right\|$, definition (1.20)

```
In[1]:= K[n_, k_] := 3!*n*k - 3!*k^2 + 1
```

Check of property (1.27)

In[1]:= $\mathbf{K}[(n^2 + n + 2)/2, 1]$

Check of expression (1.31), x^n

In[1]:= $1/2*\mathbf{Sum}[(\mathbf{K}[2*x - k, k] + \mathbf{K}[2*x - k, 0])*x^{(n - 3)}, \{k, 0, x - 1\}]$
 $1/2*\mathbf{Sum}[(\mathbf{K}[x + 1, k] + \mathbf{K}[x - 1, k])*x^{(n - 3)}, \{k, 0, x - 1\}]$
 $1/2*\mathbf{Sum}[(\mathbf{K}[(k^2 + k)/2, 1] + \mathbf{K}[(k^2 + k + 4)/2, 1])*x^{(n - 3)}, \{k, 0, x - 1\}]$
 $\mathbf{Sum}[\mathbf{K}[x, k]*x^{(n-3)}, \{k, 0, x - 1\}]$
 $\mathbf{Sum}[\mathbf{K}[(k^2 + k + 2)/2, 1]*x^{(n - 3)}, \{k, 0, x - 1\}]$

Generating formula of Triangle (1.17), Figure 3

In[1]:= $\mathbf{Column}[\mathbf{Table}[\mathbf{K}[n, k], \{n, 0, 5\}, \{k, 0, n\}], \mathbf{Center}]$

Expression (1.33), $x^n - 1$

In[1]:= $\mathbf{Sum}[\mathbf{K}[x, m]*x^{(n - 3)} + \mathbf{Sum}[x^{(n - t)}, \{t, 4, n\}], \{m, 1, x - 1\}]$

Expression (1.34), $x^n - 1$ using properties (1.27), (1.29), (1.30)

In[1]:= $\mathbf{Sum}[1/2*(\mathbf{K}[2*x - k, k] + \mathbf{K}[2*x - k, 0])*x^{(n - 3)} + \mathbf{Sum}[x^{(n - t)}, \{t, 4, n\}], \{k, 1, x - 1\}]$
 $\mathbf{Sum}[1/2*(\mathbf{K}[x + 1, k] + \mathbf{K}[x - 1, k])*x^{(n - 3)} + \mathbf{Sum}[x^{(n - t)}, \{t, 4, n\}], \{k, 1, x - 1\}]$
 $\mathbf{Sum}[1/2*(\mathbf{K}[(k^2 + k)/2, 1] + \mathbf{K}[(k^2 + k + 4)/2, 1])*x^{(n - 3)} + \mathbf{Sum}[x^{(n - t)}, \{t, 4, n\}], \{k, 1, x - 1\}]$
 $\mathbf{Sum}[\mathbf{K}[(k^2 + k + 2)/2, 1]*x^{(n - 3)} + \mathbf{Sum}[x^{(n - t)}, \{t, 4, n\}], \{k, 1, x - 1\}]$

Expression (1.35), x^n

$$\mathbf{In}[1]:= \mathbf{Sum}[\mathbf{K}[x, m]*x^{(n-3)} + \mathbf{Sum}[x^{(n-t)}, \{t, 4, n\}] - \mathbf{Sum}[x^j, \{j, 0, \mathbf{Infinity}\}], \{m, 1, x-1\}]$$

Expression (1.36), generalized version of (1.35), x^n

$$\mathbf{In}[1]:= \mathbf{Sum}[(\mathbf{K}[x, m] - 1)*x^{(n-3)} - \mathbf{Sum}[x^{(n-3+j)}, \{j, 1, \mathbf{Infinity}\}], \{m, 1, x-1\}]$$

$$\mathbf{Sum}[(\mathbf{K}[(m^2 + m + 2)/2, 1] - 1)*x^{(n-3)} - \mathbf{Sum}[x^{(n-3+j)}, \{j, 1, \mathbf{Infinity}\}], \{m, 1, x-1\}]$$

Expression (1.21), Multinomial theorem for $n = 4$

$$\mathbf{In}[1]:= \mathbf{Sum}[\mathbf{K}[u + j + m, k]*u + \mathbf{K}[u + j + m, k]*j + \mathbf{K}[u + j + m, k]*m, \{k, 0, u + j + m - 1\}]$$

Section 2, Expression (2.1), e^x representation

$$\mathbf{In}[1]:= \mathbf{Sum}[1/n!*\mathbf{Sum}[(\mathbf{K}[x, m] - 1)*x^{(n-3)} - \mathbf{Sum}[x^{(n-3+j)}, \{j, 1, \mathbf{Infinity}\}], \{m, 1, x-1\}], \{n, 0, \mathbf{Infinity}\}]$$

$$\mathbf{Sum}[1/n!*\mathbf{Sum}[(\mathbf{K}[(m^2 + m + 2)/2, 1] - 1)*x^{(n-3)} - \mathbf{Sum}[x^{(n-3+j)}, \{j, 1, \mathbf{Infinity}\}], \{m, 1, x-1\}], \{n, 0, \mathbf{Infinity}\}]$$

$$\mathbf{Sum}[1/n!*\mathbf{Sum}[(1/2*(\mathbf{K}[x + 1, m] + \mathbf{K}[x - 1, m]) - 1)*x^{(n-3)} - \mathbf{Sum}[x^{(n-3+j)}, \{j, 1, \mathbf{Infinity}\}], \{m, 1, x-1\}], \{n, 0, \mathbf{Infinity}\}]$$

$$\mathbf{Sum}[1/n!*\mathbf{Sum}[(1/2*(\mathbf{K}[2x-m, m] + \mathbf{K}[2x-m, 0]) - 1)*x^{(n-3)} - \mathbf{Sum}[x^{(n-3+j)}, \{j, 1, \mathbf{Infinity}\}], \{m, 1, x-1\}], \{n, 0, \mathbf{Infinity}\}]$$

Section 2, Expression (2.2), $e^x - e$ representation

$$\mathbf{In}[1]:= \mathbf{Sum}[1/n!*\mathbf{Sum}[\mathbf{K}[x, m]*x^{(n-3)} + \mathbf{Sum}[x^{(n-t)}, \{t, 4, n\}], \{m, 1, x-1\}], \{n, 0, \mathbf{Infinity}\}]$$

$$\mathbf{Sum}[1/n!*\mathbf{Sum}[1/2*(\mathbf{K}[x + 1, m] + \mathbf{K}[x - 1, m])*x^{(n-3)} + \mathbf{Sum}[x^{(n-t)}, \{t, 4, n\}], \{m, 1, x-1\}], \{n, 0, \mathbf{Infinity}\}]$$

$$\text{Sum}[1/n! * \text{Sum}[\mathbf{K}[(m^2 + m + 2)/2, 1] * x^{(n-3)} + \text{Sum}[x^{(n-t)}, \{t, 4, n\}], \{m, 1, x-1\}], \{n, 0, \mathbf{Infinity}\}]$$

$$\text{Sum}[1/(n!) * \text{Sum}[1/2 * (\mathbf{K}[(m^2+m+1)/2, 1] + \mathbf{K}[(m^2+m+3)/2, 1]) * x^{(n-3)} + \text{Sum}[x^{(n-t)}, \{t, 4, n\}], \{m, 1, x-1\}], \{n, 0, \mathbf{Infinity}\}]$$

Binomial Theorem based on definition (1.20), subsection (1.1)

$$\text{In}[1]:= \text{Sum}[\text{Sum}[\mathbf{K}[x+y, k] * \text{Binomial}[n-3, j] * x^{(n-3-j)} * y^j, \{j, 0, n-3\}], \{k, 0, x+y-1\}]$$

$$\text{Sum}[1/2 * \text{Sum}[(\mathbf{K}[x+y+1, k] + \mathbf{K}[x+y-1, k]) * \text{Binomial}[n-3, j] * x^{(n-3-j)} * y^j, \{j, 0, n-3\}], \{k, 0, x+y-1\}]$$

$$\text{Sum}[1/2 * \text{Sum}[(\mathbf{K}[2(x+y) - k, k] + \mathbf{K}[2(x+y) - k, 0]) * \text{Binomial}[n-3, j] * x^{(n-3-j)} * y^j, \{j, 0, n-3\}], \{k, 0, x+y-1\}]$$

$$\text{Sum}[\text{Sum}[\mathbf{K}[(k^2+k+2)/2, 1] * \text{Binomial}[n-3, j] * x^{(n-3-j)} * y^j, \{j, 0, n-3\}], \{k, 0, x+y-1\}]$$

5. APPLICATION 2. AN EXTENDED VERSION OF TRIANGLE (3)




Figure 5. Extended version of Triangle (1.19) generated from (1.15) given $n = 3$ over x from 0 to 11, sequence A287326 in OEIS, [9], [12].