

Atomic Nucleus Grimace

To some degree of approximation, atomic nuclei are spherical, though distorted to a greater or lesser extent. When the nucleus is excited, its shape may change, but only for an extremely brief moment, after which it returns to its original state. [13]

What is the mass of a proton? Scientists from Germany and Japan have made an important step toward better understanding this fundamental constant. [12]

In a paper published today in the journal Science, the ASACUSA experiment at CERN reported new precision measurement of the mass of the antiproton relative to that of the electron. [11]

When two protons approaching each other pass close enough together, they can “feel” each other, similar to the way that two magnets can be drawn closely together without necessarily sticking together. According to the Standard Model, at this grazing distance, the protons can produce a pair of W bosons. [10]

The fact that the neutron is slightly more massive than the proton is the reason why atomic nuclei have exactly those properties that make our world and ultimately our existence possible. Eighty years after the discovery of the neutron, a team of physicists from France, Germany, and Hungary headed by Zoltán Fodor, a researcher from Wuppertal, has finally calculated the tiny neutron-proton mass difference. [9]

Taking into account the Planck Distribution Law of the electromagnetic oscillators, we can explain the electron/proton mass ratio and the Weak and Strong Interactions. Lattice QCD gives the same results as the diffraction patterns of the electromagnetic oscillators, explaining the color confinement and the asymptotic freedom of the Strong Interactions.

Contents

Preface.....	2
The first light atomic nucleus with a second face.....	3
Most precise measurement of proton mass.....	4
CERN experiment improves precision of antiproton mass measurement with new innovative cooling technique	5
Exclusive production: shedding light with grazing protons.....	7

Theory of the strong interaction verified	8
Asymmetry in the interference occurrences of oscillators	9
Spontaneously broken symmetry in the Planck distribution law	10
The structure of the proton	12
The weak interaction	12
The Strong Interaction - QCD	13
Confinement and Asymptotic Freedom	13
Lattice QCD.....	14
QCD	14
Color Confinement	14
Electromagnetic inertia and mass.....	15
Electromagnetic Induction	15
The frequency dependence of mass	15
Electron – Proton mass rate	15
The potential of the diffraction pattern	15
Conclusions	16
References	17

Author: George Rajna

Preface

The fact that the neutron is slightly more massive than the proton is the reason why atomic nuclei have exactly those properties that make our world and ultimately our existence possible. Eighty years after the discovery of the neutron, a team of physicists from France, Germany, and Hungary headed by Zoltán Fodor, a researcher from Wuppertal, has finally calculated the tiny neutron-proton mass difference. The findings, which have been published in the current edition of Science, are considered a milestone by many physicists and confirm the theory of the strong interaction. As one of the most powerful computers in the world, JUQUEEN at Forschungszentrum Jülich was decisive for the simulation. [10]

The diffraction patterns of the electromagnetic oscillators give the explanation of the Electroweak and Electro-Strong interactions. [2] Lattice QCD gives the same results as the diffraction patterns which explain the color confinement and the asymptotic freedom.

The hadronization is the diffraction pattern of the baryons giving the jet of the color – neutral particles!

The first light atomic nucleus with a second face

To some degree of approximation, atomic nuclei are spherical, though distorted to a greater or lesser extent. When the nucleus is excited, its shape may change, but only for an extremely brief moment, after which it returns to its original state. A relatively permanent 'second face' of atomic nuclei has so far only been observed in the most massive elements. Recently, physicists from Poland, Italy, Japan, Belgium and Romania have for the first time registered this phenomenon in a light nucleus.

Atomic nuclei can change their shape depending on the amount of energy they possess or the speed at which they spin. Changes related only to the addition of energy (and not accounting for spin) are relatively stable only in nuclei of the most massive elements. Now, it turns out that the nuclei of much lighter elements such as nickel can also persist a little longer in their new shape.

The calculations necessary for the preparation of the experiment proved to be so complex that a computer infrastructure of about one million processors was required to perform them. The effort has been reported in the journal *Physical Review Letters*.

Constructed of protons and neutrons, atomic nuclei are generally considered to be spherical structures, but can be flattened or elongated along one, two, or sometimes three axes. What's more, atomic nuclei can change their deformation depending on the amount of energy they possess, even when they are not spinning.

"When an atomic nucleus is supplied with the right amount of energy, it can transition into a state with a different shape deformation than is typical for the basic state. However, this new deformation, illustratively speaking, is very unstable. Just like a ball returns to its original shape after squeezing it, so the nucleus returns to its original form, but it does so much, much faster—in billionths of a billionth of a second or an even shorter time. So instead of talking about the second face of the atomic nucleus, it's probably better to talk about just a grimace," explains Prof. Bogdan Fornal.

In the last few decades, evidence has accumulated confirming that relatively stable nuclei with a deformed shape can be present in a small number of elements. Measurements have shown that the nuclei of some actinides—elements with atomic numbers from 89 (actinium) to 103 (lawrencium)—are capable of maintaining their 'second face' even tens of millions of times longer than other nuclei. Actinides are quite massive, with protons and neutrons totaling well above 200. Up to now, among the non-spinning nuclei of lighter elements, an excited state with a deformed shape characterized by high stability has never been observed.

"We pointed out that two theoretical models of nuclear excitation predict the existence of relatively stable states with deformed shapes in the nuclei of light elements. Later, a third model appeared that also led to similar conclusions. Our attention was drawn to nickel-66, because it was present in the predictions of all three models," recalls Prof. Fornal.

The new experimental method proposed by Prof. Silvia Leoni (UniMi), combined with the computationally sophisticated Monte Carlo shell model developed by the Tokyo University theorists, enabled the design of appropriate, accurate measurements. The experiment was carried out at the 9

MV FN Pelletron Tandem accelerator operating in the Romanian National Institute of Physics and Nuclear Engineering (IFIN-HH).

In the experiment in Bucharest, a target of nickel-64 was fired with nuclei of oxygen-18. Relative to oxygen-16, which is the main (99.76%) isotope of atmospheric oxygen, these nuclei contain two additional neutrons. During the collisions, both the excess neutrons can be transferred to the nickel nuclei, resulting in the creation of nickel-66, the basic shape of which is almost an ideal sphere. With properly selected collision energies, a small portion of the Ni-66 nuclei thus formed achieve a certain state with a deformed shape which, as measurements showed, proved to be slightly more stable than all other excited states associated with significant deformation. In other words, the nucleus was in a local, deep minimum of potential.

"The extension of lifespan of the deformed shape of the Ni-66 nucleus is not as spectacular as that of the actinides. We recorded only five-fold growth. Nevertheless, the measurement was exceptional, because it was the first observation of its kind in light nuclei," says Prof. Fornal, who stresses that the measured delay times of return to the basic state correspond to an acceptable extent with the values provided by the new theoretical model. None of the earlier models of nuclear structure allowed for such detailed predictions. This suggests that the new theoretical approach should be helpful in describing several thousand nuclei that have not yet been discovered. [13]

Most precise measurement of proton mass

What is the mass of a proton? Scientists from Germany and Japan have made an important step toward better understanding this fundamental constant. By means of precision measurements on a single proton, they were able to improve the precision by a factor of three and also correct the existing value.

To determine the mass of a single proton more accurately, the group of physicists from the Max Planck Institute for Nuclear Physics in Heidelberg and RIKEN in Japan performed an important high-precision measurement in a greatly advanced Penning trap system, designed by Sven Sturm and Klaus Blaum from MPI-K, using ultra-sensitive single particle detectors that were partly developed by RIKEN's Ulmer Fundamental Symmetries Laboratory.

The proton is the nucleus of the hydrogen atom and one of the basic building blocks of all other atomic nuclei. Therefore, the proton's mass is an important parameter in atomic physics: it is one of the factors that affect how the electrons move around the atomic nucleus. This is reflected in the spectra, i.e., the light colours (wavelengths) that atoms can absorb and emit again. By comparing these wavelengths with theoretical predictions, it is possible to test fundamental physical theories. Further, precise comparisons of the masses of the proton and the antiproton may help in the search for the crucial difference – besides the reversed sign of the charge – between matter and antimatter.

Penning traps are well-proven as suitable "scales" for ions. In such a trap, it is possible to confine, nearly indefinitely, single charged particles such as a proton, for example, by means of electric and magnetic fields. Inside the trap, the trapped particle performs a characteristic periodic motion at a certain oscillation frequency. This frequency can be measured and the mass of the particle calculated from it. In order to reach the targeted high precision, an elaborate measurement technique was required.

The carbon isotope ^{12}C with a mass of 12 atomic mass units is defined as the mass standard for atoms. "We directly used it for comparison," says Sven Sturm. "First we stored each one proton and one carbon ion ($^{12}\text{C}^{6+}$) in separate compartments of our Penning trap apparatus, then transported each of the two ions into the central measurement compartment and measured its motion." From the ratio of the two measured values the group obtained the proton's mass directly in atomic units. The measurement compartment was equipped with specifically developed purpose-built electronics. Andreas Mooser of RIKEN's Fundamental Symmetries Laboratory explains its function: "It allowed us to measure the proton under identical conditions as the carbon ion despite its about 12-fold lower mass and 6-fold smaller charge."

The resulting mass of the proton, determined to be $1.007276466583(15)(29)$ atomic mass units, is three times more precise than the presently accepted value. The numbers in parentheses refer to the statistical and systematic uncertainties, respectively.

Intriguingly, the new value is significantly smaller than the current standard value. Measurements by other authors yielded discrepancies with respect to the mass of the tritium atom, the heaviest hydrogen isotope ($T = 3\text{H}$), and the mass of light helium (^3He) compared to the "semiheavy" hydrogen molecule HD ($D = 2\text{H}$, deuterium, heavy hydrogen). "Our result contributes to solving this puzzle, since it corrects the proton's mass in the proper direction," says Klaus Blaum.

Florian Köhler-Langes of MPIK explains how the researchers intend to further improve the precision of their measurement: "In the future, we will store a third ion in our trap tower. By simultaneously measuring the motion of this reference ion, we will be able to eliminate the uncertainty originating from fluctuations of the magnetic field." The work was published in *Physical Review Letters*. [12]

CERN experiment improves precision of antiproton mass measurement with new innovative cooling technique

In a paper published today in the journal *Science*, the ASACUSA experiment at CERN reported new precision measurement of the mass of the antiproton relative to that of the electron. This result is based on spectroscopic measurements with about 2 billion antiprotonic helium atoms cooled to extremely cold temperatures of 1.5 to 1.7 degrees above absolute zero. In antiprotonic helium atoms an antiproton takes the place of one of the electrons that would normally be orbiting the nucleus.

Such measurements provide a unique tool for comparing with high precision the mass of an antimatter particle with its matter counterpart. The two should be strictly identical.

"A pretty large number of atoms containing antiprotons were cooled below minus 271 degrees Celsius. It's kind of surprising that a 'half-antimatter' atom can be made so cold by simply placing it in a refrigerated gas of normal helium," said Masaki Hori, group leader at the ASACUSA collaboration.

Matter and antimatter particles are always produced as a pair in particle collisions. Particles and antiparticles have the same mass and opposite electric charge. The positively charged positron, for example, is an anti-electron, the antiparticle of the negatively charged electron. Positrons have been observed since the 1930s, both in natural collisions from cosmic rays and in particle accelerators. They are used today in hospital in PET scanners. However, studying antimatter particles with high-

precision remains a challenge because when matter and antimatter come into contact, they annihilate – disappearing in a flash of energy.

CERN's Antiproton Decelerator is a unique facility delivering low-energy antiproton beams to experiments for antimatter studies. In order to make measurements with these antiprotons, several experiments trap them for long periods using magnetic devices. ASACUSA's approach is different as the experiment is able to create very special hybrid atoms made of a mix of matter and antimatter: these are the antiprotonic helium atoms composed of an antiproton and an electron orbiting a helium nucleus. They are made by mixing antiprotons with helium gas. In this mixture, about 3% of the antiprotons replace one of the two electrons of the helium atom. In antiprotonic helium, the antiproton is in orbit around the helium nucleus, and protected by the electron cloud that surrounds the whole atom, making antiprotonic helium stable enough for precision measurements.

Latest precision measurement of the mass of the proton and the anti proton through the production of antiprotonic helium by the ASACUSA experiment at CERN's antimatter factory, with a beam from the Antiproton Decelerator 00:03:41.480 / 02 November 2016. Credit: CERN (License: Julien Ordan)

The measurement of the antiproton's mass is done by spectroscopy, by shining a laser beam onto the antiprotonic helium. Tuning the laser to the right frequency causes the antiprotons to make a quantum jump within the atoms. From this frequency the antiproton mass relative to the electron mass can be calculated. This method has been successfully used before by the ASACUSA collaboration to measure with high accuracy the antiproton's mass. However, the microscopic motion of the antiprotonic helium atoms introduced a significant source of uncertainty in previous measurements.

The major new achievement of the collaboration, as reported in *Science*, is that ASACUSA has now managed to cool down the antiprotonic helium atoms to temperatures close to absolute zero by suspending them in a very cold helium buffer-gas. In this way, the microscopic motion of the atoms is reduced, enhancing the precision of the frequency measurement. The measurement of the transition frequency has been improved by a factor of 1.4 to 10 compared with previous experiments. Experiments were conducted from 2010 to 2014, with about 2 billion atoms, corresponding to roughly 17 femtograms of antiprotonic helium.

According to standard theories, protons and antiprotons are expected to have exactly the same mass. To date, no difference has been found between their masses, but pushing the precision limits of this comparison is a very important test of key theoretical principles such as the CPT symmetry. CPT is a consequence of basic symmetries of space-time, such as its isotropy in all directions. The observation of even a minute breaking of CPT would call for a review of our assumptions about the nature and properties of space-time.

The ASACUSA collaboration is confident that it will be able to further improve the precision of antiproton's mass by using two laser beams. In the near future, the start of the ELENA facility at CERN will also allow the precision of such measurements to be improved. [11]

Exclusive production: shedding light with grazing protons

When two protons approaching each other pass close enough together, they can “feel” each other, similar to the way that two magnets can be drawn closely together without necessarily sticking together. According to the Standard Model, at this grazing distance, the protons can produce a pair of W bosons.

When two protons approaching each other pass close enough together, they can “feel” each other, similar to the way that two magnets can be drawn closely together without necessarily sticking together. According to the Standard Model, at this grazing distance, the protons can produce a pair of W bosons.

As its name implies, the primary mission of the Large Hadron Collider is to generate collisions of protons for study by physicists at experiments such as CMS. It may surprise you to find out that the vast majority of protons accelerated by the LHC never collide with one another. Some of these fly-by protons, however, still interact with each other in such a way as to help physicists shed light on the nature of the universe.

The LHC accelerates bunches of protons, with more than 10 billion protons in each bunch, in opposite directions around the ring. As those protons arrive at a detector, such as CMS, magnets focus the beams to increase the density of protons and thus increase the chance of a coveted collision. Despite what seems like overwhelming odds, only a few of these protons actually collide with each other: tens to hundreds per each beam “crossing.” An even smaller fraction of the remaining protons pass close enough to other protons to “feel” each other, even if they do not directly collide.

Think of two toy magnets on a tabletop: A north end and a south end moved close enough to each other will rather firmly stick to each other. However, you can also move one magnet just close enough to the other that you can make it wiggle without drawing it all the way over. This exchange of energy is mediated by the exchange of photons, the carrier particle of the electromagnetic force. Similarly, two protons in the LHC that get just the right distance from each other will exchange photons without colliding.

Now for the part that gets really interesting to particle physicists. The photons generated by these near-miss proton interactions can be billions of times more energetic than those of visible light, and as a result they carry enough energy to create particles in their own right. The Standard Model predicts the production of massive particles, such as pairs of W bosons, from these interacting photons without any of the additional activity that is seen in the messier proton-proton collision events. In a detector such as CMS, this pair of W bosons is said to be produced “exclusively.” However, “exclusive production” is an apt name in another way – creating a pair of W bosons from interacting photons is a rare occurrence in an even rarer sample of photons generated from near-miss proton interactions.

CMS scientists performed such a search for such W boson pairs emanating from interacting photons. In a data set consisting of 7- and 8-TeV collisions, 15 candidate events for this process were observed. While it may not seem like much, the expected background was considerably smaller, allowing the CMS team to claim that they have evidence of the process. (In the particle physics world, evidence is a three-standard-deviation departure from background, as explained here).

Furthermore, these results helped place stringent results on a number of models which predict a greater rate of this process. [10]

Theory of the strong interaction verified

The findings, which have been published in the current edition of Science, are considered a milestone by many physicists and confirm the theory of the strong interaction. As one of the most powerful computers in the world, JUQUEEN at Forschungszentrum Jülich was decisive for the simulation.

The existence and stability of atoms relies heavily on the fact that neutrons are slightly more massive than protons. The experimentally determined masses differ by only around 0.14 percent. A slightly smaller or larger value of the mass difference would have led to a dramatically different universe, with too many neutrons, not enough hydrogen, or too few heavier elements. The tiny mass difference is the reason why free neutrons decay on average after around ten minutes, while protons - the unchanging building blocks of matter - remain stable for a practically unlimited period.

In 1972, about 40 years after the discovery of the neutron by Chadwick in 1932, Harald Fritzsch (Germany), Murray Gell-Mann (USA), and Heinrich Leutwyler (Switzerland) presented a consistent theory of particles and forces that form the neutron and the proton known as quantum chromodynamics. Today, we know that protons and neutrons are composed of "up quarks" and "down quarks". The proton is made of one down and two up quarks, while the neutron is composed of one up and two down quarks.

Simulations on supercomputers over the last few years confirmed that most of the mass of the proton and neutron results from the energy carried by their quark constituents in accordance with Einstein's formula $E=mc^2$. However, a small contribution from the electromagnetic field surrounding the electrically charged proton should make it about 0.1 percent more massive than the neutral neutron. The fact that the neutron mass is measured to be larger is evidently due to the different masses of the quarks, as Fodor and his team have now shown in extremely complex simulations.

For the calculations, the team developed a new class of simulation techniques combining the laws of quantum chromodynamics with those of quantum electrodynamics in order to precisely determine the effects of electromagnetic interactions. By controlling all error sources, the scientists successfully demonstrated how finely tuned the forces of nature are.

Professor Kurt Binder is Chairman of the Scientific Council of the John von Neumann Institute for Computing (NIC) and member of the German Gauss Centre for Supercomputing. Both organizations allocate computation time on JUQUEEN to users in a competitive process. "Only using world-class computers, such as those available to the science community at Forschungszentrum Jülich, was it possible to achieve this milestone in computer simulation," says Binder. JUQUEEN was supported in the process by its "colleagues" operated by the French science organizations CNRS and GENCI as well as by the computing centres in Garching (LRZ) and Stuttgart (HLRS). [9]

Asymmetry in the interference occurrences of oscillators

The asymmetrical configurations are stable objects of the real physical world, because they cannot annihilate. One of the most obvious asymmetry is the proton – electron mass rate $M_p = 1840 M_e$ while they have equal charge. We explain this fact by the strong interaction of the proton, but how remember it his strong interaction ability for example in the H – atom where are only electromagnetic interactions among proton and electron.

This gives us the idea to origin the mass of proton from the electromagnetic interactions by the way interference occurrences of oscillators. The uncertainty relation of Heisenberg makes sure that the particles are oscillating.

The resultant intensity due to n equally spaced oscillators, all of equal amplitude but different from one another in phase, either because they are driven differently in phase or because we are looking at them an angle such that there is a difference in time delay:

$$(1) I = I_0 \sin^2 n \phi / 2 / \sin^2 \phi / 2$$

If ϕ is infinitesimal so that $\sin\phi = \phi$, then

$$(2) I = n^2 I_0$$

This gives us the idea of

$$(3) M_p = n^2 M_e$$




Fig. 30-3. A linear array of n equal oscillators, driven with phases $\alpha_s = s\alpha$.

Figure 1.) A linear array of n equal oscillators

There is an important feature about formula (1) which is that if the angle ϕ is increased by the multiple of 2π , it makes no difference to the formula.

So

$$(4) d \sin \theta = m \lambda$$

and we get m-order beam if λ less than d. [6]

If d less than λ we get only zero-order one centered at $\theta = 0$. Of course, there is also a beam in the opposite direction. The right chooses of d and λ we can ensure the conservation of charge.

For example

$$(5) 2(m+1) = n$$

Where $2(m+1) = N_p$ number of protons and $n = N_e$ number of electrons.

In this way we can see the H_2 molecules so that $2n$ electrons of n radiate to $4(m+1)$ protons, because $d_e > \lambda_e$ for electrons, while the two protons of one H_2 molecule radiate to two electrons of them, because of $d_e < \lambda_e$ for this two protons.

To support this idea we can turn to the Planck distribution law, that is equal with the Bose – Einstein statistics.

Spontaneously broken symmetry in the Planck distribution law

The Planck distribution law is temperature dependent and it should be true locally and globally. I think that Einstein's energy-matter equivalence means some kind of existence of electromagnetic oscillations enabled by the temperature, creating the different matter formulas, atoms molecules, crystals, dark matter and energy.

Max Planck found for the black body radiation

As a function of wavelength (λ), Planck's law is written as:

$$B_{\lambda}(T) = \frac{2hc^2}{\lambda^5} \frac{1}{e^{\frac{hc}{\lambda T}} - 1}.$$




Figure 2. The distribution law for different T temperatures

We see there are two different λ_1 and λ_2 for each T and intensity, so we can find between them a d so that $\lambda_1 < d < \lambda_2$.

We have many possibilities for such asymmetrical reflections, so we have many stable oscillator configurations for any T temperature with equal exchange of intensity by radiation. All of these configurations can exist together. At the λ_{\max} is the annihilation point where the configurations are symmetrical. The λ_{\max} is changing by the Wien's displacement law in many textbooks.

$$(7) \quad \lambda_{\max} = \frac{b}{T}$$

where λ_{\max} is the peak wavelength, T is the absolute temperature of the black body, and b is a constant of proportionality called *Wien's displacement constant*, equal to $2.8977685(51) \times 10^{-3} \text{ m} \cdot \text{K}$ (2002 CODATA recommended value).

By the changing of T the asymmetrical configurations are changing too.

The structure of the proton

We must move to the higher T temperature if we want look into the nucleus or nucleon arrive to $d < 10^{-13}$ cm. [2] If an electron with $\lambda_e < d$ move across the proton then by (5) $2(m+1) = n$ with $m = 0$ we get $n = 2$ so we need two particles with negative and two particles with positive charges. If the proton can fraction to three parts, two with positive and one with negative charges, then the reflection of oscillators are right. Because this very strange reflection where one part of the proton with the electron together on the same side of the reflection, the all parts of the proton must be quasi lepton so $d > \lambda_q$. One way dividing the proton to three parts is, dividing his oscillation by the three direction of the space. We can order $1/3$ e charge to each coordinates and $2/3$ e charge to one plane oscillation, because the charge is scalar. In this way the proton has two $+2/3$ e plane oscillation and one linear oscillation with $-1/3$ e charge. The colors of quarks are coming from the three directions of coordinates and the proton is colorless. The flavors of quarks are the possible oscillations differently by energy and if they are plane or linear oscillations. We know there is no possible reflecting two oscillations to each other which are completely orthogonal, so the quarks never can be free, however there is asymptotic freedom while their energy are increasing to turn them to orthogonal. If they will be completely orthogonal then they lose this reflection and take new partners from the vacuum. Keeping the symmetry of the vacuum the new oscillations are keeping all the conservation laws, like charge, number of baryons and leptons. The all features of gluons are coming from this model. The mathematics of reflecting oscillators show Fermi statistics.

Important to mention that in the Deuteron there are 3 quarks of $+2/3$ and $-1/3$ charge, that is three u and d quarks making the complete symmetry and because this its high stability.

The weak interaction

The weak interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and time reversal symmetry.

Another important issue of the quark model is when one quark changes its flavor such that a linear oscillation transforms into plane oscillation or vice versa, changing the charge value with 1 or -1. This kind of change in the oscillation mode requires not only parity change, but also charge and time changes (CPT symmetry) resulting a right handed anti-neutrino or a left handed neutrino.

The right handed anti-neutrino and the left handed neutrino exist only because changing back the quark flavor could happen only in reverse, because they are different geometrical constructions, the u is 2 dimensional and positively charged and the d is 1 dimensional and negatively charged. It needs also a time reversal, because anti particle (anti neutrino) is involved.

The neutrino is a $1/2$ spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $1/2$ spin. The weak interaction

changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell–Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The limit of the velocity of particles as the speed of light appropriate only for electrical charged particles, since the accelerated charges are self maintaining locally the accelerating electric force. The neutrinos are CP symmetry breaking particles compensated by time in the CPT symmetry, that is the time coordinate not works as in the electromagnetic interactions, consequently the speed of neutrinos is not limited by the speed of light.

The weak interaction T-asymmetry is in conjunction with the T-asymmetry of the second law of thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes the weak interaction, for example the Hydrogen fusion.

Probably because it is a spin creating movement changing linear oscillation to 2 dimensional oscillation by changing d to u quark and creating anti neutrino going back in time relative to the proton and electron created from the neutron, it seems that the anti neutrino fastest then the velocity of the photons created also in this weak interaction?

A quark flavor changing shows that it is a reflection changes movement and the CP- and T- symmetry breaking. This flavor changing oscillation could prove that it could be also on higher level such as atoms, molecules, probably big biological significant molecules and responsible on the aging of the life.

Important to mention that the weak interaction is always contains particles and antiparticles, where the neutrinos (antineutrinos) present the opposite side. It means by Feynman's interpretation that these particles present the backward time and probably because this they seem to move faster than the speed of light in the reference frame of the other side.

Finally since the weak interaction is an electric dipole change with $\frac{1}{2}$ spin creating; it is limited by the velocity of the electromagnetic wave, so the neutrino's velocity cannot exceed the velocity of light.

The Strong Interaction - QCD

Confinement and Asymptotic Freedom

For any theory to provide a successful description of strong interactions it should simultaneously exhibit the phenomena of confinement at large distances and asymptotic freedom at short distances. Lattice calculations support the hypothesis that for non-abelian gauge theories the two domains are analytically connected, and confinement and asymptotic freedom coexist. Similarly, one way to show that QCD is the correct theory of strong interactions is that the coupling extracted at various scales (using experimental data or lattice simulations) is unique in the sense that its variation with scale is given by the renormalization group. The data for α_s is reviewed in Section 19. In this section I will discuss what these statements mean and imply. [4]

Lattice QCD

Lattice QCD is a well-established non-perturbative approach to solving the quantum chromodynamics (QCD) theory of quarks and gluons. It is a lattice gauge theory formulated on a grid or lattice of points in space and time. When the size of the lattice is taken infinitely large and its sites infinitesimally close to each other, the continuum QCD is recovered. [6]

Analytic or perturbative solutions in low-energy QCD are hard or impossible due to the highly nonlinear nature of the strong force. This formulation of QCD in discrete rather than continuous space-time naturally introduces a momentum cut-off at the order $1/a$, where a is the lattice spacing, which regularizes the theory. As a result, lattice QCD is mathematically well-defined. Most importantly, lattice QCD provides a framework for investigation of non-perturbative phenomena such as confinement and quark-gluon plasma formation, which are intractable by means of analytic field theories.

In lattice QCD, fields representing quarks are defined at lattice sites (which leads to fermion doubling), while the gluon fields are defined on the links connecting neighboring sites.

QCD

QCD enjoys two peculiar properties:

- **Confinement**, which means that the force between quarks does not diminish as they are separated. Because of this, it would take an infinite amount of energy to separate two quarks; they are forever bound into hadrons such as the proton and the neutron. Although analytically unproven, confinement is widely believed to be true because it explains the consistent failure of free quark searches, and it is easy to demonstrate in lattice QCD.
- **Asymptotic freedom**, which means that in very high-energy reactions, quarks and gluons interact very weakly. This prediction of QCD was first discovered in the early 1970s by David Politzer and by Frank Wilczek and David Gross. For this work they were awarded the 2004 Nobel Prize in Physics.

There is no known phase-transition line separating these two properties; confinement is dominant in low-energy scales but, as energy increases, asymptotic freedom becomes dominant. [5]

Color Confinement

When two quarks become separated, as happens in particle accelerator collisions, at some point it is more energetically favorable for a new quark-antiquark pair to spontaneously appear, than to allow the tube to extend further. As a result of this, when quarks are produced in particle accelerators, instead of seeing the individual quarks in detectors, scientists see "jets" of many color-neutral particles (mesons and baryons), clustered together. This process is called hadronization,

fragmentation, or string breaking, and is one of the least understood processes in particle physics. [3]

Electromagnetic inertia and mass

Electromagnetic Induction

Since the magnetic induction creates a negative electric field as a result of the changing acceleration, it works as an electromagnetic inertia, causing an electromagnetic mass. [1]

The frequency dependence of mass

Since $E = h\nu$ and $E = mc^2$, $m = h\nu / c^2$ that is the m depends only on the ν frequency. It means that the mass of the proton and electron are electromagnetic and the result of the electromagnetic induction, caused by the changing acceleration of the spinning and moving charge! It could be that the m_0 inertial mass is the result of the spin, since this is the only accelerating motion of the electric charge. Since the accelerating motion has different frequency for the electron in the atom and the proton, they masses are different, also as the wavelengths on both sides of the diffraction pattern, giving equal intensity of radiation.

Electron – Proton mass rate

The Planck distribution law explains the different frequencies of the proton and electron, giving equal intensity to different lambda wavelengths! Also since the particles are diffraction patterns they have some closeness to each other. [2]


There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

The potential of the diffraction pattern

The force that holds protons and neutrons together is extremely strong. It has to be strong to overcome the electric repulsion between the positively charged protons. It is also of very short range, acting only when two particles are within 1 or 2 fm of each other.

1 fm (femto meter) = 10^{-15} m = 0.00000000000001 meters.

The qualitative features of the nucleon-nucleon force are shown below.



There is an extremely **strong short-range repulsion** that pushes protons and neutrons apart before they can get close enough to touch. (This is shown in orange.) This repulsion can be understood to arise because the quarks in individual nucleons are forbidden to be in the same area by the Pauli Exclusion Principle.

There is a **medium-range attraction** (pulling the neutrons and protons together) that is strongest for separations of about 1 fm. (This is shown in gray.) This attraction can be understood to arise from the exchange of quarks between the nucleons, something that looks a lot like the exchange of a pion when the separation is large.

The density of nuclei is limited by the short range repulsion. The maximum size of nuclei is limited by the fact that the attractive force dies away extremely quickly (exponentially) when nucleons are more than a few fm apart.

Elements beyond uranium (which has 92 protons), particularly the trans-fermium elements (with more than 100 protons), tend to be unstable to fission or alpha decay because the Coulomb repulsion between protons falls off much more slowly than the nuclear attraction. This means that each proton sees repulsion from every other proton but only feels an attractive force from the few neutrons and protons that are nearby -- even if there is a large excess of neutrons.

Some "super heavy nuclei" (new elements with about 114 protons) might turn out to be stable as a result of the same kind of quantum mechanical shell-closure that makes noble gases very stable chemically. [7]

Conclusions

The results of this work by Fodor's team of physicists from Bergische Universität Wuppertal, Centre de Physique Théorique de Marseille, Eötvös University Budapest, and Forschungszentrum Jülich open the door to a new generation of simulations that will be used to determine the properties of quarks, gluons, and nuclear particles. According to Professor Kálmán Szabó from Forschungszentrum Jülich, "In future, we will be able to test the standard model of elementary particle physics with a tenfold increase in precision, which could possibly enable us to identify effects that would help us to uncover new physics beyond the standard model." [9]

Lattice QCD gives the same results as the diffraction theory of the electromagnetic oscillators, which is the explanation of the strong force and the quark confinement. [8]

References

[1] The Magnetic field of the Electric current and the Magnetic induction

[http://academia.edu/3833335/The Magnetic field of the Electric current](http://academia.edu/3833335/The_Magnetic_field_of_the_Electric_current)

[2] 3 Dimensional String Theory

[http://academia.edu/3834454/3 Dimensional String Theory](http://academia.edu/3834454/3_Dimensional_String_Theory)

[3] Color confinement - Wikipedia, the free encyclopedia

http://en.wikipedia.org/wiki/Color_confinement

[4] *INTRODUCTION TO LATTICE QCD*

<http://arxiv.org/abs/hep-lat/9807028>

[5] QCD http://en.wikipedia.org/wiki/Quantum_chromodynamics

[6] http://en.wikipedia.org/wiki/Lattice_QCD

[7]

<http://www.cartage.org.lb/en/themes/sciences/physics/NuclearPhysics/WhatisNuclear/Forces/Forces.htm>

[8] Theory of Everything

http://www.academia.edu/4168202/Theory_of_Everything_-_4_Dimensional_String_Theory

[9] Theory of the strong interaction verified

<http://phys.org/news/2015-03-theory-strong-interaction.html>

[10] Exclusive production: shedding light with grazing protons

<http://news.fnal.gov/2016/06/exclusive-production-shedding-light-grazing-protons/>

[11] CERN experiment improves precision of antiproton mass measurement with new innovative cooling technique

<http://phys.org/news/2016-11-cern-precision-antiproton-mass-cooling.html>

[12] Most precise measurement of proton mass

<https://phys.org/news/2017-07-precise-proton-mass.html>

[13] The first light atomic nucleus with a second face

<https://phys.org/news/2017-07-atomic-nucleus.html>