

# Towards Soliton Computer Based on Solitary Wave Solution of Maxwell-Dirac equation: A Plausible Alternative to Manakov System

Victor Christianto\*<sup>1</sup>, Florentin Smarandache<sup>2</sup>

<sup>1</sup>Malang Institute of Agriculture (IPM), Malang, Indonesia. Founder of [www.ketindo.com](http://www.ketindo.com)

\*Email: [victorchristianto@gmail.com](mailto:victorchristianto@gmail.com). URL: [http://researchgate.net/profile/Victor\\_Christianto](http://researchgate.net/profile/Victor_Christianto)

<sup>2</sup>Dept. Mathematics and Sciences, University of New Mexico, Gallup – USA. Email: [florentin.smarandache@laposte.net](mailto:florentin.smarandache@laposte.net)

## ABSTRACT

In recent years, there are a number of proposals to consider collision-based soliton computer based on certain chemical reactions, namely *Belousov-Zhabotinsky* reaction, which leads to soliton solutions of coupled Nonlinear Schroedinger equations. They are called Manakov System. But it seems to us that such a soliton computer model can also be based on solitary wave solution of Maxwell-Dirac equation, which reduces to Choquard equation. And soliton solution of Choquard equation has been investigated by many researchers, therefore it seems more profound from physics perspective. However, we consider both schemes of soliton computer are equally possible. More researches are needed to verify our proposition.

## 1. Introduction: early development of soliton science

There is little doubt that soliton technology has recently achieved close to pop science status as its proponents seek to put it to use for transporting vast amounts of

information—the stuff that gives our era its name—farther and faster. It could well

become one of the fundamental technologies in the current communications revolution.

It was during the Industrial Revolution, however, that this phenomenon was first noticed and studied.[1]

A Scottish engineer by the name of John Scott Russell had set out to create a more

efficient hull design for canal boats (a 19th-century forerunner, perhaps, of current efforts

to speed packets of information along fiber-optic cables). One day in August of 1834, he stood beside Union Canal near Edinburgh to observe the movement of a boat being pulled by two horses. As the rope pulling the boat snapped and the boat's movement halted, its prow dropped back down and Scott Russell saw a large mass of water, a smooth, solitary wave, gather around the prow and continue rapidly down the channel. Surprised and intrigued, he followed on horseback and noticed that the wave held its shape and only very gradually diminished in height. He lost sight of it after a mile or two but was so taken with this observation that he built a 30-foot wave tank in his back yard to study the phenomenon further. Ten years later, he reported his observations to the British Association for the Advancement of Science, calling what he had observed the "wave of translation." Scott Russell considered that day back in 1834 "the happiest day of my life", but his discovery was ignored by all but one or two people, who felt compelled to prove him wrong in the scientific literature. After all, it was common knowledge that waves could not behave in this way.[1]

Vindication came from two independent camps, both of which were attempting to explain the movement of shallow water waves. Boussinesq's equation in 1872 and the Korteweg–de Vries (KdV) equation in 1895 proved that solitary waves were, indeed, theoretically possible. Many theoretical and experimental developments have been done since then.

By the early 1980s, however, fiber-optic technology had caught up somewhat, and Linn Mollenauer, Roger Stolen, and Jim Gordon were able to observe soliton propagation in the lab. As mathematical results continued to appear, this group at Bell Labs was experimenting intensively with optical solitons, looking for ways to use them in long-

distance telecommunication systems. Ironically, while their work was enthusiastically received in scientific circles, it seemed that the practical application could not be quickly realized, and they were told by the head of research, Arno Penzias, to desist.

Doggedly, they persisted until their results were so compelling that Penzias apologized and publicly praised the work. Mollenauer's group has since achieved several long-distance and speed records for optical soliton transmission.[1]

## **2. Recent development: soliton computer possibility**

It is a known fact in computer industry, that the present silicon technology cannot keep up with the Moore's law. Therefore new ways of developing unconventional computing methods are being carried out in many labs.

Since 2005, some researchers from Princeton University have reported a new concept of soliton computing. As data rates in optical communication systems continue to increase, the demand for all-optical signal processing and computing devices does as well.

Examples of such devices include the nonlinear optical loop mirror, the temporal soliton dragging gate, the spatial soliton deflection gate, and the TOAD, an asymmetric loop mirror. These devices avoid the bottleneck associated with optical-electrical conversion.[2] They describe physical state-restoring computation using collisions of optical solitons. Their work is part of a larger subject known as collision-based computing, sometimes called dynamical computation. Such constructions include ideal collisions of billiard balls, Conway's universal game of Life, and multidimensional excitable lattices. Early work on soliton computation involved soliton-like collisions in

cellular automata, which demonstrated the ability to embed computation in automata using particles. [2]

In their initial study, Rand et al, consider a Manakov system, which is essentially a system of coupled NLSE, where  $q_1(x, t)$  and  $q_2(x, t)$  are two interacting optical components,  $\mu$  is a positive parameter representing the strength of the nonlinearity, and  $x$  and  $t$  are normalized space and time, respectively. The two components can be thought of as components in two polarizations, or, as in the case of a photorefractive crystal, two uncorrelated beams.[2]

The Manakov system consists of two coupled NLSEs [18]:

$$i \frac{\partial q_1}{\partial t} + \frac{\partial^2 q_1}{\partial x^2} + 2\mu(|q_1|^2 + |q_2|^2)q_1 = 0$$
$$i \frac{\partial q_2}{\partial t} + \frac{\partial^2 q_2}{\partial x^2} + 2\mu(|q_1|^2 + |q_2|^2)q_2 = 0$$

Source: Rand et al.[2]

From a practical standpoint, successful soliton computation requires ideal interactions and error-free propagation. In this sense, it is analogous to the construction of Fredkin and Toffoli, in which ideal, elastic collisions of billiard balls were used to achieve universal and reversible computation. In reality, noise will cause variability in soliton propagation and collision, and fault tolerance based on logical state restoration would need to be implemented in order to improve system performance. In a sense, this is an analog rather than a digital computer. Rand et al. also reported more advanced work based on bistable configurations of Manakov solitons. [2]

More recent discussions of collision-based soliton computing still use that basic Manakov system. See [3][4].

In the next section we will discuss an alternative approach for soliton computer.

### 3. A plausible alternative: Soliton computer based on Choquard equation

In a recent paper, we discuss how Dirac-Maxwell equation reduces to wave equation, which can be transformed into a cellular-automaton scheme.[5]

Now, it is worthy to remark here that a recent paper shows that there is travelling wave solution of (classical) Dirac-Maxwell equation.[6]

It should be noted that numerical results showed that Dirac-Maxwell system has definitely many families of solitary waves. Here the nonnegative integer  $N$  denotes the number of nodes of the positronic component of the solution (number of zeros of the corresponding spherically symmetric solution to the Choquard equation).

It can be shown that the nonrelativistic limit of such a solitary wave solution takes the form of Choquard equation, which can be written as follows:[6]

$$\omega\phi = -\frac{1}{2m}\Delta\phi + q^2\Delta^{-1}\left(|\phi|^2\right)\phi. \quad (1)$$

Considering that the above Choquard equations is similar to the NLSE (nonlinear Schrodinger equation), and also that Manakov system is a system of coupled NLSE, then we consider it plausible to consider a Manakov-Choquard system as follows:

$$\omega\phi_1 = -\frac{1}{2m}\Delta\phi_2 + q^2\Delta^{-1}\left(|\phi_2|^2\right)\phi_1. \quad (2)$$

$$\omega\phi_2 = -\frac{1}{2m}\Delta\phi_1 + q^2\Delta^{-1}\left(|\phi_1|^2\right)\phi_2. \quad (3)$$

The solutions of (2) and (3) can be explored numerically with computer algebra system such as Mathematica.

As far as we know, such a Manakov-Choquard system has not been proposed before for studying soliton computer.

#### **4. Concluding Remarks**

Considering the everincreasing demand for better computers, some researchers have considered collision-based soliton computer. And this proposal is based on Manakov system. In this paper, we consider a new concept based on travelling wave solutions of Dirac-Maxwell equations, which we call Manakov-Choquard system.

As far as we know, such a Manakov-Choquard system has not been proposed before for studying soliton computer.

More observation and experiments are recommended to verify our propositions.

#### **Document history:**

- version 1.0: 31 august 2017, pk. 17:41

#### **References:**

[1] Bonnie and Bill Kath. Making Waves: solitons and their optical applications. From *SIAM News*, Vol. 31 No. 2

[2] Darren Rand, Ken Steiglitz and Paul R. Prucnal. Signal Standardization in Collision-based Soliton Computing. *Int. J. Unconventional Computing*, Vol.1, pp. 31-45 (2005)

[3] Darren Rand, Ken Steiglitz. Computing with solitons. July 2007

[4] M. Jakubowsky, Ken Steiglitz, Richard Squier. Computing with Classical Soliton Collisions. January 2015.

[5] Victor Christianto & Florentin Smarandache. A Cellular Automaton Molecular Model based on Wave Equation: An Alternative to Cellular Automata QM. 2017. Submitted for *Special Issue of MDPI-Mathematics journal: Beyond Quantum Physics, and Computation*. url: [www.mdpi.com](http://www.mdpi.com)

[6] Andrew Comech & David Stuart. Small amplitude solitary waves in the Dirac-Maxwell system. arXiv: 1210.7261v3 (2012)

[7] Vitaly Moroz & Jean Van Schaftingen. A guide to the Choquard equation. arXiv: 1606.02518 [math.AP]