

The distribution of primes

Ihsan Raja Muda Nasution

December 7, 2017

Abstract

In this paper, we find the axiomatic pattern of prime numbers.

MSC: 11A41

Keywords: prime numbers, distribution of primes

1 Introduction

In 1859, Riemann [Rie59] computed the distribution of primes. Our motivation is to axiomatize the structure of primes.

2 Results

These below are some patterns of number.

Let t_n denote the n th triangular number. Then

$$t_n = \binom{n+1}{2} \quad n \geq 1,$$

where $\binom{n}{k}$ is the binomial coefficients.

Let F_n be the n th Fibonacci number. Then

$$F_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2} \right)^n - \left(\frac{1-\sqrt{5}}{2} \right)^n \right],$$

where n is an integer.

Let B_n be the n th Bernoulli number. Then

$$B_n = (-1)^{n+1} n \zeta(1-n),$$

where $\zeta(1-n)$ is the Riemann zeta-function.

Postulate 2.1 (Peano Postulates). Given the number 0, the set \mathbf{N} , and the function σ . Then:

1. $0 \in \mathbf{N}$.
2. $\sigma : \mathbf{N} \rightarrow \mathbf{N}$ is a function from \mathbf{N} to \mathbf{N} .
3. $0 \notin \text{range}(\sigma)$.
4. The function σ is one-to-one.
5. If $I \subset \mathbf{N}$ such that $0 \in I$ and $\sigma(n) \in I$ whenever $n \in I$, then $I = \mathbf{N}$.

We define $1 = \sigma(0)$, $2 = \sigma(1)$, $3 = \sigma(2)$, etc. We have the following postulate.

Postulate 2.2. Given a prime number p and the function τ . Then:

1. $p \neq 0, 1$.
2. $2 \in \mathbf{Z}^+$.
3. $4 \nmid p$.
4. $(-1)^{\tau(p)} = 1$.

References

- [Rie59] B. Riemann. Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse. *Monatsber. Akad. Berlin*, pages 671–680, 1859.