

On Dark Energy and the Relativistic Bohm-Poisson Equation

Carlos Castro Perelman

October, 2017

Center for Theoretical Studies of Physical Systems, Clark Atlanta University, Atlanta,
GA. 30314. perelmanc@hotmail.com

Abstract

Recently, solutions to the *nonlinear* Bohm-Poisson (BP) equation were found with relevant cosmological applications. We were able to obtain an exact analytical expression for the observed vacuum energy density, and explain the origins of its *repulsive* gravitational nature. In this work we provide further results which include two possible extensions of the Bohm-Poisson equation to the full relativistic regime; explain how Bohm's quantum potential in four-dimensions could be re-interpreted as a gravitational potential in five-dimensions, and which explains why the presence of dark energy/dark matter in our $4D$ spacetime can only be inferred *indirectly*, but not be detected/observed directly. We conclude with some comments about the Dirac-Eddington large numbers coincidences.

1 Dark Energy and the Bohm-Poisson Equation

In physical cosmology and astronomy, dark energy is an unknown form of energy which is hypothesized to permeate all of space, tending to accelerate the expansion of the universe [1]. Assuming that the standard model of cosmology is correct, the best current measurements indicate that dark energy contributes 68.3 percent of the total energy in the present-day observable universe. The mass-energy of dark matter and ordinary (baryonic) matter contribute 26.8 and 4.9 percent respectively, and other components such as neutrinos and photons contribute a very small amount. The density of dark energy much less than the density of ordinary matter or dark matter within galaxies. However, it dominates the mass-energy of the universe because it is uniform across space [1]. Two proposed forms for dark energy are the cosmological constant, [2] representing a constant energy density filling space homogeneously, and scalar fields such as quintessence or moduli, dynamic quantities whose energy density can vary in time and space.

The nature of dark energy is more hypothetical than that of dark matter, and many things about the nature of dark energy remain matters of speculation [1]. Dark energy

is thought to be very homogeneous, not very dense and is not known to interact through any of the fundamental forces other than gravity. In the Friedmann-Lemaitre-Robertson-Walker (FLRW) metric, it can be shown that a strong constant negative pressure in all the universe causes an acceleration in universe expansion if the universe is already expanding, or a deceleration in universe contraction if the universe is already contracting. This accelerating expansion effect is sometimes labeled “gravitational repulsion”.

A major outstanding problem is that quantum field theories predict a huge cosmological constant, more than 100 orders of magnitude too large. This would need to be almost, but not exactly, cancelled by an equally large term of the opposite sign. Some supersymmetric theories require a cosmological constant that is exactly zero, which does not help because supersymmetry must be broken. Nonetheless, the cosmological constant is the most economical solution to the problem of cosmic acceleration. Thus, the current standard model of cosmology, the Lambda-CDM (cold dark matter) model, includes the cosmological constant as an essential feature [1].

The evidence for dark energy is heavily dependent on the theory of general relativity. Therefore, it is conceivable that a modification to general relativity also eliminates the need for dark energy. There are very many such theories, and research is ongoing [3], [4]. The measurement of the speed of gravity with the gravitational wave event GW170817 ruled out many modified gravity theories as alternative explanation to dark energy [1].

In quintessence models of dark energy, the observed acceleration of the scale factor is caused by the potential energy of a dynamical field, referred to as quintessence field. Quintessence differs from the cosmological constant in that it can vary in space and time. In order for it not to clump and form structure like matter, the field must be very light so that it has a large Compton wavelength. This class of theories attempts to come up with an all-encompassing theory of both dark matter and dark energy as a single phenomenon that modifies the laws of gravity at various scales. This could for example treat dark energy and dark matter as different facets of the same unknown substance, a “dark fluid” [5], or postulate that cold dark matter decays into dark energy.

The Schrödinger-Newton equation has had a long history since the 1950’s [6], [7]. It is the name given to the system coupling the Schrödinger equation to the Poisson equation. In the case of a single particle, this coupling is effected as follows: for the potential energy term in the Schrödinger equation take the gravitational potential energy determined by the Poisson equation from a matter density proportional to the probability density obtained from the wave-function. For a single particle of mass m the coupled system of equations leads to the nonlinear and nonlocal Newton-Schrödinger integro-differential equation

$$i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(\vec{r}, t) + V(\vec{r}, t) \Psi(\vec{r}, t) - \left(Gm^2 \int \frac{|\Psi(\vec{r}', t)|^2}{|\vec{r} - \vec{r}'|} d^3 r' \right) \Psi(\vec{r}, t) \quad (1)$$

Bohm’s quantum potential $V_Q = -\frac{\hbar^2}{2m} (\nabla^2 \sqrt{\rho}/\sqrt{\rho})$ has a *geometrical* description as the Weyl scalar curvature produced by an ensemble density of paths associated with one, and only one particle [8]. This geometrization process of quantum mechanics allowed to derive the Schroedinger, Klein-Gordon [8] and Dirac equations [9]. Most recently, a

related geometrization of quantum mechanics was proposed [10] that describes the time evolution of particles as geodesic lines in a curved space, whose curvature is induced by the quantum potential. This formulation allows therefore the incorporation of all quantum effects into the geometry of space-time, as it is the case for gravitation in the general relativity.

Based on these results we proposed in [11] the following nonlinear quantum-like Bohm-Poisson equation for *static* solutions $\rho = \rho(\vec{r})$

$$\nabla^2 V_Q = 4\pi G m \rho \Rightarrow -\frac{\hbar^2}{2m} \nabla^2 \left(\frac{\nabla^2 \sqrt{\rho}}{\sqrt{\rho}} \right) = 4\pi G m \rho \quad (2)$$

such that one could *replace* the nonlinear Newton-Schrödinger equation for the above non-linear quantum-like Bohm-Poisson equation (2) where the fundamental quantity is *no longer* the wave-function Ψ (complex-valued in general) but the real-valued probability density $\rho = \Psi^* \Psi$. Eq-(2) is based on Bohm's quantum potential

$$V_Q \equiv -\frac{\hbar^2}{2m} \frac{\nabla^2 \sqrt{\rho}}{\sqrt{\rho}} \quad (3)$$

If one wishes to introduce a temporal evolution to ρ via a Linblad-like equation, for instance, this would lead to an overdetermined system of differential equations for $\rho(\vec{r}, t)$. Replacing ∇^2 in eqs-(1,2) for the D'Alambertian operator $\square = \nabla_\mu \nabla^\mu, \mu = 0, 1, 2, 3$ has the caveat that in QFT $\rho(x^\mu) = \rho(\vec{r}, t)$ *no longer* has the interpretation of a probability density but is now related to the particle number current. Despite this caveat we will propose an eq-(10) below involving the D'Alambertian \square operator and a proper mass density (mass per proper four-volume).

For the time being we shall just focus on static solutions $\rho(\vec{r})$. The de Sitter space metric can be written in static coordinates in the form $g_{tt}(r) = -(1 - \frac{\Lambda}{3} r^2); g_{rr}(r) = -(g_{tt})^{-1}, \dots$, and given in terms of the cosmological constant $\Lambda = (3/R_H^2)$, where R_H is the Hubble radius. Hence, there is no inconsistency in focusing for now on static solutions $\rho_m(\vec{r})$ for the probability density.

Since almost 95 percent of the energy/mass content of the Universe is comprised of dark energy/dark matter, in a recent manuscript we envisioned the Universe's dark energy/dark matter (dark-fluid) density distribution as being proportional to a QM probability density obeying the Bohm-Poisson (BP) equation [12], in the same vain that one can view an electron orbiting the Hydrogen nucleus as an "electron probability cloud" surrounding the nucleus, permeating all of space, and whose mass density distribution is $\rho = m_e \Psi^* \Psi$, where $\Psi(\vec{r})$ are the stationary wave-function solutions to the Schrödinger equation, and m_e is the electron's mass.

The density $\rho_m = m\rho$ of dark energy/dark matter (dark-fluid) permeating all of space was postulated to be a solution to the nonlinear quantum-like Bohm-Poisson (BP) equation. It is straightforward to verify that a spherically symmetric solution to eq-(2) in a 3D spatially *flat* background ¹ is given by

¹For the time being we shall not discuss solutions in *curved* backgrounds

$$\rho_m(r) = \frac{A}{r^4}, \quad A = -\frac{\hbar^2}{2\pi G m^2} < 0 \quad (4)$$

At first glance, since $\rho_m(r) \leq 0$ one would be inclined to dismiss such solution as being unphysical. Nevertheless, we can bypass this problem by focusing instead on the *shifted* density $\tilde{\rho}_m(r) \equiv \rho_m(r) - \rho_0$ obeying the Bohm-Poisson equation

$$-\frac{\hbar^2}{2m} \nabla^2 \left(\frac{\nabla^2 \sqrt{\tilde{\rho}_m}}{\sqrt{\tilde{\rho}_m}} \right) = 4\pi G m \tilde{\rho}_m, \quad \nabla^2 f(r) \equiv r^{-2} \partial_r(r^2 \partial_r f(r)). \quad (5)$$

and whose solution for the *shifted* mass density is given by

$$\tilde{\rho}_m = A/r^4 = \rho_m(r) - \rho_0 \leq 0, \quad \Rightarrow \quad \rho_m(r) = \frac{A}{r^4} + \rho_0, \quad A = -\frac{\hbar^2}{2\pi G m^2} \quad (6)$$

One may notice that by shifting the density $\rho_m \rightarrow \rho_m - \rho_0 = \tilde{\rho}_m$, and the (radial) pressure $p \rightarrow p + p_0 = \tilde{p}$, one can eliminate the cosmological constant in the Einstein's field equations with a cosmological constant, when $\rho_0 = p_0 = (\Lambda/8\pi G)$.

For example, given a Friedman-Lemaitre-Robertson-Walker (FLRW) model with a metric $ds^2 = -(dt)^2 + a^2(t)(d\Sigma)^2$, the resulting Einstein's field equations (in units $c = 1$) lead to the Friedmann equations

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} - \frac{\Lambda}{3} = \frac{8\pi G}{3} \rho_m \quad (7a)$$

$$\frac{2\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} - \Lambda = -8\pi G p \quad (7b)$$

where $k = 1, 0, -1$ is the spatial curvature index. A simple inspection of eqs-(7a,7b) reveals that that the cosmological constant can be *reabsorbed* into a mere redefinition of ρ_m and p as follows

$$\rho_m \rightarrow \rho_m - \frac{\Lambda}{8\pi G} = \tilde{\rho}_m, \quad p \rightarrow p + \frac{\Lambda}{8\pi G} = \tilde{p} \quad (8)$$

consequently, if ρ_m, p appear in the Friedmann equations (7a,7b) with a cosmological constant Λ , then $\tilde{\rho}_m = \rho_m - \rho_0$, and $\tilde{p} = p + p_0$ (such that $\rho_0 = p_0 = \Lambda/8\pi G$), will appear in eqs-(7a, 7b) without the cosmological constant Λ . Furthermore, the cosmological equation of state associated to the dark energy permeating all of space

$$\tilde{\rho}_m = -\tilde{p} \Leftrightarrow \rho_m = -p \quad (9)$$

remains invariant.

To sum up, given an actual solution $\tilde{\rho}_m$ of the BP equation, the shifting provided in eq-(8) will allow us to focus on the *domain* of values where $\rho_m(r) \geq 0$. And, in doing so, it will permit us to show that the value of ρ_0 can be made to coincide exactly with the (extremely small) observed vacuum energy density, by simply introducing an ultraviolet

length scale l that is *very close* to the Planck scale, and infrared length scale L equal to Hubble scale R_H .

A covariant extension of the BP equation, for signature $(-, +, +, +)$, may be defined in terms of the D'Alambertian operator, and a proper mass density $\sigma(\vec{r}, t)$ of physical dimensions $(length)^{-5}$, such that $m = \int \sigma(\vec{r}, t) \sqrt{|g|} d^4x$, as follows

$$-\square \left(\frac{\square \sqrt{\sigma(\vec{r}, t)}}{\sqrt{\sigma(\vec{r}, t)}} \right) = 4\pi G m \sigma(\vec{r}, t), \quad \square \equiv \frac{1}{\sqrt{|g|}} \partial_\mu (\sqrt{|g|} g^{\mu\nu} \partial_\nu), \quad \hbar = c = 1 \quad (10)$$

Focusing for now on the static solutions (6) of the BP equation, the ultraviolet scale l is chosen at the node of $\rho_m(r)$ such that

$$\rho_m(r = l) = -\frac{\hbar^2}{2\pi G m^2} \frac{1}{l^4} + \rho_o = 0 \Rightarrow \rho_o = \frac{\hbar^2}{2\pi G m^2} \frac{1}{l^4} \quad (11)$$

The domain of physical values of r must be $r \geq l$ in order to ensure a positive-definite density $\rho_m(r) \geq 0$.

In natural units of $\hbar = c = 1$, introducing the infrared scale $L = R_H$ in the normalization condition (otherwise the mass would diverge) it yields

$$m = \int_l^{R_H} \rho(r) 4\pi r^2 dr = \int_l^{R_H} \left(\frac{A}{r^4} + \rho_0 \right) 4\pi r^2 dr = \int_l^{R_H} \left(-\frac{1}{2\pi G m^2} \frac{1}{r^4} + \rho_0 \right) 4\pi r^2 dr \quad (12)$$

Upon performing the integral in eq-(12), after plugging in the value of ρ_0 derived from eq-(11), with the provision that when $R_H \gg l$ the dominant contribution to the integral stems solely from ρ_o , one ends up with the following relationship

$$\frac{4\pi R_H^3}{3} \rho_o = \frac{4\pi R_H^3}{3} \frac{1}{2\pi G m^2 l^4} = m \Rightarrow m^3 = \frac{2}{3} \frac{R_H^3}{G l^4} \quad (13)$$

solving for m one gets

$$m = \left(\frac{2}{3 G l^4} \right)^{1/3} R_H \quad (14)$$

One can verify that when the ultraviolet scale l is chosen to be *very close* to the Planck scale, and given by

$$l^4 = \frac{4}{3} L_P^4 \Rightarrow l = \left(\frac{4}{3} \right)^{1/4} L_p = 1.0745 L_p \quad (15)$$

then upon inserting the values for m and l obtained in eqs-(14,15) into the expression for ρ_o derived in eq-(11), after setting $L_p^2 = 2G$, ² it gives in natural units of $\hbar = c = 1$

²Some authors absorb the factor of 2 inside the definition of L_p , we define the Planck scale such that the Compton wavelength coincides with the Schwarzschild radius

$$\rho_o = \frac{1}{2\pi G m^2} \frac{1}{l^4} = \frac{1}{2\pi G} \left(\frac{3}{2} \frac{G l^4}{R_H^2}\right)^{2/3} \frac{1}{l^4} = \frac{3}{8\pi G} \frac{L_p^4}{R_H^2 L_p^4} = \frac{3}{8\pi G R_H^2} \quad (16)$$

which is precisely *equal* to the observed vacuum energy density $\rho = (2\Lambda/16\pi G)$ associated with a cosmological constant $\Lambda = (3/R_H^2)$ and corresponding to a de Sitter expanding universe whose throat size is the Hubble radius R_H ($= c/H_o$, H_o is today's value of the Hubble parameter).

The physical reason behind the choice of the ultraviolet scale l in eq-(15) is based on re-interpreting ρ_o as the uniform energy (mass) density inside a black hole region of Schwarzschild radius $R = 2Gm$

$$\rho_{bh} = \frac{m}{(4\pi/3)R^3} = \frac{3}{8\pi G R^2}, \quad L_P^2 = 2G, \quad \hbar = c = 1 \quad (17)$$

In the regime $R = 2Gm \gg l$, when the dominant contribution to the integral (12) stems from the ρ_o term, we may equate the expression for ρ_o in eq-(11) to ρ_{bh} in eq-(17) giving

$$\frac{1}{2\pi G m^2 l^4} = \frac{1}{2\pi l^4} \frac{(2G)^2}{GR^2} = \frac{1}{2\pi l^4} \frac{L_p^4}{GR^2} = \frac{3}{8\pi G R^2} \Rightarrow l = \left(\frac{4}{3}\right)^{1/4} L_p, \quad \hbar = c = 1 \quad (18)$$

and leading once again to the value of $l = 1.0745 L_p$ in eq-(15). Therefore, when $R = 2Gm \gg l$, the value of l is always *very* close to the Planck scale, and *independent* of $R = 2Gm$, because the scale R has *decoupled* in eq-(18).

In this way, one can effectively view the observable universe as a “black-hole” whose Hubble radius R_H encloses a mass M_U given by $2GM_U = R_H$. From eq-(14) it follows that when $R = R_H$, the black hole density $\rho_{bh} = \rho_o = \rho_{obs}$ coincides with the observed vacuum energy density. It is well known that inside the black hole horizon region the roles of t and r are exchanged due to the switch in the signature of the g_{tt}, g_{rr} metric components. Cosmological solutions based on this $t \leftrightarrow r$ exchange were provided by the Kantowski-Sachs metric.

To sum up the results in [12] : After applying the Bohm-Poisson equation to the observable Universe as a *whole*, and by introducing an ultraviolet (very close to the Planck scale) and an infrared (Hubble) scale, one can naturally obtain a value for the vacuum energy density which coincides *exactly* with the extremely small observed vacuum energy density. It is remarkable that the Bohm-Poisson equation chooses for us a lower scale to be basically equal to the Planck scale. It was not put it in by hand, but is a direct result of the solutions to the Bohm-Poisson equation. The only assumption made was to choose the Hubble scale R_H for the infrared cutoff, and which makes physical sense since R_H is the cosmological horizon. Is it a numerical coincidence or design ? Because Bohm's formulation of QM is by construction non-local, it is this non-locality which casts light into the crucial ultraviolet/infrared entanglement of the Planck/Hubble scales which was required in order to obtain the observed values of the vacuum energy density.

Furthermore, one can also explain the origins of its *repulsive* gravitational nature. The Bohm-Poisson's (BP) equation is invariant under $\tilde{\rho}_m \rightarrow -\tilde{\rho}_m$, and $G \rightarrow -G$. Consequently $-\tilde{\rho}_m \geq 0$ is a solution to a BP equation associated to a *negative* gravitational coupling $-G < 0$ which is tantamount to *repulsive* gravity. This is perhaps the most salient feature of the results in [12].

2 Asymptotic Safety and Covariant Extensions of the BP Equation

The Renormalization Group (RG) improvement of Einstein's equations is based on the possibility that Quantum Einstein Gravity might be non-perturbatively renormalizable and asymptotically safe due to the presence of interacting ultraviolet fixed points [19]. In this program one has k (energy) dependent modifications to the Newtonian coupling $G(k)$, the cosmological constant $\Lambda(k)$ and energy-dependent spacetime metrics $g_{ij,(k)}(x)$.

In $D = 4$ there is a nontrivial interacting (non-Gaussian) ultraviolet fixed point $G_* = G(k)k^2 \neq 0$. The fixed point G_* by definition is *dimensionless* and the running gravitational coupling has the form [20], [19]

$$G(k) = G_N \frac{1}{1 + [G_N k^2 / G_*]} \quad (19)$$

The scale dependence of $\Lambda(k)$ in the de Sitter case was found to be [20]

$$\Lambda(k) = \Lambda_0 + \frac{b G(k)}{4} k^4, \quad \Lambda_0 > 0 \quad (20)$$

where b is positive numerical constant.

In $D = 4$, the dimensionless gravitational coupling has a nontrivial fixed point $G = G(k)k^2 \rightarrow G_*$ in the $k \rightarrow \infty$ limit, and the dimensionless variable $\Lambda = \Lambda(k)k^{-2}$ has also a nontrivial ultraviolet fixed point $\Lambda_* \neq 0$ [20]. The infrared limits are $\Lambda(k \rightarrow 0) = \Lambda_0 > 0$, $G(k \rightarrow 0) = G_N$. Whereas the ultraviolet limit is $\Lambda(k = \infty) = \infty$; $G(k = \infty) = 0$.

Let us choose now an actual positive-definite solution $\hat{\rho}_m \equiv -\tilde{\rho}_m = |A|/r^4 \geq 0$; $|A| = \hbar^2/2\pi G m^2$, of the BP equation associated to *repulsive* gravity $-G < 0$, as explained earlier

$$-\frac{\hbar^2}{2m} \nabla^2 \left(\frac{\nabla^2 \sqrt{-\tilde{\rho}_m}}{\sqrt{-\tilde{\rho}_m}} \right) = 4\pi (-G) m (-\tilde{\rho}_m) \quad (21)$$

The mass density solution of (21) to focus on (in $\hbar = c = 1$ units) is

$$\hat{\rho}_m(r) = -\tilde{\rho}_m(r) = \frac{1}{2\pi G m^2 r^4} \geq 0 \quad (22)$$

If one selects $m = (1/R_H)$ to coincide with the Compton mass of a particle corresponding to the Hubble scale R_H , then at the Hubble scale $r = R_H$ one has

$$\hat{\rho}_m(r = R_H) = \frac{1}{2\pi G m^2 R_H^4} = \frac{1}{2\pi G (R_H)^{-2} R_H^4} \sim (L_P R_H)^{-2} \sim 10^{-122} M_{Planck}^4 \quad (23a)$$

and which agrees with the observed vacuum energy density. It is well known (to the experts) that such extremely small value is of the same order of magnitude as $(m_{neutrino})^4$.

The problem arises when one evaluates $\hat{\rho}_m(r)$ at L_p , given $m = 1/R_H$. One gets a huge value

$$\hat{\rho}_m(r = L_p) \sim \frac{1}{G m^2 L_p^4} = (\frac{R_H}{L_p})^2 L_p^{-4} \sim 10^{122} M_p^4 \quad (23b)$$

We will see how the Asymptotic Safety scenario comes to our rescue by realizing that a Renormalization Group flow of G and m^2 solves the problem. The key idea, based on dimensional grounds, is simply to postulate that the flow of $m^2(k)$ has the *same* functional form as the flow of $\Lambda(k)$ in eq-(20)

$$m^2(k) = m_o^2 + \frac{b}{4} \frac{G(k)}{k^4}, \quad m_o^2 > 0 \quad (24)$$

The only thing remaining is to relate the scale r in eq-(17) with the energy (momentum) scale k . The authors [20] expressed k as the inverse of $d(r)$ where $d(r)$ was a proper distance derived from the Schwarzschild metric. If one opts for the simplest choice $k = 1/r$, eq-(22) can be rewritten as

$$\hat{\rho}_m(r) = \frac{1}{2\pi} \frac{1}{G(k) m^2(k) r^4} = \frac{1}{2\pi} \frac{k^4}{G(k) m^2(k)} \quad (25)$$

note that strictly speaking eq-(25) is *not* a solution to the BP equation, because if it were one must have that $G(k)m^2(k) = constant$, for all values of k , which is *not* the case. Similarly, the renormalization-group-improved black hole solutions of [20] are *not* solutions to the Einstein vacuum field equations [17]. Nevertheless, from eqs-(19,24) one learns that

$$\lim_{k \rightarrow 0} (G(k) m^2(k)) = G_N m_o^2 \quad (26a)$$

$$\lim_{k \rightarrow \infty} (G(k) m^2(k)) = \frac{b}{4} (G_*)^2 \quad (26b)$$

whereas at the Planck scale $k = M_p$

$$\lim_{k \rightarrow M_p} (G(k) m^2(k)) \sim \frac{b}{4} (G_*)^2 \quad (27)$$

Consequently, eqs-(25,27) lead to

$$\hat{\rho}_m(r = L_p) \sim \frac{1}{2\pi G(k = M_p) m^2(k = M_p) L_p^4} \sim \frac{2}{b\pi} (G_*)^{-2} L_p^{-4} \sim M_p^4 \quad (28)$$

which is the expected result for the vacuum energy density at the Planck scale.

To sum up, the renormalization group machinery (Asymptotic Safety) can be implemented such that eq-(23a) furnishes the observed vacuum energy density at the Hubble scale, while eq-(28) is the expected vacuum energy density at Planck scale. Naturally, one needs to generalize the BP equation to the fully relativistic regime as described by eq-(10). The key question is what is the “particle” represented by the mass m in the BP equation (21) ? i.e. a mass that experiences a renormalization group flow (24) similar to the flow experienced by Λ (20).

We emphasized earlier the key role that $-G < 0$ plays in all of this and which *stems* directly from the invariance of the BP equation under $\rho \rightarrow -\rho; G \rightarrow -G$. Our solutions for $\hat{\rho}_m(r) \geq 0$ correspond to $-G < 0$, thus the “particle” in question exerts a *repulsive* gravitational force which mimics “dark energy”. The RG flow behavior of G displayed in eq-(19) shows that G grows as k decreases. Meaning that G *increases* with distance, so that the magnitude of the repulsive force exemplified by $-G < 0$ becomes larger, and larger, as the universe expands. This is what is observed. Next we shall provide a different view of our findings so far.

Matter Creation from the Vacuum

The second interpretation of the solution (22) to the BP equation (21) is that involving matter creation from the vacuum, as advocated by Hoyle long ago. Imagine one pumps matter out of the vacuum in lumps/units of Planck masses. Let us assume that the Universe expands in such a way that matter is being replenished from the vacuum so that the mass at any moment is linearly proportional to the size of the Universe. As the mass of the universe grows the vacuum energy density decreases since the vacuum is being depleted. In this scenario, at the Hubble scale R_H , one has $M_U \sim R_H$.

This result is also compatible with Mach’s principle. By equating GmM_U/R_H to the rest mass m of a particle one arrives at $GM_U = R_H$, which once again is very close to the Schwarzschild radius $2GM_U$. Hence, one arrives at the scaling relation

$$\frac{M_p}{L_P} = \frac{M_U}{R_H} \quad (29)$$

which we interpreted long ago [18] as equating the proper forces (after re-introducing c) $M_U c^2/R_H = M_p c^2/L_p$ and leading to some sort of maximal/minimal acceleration duality.

Inserting the values of M_P, M_U , and $r = L_P$, into the solution (22) of the BP equation gives

$$\hat{\rho}(r = L_p; m = M_p) = \frac{1}{2\pi GM_P^2 L_p^4} \sim L_p^{-4} = M_p^4, \quad L_p^2 = 2G \quad (30a)$$

which is compatible with the large density at the Planck scale, and

$$\hat{\rho}(r = L_p; m = M_U) = \frac{1}{2\pi GM_U^2 L_p^4} \sim \frac{G}{2\pi R_H^2 L_p^4} \sim (L_p R_H)^{-2} \quad (30b)$$

which agrees with the observed vacuum energy density at the Hubble scale and obtained above in eq-(11). Before concluding we add some important remarks.

Does Dark Energy Resides in the Bulk of a 5D spacetime ?

Evaluating Bohm's quantum potential (3) for $\rho(r)$ given by eq-(4) yields

$$V_Q = - \frac{\hbar^2}{mr^2} \quad (31)$$

and which is reminiscent of an effective gravitational potential in 4 spatial dimensions (a 5D spacetime). The “quantum” force $F_Q = -\partial_r V_Q$ corresponding to V_Q in eq-(31) scales as $F_Q \sim -r^{-3}$ which has the same behavior as the gravitational force between two masses m_1, m_2 in 5D

$$F = - G_5 \frac{m_1 m_2}{r^3} \quad (32)$$

since the 5D gravitational constant G_5 has dimensions of $(length)^3$. Despite the possibility, we are not going to speculate at the moment as to whether or not the “quantum” force originating from Bohm's quantum potential is the “fifth” force. The main point is that one should consider the possibility that Bohm's quantum potential in 3 spatial dimensions (4D spacetime) mimics classical gravity in 4 spatial dimensions (5D spacetime), and for this reason one can only *indirectly* infer the gravitational effects of dark energy/dark matter in our 4D universe without directly detecting it because such dark energy/dark matter resides in 5D, which is reminiscent of the brane-world scenarios.

Finsler-Relativistic Extension of the Bohm-Poisson equation

If one wishes to introduce a temporal dependence to $\hat{\rho}_m$ we should extend the BP equation to full the relativistic regime. It is interesting that a simple exchange of $r \leftrightarrow t$ as it occurs in the Schwarzschild metric leading to the Kantowski-Sachs metric, yields $\hat{\rho}_m(t) = |A|t^{-4}$, and similar findings are obtained for the values of the vacuum energy density, simply by exchanging $L_p \leftrightarrow ct_p; R_H \leftrightarrow ct_H$ in the equations. t_p, t_H are the Planck and Hubble times, respectively. Upon doing so it leads to a Big-Bang-like singularity at $t = 0, \hat{\rho}_m(t = 0) = \infty$. This combined with the repulsive gravitational feature of our model, implies naturally that an expansion would follow.

Besides eq-(10), another relativistic generalization of the BP equation can be constructed from the Lagrange-Finsler geometrical formulation of QM recently advocated by [10]. He described the time evolution of particles as geodesic lines in a curved space, whose *curvature* is induced by the quantum potential. This formulation incorporates all quantum effects into the geometry of space-time, as it is the case for gravitation in the general relativity.

The explicit expression of the metric components in terms of the quantum potential V_Q were provided by [10]. This is not the first time where the metric components are expressed in terms of a potential. In General Relativity (GR) we learned that in the linearized weak gravity limit, and for slow moving bodies, the temporal metric component $g_{00} \sim \eta_{00} + 2V_N$ ($c = 1$), can be expressed in terms of the Newtonian potential V_N . Hence, given the explicit expressions [10] of the metric $g_{\mu\nu} = g_{\mu\nu}(V_Q)$ in terms of Bohm's quantum

potential, one can write down the curvature tensors, and the Einstein tensor leading to the field equations

$$R_{\mu\nu}(V_Q) - \frac{1}{2} g_{\mu\nu}(V_Q) R(V_Q) = 8\pi G T_{\mu\nu} \quad (33)$$

where the stress energy tensor is the one associated with a dark energy/dark matter fluid permeating all of space and given in terms of ρ, p . In particular, the BP equation (2) can be generalized to

$$R_{00}(V_Q) - \frac{1}{2} g_{00}(V_Q) R(V_Q) = 8\pi G T_{00} \quad (34)$$

where $T_{00} = \hat{\rho}_m = m\rho$. Solutions to eq-(33) will be provided in future investigations as well as the study of these equations in higher dimensions.

3 Conclusions

To conclude we add some remarks pertaining the Dirac-Eddington large numbers coincidences. Nottale [16] found long ago a direct relationship between the fine structure constant α and the cosmological constant Λ . In $\hbar = c = 1$ units, $\alpha = e^2 = 1/137$, the expression is

$$\Lambda \simeq \frac{(L_p)^4}{(r_e)^6} = \frac{(m_e)^6 (L_p)^4}{(\alpha)^6} = 10^{-56} \text{ cm}^{-2} \quad (35)$$

the classical electron radius r_e is defined in terms of the charge e , and electron mass m_e , as

$$\frac{e^2}{r_e} = \frac{\alpha}{r_e} = m_e \quad (36)$$

This important relation between Λ and α [16] warrants further investigation within the context of the Bohm-Poisson equation and the Dirac-Eddington large number coincidences.

We should mention that of the many articles surveyed in the literature pertaining the role of Bohm's quantum potential and cosmology, [13], [14], [15] we did not find any related to the Bohm-Poisson equation proposed in this work.³ The authors [14], for instance, have shown that replacing classical geodesics with quantal (Bohmian) trajectories gives rise to a quantum corrected Raychaudhuri equation (QRE). They derived the second order Friedmann equations from the QRE, and showed that this also contains a couple of quantum correction terms, the first of which can be interpreted as cosmological constant (and gives a correct estimate of its observed value), while the second as a radiation term in the early universe, which gets rid of the big-bang singularity and predicts an infinite age of our universe.

³A Google Scholar search provided the response “Bohm-Poisson equation and cosmological constant did not match any articles”

To finalize, we must say that the most attractive project is to find nontrivial solutions to the relativistic Bohm-Poisson equation (10). A careful inspection reveals that a separation of variables does not work. Solutions to eq-(33) are more difficult to find. We delegate this difficult task for future investigations as well as the study of these equations and solutions in higher dimensions.

Acknowledgments

We are indebted to M. Bowers for assistance and Tony Smith for discussions pertaining Armand Wyler's derivation of the fine structure constant and its plausible connection to the Bohm-Poisson equation.

References

- [1] “Dark Energy” http://en.wikipedia.org/wiki/Dark_energy
- [2] S. Carroll, “The cosmological constant”. Living Reviews in Relativity **4** (2001).
- [3] M. Sami and R. Myrzakulov, “Late time cosmic acceleration: ABCD of dark energy and modified theories of gravity”. International Journal of Modern Physics **D 25** (12) (2015).
- [4] A. Joyce, L. Lombriser, F. Schmidt, “Dark Energy vs. Modified Gravity”. Annual Review of Nuclear and Particle Science **66** (2016)
- [5] A. Arbey, “Is it possible to consider Dark Energy and Dark Matter as a same and unique Dark Fluid? <http://arxiv.org/abs/astro-ph/0506732>”
A. Arbey, “Dark Fluid: a complex scalar field to unify dark energy and dark matter <http://arxiv.org/abs/astro-ph/0601274>”
- [6] S.I. Pekar, “Untersuchungen über die Elektronentheorie der Kristalle” Akademie Verlag, Berlin, 1954, pp-29-34.
R. Ruffini and S. Bonazzola, “Systems of self-gravitating particles in general relativity and the concept of an equation of state”, Phys. Rev. **187** (1969) 1767
- [7] M. Bahrami, A. Grossardt, S. Donadi, and A. Bassi, “The Schrödinger-Newton equation and its foundations” New J. Phys. **16** (2014) 115007.
- [8] E. Santamato, “Geometric derivation of the Schrödinger equation from classical mechanics in curved Weyl spaces”, Phys. Rev. **D 29**, (1984) 216.
E. Santamato, “Statistical interpretation of the Klein-Gordon equation in terms of the space-time Weyl curvature”, Journal of Mathematical Physics **25**, 2477 (1984).

- [9] C. Castro, "Nonlinear Quantum Mechanics as the Weyl Geometry of a Classical Statistical Ensemble ". Found. Physics Letters, **vol.4**, no.1 (1991) 81.
C. Castro, "On Weyl Geometry, Random Processes and Geometric Quantum Mechanics ". Foundations of Physics. **vol 22**, no.4 (1992) 569- 615.
- [10] I. Tavernelli, "On the Geometrization of Quantum Mechanics" Annals of Physics **371** (2016) 239.
- [11] C. Castro, "Exact Solutions of the Newton-Schroedinger Equation, Infinite Derivative Gravity and Schwarzschild Atoms" to appear in the Physics and Astronomy International Journal.
- [12] C. Castro, "An elegant solution to the cosmological constant problem from the Bohm-Poisson equation" vixra.org :1710.0215; submitted to the AAC, Waldyr Rodrigues' memorial issue, October 2017.
- [13] A. Shojai and F. Shojai, "Constraints Algebra and Equations of Motion in Bohmian Interpretation of Quantum Gravity", arXiv : gr-qc/0311076.
D. Fiscaletti, "Bohm's Quantum Potential and the Geometry of Space", Quantum Matter **2**, 1 (2013) 45.
E. Squires, "A quantum solution to a cosmological mystery", Phys. Letts A **162**, 1 (1992) 35.
J. Vink, "The quantum potential interpretation of quantum mechanics due to Bohm is applied to the Wheeler-De Witt equation for minisuperspace" Nuc. Phys. B **369**, 3 (1992) 707.
T. Horiguchi, Mod. Phys. Lett. **A 9**, (1994).
J. Marto and P. Vargas Moniz, "de Broglie-Bohm FRW universes in quantum string cosmology", Phys. Rev. **D 65**, (2001) 023516.
- [14] A. Farag and S. Das, "Cosmology from quantum potential", PLB **741** (2015) 276.
- [15] H. Nikolic, "Time and Probability : From classical mechanics to relativistic Bohmian mechanics" arXiv : 1309.0400
H. Nikolic, "Cosmological constant, semiclassical gravity, and foundations of quantum mechanics" gr-qc/0611037.
H. Nikolic, "Interpretation miniatures" arXiv : 1703.08341
- [16] L. Nottale, "Mach's Principle, Dirac's Large Number Hypothesis and the Cosmological Constant Problem", <https://www.luth.obspm.fr/luthier/nottale/arlambda.pdf> (1993).
L. Nottale, *Scale Relativity And Fractal Space-Time: A New Approach to Unifying Relativity and Quantum Mechanics* (World Scientific 2011)
L. Nottale, *Fractal Space-Time and Micro-physics* (World Scientific 1993).

- [17] C. Castro, “Asymptotic Safety in Quantum Gravity and Diffeomorphic Non-isometric Metric Solutions to the Schwarzschild Metric” to appear in Can. J. Physics, 2017.
- [18] C. Castro, “On Maximal Proper Force, Black Hole Horizons and Matter as Curvature in Momentum Space” to appear in IJGMMP.
 C. Castro, “On Dual Phase Space Relativity, the Machian Principle and Modified Newtonian Dynamics” Progress in Physics **1** (April 2005) 20.
 C. Castro, “Solutions to the Gravitational Field Equations in Curved Phase-Spaces” to appear in the EJTP.
- [19] S. Weinberg, “Ultraviolet divergences in quantum theories of gravitation” *General Relativity : An Einstein centenary survey*, Eds. S. Hawking and W. Israel, Cambridge University Press (1979) p. 790.
 M. Reuter and F. Saueressig, “Quantum Einstein Gravity” arXiv: 1202.2274
 R. Percacci, “A short introduction to Asymptotic Safety” arXiv: 1110.6389.
 D. Litim, “Renormalization group and the Planck scale” arXiv: 1102.4624.
 S. Nagy, “Lectures on Renormalization and Asymptotic Safety”, arXiv : 1211.4151.
 M. Niedermaier, “The Asymptotic Safety Scenario in Quantum Gravity-An Introduction” , gr-qc/0610018.
- [20] A. Bonano and Reuter. “Renormalization group improved black hole spacetimes” Phys. Rev. **D62** (2000) 043008.
- [21] M. Reuter and F. Saueressig, “Asymptotic Safety, Fractals and Cosmology”, arXiv: 1205.5431.