



The article appeared in English in open access in the journal

Open Physics

Derivation method of numerous dynamics in the Special Theory of Relativity

Open Physics, Vol. 17, 2019, 153-166, ISSN: 2391-5471

Szostek Roman

<https://doi.org/10.1515/phys-2019-0016>

Derivation method of numerous dynamics in the Special Theory of Relativity

Roman Szostek

*Rzeszow University of Technology, Department of Quantitative Methods, Rzeszow, Poland
rszostek@prz.edu.pl*

Abstract:

The article presents innovative method of deriving dynamics in the Special Theory of Relativity. This method enables to derive infinitely many dynamics in relativistic mechanics. The author has shown five examples of these derivations. In this way, It is presented that the dynamics known today as the dynamics of Special Theory of Relativity is only one of infinitely of theoretically possible.

Keywords: dynamics of bodies, Special Theory of Relativity

PACS: 03.30.+p, 02.90.+p

1. Introduction

The currently accepted dynamics of STR have different experimental confirmations. Present article does not deal with these experiments. They are not analyzed, nor evaluated. The purpose of this article is to show that the concrete dynamics of STR does not only result from the kinematics of STR only. Formally and correctly mathematically, many STR dynamics, which have different properties, can be derived within the kinematics STR. Relativistic dynamics is derived based on the relativistic kinematics and one additional assumption, which allows the concept of mass, momentum and kinetic energy to be introduced into the theory. This paper is a discussion about possible assumptions and the dynamics of STR resulting from these assumptions. It presents the author's method of deriving numerous dynamics for this theory.

The decision, in which dynamic STR is correct, can only result from experiments. Available publications show that the dynamics indicated by Albert Einstein is correct, that is for $x = 3/2$ (see Section 5). However, because each experiment is fraught with errors, it is possible that more

accurate experiments carried out in the future will show that the optimal model of dynamics for $x = 3/2 \pm \Delta x$, where Δx is a noticeable correction.

Kinematics deals with the movement of bodies without taking their physical characteristics into account. The basic concepts of kinematics are: time, location, transformation, speed and acceleration.

Dynamics deals with the movement of material bodies under the action of forces. The basic concepts of dynamics are: inertial mass, force, momentum and kinetic energy.

Kinematics and dynamics are resulting in mechanics. This study deals with relativistic mechanics, i.e. the Special Theory of Relativity, which unlike classical mechanics, also applies to high-speed.

2. Kinematic assumptions of the Special Theory of Relativity

The kinematics of the Special Theory of Relativity is based on the following assumptions:

I. **All inertial systems are equivalent.**

This assumption means that there is no physical phenomenon, which distinguishes the inertial system. In a particular case, it means that there is no such phenomenon for which the absolute rest is needed to explain. Mathematically, it results from this assumption that each coordinate and time transformation have coefficients with exactly the same numerical values as inverse transformation (with the accuracy to the sign resulting from the velocity direction between the systems).

II. **Velocity of light c in vacuum is the same in every direction and in each inertial system.**

III. **Transformation of time and position coordinates between the inertial systems is linear.**

These assumptions are often written in other equivalent forms.

Based on mentioned assumptions, it is possible to derive Lorentz transformation on which the Special Theory of Relativity is based. There are many different derivation ways of this transformation. Two derivations are presented in monograph [1].

Markings adopted in Figure 1 will be convenient for our needs. Inertial systems move along their x -axis. Symbol $v_{2/1}$ stands for a velocity of U_2 system measured by the observer from U_1 system, while $v_{1/2}$ is a velocity of U_1 system measured by the observer from U_2 system. In the Special Theory of Relativity occurs that $v_{2/1} = -v_{1/2}$.




Figure 1: Relative movement of inertial systems U_1 and U_2 ($v_{2/1} = -v_{1/2}$).

Lorentz transformation from U_2 to U_1 system has a form of:

$$t_1 = \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} (t_2 + \frac{v_{2/1}}{c^2} x_2) \quad (1)$$

$$x_1 = \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} (v_{2/1} t_2 + x_2) \quad (2)$$

$$y_1 = y_2, \quad z_1 = z_2 \quad (3)$$

Lorentz transformation from U_1 to U_2 system has a form of:

$$t_2 = \frac{1}{\sqrt{1 - (v_{1/2}/c)^2}} \left(t_1 + \frac{v_{1/2}}{c^2} x_1 \right) \quad (4)$$

$$x_2 = \frac{1}{\sqrt{1 - (v_{1/2}/c)^2}} (v_{1/2} t_1 + x_1) \quad (5)$$

$$y_2 = y_1, \quad z_2 = z_1 \quad (6)$$

Transformation (1)-(3) and (4)-(6) includes complete information on the relativistic kinematics.

3. Selected properties of relativistic kinematics

In order to derive dynamics two formulas from kinematics, i.e. (20) and (23) from kinematics will be needed. They will be derived out of transformation (1)-(3).

3.1. Transformation of velocity

Determine the differentials from transformation (1)-(3)

$$dt_1 = \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} \left(dt_2 + \frac{v_{2/1}}{c^2} dx_2 \right) \quad (7)$$

$$dx_1 = \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} (v_{2/1} dt_2 + dx_2) \quad (8)$$

$$dy_1 = dy_2, \quad dz_1 = dz_2 \quad (9)$$

From the inertial system U_1 and U_2 , the moving body U_3 is observed. In U_1 system, it has a velocity of $v_{3/1}$, while in U_2 system it has a velocity of $v_{3/2}$. The components of these velocities are presented in Figure 2.




Figure 2: Movement of the body from two inertial systems U_1 and U_2 .

The coordinates of body U_3 position in U_1 system are x_1, y_1, z_1 . At the same time in U_2 system these coordinates are x_2, y_2, z_2 . Since the body U_3 moves, these coordinates change in time. When time dt_1 elapses in U_1 system then time dt_2 elapses in U_2 system. For such indications the changes of coordinates of body U_3 position in U_1 system in the time interval dt_1 are dx_1, dy_1, dz_1 . Changes of coordinates of body U_3 position in U_2 system in the time interval dt_2 are dx_2, dy_2, dz_2 .

The body velocity U_3 in inertial system U_2 has the following components:

$$v_{3/2}^x = \frac{dx_2}{dt_2}, \quad v_{3/2}^y = \frac{dy_2}{dt_2}, \quad v_{3/2}^z = \frac{dz_2}{dt_2} \quad (10)$$

The body velocity U_3 in inertial system U_1 has the following components:

$$v_{3/1}^x = \frac{dx_1}{dt_1}, \quad v_{3/1}^y = \frac{dy_1}{dt_1}, \quad v_{3/1}^z = \frac{dz_1}{dt_1} \quad (11)$$

When differentials (7)-(9) are put into Eqs. (11), one will receive

$$\left\{ \begin{array}{l} v_{3/1}^x = \frac{\frac{1}{\sqrt{1-(v_{2/1}/c)^2}}(v_{2/1}dt_2 + dx_2)}{\frac{1}{\sqrt{1-(v_{2/1}/c)^2}}(dt_2 + \frac{v_{2/1}}{c^2}dx_2)} \\ v_{3/1}^y = \frac{dy_2}{\frac{1}{\sqrt{1-(v_{2/1}/c)^2}}(dt_2 + \frac{v_{2/1}}{c^2}dx_2)} \\ v_{3/1}^z = \frac{dz_2}{\frac{1}{\sqrt{1-(v_{2/1}/c)^2}}(dt_2 + \frac{v_{2/1}}{c^2}dx_2)} \end{array} \right. \quad (12)$$

i.e.

$$\left\{ \begin{array}{l} v_{3/1}^x = \frac{v_{2/1} + dx_2/dt_2}{1 + \frac{v_{2/1}}{c^2}(dx_2/dt_2)} \\ v_{3/1}^y = \sqrt{1-(v_{2/1}/c)^2} \frac{dy_2/dt_2}{1 + \frac{v_{2/1}}{c^2}(dx_2/dt_2)} \\ v_{3/1}^z = \sqrt{1-(v_{2/1}/c)^2} \frac{dz_2/dt_2}{1 + \frac{v_{2/1}}{c^2}(dx_2/dt_2)} \end{array} \right. \quad (13)$$

On the basis of (10) the desired velocity transformation from U_2 to U_1 system is obtained

$$\left\{ \begin{array}{l} v_{3/1}^x = \frac{v_{3/2}^x + v_{2/1}}{1 + \frac{v_{3/2}^x v_{2/1}}{c^2}} \\ v_{3/1}^y = \sqrt{1-(v_{2/1}/c)^2} \frac{v_{3/2}^y}{1 + \frac{v_{3/2}^x v_{2/1}}{c^2}} \\ v_{3/1}^z = \sqrt{1-(v_{2/1}/c)^2} \frac{v_{3/2}^z}{1 + \frac{v_{3/2}^x v_{2/1}}{c^2}} \end{array} \right. \quad (14)$$

In special case, when U_3 body moves parallel to x -axis then occurs

$$v_{3/1}^x = v_{3/2}, \quad v_{3/2}^x = v_{3/2}, \quad v_{3/1}^y = v_{3/2}^y = 0, \quad v_{3/1}^z = v_{3/2}^z = 0 \quad (15)$$

Then velocity transformation (14) takes the form of formula to sum-up parallel velocities

$$v_{3/1} = \frac{v_{3/2} + v_{2/1}}{1 + \frac{v_{3/2}v_{2/1}}{c^2}} \quad (16)$$

3.2. Change of velocity seen from different inertial systems

The body at rest in U_3 system has momentary acceleration to U_3' system. The body movement is observed from U_1 and U_2 systems. The velocities of inertial systems are parallel to each other. Markings shown in Figure 3 are adopted.




Figure 3: Increases in the velocity seen in inertial systems U_1 and U_2 .

The differentials from formula (16) will be determined:

$$dv_{3/1} = \frac{d \frac{v_{3/2} + v_{2/1}}{1 + (v_{3/2}v_{2/1})/c^2}}{dv_{3/2}} dv_{3/2} = \frac{1 + \frac{v_{3/2}v_{2/1}}{c^2} - (v_{3/2} + v_{2/1})\frac{v_{2/1}}{c^2}}{\left(1 + \frac{v_{3/2}v_{2/1}}{c^2}\right)^2} dv_{3/2} \quad (17)$$

$$dv_{3/1} = \frac{1 - \frac{v_{2/1}^2}{c^2}}{\left(1 + \frac{v_{3/2}v_{2/1}}{c^2}\right)^2} dv_{3/2} \quad (18)$$

If U_3 system is U_2 system then it is necessary to replace index 3 with 2. Then,

$$dv_{3/1} = dv_{2/1}, \quad v_{3/2} = v_{2/2} = 0, \quad dv_{3/2} = dv_{2/2} \quad (19)$$

On this basis, the formula (18) takes a form of

$$dv_{2/2} = \frac{dv_{2/1}}{1 - (v_{2/1}/c)^2} \quad (20)$$

Relation (20) is related to the change of body velocity seen in the inertial system U_2 , in which the body is located ($dv_{2/2}$), and the change of velocity seen from another inertial system U_1 ($dv_{2/1}$).

3.3. Time dilatation

If motionless body is in U_2 system, then for its coordinates occurs

$$\frac{dx_2}{dt_2} = 0 \quad (21)$$

Based on time transformation (7) one receives

$$\frac{dt_1}{dt_2} = \frac{1}{\sqrt{1-(v_{2/1}/c)^2}} \left(1 + \frac{v_{2/1}}{c^2} \frac{dx_2}{dt_2}\right) \stackrel{\frac{dx_2}{dt_2}=0}{\Rightarrow} \frac{dt_1}{dt_2} = \frac{1}{\sqrt{1-(v_{2/1}/c)^2}} \quad (22)$$

On this basis we receive the formula for time dilatation of motionless body with regard to U_2 system

$$\frac{dx_2}{dt_2} = 0 \Rightarrow dt_2 = \sqrt{1-(v_{2/1}/c)^2} \cdot dt_1 \quad (23)$$

Recording of time dilatation in a form of (23) is more precise than the commonly used recording, as it has a form of implications. Such a record makes it clear that such dilatation applies only to motionless bodies in relation to U_2 system (or for events occurring in the same position in relation to U_2 system).

4. Dynamics in the Special Theory of Relativity

All dissertations will be conducted only for one-dimensional model, i.e. all analyzed vector values will be parallel to x -axis. Each derived dynamic can easily be generalized into three-dimensional cases.

In order to derive dynamics in the Special Theory of Relativity (STR), it is necessary to adopt an additional assumption, which allows the concept of mass, momentum and kinetic energy to be introduced into the theory. Depending on the assumption, different dynamics of bodies are received.

The inertial mass body resting in inertial frame of reference is determined by m_0 (rest mass). The rest mass is determined on the base unit of mass and the method of comparing any mass with this base unit. The inertial mass body at rest in U_2 , as seen from U_1 system, is determined by $m_{2/1}$ (relativistic mass). It is worth to note that the relativistic mass in this case is an inertial mass that occurs in the Newton's second law, rather than mass occurring in the formula for momentum, as assumed in the STR. In this way, a different definition of relativistic mass has been adopted, than one in the STR. Such a definition of the relativistic mass is more convenient in deriving dynamics.

The body of m_0 inertial mass is in U_2 system. It is affected by force $F_{2/2}$ that causes acceleration of $dv_{2/2}/dt_2$. Therefore, for the observer from U_2 system, the Newton's second law takes a form of

$$F_{2/2} := m_0 \cdot a_{2/2} = m_0 \frac{dv_{2/2}}{dt_2} \quad (24)$$

For the observer from U_1 system, inertial mass of the same body is $m_{2/1}$. For this observer, the force $F_{2/1}$ acts on the body, causing acceleration of $dv_{2/1}/dt_1$. Therefore, for the observer from U_1 the Newton's second law takes the form of

$$F_{2/1} := f(v_{2/1}) \cdot m_0 \cdot a_{2/1} = m_{2/1}(v_{2/1}) \cdot a_{2/1} = m_{2/1} \cdot a_{2/1} = m_{2/1} \frac{dv_{2/1}}{dt_1} \quad (25)$$

Equation (25) means that a generalized form of the Newton's second law is postulated. This generalized form contains an additional parameter $f(v)$. From the formula (24) shows that always $f(0)=1$. In classical mechanics $f(v)=1$, while in the current dynamics STR $f(v)=\gamma^3$ (formula (32)). Determining another form of parameter $f(v)$ leads to other dynamics for STR. The inertial relativistic mass $m_{2/1}$ is the product of this additional parameter $f(v)$ and the inertial mass

body at rest m_0 . In this article, the parameter $f(v)$ will not be used, only the inertial relativistic mass $m_{2/1}$.

Definitions identical as in classical mechanics apply for momentum and kinetic energy.

For the observer from U_2 system, the change of this body momentum can be recorded in the following forms

$$dp_{2/2} := F_{2/2} \cdot dt_2 = m_0 \cdot a_{2/2} \cdot dt_2 = m_0 \frac{dv_{2/2}}{dt_2} dt_2 = m_0 \cdot dv_{2/2} \quad (26)$$

For the observer from U_1 system, the change of this body momentum can be recorded in the following forms

$$dp_{2/1} := F_{2/1} \cdot dt_1 = m_{2/1} \cdot a_{2/1} \cdot dt_1 = m_{2/1} \frac{dv_{2/1}}{dt_1} dt_1 = m_{2/1} \cdot dv_{2/1} \quad (27)$$

where:

- $dp_{2/2}$ is a change of body momentum with rest mass m_0 in the inertial system U_2 , measured by the observer from the same inertial system U_2 ,
- $dp_{2/1}$ is a change of body momentum in the inertial system U_2 , measured by the observer from the same inertial system U_1 .

Kinetic energy of the body is equal of the work into its acceleration. For the observer from U_1 system, the change of kinetic energy of this body is as follows

$$dE_{2/1} := F_{2/1} \cdot dx_{2/1} = m_{2/1} \cdot a_{2/1} \cdot dx_{2/1} = m_{2/1} \frac{dv_{2/1}}{dt_1} dx_{2/1} = m_{2/1} \frac{dx_{2/1}}{dt_1} dv_{2/1} = m_{2/1} \cdot v_{2/1} \cdot dv_{2/1} \quad (28)$$

where:

- $dE_{2/1}$ is a change of kinetic energy of the body in inertial system U_2 , measured by the observer from the inertial system U_1 .

4.1. STR dynamics with constant force (STR/F)

In this section, a model of dynamics of bodies based on the assumption that the force accelerating of the body (parallel to x -axis) is the same for an observer from every inertial system will be derived (hence indication F).

4.1.1. The relativistic mass in STR/F

In the model STR/F it is assumed, that

$$F_{2/1}^F := F_{2/2} \quad (29)$$

Having introduced (24) and (25), one obtains

$$m_{2/1}^F \frac{dv_{2/1}}{dt_1} = m_0 \frac{dv_{2/2}}{dt_2} \quad (30)$$

On the base (20) and (23), one has

$$m_{2/1}^F \frac{dv_{2/1}}{dt_1} = m_0 \frac{\frac{dv_{2/1}}{1 - (v_{2/1}/c)^2}}{\sqrt{1 - (v_{2/1}/c)^2} \cdot dt_1} \quad (31)$$

Hence, a formula for relativistic mass of the body that is located in the system U_2 and is seen from the system U_1 is obtained, when assumption (29) is satisfied, as below

$$m_{2/1}^F = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^{3/2} \quad (32)$$

4.1.2. The momentum in STR/F

The body of rest mass m_0 is associated with the system U_2 . To determine the momentum of the body relative to the system U_1 a substitution of (32) to (27)

$$dp_{2/1}^F = m_{2/1}^F \cdot dv_{2/1} = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^{3/2} dv_{2/1} = m_0 c^3 \frac{1}{(c^2 - v_{2/1}^2)^{3/2}} dv_{2/1} \quad (33)$$

The body momentum is a sum of increases in its momentum, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$p_{2/1}^F = m_0 c^3 \int_0^{v_{2/1}} \frac{1}{(c^2 - v_{2/1}^2)^{3/2}} dv_{2/1} \quad (34)$$

From the work [2] (formula 72, p. 167) it is possible to read out, that

$$\int \frac{dx}{(a^2 - x^2)^{3/2}} = \frac{x}{a^2 \sqrt{a^2 - x^2}}, \quad a \neq 0 \quad (35)$$

After applying the integral (35) to (34) the formula for the body momentum in U_2 system is received and measured by the observer from U_1 system in a form of

$$p_{2/1}^F = m_0 c^3 \frac{v_{2/1}}{c^2 \sqrt{c^2 - v_{2/1}^2}} = \frac{m_0}{\sqrt{1 - (v_{2/1}/c)^2}} v_{2/1} \quad (36)$$

This formula is identical to the formula for momentum known from the STR, for the same reasons as in the case of momentum. This is because the dynamics known from the STR is derived from the assumption (29). It was adopted unconsciously, because it was considered as necessary. The awareness of this assumption allows to its change and derives other dynamics.

As already mentioned above, the definition of relativistic mass adopted is different from the definition adopted in the STR. In this case, the relativistic mass is the one, which occurs in the Newton's second law (25). In this particular case, it is expressed in terms of dependency (32). In the STR, the relativistic mass is the one, which occurs in the formula (36) per momentum.

4.1.3. The momentum in STR/F for small velocities

For small velocity $v_{2/1} \ll c$ momentum (36) comes down to the momentum from classical mechanics, because

$$v_{2/1} \ll c \Rightarrow p_{2/1}^F \approx m_0 v_{2/1} \quad (37)$$

4.1.4. The kinetic energy in STR/F

A determination of the formula for kinetic energy will be given. The dependence for the relativistic mass (32) is introduced to the formula (28)

$$dE_{2/1}^F = m_{2/1}^F \cdot v_{2/1} \cdot dv_{2/1} = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^{3/2} v_{2/1} dv_{2/1} = m_0 c^3 \frac{v_{2/1}}{(c^2 - v_{2/1}^2)^{3/2}} dv_{2/1} \quad (38)$$

The kinetic energy of body is a sum of increases in its kinetic energy, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$E_{2/1}^F = m_0 c^3 \int_0^{v_{2/1}} \frac{v_{2/1}}{(c^2 - v_{2/1}^2)^{3/2}} dv_{2/1} \quad (39)$$

From the work [2] (formula 74, p. 167) it is possible to read out, that

$$\int \frac{x dx}{(a^2 - x^2)^{3/2}} = \frac{1}{\sqrt{a^2 - x^2}} \quad (40)$$

After applying the integral (40) to (39) the formula for the kinetic energy of the body in U_2 system and measured by the observer from U_1 system in a form of

$$E_{2/1}^F = m_0 c^3 \frac{1}{\sqrt{c^2 - x^2}} \Big|_0^{v_{2/1}} = m_0 c^3 \left(\frac{1}{\sqrt{c^2 - v_{2/1}^2}} - \frac{1}{c} \right) = m_0 c^2 \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} - m_0 c^2 \quad (41)$$

This formula is identical to the formula for kinetic energy known from the STR, for the same reasons as in the case of momentum (36).

4.1.5. The kinetic energy in STR/F for small velocities

Formula (41) can be written in the form

$$E_{2/1}^F = m_0 c^2 \frac{1 - \sqrt{1 - (v_{2/1}/c)^2}}{\sqrt{1 - (v_{2/1}/c)^2}} \cdot \frac{1 + \sqrt{1 - (v_{2/1}/c)^2}}{1 + \sqrt{1 - (v_{2/1}/c)^2}} \quad (42)$$

$$E_{2/1}^F = \frac{m_0 v_{2/1}^2}{2} \frac{2}{1 - \frac{v_{2/1}^2}{c^2} + \sqrt{1 - \frac{v_{2/1}^2}{c^2}}} \quad (43)$$

On this basis, for small values $v_{2/1} \ll c$ one receives

$$v_{2/1} \ll c \Rightarrow E_{2/1}^F \approx \frac{m_0 v_{2/1}^2}{2} \frac{2}{1+1} = \frac{m_0 v_{2/1}^2}{2} \quad (44)$$

4.1.6. The force in STR/F

Due to the assumption (29) value measurement of the same force by two different observers is identical.

4.2. STR dynamics with constant momentum change (STR/ Δp)

In this section, a model of dynamics of bodies based on the assumption that the change in momentum of the body (parallel to x -axis) is the same for an observer from every inertial system will be derived (hence indication Δp).

These dynamics seem particularly interesting, because the conservation law of momentum is a fundamental law. Assumption that the change of body momentum is the same for every observer seems to be a natural extension of this law.

4.2.1. The relativistic mass in STR/ Δp

In the model STR/ Δp it is assumed, that

$$dp_{2/1}^{\Delta p} := dp_{2/2} \quad (45)$$

Having introduced (26) and (27), one obtains

$$m_{2/1}^{\Delta p} dv_{2/1} = m_0 dv_{2/2} \quad (46)$$

On the base (20), one has

$$m_{2/1}^{\Delta p} dv_{2/1} = m_0 \frac{dv_{2/1}}{1 - (v_{2/1}/c)^2} \quad (47)$$

Hence, a formula for relativistic mass of the body that is located in the system U_2 and is seen from the system U_1 is obtained, when assumption (45) is satisfied, as below

$$m_{2/1}^{\Delta p} = m_0 \frac{1}{1 - (v_{2/1}/c)^2} \quad (48)$$

4.2.2. The momentum in STR/ Δp

The body of rest mass m_0 is associated with the system U_2 . To determine the momentum of the body relative to the system U_1 a substitution of (48) to (27) is made

$$dp_{2/1}^{\Delta p} = m_{2/1}^{\Delta p} \cdot dv_{2/1} = m_0 \frac{1}{1 - (v_{2/1}/c)^2} dv_{2/1} = m_0 c^2 \frac{1}{c^2 - v_{2/1}^2} dv_{2/1} \quad (49)$$

The body momentum is a sum of increases in its momentum, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$p_{2/1}^{\Delta p} = m_0 c^2 \int_0^{v_{2/1}} \frac{1}{c^2 - v_{2/1}^2} dv_{2/1} \quad (50)$$

From the work [2] (formula 52, p. 160) it is possible to read out, that

$$\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right|, \quad a \neq 0 \quad (51)$$

After applying the integral (51) to (50) the formula for the body momentum in U_2 system and measured by the observer from U_1 system is received in a form of

$$p_{2/1}^{\Delta p} = m_0 c^2 \frac{1}{2c} \ln \left| \frac{c+x}{c-x} \right|_0^{v_{2/1}} = \frac{m_0 c}{2} \ln \left(\frac{c + v_{2/1}}{c - v_{2/1}} \right) \quad (52)$$

4.2.3. The momentum in STR/ Δp for small velocities

Formula (52) can be written in the form

$$p_{2/1}^{\Delta p} = \frac{m_0 v_{2/1}}{2} \frac{c}{v_{2/1}} \ln \left(\frac{c + v_{2/1}}{c - v_{2/1}} \right) = \frac{m_0 v_{2/1}}{2} \ln \left(\frac{(1 + v_{2/1}/c)^{c/v_{2/1}}}{(1 - v_{2/1}/c)^{c/v_{2/1}}} \right) \quad (53)$$

$$p_{2/1}^{\Delta p} = \frac{m_0 v_{2/1}}{2} \ln \left(\frac{\left(1 + \frac{1}{c/v_{2/1}}\right)^{c/v_{2/1}}}{\left(1 - \frac{1}{c/v_{2/1}}\right)^{c/v_{2/1}}} \right) \quad (54)$$

On this basis, for small values $v_{2/1} \ll c$ one receives

$$v_{2/1} \ll c \Rightarrow p_{2/1}^{\Delta p} \approx \frac{m_0 v_{2/1}}{2} \ln \left(\frac{e}{1/e} \right) = \frac{m_0 v_{2/1}}{2} \ln(e^2) = m_0 v_{2/1} \quad (55)$$

4.2.4. The kinetic energy in STR/ Δp

A determination of the formula for kinetic energy will be given. The dependence for the relativistic mass (48) is introduced to the formula (28)

$$dE_{2/1}^{\Delta p} = m_{2/1}^{\Delta p} \cdot v_{2/1} \cdot dv_{2/1} = m_0 \frac{1}{1 - (v_{2/1}/c)^2} v_{2/1} dv_{2/1} = m_0 c^2 \frac{v_{2/1}}{c^2 - v_{2/1}^2} dv_{2/1} \quad (56)$$

The kinetic energy of body is a sum of increases in its kinetic energy, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$E_{2/1}^{\Delta p} = m_0 c^2 \int_0^{v_{2/1}} \frac{v_{2/1}}{c^2 - v_{2/1}^2} dv_{2/1} \quad (57)$$

From the work [2] (formula 56, p. 160) it is possible to read out, that

$$\int \frac{x}{a^2 - x^2} dx = -\frac{1}{2} \ln|a^2 - x^2| \quad (58)$$

After applying the integral (58) to (57) the formula for the kinetic energy of the body in U_2 system and measured by the observer from U_1 system in a form of

$$E_{2/1}^{\Delta p} = -m_0 c^2 \frac{1}{2} \ln|c^2 - x^2| \Big|_0^{v_{2/1}} = -\frac{m_0 c^2}{2} \ln(c^2 - v_{2/1}^2) + \frac{m_0 c^2}{2} \ln(c^2) \quad (59)$$

$$E_{2/1}^{\Delta p} = \frac{m_0 c^2}{2} \ln \frac{c^2}{c^2 - v_{2/1}^2} = \frac{m_0 c^2}{2} \ln \frac{1}{1 - (v_{2/1}/c)^2} \quad (60)$$

4.2.5. The kinetic energy in STR/ Δp for small velocities

Formula (60) can be written in the form

$$E_{2/1}^{\Delta p} = \frac{m_0 v_{2/1}^2}{2} \frac{c^2}{v_{2/1}^2} \ln \frac{1}{1 - (v_{2/1}/c)^2} = \frac{m_0 v_{2/1}^2}{2} \ln \frac{1}{[1 - (v_{2/1}/c)^2]^{(c/v_{2/1})^2}} \quad (61)$$

$$E_{2/1}^{\Delta p} = \frac{m_0 v_{2/1}^2}{2} \ln \frac{1}{\left[1 - \frac{1}{(c/v_{2/1})^2}\right]^{(c/v_{2/1})^2}} \quad (62)$$

On this basis, for small values $v_{2/1} \ll c$ one receives

$$v_{2/1} \ll c \Rightarrow E_{2/1}^{\Delta p} \approx \frac{m_0 v_{2/1}^2}{2} \ln \frac{1}{1/e} = \frac{m_0 v_{2/1}^2}{2} \quad (63)$$

4.2.6. The force in STR/ Δp

Body with rest mass m_0 is related to U_2 system. It is affected by force that causes acceleration. For the observer from this system, the acceleration force has in accordance with (24) the following value

$$F_{2/2} = m_0 \frac{dv_{2/2}}{dt_2} \quad (64)$$

For the observer from U_1 system, acceleration force has in accordance with (25) the following value

$$F_{2/1}^{\Delta p} = m_{2/1}^{\Delta p} \frac{dv_{2/1}}{dt_1} \quad (65)$$

If to divide parties' equation (65) by (64), then on the basis of (20) and (23) one will receive

$$\frac{F_{2/1}^{\Delta p}}{F_{2/2}} = \frac{m_{2/1}^{\Delta p}}{m_0} \cdot \frac{dt_2}{dt_1} \cdot \frac{dv_{2/1}}{dv_{2/2}} = \frac{m_{2/1}^{\Delta p}}{m_0} (1 - (v_{2/1}/c)^2)^{3/2} \quad (66)$$

On the basis of (48) a relation between measurements of the same force by two different observers is obtained

$$F_{2/1}^{\Delta p} = \sqrt{1 - (v_{2/1}/c)^2} \cdot F_{2/2} \quad (67)$$

The highest value of force is measured by the observer from the inertial system in which the body is located.

4.3. STR dynamics with constant mass (STR/ m)

In this section, a model of dynamics of bodies, based on the assumption that body weight is the same for an observer from each inertial reference system, will be derived (hence indication m).

4.3.1. The relativistic mass in STR/ m

In the model STR/ m it is assumed, that

$$m_{2/1}^m := m_0 \quad (68)$$

Therefore, for the observer from inertial system U_1 , the body mass in U_2 system is the same as the rest mass.

4.3.2. The momentum in STR/ m

The body of rest mass m_0 is associated with the system U_2 . To determine the momentum of the body relative to the system U_1 a substitution of (68) to (27)

$$dp_{2/1}^m = m_{2/1}^m \cdot dv_{2/1} = m_0 dv_{2/1} \quad (69)$$

The body momentum is a sum of increases in its momentum, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$p_{2/1}^m = m_0 \int_0^{v_{2/1}} dv_{2/1} = m_0 v_{2/1} \quad (70)$$

In this relativistic dynamics the momentum is expressed with the same equation as in classical mechanics.

4.3.3. The kinetic energy in STR/m

A determination of the formula for kinetic energy will be given. The dependence for the relativistic mass (68) is introduced to the formula (28)

$$dE_{2/1}^m = m_{2/1}^m \cdot v_{2/1} \cdot dv_{2/1} = m_0 v_{2/1} dv_{2/1} \quad (71)$$

The kinetic energy of body is a sum of increases in its kinetic energy, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$E_{2/1}^m = m_0 \int_0^{v_{2/1}} v_{2/1} dv_{2/1} = \frac{m_0 v_{2/1}^2}{2} \quad (72)$$

In this relativistic dynamics the kinetic energy is expressed with the same equation as in classical mechanics.

4.3.4. The force in STR/m

Body with rest mass m_0 is related to U_2 system. It is affected by force that causes acceleration. For the observer from this system, the acceleration force has in accordance with (24) the following value

$$F_{2/2} = m_0 \frac{dv_{2/2}}{dt_2} \quad (73)$$

For the observer from U_1 system, acceleration force has in accordance with (25) the following value

$$F_{2/1}^m = m_{2/1}^m \frac{dv_{2/1}}{dt_1} = m_0 \frac{dv_{2/1}}{dt_1} \quad (74)$$

If to divide parties' equation (74) by (73), then on the basis of (20) and (23) one will receive

$$\frac{F_{2/1}^m}{F_{2/2}} = \frac{dt_2}{dt_1} \cdot \frac{dv_{2/1}}{dv_{2/2}} = (1 - (v_{2/1}/c)^2)^{3/2} \quad (75)$$

i.e.

$$F_{2/1}^m = (1 - (v_{2/1}/c)^2)^{3/2} \cdot F_{2/2} \quad (76)$$

The highest value of force is measured by the observer from the inertial system in which the body is located.

4.3.5. Discussion on the STR/m dynamics

Obtaining a relativistic dynamics, in which there is no relativistic mass, and equations for kinetic energy and momentum are identical as in classical mechanics can be surprising, because in

relativistic mechanics it is believed that the accelerated body can achieve maximum speed c . However, this dynamics is formally correct.

If the body velocity $v_{2/1}$ reaches c value, then according to (76)

$$F_{2/1}^m = (1 - v_{2/1}^2/c^2)^{3/2} \cdot F_{2/2} \approx 0 \quad (77)$$

In the inertial system U_2 , in which the body is located, can be affected by acceleration force $F_{2/2}$ of any, but finite value. However, from a perspective of the inertial system U_1 , towards which the body has c velocity, the same force is zero. This means that from a perspective of U_1 system, it is not possible to perform work on the body, which will increase its kinetic energy indefinitely. From the relation (72) it results that the kinetic energy, that a body with mass m_0 and velocity c has, a value has

$$E_{\max}^m = \frac{m_0 c^2}{2} \quad (78)$$

4.4. STR dynamics with constant force to its operation time (STR/ $F/\Delta t$)

In this section, a model of dynamics of bodies based on the assumption that the force that accelerates of the body (parallel to x -axis) divided by the time of operation of this force is the same for an observer from every inertial system will be derived (hence indication $F/\Delta t$).

4.4.1. The relativistic mass in STR/ $F/\Delta t$

In the model STR/ $F/\Delta t$ it is assumed, that

$$\frac{F_{2/1}^{F/\Delta t}}{dt_1} := \frac{F_{2/2}}{dt_2} \quad (79)$$

Having introduced (24) and (25), one obtains

$$m_{2/1}^{F/\Delta t} \frac{dv_{2/1}}{dt_1} \frac{1}{dt_1} = m_0 \frac{dv_{2/2}}{dt_2} \frac{1}{dt_2} \quad (80)$$

On the base (20) and (23), one has

$$m_{2/1}^{F/\Delta t} \frac{dv_{2/1}}{dt_1^2} = m_0 \frac{\frac{dv_{2/1}}{dt_1}}{(1 - (v_{2/1}/c)^2) dt_1^2} \quad (81)$$

Hence, a formula for relativistic mass of the body that is located in the system U_2 and is seen from the system U_1 is obtained, when assumption (79) is satisfied, as below

$$m_{2/1}^{F/\Delta t} = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^2 \quad (82)$$

4.4.2. The momentum in STR/ $F/\Delta t$

The body of rest mass m_0 is associated with the system U_2 . To determine the momentum of the body relative to the system U_1 a substitution of (82) to (27)

$$dp_{2/1}^{F/\Delta t} = m_{2/1}^{F/\Delta t} \cdot dv_{2/1} = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^2 dv_{2/1} = m_0 c^4 \frac{1}{(c^2 - v_{2/1}^2)^2} dv_{2/1} \quad (83)$$

The body momentum is a sum of increases in its momentum, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$p_{2/1}^{F/\Delta t} = m_0 c^4 \int_0^{v_{2/1}} \frac{1}{(c^2 - v_{2/1}^2)^2} dv_{2/1} \quad (84)$$

From the work [2] (formula 54, p. 160) it is possible to read out, that

$$\int \frac{dx}{(a^2 - x^2)^2} = \frac{x}{2a^2(a^2 - x^2)} + \frac{1}{4a^3} \ln \left| \frac{a+x}{a-x} \right|, \quad a \neq 0 \quad (85)$$

After applying the integral (85) to (84) the formula for the body momentum in U_2 system and measured by the observer from U_1 system in a form of

$$p_{2/1}^{F/\Delta t} = m_0 c^4 \left[\frac{x}{2c^2(c^2 - x^2)} + \frac{1}{4c^3} \ln \frac{(c+x)}{(c-x)} \right]_0^{v_{2/1}} = m_0 c \left[\frac{cv_{2/1}}{2(c^2 - v_{2/1}^2)} + \frac{1}{4} \ln \frac{(c+v_{2/1})}{(c-v_{2/1})} \right] \quad (86)$$

$$p_{2/1}^{F/\Delta t} = m_0 v_{2/1} \frac{1}{2} \left[\frac{1}{1 - (v_{2/1}/c)^2} + \ln \left(\frac{c+v_{2/1}}{c-v_{2/1}} \right)^{\frac{c}{2v_{2/1}}} \right] \quad (87)$$

4.4.3. The momentum in STR/ $F/\Delta t$ for small velocities

Formula (87) can be written in the form

$$p_{2/1}^{F/\Delta t} = m_0 v_{2/1} \left[\frac{1}{2(1 - (v_{2/1}/c)^2)} + \frac{1}{4} \ln \left(\frac{(1 + v_{2/1}/c)^{c/v_{2/1}}}{(1 - v_{2/1}/c)^{c/v_{2/1}}} \right) \right] \quad (88)$$

$$p_{2/1}^{F/\Delta t} = m_0 v_{2/1} \left[\frac{1}{2(1 - (v_{2/1}/c)^2)} + \frac{1}{4} \ln \left(\frac{\left(1 + \frac{1}{c/v_{2/1}}\right)^{c/v_{2/1}}}{\left(1 - \frac{1}{c/v_{2/1}}\right)^{c/v_{2/1}}} \right) \right] \quad (89)$$

On this basis, for small values $v_{2/1} \ll c$ one receives

$$v_{2/1} \ll c \Rightarrow p_{2/1}^{F/\Delta t} \approx m_0 v_{2/1} \left[\frac{1}{2} + \frac{1}{4} \ln \left(\frac{e}{1/e} \right) \right] = m_0 v_{2/1} \left[\frac{1}{2} + \frac{1}{4} \ln(e^2) \right] = m_0 v_{2/1} \quad (90)$$

4.4.4. The kinetic energy in STR/ $F/\Delta t$

A determination of the formula for kinetic energy will be given. The dependence for the relativistic mass (82) is introduced to the formula (28)

$$dE_{2/1}^{F/\Delta t} = m_{2/1}^{F/\Delta t} \cdot v_{2/1} \cdot dv_{2/1} = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^2 v_{2/1} dv_{2/1} = m_0 c^4 \frac{v_{2/1}}{(c^2 - v_{2/1}^2)^2} dv_{2/1} \quad (91)$$

The kinetic energy of body is a sum of increases in its kinetic energy, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$E_{2/1}^{F/\Delta t} = m_0 c^4 \int_0^{v_{2/1}} \frac{v_{2/1}}{(c^2 - v_{2/1}^2)^2} dv_{2/1} \quad (92)$$

From the work [2] (formula 58, p. 160) it is possible to read out, that

$$\int \frac{xdx}{(a^2 - x^2)^2} = \frac{1}{2(a^2 - x^2)} \quad (93)$$

After applying the integral (93) do (92) the formula for the kinetic energy of the body in U_2 system and measured by the observer from U_1 system in a form of

$$E_{2/1}^{F/\Delta t} = m_0 c^4 \frac{1}{2(c^2 - x^2)} \Big|_0^{v_{2/1}} = \frac{m_0 c^4}{2} \frac{1}{(c^2 - v_{2/1}^2)} - \frac{m_0 c^4}{2} \frac{1}{c^2} \quad (94)$$

$$E_{2/1}^{F/\Delta t} = \frac{m_0 c^2}{2} \frac{1}{1 - (v_{2/1}/c)^2} - \frac{m_0 c^2}{2} = \frac{m_0 v_{2/1}^2}{2} \frac{1}{1 - (v_{2/1}/c)^2} \quad (95)$$

The formula for kinetic energy (95) was derived from the work [3], due to the fact that the author adopted a different assumption than the one on which the dynamics known from the STR was based.

4.4.5. The kinetic energy in STR/ $F/\Delta t$ for small velocities

For small velocity $v_{2/1} \ll c$ kinetic energy (95) comes down to the kinetic energy from classical mechanics, because

$$v_{2/1} \ll c \Rightarrow E_{2/1}^{F/\Delta t} \approx \frac{m_0 v_{2/1}^2}{2} \cdot \frac{1}{1} = \frac{m_0 v_{2/1}^2}{2} \quad (96)$$

4.4.6. The force in STR/ $F/\Delta t$

Body with rest mass m_0 is related to U_2 system. It is affected by force that causes acceleration. For the observer from this system, the acceleration force has in accordance with (24) the following value

$$F_{2/2} = m_0 \frac{dv_{2/2}}{dt_2} \quad (97)$$

For the observer from U_1 system, acceleration force has in accordance with (25) the following value

$$F_{2/1}^{F/\Delta t} = m_{2/1}^{F/\Delta t} \frac{dv_{2/1}}{dt_1} \quad (98)$$

If to divide parties' equation (98) by (97), then on the basis of (20) and (23) one will receive

$$\frac{F_{2/1}^{F/\Delta t}}{F_{2/2}} = \frac{m_{2/1}^{F/\Delta t}}{m_0} \cdot \frac{dt_2}{dt_1} \cdot \frac{dv_{2/1}}{dv_{2/2}} = \frac{m_{2/1}^{F/\Delta t}}{m_0} (1 - (v_{2/1}/c)^2)^{3/2} \quad (99)$$

On the basis of (82) relation between measurements of the same force by two different observers is obtained

$$F_{2/1}^{F/\Delta t} = \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} \cdot F_{2/2} \quad (100)$$

The lowest value of force is measured by the observer from the inertial system in which the body is located.

4.5. STR dynamics with constant mass to elapse of observer's time (STR/ $m/\Delta t$)

In this subchapter a model of body dynamics will be derived based on the assumption that the body mass divided by the elapse of time in observer system is the same for the observer from each inertial frame of reference (hence indication $m/\Delta t$).

4.5.1. The relativistic mass in STR/ $m/\Delta t$

In the model STR/ $m/\Delta t$ it is assumed, that

$$\frac{m_{2/1}^{m/\Delta t}}{dt_1} := \frac{m_0}{dt_2} \quad (101)$$

On the base (23), one obtains

$$\frac{m_{2/1}^{m/\Delta t}}{dt_1} = \frac{m_0}{\sqrt{1 - (v_{2/1}/c)^2} \cdot dt_1} \quad (102)$$

Hence, a formula for relativistic mass of the body that is located in the system U_2 and is seen from the system U_1 is obtained, when assumption (101) is satisfied, as below

$$m_{2/1}^{m/\Delta t} = m_0 \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} \quad (103)$$

4.5.2. The momentum in STR/ $m/\Delta t$

The body of rest mass m_0 is associated with the system U_2 . To determine the momentum of the body relative to the system U_1 a substitution of (103) to (27)

$$dp_{2/1}^{m/\Delta t} = m_{2/1}^{m/\Delta t} \cdot dv_{2/1} = m_0 \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} dv_{2/1} = m_0 c \frac{1}{\sqrt{c^2 - v_{2/1}^2}} dv_{2/1} \quad (104)$$

The body momentum is a sum of increases in its momentum, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$p_{2/1}^{m/\Delta t} = m_0 c^2 \int_0^{v_{2/1}} \frac{1}{\sqrt{c^2 - v_{2/1}^2}} dv_{2/1} \quad (105)$$

From the work [2] (formula 71, p. 167) it is possible to read out, that

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a}, \quad a > 0 \quad (106)$$

After applying the integral (106) to (105) the formula for the body momentum in U_2 system and measured by the observer from U_1 system in a form of

$$p_{2/1}^{m/\Delta t} = m_0 c \cdot \arcsin \frac{v_{2/1}}{c} \Big|_0^{v_{2/1}} = m_0 c \cdot \arcsin \frac{v_{2/1}}{c} \quad (107)$$

4.5.3. The momentum in STR/ $m/\Delta t$ for small velocities

Formula (107) can be written in the form

$$p_{2/1}^{m/\Delta t} = m_0 v_{2/1} \frac{\arcsin \frac{v_{2/1}}{c}}{\frac{v_{2/1}}{c}} \quad (108)$$

On this basis, for small values $v_{2/1} \ll c$ one receives

$$v_{2/1} \ll c \Rightarrow p_{2/1}^{m/\Delta t} \approx m_0 v_{2/1} \quad (109)$$

4.5.4. The kinetic energy in STR/ $m/\Delta t$

A determination of the formula for kinetic energy will be given. The dependence for the relativistic mass (103) is introduced to the formula (28)

$$dE_{2/1}^{m/\Delta t} = m_{2/1}^{m/\Delta t} \cdot v_{2/1} \cdot dv_{2/1} = m_0 \frac{1}{\sqrt{1-(v_{2/1}/c)^2}} v_{2/1} dv_{2/1} = m_0 c \frac{v_{2/1}}{\sqrt{c^2 - v_{2/1}^2}} dv_{2/1} \quad (110)$$

The kinetic energy of body is a sum of increases in its kinetic energy, when the body is accelerated from the inertial system U_1 (the body has velocity 0) to the inertial system U_2 (the body has velocity $v_{2/1}$), i.e.

$$E_{2/1}^{m/\Delta t} = m_0 c \int_0^{v_{2/1}} \frac{v_{2/1}}{\sqrt{c^2 - v_{2/1}^2}} dv_{2/1} \quad (111)$$

From the work [2] (formula 73, p. 167) it is possible to read out, that

$$\int \frac{x}{\sqrt{a^2 - x^2}} dx = -\sqrt{a^2 - x^2} \quad (112)$$

After applying the integral (112) do (111) the formula for the kinetic energy of the body in U_2 system and measured by the observer from U_1 system in a form of

$$E_{2/1}^{m/\Delta t} = -m_0 c \sqrt{c^2 - v_{2/1}^2} \Big|_0^{v_{2/1}} = -m_0 c \sqrt{c^2 - v_{2/1}^2} + m_0 c \sqrt{c^2} \quad (113)$$

$$E_{2/1}^{m/\Delta t} = m_0 c^2 - m_0 c \sqrt{c^2 - v_{2/1}^2} = m_0 c^2 (1 - \sqrt{1 - (v_{2/1}/c)^2}) \quad (114)$$

4.5.5. The kinetic energy in STR/ $m/\Delta t$ for small velocities

Formula (114) can be written in the form

$$E_{2/1}^{m/\Delta t} = \frac{m_0 v_{2/1}^2}{2} \cdot \frac{2c^2}{v_{2/1}^2} \cdot \frac{(1 - \sqrt{1 - (v_{2/1}/c)^2})(1 + \sqrt{1 - (v_{2/1}/c)^2})}{1 + \sqrt{1 - (v_{2/1}/c)^2}} \quad (115)$$

$$E_{2/1}^{m/\Delta t} = \frac{m_0 v_{2/1}^2}{2} \cdot \frac{2c^2}{v_{2/1}^2} \cdot \frac{1 - (1 - (v_{2/1}/c)^2)}{1 + \sqrt{1 - (v_{2/1}/c)^2}} = \frac{m_0 v_{2/1}^2}{2} \frac{2}{1 + \sqrt{1 - (v_{2/1}/c)^2}} \quad (116)$$

On this basis, for small values $v_{2/1} \ll c$ one receives

$$v_{2/1} \ll c \Rightarrow E_{2/1}^{m/\Delta t} \approx \frac{m_0 v_{2/1}^2}{2} \cdot \frac{2}{2} = \frac{m_0 v_{2/1}^2}{2} \quad (117)$$

4.5.6. The force in STR/ $m/\Delta t$

Body with rest mass m_0 is related to U_2 system. It is affected by force that causes acceleration. For the observer from this system, the acceleration force has in accordance with (24) the following value

$$F_{2/2} = m_0 \frac{dv_{2/2}}{dt_2} \quad (118)$$

For the observer from U_1 system, acceleration force has in accordance with (25) the following value

$$F_{2/1}^{m/\Delta t} = m_{2/1}^{m/\Delta t} \frac{dv_{2/1}}{dt_1} \quad (119)$$

If to divide parties' equation (119) by (118), then on the basis of (20) and (23) one will receive

$$\frac{F_{2/1}^{m/\Delta t}}{F_{2/2}} = \frac{m_{2/1}^{m/\Delta t}}{m_0} \cdot \frac{dt_2}{dt_1} \cdot \frac{dv_{2/1}}{dv_{2/2}} = \frac{m_{2/1}^{m/\Delta t}}{m_0} (1 - (v_{2/1}/c)^2)^{3/2} \quad (120)$$

On the basis of (103) relation between measurements of the same force by two different observers is obtained

$$F_{2/1}^{m/\Delta t} = (1 - (v_{2/1}/c)^2) \cdot F_{2/2} \quad (121)$$

The highest value of force is measured by the observer from the inertial system in which the body is located.

5. The general form of dynamics

In presented examples, assumptions have been adopted which can be written in forms (30), (46), (68), (80) and (101). On this basis, it can be seen that the assumption for relativistic dynamics is as follows

$$m_{2/1}^{\{a,b\}} \frac{dv_{2/1}^a}{dt_1^b} = m_0 \frac{dv_{2/2}^a}{dt_2^b}, \quad a, b \in R \quad (122)$$

The physical meaning of the formula (122) depends on the value of the parameters a and b to be determined. For example, if $a = b = 1$, then this formula takes the form (29), equivalent to the form (30), from the first example.

On the basis of (20) and (23) one receives

$$m_{2/1}^{\{a,b\}} \frac{dv_{2/1}^a}{dt_1^b} = m_0 \frac{\frac{dv_{2/1}^a}{(1 - (v_{2/1}/c)^2)^a}}{(1 - (v_{2/1}/c)^2)^{b/2} \cdot dt_1^b} = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^{a+b/2} \frac{dv_{2/1}^a}{dt_1^b} \quad (123)$$

Markings are adopted

$$\{x\} \equiv \{a, b\} \quad \wedge \quad x = a + \frac{b}{2} \in R \quad (124)$$

Now on the basis of (123) the relativistic inertial mass of body in U_2 system, seen from U_1 system, when an assumption is fulfilled (122), is expressed in dynamics $\{x\}$ by the following formula

$$m_{2/1}^{\{x\}} = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^x \quad (125)$$

Each such relativistic mass defines a different relativistic dynamics.

According to presented examples, based on formulas (27) and (125), the momentum in dynamics $\{x\}$ is expressed by the following formula

$$p_{2/1}^{\{x\}} = \int_0^{v_{2/1}} dp_{2/1}^{\{x\}} = \int_0^{v_{2/1}} m_{2/1}^{\{x\}} \cdot dv_{2/1} = m_0 \int_0^{v_{2/1}} \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^x dv_{2/1} \quad (126)$$

$$p_{2/1}^{\{x\}} = m_0 c^{2x} \int_0^{v_{2/1}} \frac{1}{(c^2 - v_{2/1}^2)^x} dv_{2/1} \quad (127)$$

According to presented examples, based on formulas (28) and (125), the kinetic energy in dynamics $\{x\}$ is expressed by the following formula

$$E_{2/1}^{\{x\}} = \int_0^{v_{2/1}} dE_{2/1}^{\{x\}} = \int_0^{v_{2/1}} m_{2/1}^{\{x\}} \cdot v_{2/1} \cdot dv_{2/1} = m_0 \int_0^{v_{2/1}} \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^x v_{2/1} dv_{2/1} \quad (128)$$

$$E_{2/1}^{\{x\}} = m_0 c^{2x} \int_0^{v_{2/1}} \frac{v_{2/1}}{(c^2 - v_{2/1}^2)^x} dv_{2/1} \quad (129)$$

According to presented examples, based on formulas (24), (25) and (20), (23), the relation between forces in dynamics $\{x\}$ is expressed by the following formula

$$\frac{F_{2/1}^{\{x\}}}{F_{2/2}} = \frac{m_{2/1}^{\{x\}} \frac{dv_{2/1}}{dt_1}}{m_0 \frac{dv_{2/2}}{dt_2}} = \frac{m_{2/1}^{\{x\}} \frac{dv_{2/1}}{dt_1}}{m_0 \frac{dv_{2/1}}{1 - (v_{2/1}/c)^2} \cdot \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} \cdot dt_1}} = \frac{m_{2/1}^{\{x\}}}{m_0} (1 - (v_{2/1}/c)^2)^{3/2} \quad (130)$$

On the basis of (125) one receives

$$\frac{F_{2/1}^{\{x\}}}{F_{2/2}} = \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^x (1 - (v_{2/1}/c)^2)^{3/2} = \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^{x - \frac{3}{2}} \quad (131)$$

On the basis of (25) and (125) the Newton's second law for dynamics $\{x\}$ is obtained

$$F_{2/1}^{\{x\}} = m_0 \left[\frac{1}{1 - (v_{2/1}/c)^2} \right]^x a_{2/1} \quad (132)$$

6. Summary of dynamics

Summary derived formulas for momentum and kinetic energy:

Dynamics $x = 0$

$$p_{2/1}^m = m_0 v_{2/1} \quad (133)$$

$$E_{2/1}^m = \frac{m_0 v_{2/1}^2}{2} \quad (134)$$

Dynamics $x = 1/2$

$$p_{2/1}^{m/\Delta t} = m_0 c \cdot \arcsin \frac{v_{2/1}}{c} = m_0 v_{2/1} \frac{\arcsin(v_{2/1}/c)}{v_{2/1}/c} \quad (135)$$

$$E_{2/1}^{m/\Delta t} = m_0 c^2 (1 - \sqrt{1 - (v_{2/1}/c)^2}) = \frac{m_0 v_{2/1}^2}{2} \frac{2}{1 + \sqrt{1 - (v_{2/1}/c)^2}} \quad (136)$$

Dynamics $x = 1$

$$p_{2/1}^{\Delta p} = \frac{m_0 c}{2} \ln \left(\frac{c + v_{2/1}}{c - v_{2/1}} \right) = m_0 v_{2/1} \ln \left(\frac{c + v_{2/1}}{c - v_{2/1}} \right)^{\frac{c}{2v_{2/1}}} \quad (137)$$

$$E_{2/1}^{\Delta p} = \frac{m_0 c^2}{2} \ln \frac{1}{1 - (v_{2/1}/c)^2} = \frac{m_0 v_{2/1}^2}{2} \ln \frac{1}{[1 - (v_{2/1}/c)^2]^{(c/v_{2/1})^2}} \quad (138)$$

Dynamics $x = 3/2$

(recognized STR dynamics)

$$p_{2/1}^F = m_0 v_{2/1} \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} \quad (139)$$

$$E_{2/1}^F = m_0 c^2 \frac{1}{\sqrt{1 - (v_{2/1}/c)^2}} - m_0 c^2 = \frac{m_0 v_{2/1}^2}{2} \frac{2}{\sqrt{1 - \frac{v_{2/1}^2}{c^2}} \left(1 + \sqrt{1 - \frac{v_{2/1}^2}{c^2}} \right)} \quad (140)$$

Dynamics $x = 2$

$$p_{2/1}^{F/\Delta t} = m_0 v_{2/1} \frac{1}{2} \left[\frac{1}{1 - (v_{2/1}/c)^2} + \ln \left(\frac{c + v_{2/1}}{c - v_{2/1}} \right)^{\frac{c}{2v_{2/1}}} \right] \quad (141)$$

$$E_{2/1}^{F/\Delta t} = \frac{m_0 c^2}{2} \frac{1}{1 - (v_{2/1}/c)^2} - \frac{m_0 c^2}{2} = \frac{m_0 v_{2/1}^2}{2} \frac{1}{1 - (v_{2/1}/c)^2} \quad (142)$$

Figure 4 shows compared momentums from derived relativistic dynamics.




Figure 4: Module of the momentum in dynamics:
STR/m ($x=0$), STR/m/ Δt ($x=1/2$), STR/ Δp ($x=1$), STR/F ($x=3/2$) and STR/F/ Δt ($x=2$).

Figure 5 shows compared kinetic energies from derived relativistic dynamics.




Figure 5: Kinetic energies in dynamics:
STR/m ($x=0$), STR/m/ Δt ($x=1/2$), STR/ Δp ($x=1$), STR/F ($x=3/2$) and STR/F/ Δt ($x=2$).

Figure 6 shows relation between measurements of the same force.




Figure 6: Relation between measurements of the same force by two different observers in dynamics:
STR/m ($x=0$), STR/m/ Δt ($x=1/2$), STR/ Δp ($x=1$), STR/F ($x=3/2$) and STR/F/ Δt ($x=2$).

7. Even more general form of dynamics

Relation (125) to the relativistic mass can be even more generalized. In the general case, it is possible to assume that the relativistic mass is expressed by the following formula

$$m_{2/1}^{\{f\}} = m_0 \cdot f(v_{2/1}) \quad (143)$$

Where $f(v_{2/1})$ is any continuous function with the following properties

$$f(v_{2/1}) \geq 0 \quad (144)$$

$$f(0) = 1 \quad (145)$$

$$f(v_{2/1}) = f(-v_{2/1}) \quad (146)$$

Each function $f(v_{2/1})$ defines a different dynamics of the STR.

8. Final conclusions

This study presents the author's method of deriving dynamics in the Special Theory of Relativity (STR). Five examples of such deriving were shown.

Derivation of dynamics is based on two formulas applicable in the kinematics of STR, i.e. (20) and (23). In order to derive the dynamics of STR, it is necessary to adopt an additional assumption in kinematics, which allows the concept of mass, kinetic energy and momentum to be introduced into the theory.

The dynamics of STR/ F ($x = 3/2$) is nowadays recognized as the dynamics of the STR. It is based on the assumption that each force parallel to x -axis has the same value for the observer from each inertial frame of reference. Formally, however, other dynamics are possible in accordance with the kinematics of the STR. In order to derive them, it is necessary to base on a different assumption.

The currently accepted dynamics of STR has numerous experimental confirmations. However, it is not excluded that more accurate experiments designed specifically for this purpose will show that the optimal model is the dynamics for $x = 3/2 \pm \Delta x$, where Δx is a noticeable correction. A calorimeter can be useful for verification of different dynamics. This device can measure the amount of heat released when stopping particles to high speed. On this basis, it is possible to determine graphs of the kinetic energy of accelerated particles (for example in accelerators of elementary particles) as a function of their velocity, analogous to those presented in Figure 5. On this basis, it is possible to indicate the dynamics in which the kinetic energy of particles is compatible with experiments.

The presented method of dynamism derivation can also be used in other theory of body kinematics. In the monograph [1] this method was used to derive four dynamics in the Special Theory of Ether, which are allowed for kinematics derived in the articles [4, 5]. The presented method is analogous to that used in another area, in the article [6].

Reference list

- [1] Szostek K., Szostek R., *Special Theory of Ether*, Publishing house AMELIA, Rzeszów, Poland, (www.ste.com.pl), 2015, ISBN 978-83-63359-81-2 (in English)
Szostek K., Szostek R., *Szczególna Teoria Eteru*, Wydawnictwo Amelia, Rzeszów, Polska, (www.ste.com.pl), 2015, ISBN 978-83-63359-77-5 (in Polish)

- [2] Воднев В., Наумович А., Наумович Н., *Основные математические формулы. Справочник*, 1988, Минск, Издательство «Вышэйшая школа» Государственного комитета БССР, ISBN 5-339-00083-4
- [3] Osiak Z., *Energy in Special Relativity*, Theoretical Physics, Isaac Scientific Publishing, 2019, 4 (1), 22-25, ISSN 2519-9625
- [4] Szostek K., Szostek R., *The derivation of the general form of kinematics with the universal reference system*, Results in Physics, 2018, 8, 429-437, ISSN: 2211-3797
- [5] Szostek K., Szostek R., *Kinematics in Special Theory of Ether*, Moscow University Physics Bulletin, 2018, 73 (4), 413-421, ISSN 0027-1349
- Szostek K., Szostek R., *Кинематика в Специальной Теории Эфира*, Вестник Московского Университета. Серия 3. Физика и Астрономия, 2018, 73 (4), 413-421, ISSN 0579-9384 (in Russian)
- [6] Szostek R., *An estimation of the geothermal energy sources for generating electricity*, Springer, Lecture Notes in Electrical Engineering 324, Analysis and Simulation of Electrical and Computer, 2015, 127-133, ISSN 1876-1100

* * *

Additional author comment

There is no reason to treat relativistic dynamics in Special Theory of Relativity as exceptional, either for experimental or theoretical reasons. Therefore, determination of which possible dynamics of relativistic mechanics is a correct model of reality remains an open problem of physics and should be one of the most important tasks of future physics.

The fact that as a part of the Special Theory of Relativity, numerous dynamics can be derived greatly undermines the truth of the formula $E = mc^2$. According to my research, on the basis of relativistic mechanics, it is impossible to derive a formula expressing the internal energy of matter. All derivations of this formula are wrong. The relation between mass and energy ($E = mc^2$) can be introduced into the STR as an independent assumption, but it does not result from Lorentz transformation, nor from the assumption (29) on which the dynamics of STR is based. But then there is a need to experimentally show what exactly is the form of such a dependency (e.g. why not $E = mc^2/2$) and experimentally investigate whether sometimes the form of such a dependency does not depend on the type of matter that this formula regards.