

General Relativity Test by Investigating the Conservation of Energy in a Relativistic Free Fall in the Uniform Gravitational Field

By Jaroslav Hynecek¹

Abstract: This paper investigates the General Relativity Theory (GRT) by studying the relativistic free fall of a small test body in a uniform gravitational field. The paper compares the predictions of energy loss, perhaps by radiation, in a free fall obtained from the GRT and the Metric Theory of Gravity (MTG). It is found that the gravitational mass dependence on velocity in GRT is not correct, because it predicts a negative loss of energy while the MTG predicts correctly a positive loss.

Introduction: The theories describing the free fall are well understood in both; the GRT and the MTG. In the GRT the inertial mass and the gravitational mass are assumed identical with identical dependencies on velocity. In the MTG, on the other hand, the gravitational mass depends on velocity differently than the inertial mass ^[1, 2]. It is thus simple for both theories to derive equations describing the free fall velocity and from that the energy loss of a small test body that falls in a uniform gravitational field.

Theories: In the GRT the relation between the velocity v and time is somewhat more complicated but can be easily derived as follows:

$$\frac{d}{dt} \left[\frac{m_0 v}{\sqrt{1-v^2/c^2}} \right] = g \frac{m_0}{\sqrt{1-v^2/c^2}} \quad (1)$$

where m_0 is the rest mass and c the speed of light in a vacuum. The left hand side of Eq.1 is the relativistic formula for the inertial force and the right hand side is the formula for the gravitational force that includes the gravitational force dependence on velocity. The formula in Eq.1 can be rearranged and simplified resulting in the following relation for the small body acceleration:

$$\frac{dv}{dt} = g(1-v^2/c^2) \quad (2)$$

The energy loss will be derived from a comparison to the potential energy that is obtained by lifting the test body very slowly in the uniform field by a distance z . This energy is simply expressed as follows:

$$E = m_0 g \cdot z \quad (3)$$

For the falling body the energy incremental gain during the fall can be expressed in terms of the velocity v and the gravitational force F as follows:

$$dE = F \cdot v \cdot \frac{dv}{\left(\frac{dv}{dt}\right)} = \frac{g \cdot m_0}{\sqrt{1-\frac{v^2}{c^2}}} \frac{v \cdot dv}{\left(1-\frac{v^2}{c^2}\right)} \quad (4)$$

After rearrangements and integration the expression for the energy as a function of velocity becomes:

¹ Email: jhynecek@netscape.net ; © 2018 Isetex, Inc.

$$E = \frac{m_0 c^2}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (5)$$

This result is expected and it is a nice confirmation of methodology used in Eq.1.

In the next step we will evaluate the energy difference ΔE given by Eq.3 and Eq.5. However, for the convenience of calculations it will be beneficial to first evaluate the differential of this difference.

$$\frac{d\Delta E}{dt} = m_0 g \cdot v - \frac{m_0 v}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{3}{2}}} \frac{dv}{dt} \quad (6)$$

In this formula the variable v in the first term was substituted for the differential of z . By substituting for the acceleration from Eq.2 the result becomes:

$$\frac{d\Delta E}{dt} = m_0 g \cdot v \left(1 - \frac{1}{\sqrt{1 - v^2/c^2}}\right) \cong -\frac{1}{2} m_0 g \cdot \frac{v^3}{c^2} \quad (7)$$

By integrating this result in the velocity domain, similarly as it was shown in Eq.4, we obtain the expression for the energy difference loss:

$$\Delta E \cong -\frac{m_0 c^2}{8} \left(\frac{v}{c}\right)^4 \quad (8)$$

This is a very strange result. It seems that the falling body is gaining some additional energy on top of the energy that is predicted from the free fall by Eq.5. This is not reasonable and it is pointing to a problem that exists in the GRT for a long time. The gravitational mass cannot depend on velocity the same way as the inertial mass. This problem will become clear from the result presented next.

The similar expression introduced in Eq.1 is used, but with the gravitational mass depending on velocity as follows ^[1, 2]:

$$m_g = m_0 \sqrt{1 - v^2 / c^2} \quad (9)$$

This leads to the following formula:

$$\frac{d}{dt} \left[\frac{m_0 v}{\sqrt{1 - v^2 / c^2}} \right] = g \cdot m_0 \sqrt{1 - \frac{v^2}{c^2}} \quad (10)$$

After rearrangements the formula is simplified with the result as follows:

$$\frac{dv}{dt} = g(1 - v^2 / c^2)^2 \quad (11)$$

Following the same procedure as above for the GRT case the differential of energy will be:

$$dE = g \cdot m_0 \sqrt{1 - \frac{v^2}{c^2}} \frac{v \cdot dv}{g \left(1 - \frac{v^2}{c^2}\right)^2} \quad (12)$$

This becomes, after integration, identical to formula in Eq.5. Both theories, the GRT and MTG, thus give the same expression for the energy, which is expected and confirms once more that the calculating procedure is correct. For the energy loss the same procedure is also followed with the result:

$$\frac{d\Delta E}{dt} = m_0 g \cdot v - \frac{m_0 v}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{3}{2}}} \frac{dv}{dt} = m_0 g \cdot v \left(1 - \sqrt{1 - \frac{v^2}{c^2}}\right) \cong \frac{1}{2} m_0 g \frac{v^3}{c^2} \quad (13)$$

This is a similar result as in Eq.7 except that the energy loss differential is now positive as it should be. This confirms the correctness of the gravitational mass dependence on velocity and therefore disproves the validity of GRT. The energy loss is likely due to the gravitational radiation, because the falling body is accelerated. By integrating the result from Eq.13 the energy loss is equal to:

$$\Delta E \cong \frac{m_0 c^2}{8} \left(\frac{v}{c}\right)^4 \quad (14)$$

For a better understanding of the magnitude of radiated energy the graph of the energy loss as a function of the fall time for a mass of 100kg is shown in FIG.1.

Energy Conservation Test During a Test Body Free Fall in Uniform Gravitational Field g

$$g = 9.80665 \frac{\text{m}}{\text{s}^2} \quad c = 2.997925 \times 10^8 \frac{\text{m}}{\text{s}} \quad t := 0, 0.1..30 \quad \text{vnt}(t) := g \cdot t \quad m_0 := 100.0 \text{kg}$$

$$\Delta E(t) := \frac{m_0 c^2}{8} \cdot \left(\frac{g \cdot t}{c}\right)^4$$




FIG.1 the dependency of energy loss due to radiation for a 100kg test body free falling in a uniform gravitational field equal to Earth's gravity. The loss for a 30 sec fall is about 1.0 micro Joules.

Conclusions: The paper derived simple expressions for the energy loss during the small body free fall in a uniform gravitational field. It was shown that the loss derived according to the GRT is negative. This is unacceptable and this fact thus disproves the validity of GRT. This problem has its root cause in the identical dependency of inertial mass and gravitational mass on velocity. When the correct dependency of gravitational mass on velocity, as derived in the MTG, is used the correct positive energy loss is calculated.

This result has fatal consequences for the GRT, because unquestionably proves its incorrectness. This finding thus has a significant impact on all the theories based on the GRT such as the Big Bang and similar ridiculous models of the Universe.

The author hopes that the main stream relativists finally recognize this problem and abandon the GRT with all its ridiculous claims of existence of Black Holes, Event Horizons, and the Big Bang Universe with its accelerating expansion to infinity from nothing.

References:

- [1] <http://gsjournal.net/Science-Journals/Research%20Papers/View/7070>
- [2] <http://physicsessays.org/browse-journal-2/product/904-8-jaroslav-hynecek-remarks-on-the-equivalence-of-inertial-and-gravitational-masses-and-on-the-accuracy-of-einstein-s-theory-of-gravity.html>