

# **New coordinate vacuum solution in general relativity theory**

**Sangwha-Yi**

**Department of Math , Taejon University 300-716**

## **ABSTRACT**

In the general relativity theory, we discover new vacuum solution by Einstein's gravity field equation in general relativity theory. We investigate the new coordinate in general relativity theory.

**PACS Number:04,04.90.+e,98.80,98.80.E**

**Key words:General relativity theory,**

**Gravity field equation**

**New coordinate solution**

**e-mail address:sangwha1@nate.com**

**Tel:051-624-3953**

## 1. Introduction

We solve new vacuum solution by gravity field equation in general relativity theory.

New spherical coordinate is

$$d\tau^2 = dt^2 - \frac{1}{c^2} [dr^2 + V(t,r)\{d\theta^2 + \sin^2 \theta d\phi^2\}]$$

$$V(t,r) = C_1(act + br)^2, \quad C_1 = \frac{1}{b^2 - a^2}$$

$$a, b, C_1 \text{ is constant, } c \text{ is light's velocity.} \quad (1)$$

In this time, Einstein's gravity equation is

$$\begin{aligned} R_{tt} &= \frac{1}{2} \frac{\ddot{U}}{U} - \frac{\dot{U}^2}{4U^2} \\ &= \frac{2a^2}{(act + br)^2} - \frac{1}{2} \frac{4a^2}{(act + br)^2} = 0 \end{aligned} \quad (2)$$

$$\begin{aligned} R_{rr} &= \frac{V''}{V} - \frac{1}{2} \frac{V'^2}{V^2} \\ &= \frac{2b^2}{(act + br)^2} - \frac{1}{2} \frac{4b^2}{(act + br)^2} = 0 \end{aligned} \quad (3)$$

$$\begin{aligned} R_{\theta\theta} &= -\frac{\ddot{V}}{2} + \frac{V''}{2} - 1 \\ &= -C_1 a^2 + C_1 b^2 - 1 = 0 \end{aligned} \quad (4)$$

$$R_{\phi\phi} = \sin^2 \theta R_{\theta\theta} = 0 \quad (5)$$

$$\begin{aligned} R_{tr} &= \frac{\dot{V}'}{V} - \frac{\dot{V}V'}{2V^2} \\ &= \frac{2C_1 ab}{(act + br)^2} - \frac{1}{2} \frac{4C_1 ab}{(act + br)^2} = 0 \end{aligned} \quad (6)$$

In this time,

$$V' = 2C_1 b(act + br), \dot{V} = 2C_1 a(act + br), V'' = 2C_1 a^2, \ddot{V} = 2C_1 b^2$$

$$A' = \frac{\partial A}{\partial r}, \dot{A} = \frac{1}{c} \frac{\partial A}{\partial t}$$

## 2. New vacuum solution in general relativity theory

Hence, new vacuum solution is

$$d\tau^2 = dt^2 - \frac{1}{c^2} \left[ dr^2 + \frac{1}{b^2 - a^2} (act + br)^2 \{d\theta^2 + \sin^2 \theta d\phi^2\} \right]$$

$a, b, C_1$  are constant,  $C$  is light's velocity. (7)

In this time, if  $r'$  is

$$r' = \frac{1}{\sqrt{b^2 - a^2}} (act + br)$$

As

$$dr' = \frac{1}{\sqrt{b^2 - a^2}} (acdt + bdr)$$

Or

$$dr = \frac{\sqrt{b^2 - a^2}}{b} dr' - \frac{a}{b} cdt \quad (8)$$

If new solution Eq(7) is inserted by transformation Eq(8),

$$dr^2 = \frac{b^2 - a^2}{b^2} dr'^2 - 2 \frac{a}{b^2} \sqrt{b^2 - a^2} dr' cdt + \frac{a^2}{b^2} c^2 dt^2 \quad (9)$$

In this time, if  $\alpha_0$  is

$$\alpha_0 = \frac{a}{b} \quad (10)$$

Hence, proper time  $d\tau$  of new solution is

$$d\tau^2 = (1 - \alpha_0^2) dt^2 + 2\alpha_0 \sqrt{1 - \alpha_0^2} dr' \frac{dt}{c} - \frac{1}{c^2} [(1 - \alpha_0^2) dr'^2 + r'^2 \{d\theta^2 + \sin^2 \theta d\phi^2\}] \quad (11)$$

In this time, if  $dt'$  is

$$dt' = \sqrt{1 - \alpha_0^2} dt \quad (12)$$

Therefore, new solution is

$$d\tau^2 = dt'^2 + 2\alpha_0 dr' \frac{dt'}{c} - \frac{1}{c^2} [(1 - \alpha_0^2) dr'^2 + r'^2 \{d\theta^2 + \sin^2 \theta d\phi^2\}] \quad (13)$$

If we rewrite  $dt, dr$  instead of  $dt', dr'$ , the proper time  $d\tau$  of new solution is

$$d\tau^2 = dt^2 + 2\alpha_0 dr \frac{dt}{c} - \frac{1}{c^2} [(1 - \alpha_0^2) dr^2 + r^2 \{d\theta^2 + \sin^2 \theta d\phi^2\}] \quad (14)$$

### 3. Conclusion

Therefore, new spherical solution in general relativity theory is

$$d\tau^2 = dt^2 + 2\alpha_0 dr \frac{dt}{c} - \frac{1}{c^2} [(1 - \alpha_0^2) dr^2 + r^2 \{d\theta^2 + \sin^2 \theta d\phi^2\}]$$

$\alpha_0$  is constant

(15)

## Reference

- [1]S.Weinberg,Gravitation and Cosmology(John wiley & Sons,Inc,1972)
- [2]P.Bergman,Introduction to the Theory of Relativity(Dover Pub. Co.,Inc., New York,1976),Chapter V
- [3]C.Misner, K,Thorne and J. Wheeler, Gravitation(W.H.Freedman & Co.,1973)
- [4]S.Hawking and G. Ellis,The Large Scale Structure of Space-Time(Cam-bridge University Press,1973)
- [5]R.Adler,M.Bazin and M.Schiffer,Introduction to General Relativity(McGraw-Hill,Inc.,1965)
- [6]E.Kasner, Am. J. Math. 43, 217(1921)
- [7]G.Birkoff,Relativity and Modern Physics(Harvard University Press,1923),p.253
- [8]T.Kaluza, Berl. Ber. 996(1921); O. Klein, Z. Phys. 37, 895(1926)
- [9]Y. Cho, J. Math. Phys. 16, 2029(1975); Y. Cho and P. Freund, Phys. Rev. D12, 1711(1975)
- [10]P. van Nieuwenhuizen, Phys. Rep. 68. 189(1981)