

**Waves Generate Electrons  
And Both are Quantized into Phosons  
(New Definition of Relativistic Mass)**

Author: Yaseen Ali Mohammad Al Azzam  
Email : [vaseenalazzam000@gmail.com](mailto:vaseenalazzam000@gmail.com)

## Index

Abstract, P (2)

Chapter 1 The Phoson, P (3-5)

1.1 Introduction

1.2 What is the Phoson

Chapter 2 Particles at the Speed of Light, P (5-9)

2.1 Introduction

2.2 Relativistic Mass at the Speed of Light

2.3 Relativistic Mass at Speeds less than C

2.4 Realistic Relativistic Mass

Chapter 3 Phoson's Model P (9-13)

3.1 Introduction

3.2 Phoson's Model

3.3 Energy and Momentum

3.4 Force

Chapter 4 Electron Generation and Mass Transfer, P (14-16)

4.1 Electron Generation

4.2 Electron Motion

Conclusions, P (17-29)

References, P (20)

## Abstract

This paper is a study of the behavior of light waves from a purely particles' point of view.

I started with showing that waves consist of fundamental units of mass (I called phosons) which are identical in all waves, don't depend on any wave's parameter, have the same mass and carry the same energy.

By interpreting Compton's effect experiment from a different point of view, I deduced the mass of the photon and explained how waves and electrons are quantized into photons and how the electron's mass and energy are the summation of masses and energies of the photons comprising its mass.

Since my work contradicts with the theory of relativity, I found it mandatory to find an alternative which works at all speeds and give more logical results.

After finding the nature of the photon's mass with the new definition of the relativistic mass at the speed of light, everything became ready to propose a model to describe the photons behavior and propagation as a continuous energy transformation between two forms of kinetic energies and a continuous mass variation between two levels.

The model explains the actual meaning of  $mc^2$  and how even if we believe in mass energy equivalency, both are conserved individually.

At last I proposed how electrons are generated by waves' and how these photons shape the electron.

## Chapter 1 The Phoson

### 1.1 Introduction

This discussion assumes that any beam of light consists of rays where each ray is a stream of particles (I named phosons).

The phoson which will be introduced in this chapter is the fundamental unit of energy carrying mass where it carries always one  $h$  (J.s).

All waves consist of discrete identical phosons which do not change its energy or mass with frequency or any other wave's parameter.

After defining the phoson, it will be obvious how waves and electrons are quantized into phosons.

The name phosons I gave to the waves' particles stands for (photon son)


### 1.2 what is the Phoson

This section is to define the phoson and calculate its mass using the data we have about Compton effect experiment.

Compton famous equation for the change wave length is (figure 1.1)

$$\Delta\lambda = \frac{h}{m.c} (1 - \cos\theta) \text{ where } m \text{ is the electron's mass.}$$

This experiment was explained as a collision and scattering physical event using the principle of energy and momentum conservation to prove that light consists of particles which can scatter the wave and eject electrons.



The part ( $\Delta\lambda = \frac{h}{m.c}$ ) in the equation above consists of constants and represents a full value of  $\Delta\lambda$  when ignoring the fraction caused by the other part of the equation.

If each phoson occupies one wave length, then the frequency of the wave corresponds to the number of phosons in one second of the wave's ray.

Accordingly, an absence of phosons represents an increase in wave length and a decrease in frequency proportional to the number of missing phosons.

The results of Compton experiment gave two peaks of scattered waves, one for the part of the wave which is scattered directly without interaction and one for the part of the wave after losing some of its phosons in the interaction at specific scattering angles.

The second peak at  $90^\circ$  and  $180^\circ$  scattering angles corresponds to a full Compton wave length and consequently a full interaction.

Thus, the interactions in this experiment are one of four types, the first is scattering without wave length alteration where phosons are not involved in the interaction, the second with increased wave length which is a fraction of  $\lambda_c$  where the wave loses part of its phosons in a partial

interaction, the third and fourth at scattering angles  $90^\circ$  and  $180^\circ$  which represents a full interaction where the wave length increment equals to  $\lambda_c$  or twice  $\lambda_c$ .

The interpretation of the latter two cases can have other possibilities than what Compton gave. The first is the possibility to have a newly generated electron and the second is a full interaction when both the wave and electron are composed of the same number of identical particles, a full one to one interaction will give the number of photons in the electron and consequently the number of photons involved from the wave.

The increase in wave length represents a decrease of frequency which corresponds to the number of reduced photons contributed in producing new electrons in a full interaction, Compton frequency denoted by  $f_c$  can be defined as:

*$f_c$  is the number of missing photons in the scattered wave when the increment in wave length is equal to  $(h/m.c)$  which contributed in generating a new electron or involved in a full interaction.*

If the interaction is one to one, then the number of photons in the electron and the number of photons the wave lost are the same and if the ejected electron is a newly generated one, then the mass of the electron is equal to summation of the masses of the involved photons.

But the number of photons involved equals to the increase in frequency of the scattered wave

$$f_c = c / \lambda_c$$

$$f_c = \frac{m.c^2}{h}$$

$$f_c = 1.235589965 \times 10^{20} \text{ Hz}$$

$$m_{phs} = m/f_c \quad 1.1$$

$$m_{phs} = 7.372497201 \times 10^{-51} \text{ Kg. s} \quad 1.2$$

Using the famous equation ( $E = m.c^2$ ) we can find the energy and mass of the photon easily in an equivalent way where

$$E = h \quad 1.3$$

$$m_{phs} = h/c^2 \quad 1.4$$

*Therefore, the electron is generated by  $f_c$  number of photons and if this electron is emitted fully as a wave (not ejected as an electron) will produce a wave of frequency  $f_c$  and a wave length  $\lambda_c$ .*

*It should be noted that  $f_c$  can be a frequency with units ( $s^{-1}$ ) or it can be just a figure representing the number of photons in the electron.*

*Consequently, this implies that waves and electrons are quantized into photons and waves are just discrete identical photons.*

In the photoelectric effect experiment, it is our measurement units which are quantized into values per second not the waves and saying that waves are quantized into photons of energy  $E = hf$  is just as saying that nature follows our manmade measurement units.

Thus, the photon is just a group of photons involved in an electron generation (or in an interaction).

## Chapter 2 – Particles at the Speed of Light

### 2.1 Introduction

As we have seen in the previous chapter, waves' particles have mass and can generate electrons which contradicts with theory of relativity, but attention should be paid to that any existing particle which can carry energy should have a mass regardless of what type of mass it has or our capability to measure or detect it.

Since experiments proved that waves' particles have momentum which is an exclusive property of mass, it is essential to find what are the characteristics of this mass.

All the coming discussion will be to show that light particles do have invariant mass and a determined mass at the speed of light.

I will use a different approach in this section to describe the behavior of mass particles travelling at the speed of light.

### 2.2 Relativistic Mass at the Speed of Light.

Referring to figure 2.1, if an external force  $F$  is applied on a particle travelling at the speed of light to cross a distance  $S$ , then the change in kinetic energy can be found from

$$\Delta k = \Delta W = F \cdot \Delta s$$

where  $F$  is force and  $W$  is the work done in the distance  $S$ .




Figure 2.1

$$F = \partial P / \partial t = \partial / \partial t (mv)$$

$$F = m \partial v / \partial t + v \partial m / \partial t \quad 2.1$$

But at the speed of light, acceleration is zero i.e.  $\partial v = 0$ , and  $\partial s / \partial t = v = c$  then

$$\Delta k = c \partial m / \partial t \cdot \Delta s$$


$$\Delta k = c (\partial s / \partial t) \cdot \Delta m.$$

$$\Delta k = c^2 \cdot \Delta m.$$

$$\Delta k = \Delta m \cdot c^2 \quad 2.2$$

Referring to figure 2.2, if a particle is travelling at the speed of light from point A to point B where its kinetic energy and mass at point A are  $k_0$  and  $m_0$  and its kinetic energy and mass at point B are  $k$  and  $m$  respectively then, the total energy at points A and B are equal because there is no external source of energy or force

$$\frac{1}{2}mc^2 = \frac{1}{2}m_0c^2 + Ep.$$



Where  $Ep.$  is additional energy carried by the particle at point A in another form of energy which works as a potential energy.

$$\frac{1}{2}mc^2 - \frac{1}{2}m_0c^2 = Ep. = \Delta k \quad 2.3$$

If we define  $Ep.$  in a translational kinetic energy scale to be equivalent to the energy required to accelerate the particle from rest to speed  $v$  (maximum value of  $v = c$ ) with constant mass  $m_0$  or to accelerate the particle from speed  $c$  to  $(c + v)$  with constant mass  $m_0$  (impossible case), then

$$Ep = \frac{1}{2}m_0v^2$$

Substituting in equation 2.3 we get

$$\frac{1}{2}mc^2 - \frac{1}{2}m_0c^2 = \frac{1}{2}m_0v^2 \quad 2.4$$

$$mc^2 - m_0c^2 = m_0v^2 \quad 2.5$$

$$m = m_0 \left(1 + \frac{v^2}{c^2}\right) \quad 2.6$$

Since the maximum value of  $v$  is  $C$ , then substituting  $C$  for  $v$  in equation 2.6 gives

$$m = 2m_0 \quad 2.7$$

also, equation 2.4 with  $v$  equals to its peak value  $C$  is given as

$$\Delta k = \frac{1}{2}m_0c^2 \quad 2.8$$

$$\frac{\Delta m}{m_0} = \frac{v^2}{c^2} \quad 2.9$$

If  $Ep = \frac{1}{2}m_0c^2$  at point A in its peak, the total carried energy is

$$k = k_0 + Ep = \frac{1}{2}m_0c^2 + \frac{1}{2}m_0c^2 = m_0c^2 \quad 2.10$$

And at point B with  $m = 2m_0$ , the total carried energy is

$$E = \frac{1}{2}(2m_0)c^2 = m_0c^2 \quad 2.11$$

Where all the energy  $Ep$  is converted to translational kinetic energy.

The first case is theoretical because waves' particles do not need an external force to propagate while the second case is a description of the behavior of these waves' particles

If the particle's translational kinetic energy is  $\frac{1}{2}m_0c^2$ , then it can carry another  $\frac{1}{2}m_0c^2$  as a maximum in another form of energy.

While the particle travels at the speed of light, it tends to resist motion by increasing its mass and converting the potential kinetic energy to translational kinetic energy until all this energy is consumed to reach to a translational kinetic energy equal to  $m_0 c^2$ .

Equation 2.1 states that the change in kinetic energy is due to the change in velocity and mass, the question which will be answered in the next section is whether both mass and speed can increase together in the same time.

### 2.3 Relativistic Mass at Speeds less than C.

To discuss what happens to mass at speeds below the speed of light, I will start with equation 2.1 in full to find the change in kinetic energy

$$F = m \frac{\partial v}{\partial t} + v \frac{\partial m}{\partial t}$$

$$\partial k = \partial s \cdot m \frac{\partial v}{\partial t} + \partial s \cdot v \frac{\partial m}{\partial t} \quad (\text{but } \partial s/\partial t = v \text{ then})$$

$$\partial k = mv\partial v + v^2\partial m \quad 2.12$$

Also  $\gamma$  is expressed as

$$m = m_0 / \sqrt{1 - \frac{v^2}{c^2}} \quad 2.13$$

$$m^2 = m_0^2 / (1 - v^2/c^2)$$

$$m^2 c^2 - m^2 v^2 = m_0^2 c^2 \quad 2.14$$

$$2mc^2\partial m - 2mv^2\partial m - 2m^2v\partial v = 0 \quad (\text{deriving equation 2.14}) \quad 2.15$$

$$c^2\partial m = v^2\partial m + mv\partial v \quad (\text{Dividing equation 2.15 by } 2m) \quad 2.16$$

Comparing equation 2.16 with 2.12 we get

$$\partial k = c^2\partial m \quad 2.17$$

If we try to integrate equation 2.16 going back to equation 2.14 we get the following:

Multiply equation 2.16 by  $2m$  we get

$$2mc^2\partial m = 2mv^2\partial m + 2m^2v\partial v \quad 2.18$$

Integrate equation 2.18 with  $m$  ranges from  $m_0$  to  $m$  and  $v$  ranges from 0 to  $v$ , we get

$$\int_{m_0}^m 2mc^2\partial m = \int_{m_0}^m 2mv^2\partial m + \int_0^v 2m^2v\partial v \quad 2.19$$

$$m^2 c^2 - m_0^2 c^2 = m^2 v^2 - m_0^2 v^2 + m^2 v^2 \quad 2.20$$

It is obvious that equation 2.14 can't be recovered from equation 2.20 unless we consider  $m$  as constant equal to  $m_0$  in the second term of equation 2.14 which leads us to equation 2.6 but with squared masses.

If we integrate equation 2.16 directly we get

$$\int_{m_0}^m c^2 \partial m = \int_{m_0}^m v^2 \partial m + \int_0^v mv \partial v$$

$$c^2(m - m_0) = v^2(m - m_0) + \frac{1}{2} m(v^2 - 0)$$

$$c^2(m - m_0) = c^2(m - m_0) + \frac{1}{2} mc v^2$$

$c^2(m - m_0)$  appears in both sides which implies that the particle either remained at rest or has a zero-relative mass.

Thus,  $\gamma$  is not suitable to describe the relativistic mass because it compressed two ranges of two separate effects and overlapped it in one range.

From section 2.2 and the above discussion we can conclude that in equation 2.12 which is  $\partial k = mv \partial v + v^2 \partial m$ , the plus sign works as an (OR) rather than addition.

The kinetic energy and consequently the momentum increases either with increasing velocity for speeds below the speed of light or by increasing mass at the speed of light but not both in the same time.

Thus, if speed can be increased for speeds below the speed of light, the accelerated object will maintain its rest mass without any amplification.

At the constant speed of light, mass increases to maintain the proportionality with the translational kinetic energy.

Accelerated objects should be destructed to molecules or compounds then to atoms then subatomic particles to have the possibility to be accelerated to the speed of light where no longer the original object exists.

#### 2.4. Realistic Relativistic Mass

At speeds below the speed of light the mass is constant and equal to the rest mass  $m_0$

$$F = m_0(\partial v / \partial t)$$

$$\partial k = F \cdot \partial s$$

$$\partial k = m_0 \cdot v \cdot \partial v \quad 2.21$$

And the new relativistic mass equation is

$$m = m_0(1 + v^2/c^2)$$

$$mc^2 = m_0c^2 + m_0v^2 \text{ (deriving this equation)}$$

$$c^2 \partial m = 2m_0 v \partial v \quad 2.22$$

comparing equations 2.21 and 2.22 we get

$$c^2 \partial m = 2 \partial k$$

If mass changed from  $m_0$  to  $m$  speed from zero to  $v$ , then integration we give

$$c^2(m-m_0) = 2(\frac{1}{2} m_0 v^2 - 0)$$

$$mc^2 - m_0 c^2 = m_0 v^2$$

$$\frac{1}{2} mc^2 - \frac{1}{2} m_0 c^2 = \frac{1}{2} m_0 v^2$$

*This equivalency means that the change in kinetic energy caused by increasing the mass of a particle travelling at the speed of light is equivalent to the change in kinetic energy caused by increasing the velocity of the same particle when accelerated from rest to a specific speed v.*

*As a conclusion, any object travelling with a speed below the speed of light will not experience any change in mass but for particle travelling at the speed of light mass varies in proportion to the translational kinetic energy.*

## Chapter 3 – The Phoson's Model

### 3.1 Introduction to Phoson's Model

The following points are fundamental in this section:

- At the speed of light, the source of mass increase is not the energy involved, mass and energy are conserved separately.
- Phosons mass works as the energy carrier.
- Each Phoson carries  $h$  (J.s) energy and have mass  $m_{phs}$  (Kg. s) where both are constants and do not vary in normal conditions with frequency or other wave's parameters.

Figure 3.1 shows a sketch of the proposed behavior of the phoson while travelling in a wave. It has two peak states, state 1 with mass  $m_0$  at point A and state 2 with mass  $m$  at point B.

### 3.2 Phoson's Model

In figure 3.1, I assumed that the phoson is a ring of mass which collects and releases mass during travelling each wave length and has one of two states as shown in the figure.

In state 1: The phoson has minimum mass, minimum translational kinetic energy and maximum spinning kinetic energy.

In state 2: The phoson has maximum mass, maximum translational kinetic energy and zero spinning kinetic energy.

The phoson goes from state 1 to state 2 in half wave length and back to state 1 in the other half.




Figure 3.1

Usually each wave length is occupied by one photon but here different stages of one photon is shown in one wave length travel for clarity.

While traveling from point A to point B the photon translational kinetic energy  $K$  increases to maximum and its spinning kinetic energy  $S$  reduces to zero maintaining a total kinetic energy equal to  $(h)$  (J.s) always, its mass always follows the rate of change of its translational kinetic energy.

$$K + S = h \quad 3.1$$

If this equation is derived it gives that the rate of increase of any of these two energies equals to the rate of decrease of the other.

$$\partial K / \partial t = - \partial S / \partial t \quad 3.2$$

At point A, the photon tends to work against its translational kinetic energy by increasing its mass with constant velocity  $C$ , the increase in translational kinetic energy is supplied by the spinning kinetic energy until it is consumed fully at point B.

At point B, the photon's energy is fully translational which is an unstable state equivalent to a nonexistence state of the photon, so it starts to decrease its translational kinetic energy again and reduce its mass to suit this decrease with restoring its spinning energy back.

**At point A** the photon is in state 1 and has a ring shape because of its high spinning, its moment of inertia is  $I = m \cdot r^2$ .

The photon mass here is  $m_o = 7.372497201 \times 10^{-51}$  Kg. s

The translational kinetic energy  $K$  and the spinning kinetic energy  $S$  are both in (J.s) and equal to

$$K = \frac{1}{2} m_o \cdot c^2 \quad 3.3$$

$$S = \frac{1}{2} I \cdot \omega^2 = \frac{1}{2} m_o \cdot r^2 \cdot \omega^2 \quad 3.4$$

At this state, the two energies are equal because  $(r \cdot \omega = c)$  which occurs at

$(0, 2\pi, 4\pi, 6\pi \dots$  in figure 3.2)

$$E_T = h = K + S = \frac{1}{2} m_o c^2 + \frac{1}{2} m_o c^2$$

$$E_T = h = m_o \cdot c^2 \quad 3.5$$

**At point B** the photon is in state 2 and its energy is a fully translational kinetic energy given by

$$E_T = K = \frac{1}{2} m \cdot c^2$$

$$E_T = \frac{1}{2} (2 \cdot m_o) \cdot c^2 \quad (m=2m_o)$$

$$E_T = h = m_o \cdot c^2 \quad 3.6$$

These points can be seen in figure 3.2 at  $(\pi, 3\pi, 5\pi \dots)$

So, while the photon is moving a full wave length its angular speed  $\omega$  reduces from its maximum value to zero in one half wave length and back to maximum in the other half.

### From point A to Point B

The energy stored as spinning energy is converted to translational kinetic energy during motion and mass increases to suit the new translational kinetic energy with constant speed.

At point B, the photon's energy is completely translational kinetic energy but it can't keep this situation without spinning, so it starts again to spin and go back to a new point A, then repeat this process each wave length.




Figure 3.2

During motion, the photon maintains a constant translational kinetic energy equal to ( $h/2$ ) beside the other ( $h/2$ ) exchanged with the spinning energy.

The motion of the photon during its travel from point A to point B can be described by

$$\frac{1}{2}m \cdot c^2 + \frac{1}{2}I \cdot \omega^2 = h \quad 3.7$$

$$h = \frac{1}{2}m \cdot c^2 \left( 1 + \frac{r^2 \omega^2}{c^2} \right) \quad 3.8$$

$$\frac{h}{c^2} = \frac{1}{2} \cdot m \left( 1 + \frac{r^2 \omega^2}{c^2} \right) \quad 3.9$$

Knowing that  $\frac{h}{c^2} = m_0$  and  $m$  in the above equations is the relativistic mass at any time between point A and B

$$A_m = \frac{m}{m_0} = \frac{2}{1 + \frac{r^2 \omega^2}{c^2}} \quad 3.10$$

Where  $A_m$  is the mass amplification and here the mass is doubled at point B.

In trigonometric forms, we can express both energies as

$$K = h/4 \{ \cos((kx - \omega t) - \pi) + 3 \} = h/4 \{ 3 - \cos(kx - \omega t) \} \quad 3.11$$

$$S = h/4 \{ \cos(kx - \omega t) + 1 \} \quad 3.12$$

For waves  $m = m_{phs}$ , and thus

$$f \cdot \lambda = c = (m \cdot c^2) / (m \cdot c)$$

$$\lambda = h / p f = c / f$$

$$\lambda = \frac{h}{p f} \quad 3.13$$

$$k = 2\pi / \lambda$$

$$k = \omega \cdot \frac{p}{h} = f \cdot \frac{p}{\hbar} \quad 3.14$$

### 3.3 Energy and Momentum.

If we look at point 1 and 2 in figure 3.1, the total energy at both points is

$$\frac{1}{2} m.c^2 + \frac{1}{2} I_1 \cdot \omega_1 = \frac{1}{2} (m + \Delta m).c^2 + \frac{1}{2} I_2 \cdot \omega_2$$

$$\frac{1}{2} (\Delta m \cdot c^2) = \frac{1}{2} I_1 \cdot \omega_1 - \frac{1}{2} I_2 \cdot \omega_2$$

$$\Delta S = \frac{1}{2} \Delta m \cdot c^2 \quad 3.15$$

But  $\Delta S = - \Delta K$  then

$$\Delta K = - \frac{1}{2} \Delta m \cdot c^2 \quad 3.16$$

If the total change in mass  $\Delta m = m_o$  in half wave length, then

$$\Delta S = \frac{1}{2} m_o \cdot c^2 \quad 3.17$$

$$\Delta K = - \frac{1}{2} m_o \cdot c^2 \quad 3.18$$

Accordingly, the total rate of change of energy between any two points is equal to zero

$$\Delta E_T = 0.0 \quad 3.19$$

Since the change in spinning and translational momentum is generated by the change in mass then similarly we can find that the change in translational momentum P is

$$\Delta P = m_o \cdot c \quad 3.20$$

### 3.4 Force

With the phason's mass increment to double in one half wave length, it generates a force which causes the translational momentum to increase in the same rate.

Usually external forces make a change in velocity and consequently a change in momentum with constant mass.

In the phason's case, the variable is the mass with constant speed, this mass variation produces a force which generates momentum.

The force produced by one phason is

$$F = m \cdot a \left( Kg \cdot s \cdot \frac{m}{s^2} \right) \text{ this unit is equivalent to } \left( \frac{Kg \cdot s}{s} \cdot \frac{m}{s} \right) \text{ and we can rewrite the equation as}$$

$$(F = \frac{\Delta m}{\Delta t} \cdot c) \text{ where } c \text{ is constant, and I will call } \frac{\Delta m}{\Delta t} \text{ as } M_m \text{ the rate of change of mass in (Kg. s / s).}$$

$$F = M_m \cdot c \quad (N.s) \quad 3.21$$

We can find this equation also by deriving  $p = m.v$  with  $\partial v=0$ .

So, while the phoson travels half wave length the mass increment generates a hammering force given by equation 3.21 and

$$M_m = (m - m_0) / t \quad 3.22$$

$$m = m_0 + M_m \cdot t \quad 3.23$$

$$M_m = \Delta m / t = 2 m_0 \cdot f \quad (t = T/2 \text{ where } T \text{ is the wave's period}) \quad 3.24$$

$$F = 2 \cdot m_0 \cdot f \cdot c \quad (\text{From equation 3.21})$$

$$F = 2 \cdot P_0 \cdot f \quad 3.25$$

$$F = P \cdot f \quad (\text{Since } P = 2p_0) \quad 3.26$$

To find the energy

$$E = F \cdot x \text{ with } x = \lambda/2 \text{ we get}$$

$$E = P \cdot f \cdot \lambda/2$$

$$E = m \cdot c \cdot f \cdot \lambda / 2$$

$$E = \frac{1}{2} mc^2$$

$$E = m_0 c^2 \quad (m = 2m_0)$$

## Chapter 4 – Electron Generation and Mass Transfer

### 4.1 Electron Generation

In certain conditions when phasons with proper frequency and direction inter the atom's field, it forms a new electron. The rays of phasons are bent over to orbits such that each orbit circumference is  $\alpha\lambda_c$  and each orbit is occupied by one phason.

Figure 4.1 shows one phason forming the orbit ( $\alpha^{-1} = 137.03587$  is the fine structure constant reciprocal).




Figure 4.1

Figure 4.2 shows how  $f$  number of phasons are bent over to circular orbits with maintaining its spinning and translational motion with the speed of light to form a string of phasons spinning around an axis.

Each orbit circumference is one wave length of the incident wave reduced by the  $\alpha$  factor.

$$2\alpha r_e = \lambda_c \cdot \alpha \quad (r_e \text{ is the electron's radius})$$




Figure 4.2

Also because of the force applied by the nucleus magnetic field, the orbits are stacked to look like a cylinder of rings.

*The electron's mass and energy are those complying with the equation  $E=mc^2$  and are the summations of masses and energies of the photons comprising it.*

As an example, if an electron of mass  $m$  consists of  $f_c$  photons, each photon has a translational kinetic energy equal to  $h/2$  and spinning with an energy equal to  $h/2$ , then the rest mass and rest energy of the electron are

$$m_{rest} = f_c \cdot m_{phs} \quad 4.1$$

$$E_{rest} = f_c \cdot (h/2 + h/2) = f_c \cdot h \quad 4.2$$

*The total translational kinetic energies of all the photons comprising the electron equals to the spinning motion energy of the whole electron mass given by*

$$S = f_c \cdot h/2 \quad 4.3$$

$$S_e = f_c \cdot h/2 = \frac{1}{2} f_c \cdot m_{phs} \cdot c^2 = \frac{1}{2} m_e \cdot c^2 \quad 4.4$$

This  $S_e$  is half the rest energy of the electron and after shrinkage each ring circumference becomes

$$2\pi r_e = 1.770538 \times 10^{-14} \text{ m}$$

When  $f_c$  number of photons in the wave front enters the atom's energy field, it is supposed to have an angular speed and radius complying with ( $c = \omega \cdot r$ ) such that one wavelength  $\lambda_c$  is rounded to a circle with one photon rotating at the speed of light

$$t = \lambda_c/c$$

$$f = c/\lambda_c$$

$$\omega = 2\pi c/\lambda_c$$

$$\omega_c = 2\pi f_c = 7.763441 \times 10^{20} \text{ rad/s}$$

$$r_c = 3.8616 \times 10^{-13} \text{ m}$$

Turning one wavelength to circular shape gives a radius equals to  $\lambda_c/2\pi$  and consequently an angular speed  $\omega_c$  but the shrinkage reduces the radius and accelerates the angular speed to

$$r_e = r_c / \alpha^{-1} = 2.8179 \times 10^{-15} \text{ m}$$

$$\omega = \alpha^{-1} \cdot \omega_c = 1.06389 \times 10^{23} \text{ rad/s}$$

When the electron's photons are emitted as a wave, it will have a wavelength equal to one orbit circumference multiplied by the fine structure constant factor  $\alpha^{-1}$ .

$$\lambda_c = 2\pi r \alpha^{-1}$$

If a wave's ray happens to fall on an electron in a proper direction and its frequency is high enough to provide the required photons in a time equal or shorter than the time required by the electron to escape from the interaction area because of its orbital motion, the wave's photons will replace the electron's photons and the original photons will be ejected as an electron. The same condition may happen if the intensity of the wave is enough to replace the electron's photons in the electron's availability time during motion.

Because the speed of rotating of the photons is much faster than the electron motion caused by the nucleus, it takes the shape of moving rings in the direction of motion of the electron, but if we slow down this view, the electron shape will look like a helical shape figure 4.3




Figure 4.3

## 4.2 Electron's Motion.

In figure 4.5 if the generated electron has a pitch  $P = \lambda_p$ , and speed  $v$ , then the time required by one photon to travel across the pitch horizontally in parallel to the axis of rotation is equal to

$$t = \lambda_p/v \quad 4.5$$

The same time  $t$  is required to travel one ring orbit following the motion route and is given by

$$t = \lambda_c/c \quad 4.6$$

equalizing equation 4.4 and 4.5 we get

$$\begin{aligned} \lambda_p/v &= \lambda_c/c \\ \lambda p &= \lambda_c (v/c) \quad (v = \alpha c) \end{aligned} \quad 4.7$$

$$\lambda p = \alpha \cdot \lambda_c \quad 4.8$$

This means that  $\lambda_p$  is constant if  $\alpha$  is constant, that's why we should see what is  $\alpha$ .

As discussed earlier, the electron spinning energy is the summation of the translational energies of the photons comprising it.

Figure 4.4 shows one photon in its orbit which is inclined by an angle  $\theta$  from the x-axis because of the helical route it follows and running with the speed of light  $c$ .

Its tangential speed  $C$  has two components, one in the direction of motion of the electron (the y-axis in the graph) and equal to  $\alpha c$  and the other is perpendicular to the motion of the electron.

If the speed in the x direction is  $V_x$  then

$$V_x^2 = c^2 - \alpha^2 c^2$$

$$\alpha^2 c^2 = c^2 - V_x^2$$




Figure 4.4

$$\alpha^2 = 1 - v_x^2/c^2$$

$$\alpha = \sqrt{1 - \frac{v_x^2}{c^2}} \quad 4.9$$

If the speed of the photons in the y-axis direction which is the electron motion direction is  $v = \alpha c$  then substitute for  $v_x$  we get

$$\alpha = \sqrt{1 - \frac{(c^2 - v^2)}{c^2}}$$

$$\alpha = v/c$$

this shows that the inclination which came from the helical shape determines the value of  $\alpha$  and that  $\alpha$  is proportional to the speed of the electron.

Thus, as the velocity is increased up to the speed of light, the electron will elongate until it becomes a string of photons acting as a wave.

The kinetic energy at ( $v = \alpha c$ ) is

$$K = \frac{1}{2} m (\alpha c)^2 \approx 13.6 \text{ eV}$$

which equals to the atom's energy level number 1 or Bohr's hydrogen atom electron's energy level.




Figure 4.5

## Conclusions

1. Waves and electrons are quantized into discrete identical fundamental energy carrying mass particles (which I called photons).
2. Each has a mass equal to  $7.372497201 \times 10^{-51} \text{ Kg. s}$  and carry energy equal to  $h$  ( $6.626 \times 10^{-34} \text{ J.s}$ ).
3. Waves can generate electrons where the generated electron's mass and energy are

$$m_e = f_c \cdot m_{phs}$$

$$E_e = f_c \cdot h$$

$$m_{phs} = h / c^2 = m_e / f_c$$

where  $f_c$  is Compton frequency and  $m_{phs}$  is the photon mass, this number of photons is applicable to free electrons, electrons in a one electron atom, ready to leave the atom electrons or generated electrons.

4. Phoson,  $h$  or  $mc^2$  are faces of the same particle, any object which complies with  $E = mc^2$  should be comprised of phasons.
5. If a generated electron is emitted fully as a wave it will produce a wave with the same characteristic of the wave which generated it.
6. Particles travelling at speeds below the speed of light do not experience a change in mass, but this change happen exclusively at the speed of light.
7. The potential energy carried by particles travelling at the speed of light works as the external force at speeds below the speed of light.
8. At the speed of light, the potential energy increases the translational kinetic energy by increasing the mass while at speeds below the speed of light the translational kinetic energy is increased by increasing the velocity.
9. For waves the potential energy is an additional energy to the translational kinetic energy and is carried in another form of energy like spinning.
10. Particles at the speed of light can have a maximum translational kinetic energy  $k = m_0 c^2$
11. As a general form, the change in translational kinetic energy at the speed of light is

$$\frac{1}{2} mc^2 - \frac{1}{2} m_0 c^2 = \frac{1}{2} m_0 v^2$$

The change in mass at the speed of light follows the equation

$$m = m_0 (1 + v^2/c^2)$$

For light and electromagnetic waves, the equation becomes

$$\frac{1}{2} mc^2 - \frac{1}{2} m_0 c^2 = \frac{1}{2} m_0 c^2$$

$$\Delta k = \frac{1}{2} m_0 c^2$$

$$\Delta m = m_0$$

$$m = 2m_0$$

12. The proposed a model for phasons based on the following:

- a) One phoron occupies one wave length and has two peak states, one with minimum translational kinetic energy, maximum angular kinetic energy and minimum mass and the other is with maximum translational kinetic energy, zero angular kinetic energy and maximum mass.
- b) The phoron travels between the two states in half wave length.
- c) In the second state the energy is purely translational, and the mass is doubled.
- d) The summation of the translational and angular kinetic energies is equal to  $h$  always.

$E_T = h = m_0 c^2 = K + S$  where  $E_T$  is the total energy,  $K$  and  $S$  are the translational and angular kinetic energies respectively.

- e) The rate of change of each of the energies is the opposite of the other making a total of zero

$$\partial K/\partial t + \partial S/\partial t = 0$$

- f) At the state with minimum mass  $c = (\omega.r)$ ,  $k = h/2$  and  $S = h/2$

- g) The mass amplification between the two states is given by

$$A_m = m / m_0 = \frac{2}{1 + \frac{r^2 \omega^2}{c^2}}$$

- h) S and K energies can be expressed in trigonometric forms as

$$K = h/4 \{ \cos((kx - \omega t) - \pi) + 3 \} = h/4 \{ 3 - \cos(kx - \omega t) \}$$

$$S = h/4 \{ \cos(kx - \omega t) + 1 \}$$

- i) In terms of phonon's momentum and wave frequency, the wave length can be expressed by the equation

$$\lambda = \frac{h}{pf}$$

And the wave number is

$$k = \omega \cdot \frac{p}{h} = f \cdot \frac{p}{\hbar}$$

- j) knowing that the mass change is  $m_0$  we get the change in energies as

$$\Delta S = \frac{1}{2} m_0 \cdot c^2$$

$$\Delta K = -\frac{1}{2} m_0 \cdot c^2$$

- k) During motion, the phonon generates a force expressed as

$F = P \cdot f$  where  $p$  is the phonon's momentum in its maximum mass state and  $f$  is the wave frequency.

13. When an electron is generated by a wave, the phonon rays of the wave are bent over to form circular orbits with one phonon per orbit and the orbit radius and angular velocity are given by

$$r_e = r_c / \alpha^{-1} = 2.8179 \times 10^{-15} \text{ m}$$

$$\omega = \alpha^{-1} \cdot \omega_c = 1.06389 \times 10^{23} \text{ rad/s}$$

14. The generated electron will be a string of phonons moving in ring orbits around a common axis and when the electron moves it takes a helical shape.

15. The phonons comprising the electron are in a state where each carries a translational kinetic energy  $h/2$  and an angular kinetic energy  $h/2$

16. The summation of the translational kinetic energies of the photons comprising the electron acts as the electron's spinning energy.

$$S_e = f_c \cdot (h/2) = \frac{1}{2} m_e \cdot c^2$$

17. The summation of the translational and angular energies of the photons comprising the electron is equal to its rest energy

$$E_{rest} = f_c \cdot (h/2 + h/2) = f_c \cdot h = mc^2$$

18. A moving electron will elongate in proportion to its velocity  $v$ , at the speed of light it becomes a string of photons travelling as a wave, the value of one-unit of elongation is

$$\lambda_p = \alpha \cdot \lambda_c (v/c)$$

where  $\lambda_p$  is the pitch between any two successive photons and  $\lambda_c$  is Compton wave length.

19. The inclination which came from the helical shape determines the value of  $\alpha$  such that  $\alpha$  is proportional to the speed of the electron given by

$$\alpha = \sqrt{1 - \frac{v_x^2}{c^2}}$$

$$\alpha = (v/c)$$

Where  $v$  is the electron velocity which is the component of the tangential speed of the photon in the direction of motion of the electron.

And  $v_x$  is the component of the tangential speed of the rotating photon (equals to  $c$ ) in a direction perpendicular to the electrons motion.

#### References -----

- 1- Elementary Quantum Mechanics, David S. Saxon, McGraw - Hill book co,1968, PP 1 – 12,25
- 2- Modern Physics, Hans C. Ohanian, Prentice-Hall Inc., New Jersey, 1987, PP.46, PP.65-99.
- 3- Elements of Quantum Mechanics, Binayak Dutta-Roy, New Age Science Limited, UK, 2010, PP 1-17, 82,187.
- 4- Physics for Scientists and Engineers with Modern Physics- Third edition, Raymond A. Serway, Saunders college publishing- Philadelphia, 1990, pages 1102-1136, page 1147-1164, PP. 1170-1176, ,1150-1153.
- 5- A first course in quantum mechanics, Clark, H., Nostrand Reinhold co ltd, New York 1982. PP 20-42, 54-55,236-245.
- 6- Fundamentals of Physics – Second Edition, David Halliday & Robert Resnick, John Wiley & Sons-New York, pages 169-173, PP 178-193, 773-792,798-810.