



32 The paper is organized as follows. In [section 2](#) we define terms that will be  
 33 used in the rest of the paper. These terms are necessary to state our main results.  
 34 In [section 3](#), we state our 4 main theorems. [section 4](#) provides a proof of the first  
 35 theorem, [section 5](#) provides a proof of the second, [section 6](#) provides a proof of the  
 36 third and [section 7](#) provides a proof of the fourth. In [section 8](#), we state 2 conjectures.  
 37 In [section 9](#), we conclude.

## 38 2. Terms and definitions.

### 39 2.1. Boundary matrix.

- 40 • Given a point  $P$  on  $\partial\Delta$  (the boundary of  $\Delta$ ) and given a unitary matrix  $U$   
 41 such that  $R_m(U) = P$ , we call  $U$  a **boundary matrix** of  $\Delta$ . See [\(1.3\)](#).
- 42 • Given a boundary matrix  $U$ . If  $\partial\Delta$  is smooth at  $R_m(U)$  and  $U$  is not the  
 43 product of a unitary diagonal matrix and a permutation matrix, we say  $U$  is  
 44 a **regular boundary matrix**.

45 **2.2. Properties of unitary matrices given  $A_0$  and  $B_0$ .** In this section, we  
 46 define four properties of unitary matrices that will be very useful when examining  
 47 boundary matrices of  $\Delta$ .

48 The first three of these properties are matrices related to  $U$ . These matrices are  
 49 defined in [\[1\]](#), p.27. They provide a language to talk about unitary matrices within  
 50 the context of the determinantal conjecture.

#### 51 **B-matrix**

$$52 \quad B = UB_0U^* \quad (2.1)$$

#### 53 **C-matrix**

$$54 \quad C = A_0 + UB_0U^* \quad (2.2)$$

55 Using [\(1.3\)](#),  $R_m(U) = \det(C)$

#### 56 **F-matrix**

$$57 \quad F = BC^{-1} - C^{-1}B$$

58 We can change the F-matrix into a more useful form:

$$59 \quad F = (C - A_0)C^{-1} - C^{-1}(C - A_0)$$

$$60 \quad 61 \quad F = C^{-1}A_0 - A_0C^{-1} \quad (2.3)$$

62 The F-matrix is only defined when  $C$  is invertible or equivalently  $R_m(U) \neq 0$ .

63 Since  $A_0$  is diagonal, we see that  $F$  is a zero-diagonal matrix.

64 As demonstrated in [\[1\]](#), p.27, the F-matrix is 0 if and only if  $U$  is the product of  
 65 a unitary diagonal matrix and a permutation matrix.

66 The fourth property is conditional. Given a unitary matrix  $U$  with  $R_m(U) \neq 0$   
 67 and with F-matrix  $F$ . Suppose there exist two skew-hermitian matrices  $Z_1$  and  $Z_2$  such

68 that  $\text{tr}(Z_1 F)$  and  $\text{tr}(Z_2 F)$  are both non-zero and non-collinear vectors in the complex  
 69 plane. Then we say that  $U$  is a **multidirectional** matrix. A multidirectional matrix  
 70 must have a non-zero  $F$ -matrix to allow those non-zero traces. So a permutation  
 71 matrix cannot be multidirectional because its  $F$ -matrix is 0.

72 Note that these properties require an  $A_0$  and  $B_0$  to be defined. Throughout the  
 73 paper we will assume there's a defined  $A_0$  and  $B_0$  in the background. We will not  
 74 mention them explicitly in order to simplify our language. For example when we  
 75 say "the  $C$ -matrix of a unitary matrix  $U$ ", it is clear that there's an unmentioned  
 76  $A_0$  and  $B_0$  according to which the  $C$ -matrix of  $U$  is defined. It is the same thing  
 77 with the terms "boundary matrix" and "regular boundary matrix". Obviously it is  
 78 meaningless for a unitary matrix to be a boundary matrix "in general". These terms  
 79 only make sense in the context of  $A_0$ ,  $B_0$  and the corresponding  $\Delta$ . So we'll assume  
 80 this context has been defined.

### 81 3. Main Theorems.

82 **THEOREM 3.1.** *Given  $U$  is a unitary matrix that cannot be written as the product*  
 83 *of a unitary diagonal matrix and a permutation matrix. Given  $R_m(U) \neq 0$  and its*  
 84  *$F$ -matrix is  $F$ . Given an arbitrary skew-hermitian matrix  $Z$ . There exists a curve*  
 85  *$R_f(t) \subseteq \Delta$ , where  $t$  is real, such that  $R_f(0) = R_m(U)$  and  $R'_f(0) = R_m(U)\text{tr}(ZF)$ .*

86 **THEOREM 3.2.** *If  $U$  is a boundary matrix, then  $U$  is not multidirectional.*

87 **THEOREM 3.3.** *Given a boundary matrix  $U$  such that  $R_m(U) \neq 0$ . Then its  $F$ -*  
 88 *matrix has the form  $F = e^{i\theta} H$  where  $H$  is a zero-diagonal hermitian matrix.*

89 **THEOREM 3.4.** *Given a regular boundary matrix  $U$  such that  $R_m(U) \neq 0$ . Let*  
 90  *$F = e^{i\theta} H$  be the  $F$ -matrix of  $U$ . let  $l$  be the tangent line to  $\partial\Delta$  at the boundary point.*  
 91 *Then  $l$  makes an angle  $\arg(R_m(U)) + \theta + \pi/2$  with the positive real axis.*

92 **4. Proof of Theorem 3.1.** This theorem is apparent from [1], p.27, but it is  
 93 not stated explicitly there. It is worth proving explicitly here as it will be used for  
 94 the other theorems.

95 Before we can prove the theorem we need to set up some tools. Our aim is to  
 96 examine boundary matrices of  $\Delta$ . Towards this aim, it is useful to consider smooth  
 97 functions of unitary matrices going through these boundary matrices and see how  
 98 they behave under (1.3). For this reason, we introduce the functional form of (1.3).

$$99 \quad R_f(t) = \det(A_0 + U_f(t)B_0U_f^*(t)) \quad (4.1)$$

100 where  $t$  is real and  $U_f(t)$  is some smooth function of unitary matrices.

101 Every unitary matrix can be written as an exponential of a skew-hermitian matrix.  
 102 So we can write:

$$103 \quad U_f(t) = e^{S_f(t)}. \quad (4.2)$$

104 where  $S_f(t)$  is a smooth function of skew hermitian matrices

105 For small  $\Delta t$ ,

$$106 \quad U_f(t + \Delta t) = (e^{S_f(t+\Delta t)})$$

$$107 \quad U_f(t + \Delta t) = (e^{S_f(t)+(\Delta t)S'_f(t)})$$

$$108 \quad U_f(t + \Delta t) = (e^{(\Delta t)S'_f(t)})U_f(t)$$

109 If we take the above function and plug it into  $R_f(t)$  we'll get  $R_f(t + \Delta t)$ , but it  
 110 won't be in a form useful to us. We use a result from [1], p.27 for this purpose. In  
 111 order to state this result within the context of this paper, we first need the functional  
 112 forms of the B-matrix, C-matrix, F-matrix (these were defined in [section 2](#)):

$$113 \quad B_f(t) = U_f(t)B_0U_f^*(t) \quad (4.3)$$

$$114 \quad C_f(t) = A_0 + B_f(t) \quad (4.4)$$

$$115 \quad F_f(t) = C_f^{-1}(t)A_0 - A_0C_f^{-1}(t) \quad (4.5)$$

116 Note,  $F_f(t)$  is only defined if  $R_f(t) \neq 0$ . Also  $F_f(t) = 0$  only when  $U_f(t)$  is the  
 117 product of a unitary diagonal matrix and a permutation matrix.

118 Now we can state the result from [1]:

119 When  $F_f(t) \neq 0$ ,

$$120 \quad R_f(t + \Delta t) = R_f(t) + (\Delta t) \det(C_f(t)) \text{tr}(S'_f(t)F_f(t)) + O((\Delta t)^2) \quad (4.6)$$

$$121 \quad R'_f(t) = \det(C_f(t)) \text{tr}(S'_f(t)F_f(t)) \quad (4.7)$$

123 Now we have the tools needed to prove [Theorem 3.1](#).

124 *Proof.* Given a unitary matrix  $U$  that cannot be written as the product of a  
 125 diagonal unitary matrix with a permutation matrix. Given  $R_m(U) \neq 0$ . let  $C$  be the  
 126 C-matrix of  $U$ . let  $F$  be the F-matrix of  $U$ . Given  $Z$  is some arbitrary skew-hermitian  
 127 matrix. We can find a skew-hermitian matrix  $S$  such that  $U = e^S$ .

128 We choose:

$$129 \quad S_f(t) = S + tZ \quad (4.8)$$

130 Note that  $S_f(t)$  is a smooth function of skew-hermitian matrices. We use it with  
 131 [\(4.1\)](#),[\(4.2\)](#),[\(4.4\)](#),[\(4.5\)](#) and [\(4.7\)](#) to get  $R_f(t)$ ,  $U_f(t)$ ,  $C_f(t)$ ,  $F_f(t)$  and  $R'_f(t)$ . Note that  
 132  $U_f(0) = U$ , the unitary matrix we're originally given. The choice of  $t = 0$  is merely  
 133 for simplicity and has no special significance. We could time-shift  $S_f(t)$  to the right  
 134 by  $t_1$  to make  $U_f(t_1) = U$  instead.

135 Note that  $C_f(0) = C$

136 Note that  $F_f(0) = F$

137 Note that  $R_f(0) = R_m(U)$ . See [\(1.3\)](#) and [\(4.1\)](#).

$$138 \quad R'_f(t) = \det(C_f(t))tr(ZF_f(t))$$

$$139 \quad R'_f(0) = \det(C_f(0))tr(ZF_f(0))$$

$$140 \quad R'_f(0) = \det(C)tr(ZF)$$

141 therefore

$$142 \quad R'_f(0) = R_m(U)tr(ZF) \quad (4.9)$$

143 This proves [Theorem 3.1](#).  $\square$

144 **5. Proof of [Theorem 3.2](#).** We will prove the contrapositive. ie: We'll start  
145 with a multidirectional matrix U, and prove that it is not a boundary matrix.

146 *Proof.* Given we have a multidirectional matrix U. Let F be its F-matrix and  
147 C-matrix C. We know  $R_m(U) = \det(C) \neq 0$  and we know F is non-zero. See the  
148 discussion on multidirectional matrices in the second last paragraph of [section 2](#).

149 There exist two skew-hermitian matrices  $Z_1$  and  $Z_2$  such that

$$150 \quad T_1 = tr(Z_1F) \quad (5.1)$$

$$151 \quad T_2 = tr(Z_2F) \quad (5.2)$$

152 are both non-zero and non-collinear.

153 By [Theorem 3.1](#), there exist two functions  $R_1(t)$  and  $R_2(t)$  such that  $R_1(0) =$   
154  $R_2(0) = R_m(U)$  and such that

$$155 \quad R'_1(0) = R_m(U)tr(Z_1F)$$

$$156 \quad R'_2(0) = R_m(U)tr(Z_2F)$$

157 substitute in [\(5.1\)](#) and [\(5.2\)](#),

$$158 \quad R'_1(0) = R_m(U)T_1$$

$$159 \quad R'_2(0) = R_m(U)T_2$$

160 Since we know  $T_1$  and  $T_2$  are non-collinear,  $R'_1(0)$  and  $R'_2(0)$  are non-collinear.  
161 They are also non-zero. Therefore they form a linear basis for all the complex numbers  
162 over the real numbers. Let Q be an arbitrary complex number.

$$163 \quad Q = aR'_1(0) + bR'_2(0) \text{ where } a \text{ and } b \text{ are real.}$$

$$164 \quad Q = a(R_m(U))T_1 + b(R_m(U))T_2$$

$$165 \quad Q = R_m(U)(aT_1 + bT_2)$$

166 substitute in [\(5.1\)](#) and [\(5.2\)](#),

$$167 \quad Q = R_m(U)(tr(aZ_1F) + tr(bZ_2F))$$

$$168 \quad Q = R_m(U)tr((aZ_1 + bZ_2)F)$$

$$169 \quad \text{let } Z_3 = aZ_1 + bZ_2$$

$$170 \quad Q = R_m(U)tr(Z_3F)$$

171 Note that  $Z_3$  is also a skew-hermitian matrix.

172 Again by [Theorem 3.1](#), there exists a function  $R_3(t)$  such that

$$173 \quad R_3(0) = R_m(U)$$

174 and

$$175 \quad R'_3(0) = R_m(U)tr(Z_3F) = Q$$

176 Therefore  $R_3(t)$  goes through  $R_m(U)$  in a direction parallel to  $Q$ .  $Q$  was chosen  
 177 arbitrarily. So through  $R_m(U)$  there exists curves  $R_3(t) \subseteq \Delta$  going in all directions.  
 178 Therefore  $R_m(U)$  is an internal point of  $\Delta$ . So it's not a boundary point. Therefore  
 179  $U$  is not a boundary matrix. That gives us [Theorem 3.2](#).  $\square$

180 **6. Proof of [Theorem 3.3](#).** For  $n = 3$ , we define the following 12 skew-hermitian  
 181 matrices with zero diagonal:

$$182 \quad Z_{12} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad Z_{13} = \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \quad Z_{23} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}$$

$$183 \quad Z_{21} = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad Z_{31} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix} \quad Z_{32} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}$$

$$184 \quad Z_{12,i} = Z_{21,i} = \begin{bmatrix} 0 & i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad Z_{13,i} = Z_{31,i} = \begin{bmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{bmatrix} \quad Z_{23,i} = Z_{32,i} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & i \\ 0 & i & 0 \end{bmatrix} \blacksquare$$

185 Note that the commas do not indicate tensors. They're just used here as a label  
 186 to distinguish imaginary and real matrices.

187 We define  $Z_{ab}$  and  $Z_{ab,i}$  similarly for all  $n > 3$ , where  $a \neq b$ . For a given  $n$  we  
 188 have  $n(n-1)$  real matrices and  $n(n-1)$  imaginary matrices.

189 *Proof.* Given a boundary matrix  $U$  with  $R_m(U) \neq 0$ . Let  $F$  be its  $F$ -matrix. We  
 190 know that  $F$  is zero-diagonal by [\(4.5\)](#).

191 Suppose  $F_{ab} = F_{ab,r} + iF_{ab,i}$  where  $F_{ab,r}$  and  $F_{ab,i}$  are real numbers.

$$192 \quad tr(Z_{ab}F) = F_{ab} - F_{ba}$$

$$193 \quad tr(Z_{ab,i}F) = (F_{ab} + F_{ba})i$$

194 Substitute in for  $F_{ab}$  and  $F_{ba}$

$$195 \quad \text{tr}(Z_{ab}F) = (F_{ab,r} - F_{ba,r}) + i(F_{ab,i} - F_{ba,i}) \quad (6.1)$$

$$196 \quad \text{tr}(Z_{ab,i}F) = (-F_{ab,i} - F_{ba,i}) + i(F_{ab,r} + F_{ba,r}) \quad (6.2)$$

197 By [Theorem 3.2](#), we know that  $U$  is not multidirectional.

198 Therefore

$$199 \quad (F_{ab,i} - F_{ba,i})(-F_{ab,i} - F_{ba,i}) = (F_{ab,r} + F_{ba,r})(F_{ab,r} - F_{ba,r})$$

200 We can simplify this to get:

$$201 \quad F_{ab,r}^2 + F_{ab,i}^2 = F_{ba,r}^2 + F_{ba,i}^2$$

$$202 \quad |F_{ab}| = |F_{ba}|$$

203 We can write:

$$204 \quad F_{ab} = |F_{ab}| \angle \theta_{ab}$$

$$205 \quad F_{ba} = |F_{ab}| \angle \theta_{ba}$$

206 There are multiple cases we need to deal with.

207 **Case 1: F-matrix is 0**

208  $F=0$  is hermitian so we're finished.

209 **Case 2:  $|F_{ab}|$  is non-zero for only one pair (a,b) where  $a \neq b$**

210 In this case,

211  $H = e^{-(\theta_{ab} + \theta_{ba})/2} F$  is a hermitian matrix, and we're finished.

212 **Case 3:  $|F_{ab}|$  is non-zero for multiple pairs (a,b) where  $a \neq b$ . For an**  
 213 **arbitrary skew-hermitian  $Z$ , when  $\text{tr}(ZF)$  is non-zero, it is imaginary.**

214 If  $|F_{ab}| \neq 0$ , then by [\(6.1\)](#) and [\(6.2\)](#),  $\theta_{ab} = -\theta_{ba}$ . So our  $F$ -matrix is already  
 215 hermitian, and we're done.

216 **Case 4:  $|F_{ab}|$  is non-zero for multiple pairs (a,b) where  $a \neq b$ . For an**  
 217 **arbitrary skew-hermitian  $Z$ , when  $\text{tr}(ZF)$  is non-zero, it is real.**

218 If  $|F_{ab}| \neq 0$ , then by [\(6.1\)](#) and [\(6.2\)](#),  $\theta_{ab} = \pi - \theta_{ba}$ .

219  $H = e^{-i(\frac{\pi}{2})} F$  is hermitian and we're done.

220 **Case 5:  $|F_{ab}|$  is non-zero for multiple pairs (a,b) where  $a \neq b$ . For**  
 221 **an arbitrary skew-hermitian  $Z$ , when  $\text{tr}(ZF)$  is non-zero, it isn't real or**  
 222 **imaginary.**

223 Suppose  $|F_{ab}| \neq 0$  and  $|F_{cd}| \neq 0$

224 if  $\text{tr}(Z_{ab}F) \neq 0$ , then

$$225 \quad \text{slope of } \text{tr}(Z_{ab}F) = \frac{\sin(\theta_{ab}) - \sin(\theta_{ba})}{\cos(\theta_{ab}) - \cos(\theta_{ba})} = -\cot\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right)$$

226 if  $\text{tr}(Z_{ab,i}F) \neq 0$ :

$$227 \quad \text{slope of } \text{tr}(Z_{ab,i}F) = \frac{\cos(\theta_{ab}) + \cos(\theta_{ba})}{-\sin(\theta_{ab}) - \sin(\theta_{ba})} = -\cot\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right)$$

228 similarly,

$$229 \quad \text{slope of } \text{tr}(Z_{cd}F) = -\cot\left(\frac{\theta_{cd} + \theta_{dc}}{2}\right)$$

230 or

$$231 \quad \text{slope of } \text{tr}(Z_{cd,i}F) = -\cot\left(\frac{\theta_{cd} + \theta_{dc}}{2}\right)$$

$$232 \quad \cot\left(\frac{\theta_{cd} + \theta_{dc}}{2}\right) = \cot\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right)$$

233 therefore either:

$$234 \quad \frac{\theta_{cd} + \theta_{dc}}{2} = \frac{\theta_{ab} + \theta_{ba}}{2}$$

235 or,

$$236 \quad \frac{\theta_{cd} + \theta_{dc}}{2} = \frac{\theta_{ab} + \theta_{ba}}{2} + \pi$$

237 For some specific  $x, y$  where  $x \neq y$  and  $|F_{xy}| \neq 0$

$$238 \quad \text{let } \beta = \frac{\theta_{xy} + \theta_{yx}}{2}$$

$$239 \quad \text{let } H = e^{-i\beta} F$$

240 For any  $a \neq b$ ,

$$241 \quad H_{ab} = |H_{ab}| \angle \alpha_{ab}$$

$$242 \quad \frac{\alpha_{ab} + \alpha_{ba}}{2} = 0 \text{ or } \pi$$

243 Therefore H is zero-diagonal, with transpositional elements of equal magnitude  
244 and opposite arguments. Therefore H is hermitian.

245 So in all 5 cases we can write  $F = e^{i\beta} H$  for some hermitian matrix H and some  
246 real  $\beta$ .

247 This completes our proof of [Theorem 3.3](#). □

248 **7. Proof of [Theorem 3.4](#).** Given a regular boundary matrix U. Let F be the  
249 F-matrix of U.

250 *Proof.* Therefore by [Theorem 3.3](#) we know that

$$251 \quad F = e^{i\theta} H \tag{7.1}$$

252 for some real  $\theta$  and some zero-diagonal hermitian matrix H.

253 We can substitute (7.1) into (6.1) and (6.2) and simplify to get:

$$254 \quad \text{tr}(Z_{ab}F) = 2H_{ab,i} e^{i(\theta + \pi/2)} \tag{7.2}$$

$$255 \quad \text{tr}(Z_{ab,i}F) = 2H_{ab,r}e^{i(\theta+\pi/2)} \quad (7.3)$$

256 As expected the vectors are collinear.

257 Since  $U$  is a regular boundary matrix,  $\partial\Delta$  is smooth at  $R_m(U)$  ie: the tangent to  
258 the curve exists at  $R_m(U)$ .

259 So using [Theorem 3.1](#), we see that the tangent line forms an angle  $\arg(R_m(U)) +$   
260  $\theta + \pi/2$  with the positive real axis. This completes our proof of [Theorem 3.4](#).  $\square$

261 **8. Conjectures.** Before we state our conjectures we define a region  $\Delta_S$  which  
262 is a restriction of  $\Delta$ . See [\(1.1\)](#).

$$263 \quad \Delta_S = \{ \det(A_0 + OB_0O^*) : O \in O(n) \} \quad (8.1)$$

264 where  $O(n)$  is the set of  $n \times n$  real orthogonal matrices.

265 As proven in [\[3\]](#), p.207, theorem 4.4.7, a matrix is normal and symmetric if and  
266 only if it is diagonalizable by a real orthogonal matrix.

267 Therefore  $\Delta_S$  is the set of determinants of sums of normal, symmetric matrices  
268 with prescribed eigenvalues. We know  $\Delta_S$  contains all the permutation points.

269 CONJECTURE 8.1 (Restricted Marcus-de Oliveira Conjecture).

$$270 \quad \Delta_S \subseteq \text{co} \left\{ \prod (a_i + b_{\sigma(i)}) \right\}$$

271 CONJECTURE 8.2 (Boundary Conjecture).

$$272 \quad \partial\Delta \subseteq \partial\Delta_S$$

273 **THEOREM 8.3.** *If the boundary conjecture is true, the restricted Marcus-de Oliveira*  
274 *conjecture implies the full Marcus-de Oliveira conjecture.*

275 *Proof.* Suppose we know [Conjecture 8.1](#) is true. Then  $\Delta_S$  along with its boundary  
276 is within the convex-hull. Suppose we also know that [Conjecture 8.2](#) is true. Then we  
277 know that  $\partial\Delta$  is inside the convex-hull. Can we have a unitary matrix  $U$  such that  
278  $R_m(U)$  is outside the convex-hull? No, because that would mean we have points of  
279  $\Delta$  on both the inside and outside of  $\partial\Delta$ . This is impossible since  $\Delta$  is a closed set  
280 (See the second last paragraph of [section 1](#)). So  $\Delta$  is within the convex hull proving  
281 [Conjecture 1.1](#).  $\square$

282 **9. Conclusion.** We hope that further analysis on boundary matrices of  $\Delta$ , either  
283 by expanding on the results in this paper, or novel research, leads to a proof of the  
284 Boundary Conjecture. Then proving the full Marcus-de Oliveira conjecture would  
285 amount to proving the restricted conjecture. Whether the restricted conjecture is any  
286 easier to prove is unknown, but it's an avenue worth exploring.

287 REFERENCES

- 288 [1] N. BEBIANO AND J. QUERÍO, *The determinant of the sum of two normal matrices with prescribed*  
289 *eigenvalues*, *Linear Algebra and its Applications*, 71 (1985), pp. 23–28.
- 290 [2] G. N. DE OLIVEIRA, *Research problem: Normal matrices*, *Linear and Multilinear Algebra*, 12  
291 (1982), pp. 153–154.
- 292 [3] R. HORN AND C. JOHNSON, *Matrix Analysis*, Cambridge University Press, 1990.
- 293 [4] M. MARCUS, *Derivations, plücker relations and the numerical range*, *Indiana University Math*  
294 *Journal*, 22 (1973), pp. 1137–1149.