

The permanent and diagonal products on the set of nonnegative matrices with bounded rank

Yair Lavi

ABSTRACT

We formulate conjectures regarding the maximum value and maximizing matrices of the permanent and of diagonal products on the set of stochastic matrices with bounded rank. We formulate equivalent conjectures on upper bounds for these functions for nonnegative matrices based on their rank, row sums and column sums. In particular we conjecture that the permanent of a singular nonnegative matrix is bounded by $\frac{1}{2}$ times the minimum of the product of its row sums and the product of its column sums, and that the product of the elements of any diagonal of a singular nonnegative matrix is bounded by $\frac{1}{4}$ times the minimum of the product of its row sums and the product of its column sums.

1. Introduction

The permanent of a real $n \times n$ matrix is defined by $per(A) = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i\sigma(i)}$

Consider the familiar inequality for the permanent of an $n \times n$ (row or column) stochastic matrix A [1]:

$$per(A) \leq 1 \tag{1}$$

Equality is obtained iff A is a permutation matrix, and in particular a necessary condition for equality is that $rank(A) = n$. One could naturally ask then:

What is the maximum value of the permanent function and what are the maximizing matrices in the set of stochastic matrices with rank bounded from above by some $k < n$?

In this note we conjecture an answer to this question.

An equivalent result to inequality 1 is the following bound for the permanent of a nonnegative matrix A :

$$per(A) \leq \min\left(\prod_{i=1}^n r_i, \prod_{i=1}^n c_i\right) \tag{2}$$

where (r_1, \dots, r_n) and (c_1, \dots, c_n) are the row sums and column sums of A respectively.

We formulate our conjecture in the equivalent "nonnegative formulation" as well.

Let $\sigma \in S_n$ be any permutation. We call $\prod_{i=1}^n a_{i\sigma(i)}$ a diagonal product of A . Similarly to the permanent conjecture we formulate a conjecture for the maximum of any diagonal product on the set of stochastic matrices with bounded rank and also present it in the equivalent "nonnegative formulation".

We begin in Section 2 with notations and definitions. In Section 3, we formulate the conjecture regarding the permanent function and in Section 4 we formulate the conjecture regarding the diagonal products.

2. Preliminaries

We denote by \mathcal{M}_n the set of real $n \times n$ matrices and by \mathcal{M}_n^+ the set of nonnegative matrices in \mathcal{M}_n .

We denote the set of nonnegative matrices of rank less than or equal to k by

$$\mathcal{M}_n^{+\bar{k}} := \{A \in \mathcal{M}_n^+ | \text{rank}(A) \leq k\}$$

Let R_n denote the set of row stochastic matrices in \mathcal{M}_n^+ , so R_n is the set of $n \times n$ real matrices $A = (a_{ij})$ satisfying:

$$\sum_{j=1}^n a_{ij} = 1 \quad i = 1, \dots, n$$

and

$$0 \leq a_{ij} \quad 1 \leq i, j \leq n$$

we denote the set of row stochastic matrices of rank less than or equal to k by

$$R_n^{\bar{k}} := \{A \in R_n | \text{rank}(A) \leq k\}$$

Similarly, we will write L_n , and $L_n^{\bar{k}}$ for the sets of column stochastic matrices and column stochastic matrices of rank less than or equal to k , respectively.

For a matrix $A \in \mathcal{M}_n^+$ with rows A_1, \dots, A_n and row sums $r_i > 0$, $i = 1, \dots, n$, we will denote by \bar{A}^r the row stochastic matrix whose rows are $\frac{A_1}{r_1}, \frac{A_2}{r_2}, \dots, \frac{A_n}{r_n}$. Similarly, for a matrix $A \in \mathcal{M}_n^+$ with columns $\tilde{A}_1, \dots, \tilde{A}_n$ and column sums $c_j > 0$, $j = 1, \dots, n$, we will denote by \bar{A}^c the column stochastic matrix whose columns are $\frac{\tilde{A}_1}{c_1}, \frac{\tilde{A}_2}{c_2}, \dots, \frac{\tilde{A}_n}{c_n}$.

Notation 1 (The Set of Compositions of n with s parts). We denote by $P_{n;s} \subset \mathbb{R}^s$ the set of all positive integer compositions of n , with exactly s parts, so $\vec{r} \in P_{n;s}$ iff $\sum_{i=1}^s r_i = n$, $r_i \in \mathbb{N}, i = 1, 2, \dots, s$.

Let J_r be the $r \times r$ doubly stochastic matrix whose entries all equal to $\frac{1}{r}$.

Definition 2.1 (s -composition matrix). Given a composition $\vec{r} = (r_1, r_2, \dots, r_s) \in$

$P_{n;s}$ we shall call the doubly stochastic matrix $J_{r_1} \oplus J_{r_2} \dots \oplus J_{r_s}$ an s -composition matrix and denote it by $J_{\vec{r}}$

Notation 2. Let $\pi : S_n \rightarrow \mathcal{M}_n$ be the natural permutation representation of the symmetric group S_n defined by $\pi(\sigma)_{ij} = \delta_{\sigma(i),j}$ and denote its image, the set of permutation matrices, by $\mathfrak{S}_n = \pi(S_n)$.

3. Conjectures for the permanent function

We formulate our conjecture in two equivalent forms, starting with the stochastic matrices formulation:

Conjecture 1. Maximum of the permanent function on the set of stochastic matrices with bounded rank. For $k, n \in \mathbb{N}$, $1 \leq k \leq n$, let $R_n^{\bar{k}}$ ($L_n^{\bar{k}}$) be the set of $n \times n$ row (column) - stochastic matrices of rank less than or equal to k , and let $r, s \in \mathbb{N}$ be the unique integers such that $n = rk + s$, $0 \leq s < k$. Then

$$A \in R_n^{\bar{k}} \cup L_n^{\bar{k}} \implies \text{per}(A) \leq \left(\frac{r!}{r^r}\right)^{k-s} \times \left(\frac{(r+1)!}{(r+1)^{r+1}}\right)^s$$

with equality iff $A = PJ_{\vec{r}}Q$ where $\vec{r} = (\underbrace{r, r, \dots, r}_{(k-s)\text{-times}}, \underbrace{r+1, r+1, \dots, r+1}_{s\text{-times}})$ and $P, Q \in \mathfrak{S}_n$ are any two permutation matrices.

In particular the maximizing matrices are doubly stochastic matrices of rank $= k$.

We now present the equivalent formulation for nonnegative matrices. The equivalence of the two formulations is trivial.

Conjecture 2. Permanent function of nonnegative matrices with bounded rank. For $k, n \in \mathbb{N}$, $1 \leq k \leq n$, let $M_n^{+\bar{k}}$ be the set of $n \times n$ nonnegative matrices of rank less than or equal to k , let $r, s \in \mathbb{N}$ be the unique integers such that $n = rk + s$, $0 \leq s < k$ and let $\vec{r} = (\underbrace{r, r, \dots, r}_{(k-s)\text{-times}}, \underbrace{r+1, r+1, \dots, r+1}_{s\text{-times}})$. Then

$$A \in M_n^{+\bar{k}} \implies \text{per}(A) \leq \min\left(\prod_{i=1}^n r_i, \prod_{i=1}^n c_i\right) \left(\frac{r!}{r^r}\right)^{k-s} \times \left(\frac{(r+1)!}{(r+1)^{r+1}}\right)^s$$

where (r_1, \dots, r_n) and (c_1, \dots, c_n) are the row sums and column sums of A respectively, with equality iff one of the following holds:

- (1) A has a zero row or a zero column
- (2) $0 < \prod_{i=1}^n r_i < \prod_{i=1}^n c_i$ and $\bar{A}^r = PJ_{\vec{r}}Q$ for some $P, Q \in \mathfrak{S}_n$
- (3) $0 < \prod_{i=1}^n c_i < \prod_{i=1}^n r_i$ and $\bar{A}^c = PJ_{\vec{r}}Q$ for some $P, Q \in \mathfrak{S}_n$
- (4) $0 < \prod_{i=1}^n r_i = \prod_{i=1}^n c_i$ and $\bar{A}^r = PJ_{\vec{r}}Q$ and $\bar{A}^c = P'J_{\vec{r}}Q'$ for some $P, Q, P', Q' \in \mathfrak{S}_n$

In particular for singular nonnegative matrices we make the following conjecture:

Conjecture 3. Permanent function of nonnegative singular matrices. Let A be a singular nonnegative matrix and let $\vec{r} = \underbrace{(1, 1, \dots, 1)}_{(n-2)\text{-times}}, 2$. then

$$\text{per}(A) \leq \frac{\min(\prod_{i=1}^n r_i, \prod_{i=1}^n c_i)}{2}$$

where (r_1, \dots, r_n) and (c_1, \dots, c_n) are the row sums and column sums of A respectively, with equality iff one of the following holds:

- (1) A has a zero row or a zero column
- (2) $0 < \prod_{i=1}^n r_i < \prod_{i=1}^n c_i$ and $\overline{A}^r = PJ_{\vec{r}}Q$ for some $P, Q \in \mathfrak{S}_n$
- (3) $0 < \prod_{i=1}^n c_i < \prod_{i=1}^n r_i$ and $\overline{A}^c = PJ_{\vec{r}}Q$ for some $P, Q \in \mathfrak{S}_n$
- (4) $0 < \prod_{i=1}^n r_i = \prod_{i=1}^n c_i$ and $\overline{A}^r = PJ_{\vec{r}}Q$ and $\overline{A}^c = P'J_{\vec{r}}Q'$ for some $P, Q, P', Q' \in \mathfrak{S}_n$

4. Conjectures for diagonal products

We formulate our conjecture in two equivalent forms, starting with the stochastic matrices formulation:

Conjecture 4. Maximum of diagonal products on the set of stochastic matrices with bounded rank. Let $\sigma \in S_n$ be any permutation. For $k, n \in \mathbb{N}$, $1 \leq k \leq n$, let $R_n^{\bar{k}}$ ($L_n^{\bar{k}}$) be the set of $n \times n$ row (column) - stochastic matrices of rank less or equal to k , and let $r, s \in \mathbb{N}$ be the unique integers such that $n = rk + s$, $0 \leq s < k$. Then

$$A \in R_n^{\bar{k}} \cup L_n^{\bar{k}} \implies \prod_{i=1}^n a_{i\sigma(i)} \leq \left(\frac{1}{r}\right)^{k-s} \times \left(\frac{1}{(r+1)}\right)^s$$

with equality iff $A = P^t J_{\vec{r}} P \pi(\sigma)$ where $\vec{r} = \underbrace{(r, r, \dots, r)}_{(k-s)\text{-times}}, \underbrace{(r+1, r+1, \dots, r+1)}_{s\text{-times}}$ and

$P \in \mathfrak{S}_n$ is any permutation matrix.

We now present the equivalent formulation for nonnegative matrices. The equivalence of the two formulations is trivial.

Conjecture 5. Diagonal products of nonnegative matrices with bounded rank.

Let $\sigma \in S_n$ be any permutation. For $k, n \in \mathbb{N}$, $1 \leq k \leq n$, let $M_n^{+\bar{k}}$ be the set of $n \times n$ nonnegative matrices of rank less or equal to k , let $r, s \in \mathbb{N}$ be the unique integers such that $n = rk + s$, $0 \leq s < k$ and let $\vec{r} = \underbrace{(r, r, \dots, r)}_{(k-s)\text{-times}}, \underbrace{(r+1, r+1, \dots, r+1)}_{s\text{-times}}$. Then

$$A \in M_n^{+\bar{k}} \implies \prod_{i=1}^n a_{i\sigma(i)} \leq \min\left(\prod_{i=1}^n r_i, \prod_{i=1}^n c_i\right) \left(\frac{1}{r}\right)^{k-s} \times \left(\frac{1}{(r+1)}\right)^s$$

where (r_1, \dots, r_n) and (c_1, \dots, c_n) are the row sums and column sums of A respectively, with equality iff one of the following holds:

- (1) A has a zero row or a zero column
- (2) $0 < \prod_{i=1}^n r_i < \prod_{i=1}^n c_i$ and $\overline{A}^r = P^t J_{\vec{r}} P \pi(\sigma)$ for some $P \in \mathfrak{S}_n$
- (3) $0 < \prod_{i=1}^n c_i < \prod_{i=1}^n r_i$ and $\overline{A}^c = P^t J_{\vec{r}} P \pi(\sigma)$ for some $P \in \mathfrak{S}_n$
- (4) $0 < \prod_{i=1}^n r_i = \prod_{i=1}^n c_i$ and $\overline{A}^r = P^t J_{\vec{r}} P \pi(\sigma)$ and $\overline{A}^c = P'^t J_{\vec{r}} P' \pi(\sigma)$ for some $P, P' \in \mathfrak{S}_n$

In particular for singular nonnegative matrices we make the following conjecture:

Conjecture 6. Diagonal products of nonnegative singular matrices. Let $\sigma \in S_n$ be any permutation, let A be a singular nonnegative matrix and let $\vec{r} = \underbrace{(1, 1, \dots, 1, 2)}_{(n-2)\text{-times}}$ then

$$\prod_{i=1}^n a_{i\sigma(i)} \leq \frac{\min(\prod_{i=1}^n r_i, \prod_{i=1}^n c_i)}{4}$$

where (r_1, \dots, r_n) and (c_1, \dots, c_n) are the row sums and column sums of A respectively, with equality iff one of the following holds:

- (1) A has a zero row or a zero column
- (2) $0 < \prod_{i=1}^n r_i < \prod_{i=1}^n c_i$ and $\overline{A}^r = P^t J_{\vec{r}} P \pi(\sigma)$ for some $P \in \mathfrak{S}_n$
- (3) $0 < \prod_{i=1}^n c_i < \prod_{i=1}^n r_i$ and $\overline{A}^c = P^t J_{\vec{r}} P \pi(\sigma)$ for some $P \in \mathfrak{S}_n$
- (4) $0 < \prod_{i=1}^n r_i = \prod_{i=1}^n c_i$ and $\overline{A}^r = P^t J_{\vec{r}} P \pi(\sigma)$ and $\overline{A}^c = P'^t J_{\vec{r}} P' \pi(\sigma)$ for some $P, P' \in \mathfrak{S}_n$

References

- [1] Minc H, Permanents, Encyclopedia Math. Appl., Addison-Wesley, Reading, MA, 1978.