

Multi-criteria decision making method based on similarity measures under single-valued neutrosophic refined and interval neutrosophic refined environments

Faruk Karaaslan

Department of Mathematics, Faculty of Sciences,
Çankırı Karatekin University, 18100, Çankırı, Turkey,
fkaraaslan@karatekin.edu.tr

Abstract

In this paper, we propose three similarity measure methods for single-valued neutrosophic refined sets and interval neutrosophic refined sets based on Jaccard, Dice and Cosine similarity measures of single-valued neutrosophic sets and interval neutrosophic sets. Furthermore, we suggest two multi-criteria decision making methods under single-valued neutrosophic refined environment and interval neutrosophic refined environment, and give applications of proposed multi-criteria decision making methods. Finally we suggest a consistency analysis method for proposed similarity measures between interval neutrosophic refined sets and give an application to demonstrate process of the method.

Key words: Neutrosophic set, single-valued neutrosophic set, interval neutrosophic set, neutrosophic refined set, single-valued neutrosophic refined set, interval neutrosophic refined set, similarity measure, decision making.

1. Introduction

Overcoming of situations containing uncertainty and inconsistency data has been very important matter for researchers that study on mathematical modeling and decision making which are very important in some areas such as operations research, social science, economics and management science, etc. From past to present many studies on mathematical modeling have been performed. Some of well-known approximations are fuzzy set (FS) theory proposed by Zadeh [30], intuitionistic fuzzy set (IFS) theory introduced by Atanassov [1] and interval valued intuitionistic fuzzy set theory suggested by Atanassov and Gargov [2]. A FS is identified by membership function, IFS, which is a generalization of the FSs, is characterized by membership and nonmembership functions. Even though these set theories are very successful in order to model some decision making problems containing uncertainty and incomplete information, sometimes they may not suffice to model indeterminate and inconsistent information encountered in real world. Therefore, Smarandache [23] introduced the concept of neutrosophic set, which is very useful to model problems containing indeterminate and inconsistent information, based on neutrosophy which is a branch of philosophy. A neutrosophic set is characterized by three functions called truth-membership function ($T(x)$), indeterminacy-membership function ($I(x)$) and falsity membership function ($F(x)$). These functions are real standard or nonstandard subsets of $]^{-}0, 1^{+}[$, i.e., $T(x) : X \rightarrow]^{-}0, 1^{+}[$, $I(x) : X \rightarrow]^{-}0, 1^{+}[$, and $F(x) : X \rightarrow]^{-}0, 1^{+}[$. Basic of the neutrosophic set stands up to the non-standard analysis introduced by Abraham Robinson in 1960s [21]. Smarandache [24] discussed comparisons between neutrosophic set, paraconsistent set and intuitionistic

fuzzy set, and he showed that the neutrosophic set is a generalization of paraconsistent set and intuitionistic fuzzy set. In some areas such as engineering and real scientific fields, modeling of problems by using real standard or nonstandard subsets of $]^{-0, 1^{+}[$ may not be easy sometimes, to cope with this issue concepts of single-valued neutrosophic set (SVN-set) and interval neutrosophic set (IN-set) were defined by Wang et al. in [27] and [28], respectively. Zhang et al. [31] presented an application of IN-set in multi criteria decision making problems. Some novel operations of interval neutrosophic sets were defined by Broumi and Smarandache [8]. Bhowmik and Pal [6] defined concept of intuitionistic neutrosophic set by combining intuitionistic fuzzy set with neutrosophic set, and gave some set theoretical operations of the intuitionistic neutrosophic set such as complement, union and intersection. Ansari et al. [3] made an application of neutrosophic set theory to medical AI. Ye [37] introduced concept of simplified neutrosophic sets and proposed some aggregation operators. He also developed a multi-criteria decision making method based on these aggregation operators and cosine similarity measures for simplified neutrosophic sets. Ye [41] proposed concept of trapezoidal neutrosophic set by combining trapezoidal fuzzy set with single-valued neutrosophic set. He also presented some operational rules related to this novel sets, and proposed score and accuracy function for trapezoidal neutrosophic numbers. Peng et al. [19] suggested an outranking method for multi-criteria decision making under the simplified neutrosophic environment. They also proposed a multi-criteria group decision making method using simplified neutrosophic sets [20]. Peng et al. [26] put forward multi-criteria decision making method based on a cross-entropy with interval neutrosophic sets. Ma et al. [14] developed a decision making method for a medical treatment selection based on prioritized harmonic mean operators under an interval neutrosophic linguistic environment and gave application of the proposed method, and compared accuracy of proposed method with traditional methods. Also you can find other important studies on decision making in references [29, 42].

Set theories mentioned above are based on idea which each element of a set appear only one time in the set. However, in some situations, a structure containing repeated elements may be need. For instance, while search in a dad name-number of children-occupation relational basis. To model such cases, a structure called bags was defined by Yager [32]. In 1998, Baowen [5] defined concepts of fuzzy bags and their operations based on Peizhuang's theory of set-valued statistics [18] and Yager's bags theory [32]. Concept of intuitionistic fuzzy bags (multi set) and its operations were defined by Shinoj and Sunil [22], and they gave an application in medical diagnosis under intuitionistic fuzzy multi environment.

In 2013, Smarandache [25] put forward n -symbolic (respectively i.e numerical valued) neutrosophic logic which is a refined symbolic and respectively numerical neutrosophic logic. Although existing set theoretical approximations are generally successful in order to model some problems encountered in real world, in some cases they may not allow for modeling of problems. For example, when elements in a set are evaluated by SVN-values in different times as t_1, t_2, \dots, t_p , SVN-set may not be sufficient in order to express such a case. Therefore, Ye and Ye [35] defined concept of single-valued neutrosophic multiset (refined) (SVNR-set) as a generalization of single-valued neutrosophic sets, and gave operational rules for proposed novel set. In a SVNR-set, each of truth membership values, indeterminacy membership values and falsity membership values are expressed sequences called truth membership sequence, indeterminacy membership sequence and falsity membership sequence, respectively. SVNR-set allows for modeling of problems containing changing values with respect to times under SVN-environment. In this regard, SVNR-set is an important tool to model some problems. Broumi et al. [12] proposed concept of n -valued interval neutrosophic set and set theoretical operations on n -valued interval neutrosophic set (or interval neutrosophic set) such as union, intersection, addition, multiplication, scalar multiplication, scalar

division, truth-favorite. Also they developed a multi-criteria group decision making method and gave its an application in medical diagnosis.

Similarity measure has an important role in many areas such as medical diagnosis, pattern recognition, clustering analysis, decision making and so on. There are many studies on similarity measures of neutrosophic sets and IN-sets. For example, Broumi and Smarandache [7] developed some similarity measure methods between two neutrosophic sets based on Hausdorff distances and they used these methods to calculate similarity degree between two neutrosophic sets. Ye [34] proposed three similarity measure methods based on simplified neutrosophic sets (SN-sets) which is a subclass of neutrosophic set that is more useful than neutrosophic set in some applications in engineering and real sciences. He also applied the these methods to decision making problem under SN-environment. Ye and Zhang [33] suggested similarity measure between two SVN-sets based on minimum and maximum operators. They also developed a multi-attribute decision making method based on weighted similarity measure of SVN-sets, and gave applications to demonstrate effectiveness of the proposed methods. Ye [38] proposed two similarity measures between SVN-sets by defining a generalized distance measure, and presented a clustering algorithm based on proposed similarity measure. In 2015, Ye [41] pointed out some drawbacks of similarity measures given in [34] and proposed improved cosine similarity measures of simplified neutrosophic sets (SN-sets) based on cosine function. Moreover, he defined weighted cosine similarity measures of SN-sets and gave an application in medical diagnosis problem containing SN-information. Ye and Fub [39] proposed a similarity measure of SVN-sets based on tangent function and put forward a medical diagnosis method called multi-period medical diagnosis method based on suggested similarity measure and weighted aggregation of multi-period information. They also made a comparison between tangent similarity measures of SVN-sets and existing similarity measures of SVN-sets. Furthermore, Ye [40] introduced a similarity measure of SVN-sets based on cotangent function and gave an application in the fault diagnosis of steam turbine, and he gave comparative analysis between cosine similarity measure and cotangent similarity measure in the fault diagnosis of steam turbine. Majumdar and Samanta [15] defined notion of distance between two SVN-sets and investigated its some properties. They also put forward a measure of entropy for a SVN-set. Aydođdu [4] introduced a similarity measure between two SVN-sets and developed an entropy of SVN-sets. Broumi and Smarandache [11] extended similarity measures proposed in [38] to IN-sets. Ye [36] proposed a similarity measure between two IN-sets based on Hamming and Euclidian distances and gave a multi-criteria decision making method.

Similarity measure on the NR-sets was studied Broumi and Smarandache [9]. Broumi and Smarandache extended improved cosine similarity measure of SVN-sets to NR-sets and gave its an application in medical diagnosis. Mondal and Pramanik [16] introduced cotangent similarity measure of NR-sets and studied on its properties, and applied cotangent similarity measure to educational stream selection. Also they proposed a similarity measure method [17] for NR-sets based on tangent function and gave an application in multi-attribute decision making. In 2015. Broumi and Smarandache [10] presented a new similarity measure method by extending the Hausdorff distance to NR-sets, and gave an application of proposed method in medical diagnosis.

In this paper, we propose three similarity measure methods for single-valued neutrosophic refined sets (SVNR-sets) and interval neutrosophic refined sets (INR-sets) by extending Jaccard, Dice and Cosine similarity measures under SVN-value and IN-value given by Ye in [34]. Also we give two multi-criteria decision making methods by defining ideal solutions for best and cost criteria under SVNR-environment and INR-environment. Furthermore, to determine which similarity measure under INR-environment is more appropriate for considered problems, we give a consistency analysis method based on developed similarity measure methods. To demonstrate processes

of the similarity measure methods and consistency analysis method, we present real examples based on criteria and attributes given in [34]. The rest of the article is organized as follows. In section 2, some definitions related to the SVN-sets and IN-sets and formulas of Jaccard, Dice and Cosine similarity measures under SVN-environment are given. In section 3, for SVN-sets and IN-sets similarity measures methods are developed as an extension of vector similarity measures between two SVN-sets and between two IN-sets given in [34]. In section 4, multi criteria decision making methods are developed under SVN-environment and IN-environment, and given examples related to the developed methods, and proposed similarity measure methods are compared with existing similarity measures methods. In section 5 for similarity measures between two IN-sets, a consistency analysis method is suggested and an application of this method is given. In section 6, conclusions of the paper and studies that can be made in future are presented.

2. Preliminary

In this section, concepts of SVN-set, IN-set, SVN-sets and IN-sets and some set theoretical operations required in subsequent sections of them are presented .

Throughout the paper, X denotes initial universe, E is a set of parameters and $I_p = \{1, 2, \dots, p\} (p \in N)$ is an index set.

Definition 1. [28] Let X be a nonempty set (initial universe), with a generic element in X denoted by x . A single-valued neutrosophic set (SVN-set) A is characterized by three functions called truth membership function $t_A(x)$, indeterminacy membership function $i_A(x)$ and falsity membership function $f_A(x)$ such that $t_A(x), i_A(x), f_A(x) \in [0, 1]$ for all $x \in X$, as follows:

If X is continuous, a SVN-set A can be written as follows:

$$A = \int_X \langle t_A(x), i_A(x), f_A(x) \rangle / x, \text{ for all } x \in X.$$

If X is crisp set, a SVN-set A can be written as follows:

$$A = \sum_x \langle t_A(x), i_A(x), f_A(x) \rangle / x, \text{ for all } x \in X.$$

Also, finite SVN-set A can be presented as follows:

$$A = \{ \langle x_1, t_A(x_1), i_A(x_1), f_A(x_1) \rangle, \dots, \langle x_M, t_A(x_M), i_A(x_M), f_A(x_M) \rangle \}$$

for all $x \in X$. Here $0 \leq t_A(x) + i_A(x) + f_A(x) \leq 3$ for all $x \in X$.

Throughout this paper, initial universe will be considered as a finite and crisp set, and set of all SVN-sets over X will be denoted by SVN_X .

Operations on SVN-sets are given by Wang et. al in [28] as follows:

Definition 2. [28] Let $A, B \in SVN_X$. Then,

1. $A \subseteq B$ if and only if $t_A(x) \leq t_B(x), i_A(x) \geq i_B(x), f_A(x) \geq f_B(x)$ for all $x \in X$.
2. $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$ for all $x \in X$.
3. $A^c = \{ \langle x, f_A(x), 1 - i_A(x), t_A(x) \rangle : x \in X \}$.
4. $A \cup B = \{ \langle x, (t_A(x) \vee t_B(x)), (i_A(x) \wedge i_B(x)), (f_A(x) \wedge f_B(x)) \rangle : x \in X \}$

5. $A \cap B = \{ \langle x, (t_A(x) \wedge t_B(x)), (i_A(x) \vee i_B(x)), (f_A(x) \vee f_B(x)) \rangle : x \in X \}$.

Definition 3. [35] Let X be a nonempty set with generic elements in X denoted by x . A single-valued neutrosophic refined set (SVNR-set), denoted by \tilde{A} , is defined as follows:

$$\tilde{A} = \left\{ \left\langle x, (t_A^1(x), t_A^2(x), \dots, t_A^p(x)), (i_A^1(x), i_A^2(x), \dots, i_A^p(x)), (f_A^1(x), f_A^2(x), \dots, f_A^p(x)) \right\rangle : x \in X \right\}.$$

Here, $t_A^1, t_A^2, \dots, t_A^p : X \rightarrow [0, 1]$, $i_A^1, i_A^2, \dots, i_A^p : X \rightarrow [0, 1]$ and $f_A^1, f_A^2, \dots, f_A^p : X \rightarrow [0, 1]$ such that $0 \leq t_A^r(x) + i_A^r(x) + f_A^r(x) \leq 3$ for all $x \in X$ and $r \in I_p$. $(t_A^1(x), t_A^2(x), \dots, t_A^p(x))$, $(i_A^1(x), i_A^2(x), \dots, i_A^p(x))$ and $(f_A^1(x), f_A^2(x), \dots, f_A^p(x))$ are called truth-membership sequence, indeterminacy-membership sequence and falsity-membership sequence of the element x . These sequences may be in decreasing or increasing order. Also p is called the dimension of single-valued neutrosophic refined set \tilde{A} .

A SVNR-set \tilde{A} can be represented as follows:

$$\tilde{A} = \left\{ \left\langle x, t_A^r(x), i_A^r(x), f_A^r(x) \right\rangle : x \in X, r \in I_p \right\}.$$

From now on, set of all single-valued neutrosophic refined sets with p dimension over X will be denoted by $SVNR_X^p$.

Definition 4. [35] Let $\tilde{A}, \tilde{B} \in SVNR_X^p$. Then,

1. If $t_A^r(x) \leq t_B^r(x)$, $i_A^r(x) \geq i_B^r(x)$, $f_A^r(x) \geq f_B^r(x)$ for all $r \in I_p$ and $x \in X$, then \tilde{A} is said to be SVNR-subset of \tilde{B} and this relation is denoted by $\tilde{A} \subseteq \tilde{B}$.
2. $\tilde{A} \subseteq \tilde{B}$ and $\tilde{B} \subseteq \tilde{A}$ if and only if $\tilde{A} = \tilde{B}$;
3. The complement of \tilde{A} , denoted by \tilde{A}^c , is defined by

$$\tilde{A}^c = \left\{ \left\langle x, f_A^r(x), 1 - i_A^r(x), t_A^r(x) \right\rangle : x \in X, r \in I_p \right\}.$$

Definition 5. [13] Let $\tilde{A} \in SVNR_X^p$. Then,

1. if $t_A^r(x) = 0$, $i_A^r(x) = 1$ and $f_A^r(x) = 1$ for all $r \in I_p$ and $x \in X$, \tilde{A} is called a null SVNR-set, and denoted by $\tilde{\Phi}$,
2. if $t_A^r(x) = 1$, $i_A^r(x) = 0$ and $f_A^r(x) = 0$ for all $r \in I_p$ and $x \in X$, \tilde{A} is called universal SVNR-set, and denoted by \tilde{X} .

Definition 6. [35] Let $\tilde{A}, \tilde{B} \in SVNR_X^p$. Then,

1. union:

$$\tilde{A} \tilde{\cup} \tilde{B} = \left\{ \left\langle x, t_A^r(x) \vee t_B^r(x), i_A^r(x) \wedge i_B^r(x), f_A^r(x) \wedge f_B^r(x) \right\rangle : x \in X, r \in I_p \right\},$$

2. intersection:

$$\tilde{A} \tilde{\cap} \tilde{B} = \left\{ \left\langle x, t_A^r(x) \wedge t_B^r(x), i_A^r(x) \vee i_B^r(x), f_A^r(x) \vee f_B^r(x) \right\rangle : x \in X, r \in I_p \right\}.$$

Example 1. Consider SVN-sets \tilde{A} , \tilde{B} and \tilde{C} are given as follows:

$$\tilde{A} = \left\{ \begin{array}{l} \langle x_1(.1, .2, .4), (.1, .4, .6), (.0, .3, .3) \rangle, \\ \langle x_2(.3, .3, .5), (.2, .3, .7), (.1, .5, .6) \rangle, \\ \langle x_3(.2, .4, .8), (.1, .3, .3), (.5, .6, .9) \rangle \end{array} \right\}, \quad \tilde{B} = \left\{ \begin{array}{l} \langle x_1(.5, .6, .7), (.4, .6, .7), (.3, .3, .4) \rangle, \\ \langle x_2(.2, .4, .4), (.2, .5, .8), (.2, .6, .7) \rangle, \\ \langle x_3(.1, .6, .6), (.1, .5, .5), (.3, .4, .7) \rangle \end{array} \right\}$$

and

$$\tilde{C} = \left\{ \begin{array}{l} \langle x_1(.3, .3, .5), (.4, .5, .6), (.1, .3, .4) \rangle, \\ \langle x_2(.0, .1, .3), (.2, .3, .6), (.1, .4, .6) \rangle, \\ \langle x_3(.1, .4, .7), (.1, .3, .4), (.3, .3, .5) \rangle \end{array} \right\}.$$

$$\text{Then, } \tilde{A} \tilde{\cup} \tilde{B} = \left\{ \begin{array}{l} \langle x_1(.5, .6, .7), (.4, .6, .7), (.3, .3, .4) \rangle, \\ \langle x_2(.2, .3, .4), (.2, .3, .7), (.1, .5, .6) \rangle, \\ \langle x_3(.1, .4, .6), (.1, .3, .3), (.3, .4, .7) \rangle \end{array} \right\}, \quad \tilde{A} \tilde{\cap} \tilde{B} = \left\{ \begin{array}{l} \langle x_1(.1, .2, .4), (.1, .4, .6), (.0, .3, .3) \rangle, \\ \langle x_2(.3, .4, .5), (.2, .5, .8), (.2, .6, .7) \rangle, \\ \langle x_3(.2, .6, .8), (.1, .5, .7), (.5, .6, .9) \rangle \end{array} \right\},$$

and $\tilde{C} \subseteq \tilde{B}$.

Definition 7. [27] Let $D[0, 1]$ be the set of all closed sub-intervals of the interval $[0, 1]$ and X be an ordinary finite non-empty set. An IN-set \hat{A} over X is a set of quadruple given as follows:

$$\hat{A} = \{ \langle x, t_A(x), i_A(x), f_A(x) \rangle \mid x \in X \},$$

where, $t_A(x) \in D[0, 1]$, $i_A(x) \in D[0, 1]$, and $f_A(x) \in D[0, 1]$ with the relation

$$0 \leq \sup t_A(x) + \sup i_A(x) + \sup f_A(x) \leq 3, \text{ for all } x \in X.$$

Here intervals $t_A(x) = [t_A^L(x), t_A^U(x)] \subset [0, 1]$, $i_A(x) = [i_A^L(x), i_A^U(x)] \subset [0, 1]$ and $f_A(x) = [f_A^L(x), f_A^U(x)] \subset [0, 1]$ denote the degree of truth, indeterminacy, and falsity membership of $x \in X$ in \hat{A} , respectively; moreover

$$\begin{aligned} t_A^L(x) &= \inf t_A(x), t_A^U(x) = \sup t_A(x) \\ i_A^L(x) &= \inf i_A(x), i_A^U(x) = \sup i_A(x) \\ f_A^L(x) &= \inf f_A(x), f_A^U(x) = \sup f_A(x) \end{aligned}$$

for every $x \in X$. The interval neutrosophic set \hat{A} can be expressed in the following interval format:

$$\hat{A} = \{ \langle x, [t_A^L(x), t_A^U(x)] [i_A^L(x), i_A^U(x)] [f_A^L(x), f_A^U(x)] \rangle \mid x \in X \}$$

where, $0 \leq \sup t_A^U(x) + \sup i_A^U(x) + \sup f_A^U(x) \leq 3$, $T_A^L(x) \geq 0$, $I_A^L(x) \geq 0$ and $F_A^L(x) \geq 0$ for all $x \in X$. Henceforth set of all IN-sets over X will be denoted by IN_X .

Definition 8. [12] Let X be a nonempty initial universe whose elements are discrete. An n -valued interval neutrosophic refined set (or interval neutrosophic refined set) \ddot{A} is defined as follows:

$$\ddot{A} = \left\{ \left\langle x, ([t_A^{rL}(x), t_A^{rU}(x)]), ([i_A^{rL}(x), i_A^{rU}(x)]), ([f_A^{rL}(x), f_A^{rU}(x)]) \right\rangle : x \in X \text{ and } r \in I_p \right\},$$

where $t_A^{rL}(x) \leq t_A^{rU}(x)$, $i_A^{rL}(x) \leq i_A^{rU}(x)$ and $f_A^{rL}(x) \leq f_A^{rU}(x)$ for all $x \in X$.

Here, $0 \leq t_A^{rL}(x) + i_A^{rL}(x) + f_A^{rL}(x) \leq 3$ and $0 \leq t_A^{rU}(x) + i_A^{rU}(x) + f_A^{rU}(x) \leq 3$ for all $x \in X$ and $r \in I_p$.

$$([t_A^{1L}(x), t_A^{1U}(x)], [t_A^{2L}(x), t_A^{2U}(x)], \dots, [t_A^{pL}(x), t_A^{pU}(x)]),$$

$$([i_A^{1L}(x), i_A^{1U}(x)], [i_A^{2L}(x), i_A^{2U}(x)], \dots, [i_A^{pL}(x), i_A^{pU}(x)])$$

and

$$([f_A^{1L}(x), f_A^{1U}(x)], [f_A^{2L}(x), f_A^{2U}(x)], \dots, [f_A^{pL}(x), f_A^{pU}(x)])$$

are the truth-membership sequence, indeterminacy-membership sequence and falsity-membership sequence of the element x , respectively. p is called the dimension of n -valued neutrosophic sets A .

Henceforth set of all n -valued interval neutrosophic set with p dimension over X will be denoted by INR_X^p . Also notion of interval neutrosophic refined set (INR-set) will be used instead of notion of n -valued interval neutrosophic set.

2.1. Similarity measures of SVN-sets and IN-sets

In this subsection, Jaccard, Dice, and Cosine similarity measures between two SVN-sets and between two IN-sets defined in [34] are given.

Definition 9. Let A and B be two SVN-sets over universe $X = \{x_1, x_2, \dots, x_n\}$. Then the Jaccard similarity measure between SVN-sets A and B in the vector space is defined as follows:

$$(A, B)_J = \frac{1}{n} \sum_{i=1}^n \frac{t_A(x_i)t_B(x_i) + i_A(x_i)i_B(x_i) + f_A(x_i)f_B(x_i)}{\left(\begin{array}{l} (t_A^2(x_i) + i_A^2(x_i) + f_A^2(x_i)) + (t_B^2(x_i) + i_B^2(x_i) + f_B^2(x_i)) \\ - (t_A(x_i)t_B(x_i) + i_A(x_i)i_B(x_i) + f_A(x_i)f_B(x_i)) \end{array} \right)}. \quad (1)$$

Definition 10. Let A and B be two SVN-sets over universe $X = \{x_1, x_2, \dots, x_n\}$. Then the Dice similarity measure between SVN-sets A and B in the vector space is defined as follows:

$$(A, B)_D = \frac{1}{n} \sum_{i=1}^n \frac{2(t_A(x_i)t_B(x_i) + i_A(x_i)i_B(x_i) + f_A(x_i)f_B(x_i))}{[(t_A^2(x_i) + i_A^2(x_i) + f_A^2(x_i)) + (t_B^2(x_i) + i_B^2(x_i) + f_B^2(x_i))]} \quad (2)$$

Definition 11. Let A and B be two SVN-sets in a universe of discourse $X = \{x_1, x_2, \dots, x_n\}$. Then the cosine similarity measure between SVN-sets A and B in the vector space is defined as follows:

$$(A, B)_C = \frac{1}{n} \sum_{i=1}^n \frac{(t_A(x_i)t_B(x_i) + i_A(x_i)i_B(x_i) + f_A(x_i)f_B(x_i))}{\left[\sqrt{(t_A^2(x_i) + i_A^2(x_i) + f_A^2(x_i))} \cdot \sqrt{(t_B^2(x_i) + i_B^2(x_i) + f_B^2(x_i))} \right]}. \quad (3)$$

In some applications, each element $x_i \in X (i \in I_n)$ may have different weights. Let w_1, w_2, \dots, w_n be the weights of elements $x_1, x_2, \dots, x_n \in X$ such that $w_i \geq 0 (\forall i \in I_n)$ and $\sum_{i=1}^n w_i = 1$, respectively. Then, formulas of Jaccard, Dice and Cosine similarity measures between A and B can be extended to weighted Jaccard, Dice and Cosine similarity measures are defined as follows: defined as follows:

$$W(A, B)_J = \sum_{i=1}^n w_i \frac{t_A(x_i)t_B(x_i) + i_A(x_i)i_B(x_i) + f_A(x_i)f_B(x_i)}{\left(\begin{array}{l} (t_A^2(x_i) + i_A^2(x_i) + f_A^2(x_i)) + (t_B^2(x_i) + i_B^2(x_i) + f_B^2(x_i)) \\ - (t_A(x_i)t_B(x_i) + i_A(x_i)i_B(x_i) + f_A(x_i)f_B(x_i)) \end{array} \right)}, \quad (4)$$

$$W(A, B)_D = \sum_{i=1}^n w_i \frac{2(t_A(x_i)t_B(x_i) + i_A(x_i)i_B(x_i) + f_A(x_i)f_B(x_i))}{[(t_A^2(x_i) + i_A^2(x_i) + f_A^2(x_i)) + (t_B^2(x_i) + i_B^2(x_i) + f_B^2(x_i))]} \quad (5)$$

and

$$W(A, B)_C = \sum_{i=1}^n w_i \frac{(t_A(x_i)t_B(x_i) + i_A(x_i)i_B(x_i) + f_A(x_i)f_B(x_i))}{\left[\sqrt{(t_A^2(x_i) + i_A^2(x_i) + f_A^2(x_i))} \cdot \sqrt{(t_B^2(x_i) + i_B^2(x_i) + f_B^2(x_i))} \right]}, \quad (6)$$

respectively.

3. Similarity measures under SVNR and INR-environments

In this section, similarity measures between two SVNR-sets and between two INR-sets are defined based on similarity measures between two SVN-sets and similarity measures between two IN-sets given in [34].

Definition 12. Let $\tilde{A}, \tilde{B} \in SVNR_X^p$. Then, the Jaccard similarity measure between SVNR-sets \tilde{A} and \tilde{B} is defined as follows:

$$(\tilde{A}, \tilde{B})_J = \frac{1}{p.n} \sum_{j=1}^n \sum_{r=1}^p \frac{(t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j))}{\left(\begin{aligned} &([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2) + ([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2) \\ &- [t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j)] \end{aligned} \right)}, \quad (7)$$

Definition 13. Let $\tilde{A}, \tilde{B} \in SVNR_X^p$. Then the Dice similarity measure between SVNR-sets \tilde{A} and \tilde{B} is defined as follows:

$$(\tilde{A}, \tilde{B})_D = \frac{1}{p.n} \sum_{j=1}^n \sum_{r=1}^p \frac{2(t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j))}{\left([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2 + ([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2) \right)}. \quad (8)$$

Definition 14. Let $\tilde{A}, \tilde{B} \in SVNR_X^p$. Then, the cosine similarity measure between SVNR-sets \tilde{A} and \tilde{B} is defined as follows:

$$(\tilde{A}, \tilde{B})_C = \frac{1}{p.n} \sum_{j=1}^n \sum_{r=1}^p \frac{(t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j))}{\left(\sqrt{([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2)} \cdot \sqrt{([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2)} \right)}. \quad (9)$$

If $w_j \in [0, 1]$ be the weight of each element x_j for $j = 1, 2, \dots, n$ such that $\sum_{j=1}^n w_j = 1$, then the weighted Jaccard, Dice and Cosine similarity measures between SVNR-sets \tilde{A} and \tilde{B} are defined as follows:

$$W(\tilde{A}, \tilde{B})_J = \frac{1}{p} \sum_{j=1}^n \sum_{r=1}^p w_j \frac{(t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j))}{\left(\begin{aligned} &([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2) + ([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2) \\ &- [t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j)] \end{aligned} \right)}, \quad (10)$$

$$W(\tilde{A}, \tilde{B})_D = \frac{1}{p} \sum_{j=1}^n \sum_{r=1}^p w_j \frac{2(t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j))}{\left(([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2) + ([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2) \right)} \quad (11)$$

and

$$W(\tilde{A}, \tilde{B})_C = \frac{1}{p} \sum_{j=1}^n \sum_{r=1}^p w_j \frac{(t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j))}{\left(\sqrt{([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2)} \cdot \sqrt{([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2)} \right)}, \quad (12)$$

respectively.

Example 2. Consider SVNR-sets \tilde{A} and \tilde{B} given in Example 1. Then, by using Eqs.(7),(8) and (9), similarity measures between SVNR-sets \tilde{A} and \tilde{B} are obtained as in Table 1.

If weights of the x_1, x_2 and x_3 are taken as $w_1 = .1$ $w_2 = .65$ and $w_3 = .25$, respectively. Then, by using Eqs.(10),(11) and (12), weighted similarity measures are obtained as in Table 3:

Table 1: Similarity measure under SVN R -environment

Similarity measures	Values
$(\tilde{A}, \tilde{B})_J$	0.784
$(\tilde{A}, \tilde{B})_D$	0.856
$(\tilde{A}, \tilde{B})_C$	0.955

Table 2: Weighted similarity measures under SVN R -environment

Similarity measure	Values
$W(\tilde{A}, \tilde{B})_J$	0.683
$W(\tilde{A}, \tilde{B})_D$	0.773
$W(\tilde{A}, \tilde{B})_C$	0.946

Proposition 1. Let $\tilde{A}, \tilde{B} \in SVN\mathcal{R}_X^p$. Then, each similarity measure $(\tilde{A}, \tilde{B})_\Lambda$ ($\Lambda = J, D, C$) satisfies the following properties:

1. $0 \leq (\tilde{A}, \tilde{B})_\Lambda \leq 1$
2. $(\tilde{A}, \tilde{B})_\Lambda = (\tilde{B}, \tilde{A})_\Lambda$;
3. $(\tilde{A}, \tilde{B})_\Lambda = 1$ if $\tilde{B} = \tilde{A}$ i.e. $t_A^r(x_j) = t_B^r(x_j)$, $i_A^r(x_j) = i_B^r(x_j)$, and $f_A^r(x_j) = f_B^r(x_j)$ for every $x_j \in X$ and $r \in I_p$.

PROOF. 1. For $p = 1$ Eq. (7), (8) and (9) are reduce to Eq. (1), (2) and (3), respectively. For all $r \in I_p$ ($p > 1$) according to inequality $x^2 + y^2 \geq 2xy$, for any $x_j \in X$ we know that

$$\begin{aligned} \sum_{r=1}^p ([t_A^r(x_j)]^2 + [t_B^r(x_j)]^2) &\geq 2 \sum_{r=1}^p ([t_A^r(x_j)] \cdot [t_B^r(x_j)]), \\ \sum_{r=1}^p ([i_A^r(x_j)]^2 + [i_B^r(x_j)]^2) &\geq 2 \sum_{i=1}^p ([i_A^r(x_j)] \cdot [i_B^r(x_j)]), \\ \sum_{r=1}^p ([f_A^r(x_j)]^2 + [f_B^r(x_j)]^2) &\geq 2 \sum_{r=1}^p ([f_A^r(x_j)] \cdot [f_B^r(x_j)]), \end{aligned}$$

and

$$\sum_{i=1}^p ([t_A^r(x_j)]^2 + [t_B^r(x_j)]^2 + [i_A^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_A^r(x_j)]^2 + [f_B^r(x_j)]^2) \geq 2 \sum_{i=1}^p ([t_A^r(x_j)] \cdot [t_B^r(x_j)] + [i_A^r(x_j)] \cdot [i_B^r(x_j)] + [f_A^r(x_j)] \cdot [f_B^r(x_j)]). \text{ Thus,}$$

$$\frac{1}{p} \sum_{r=1}^p \left(\frac{(t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j))}{([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2) + ([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2)} - \frac{[t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j)]}{([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2) + ([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2)} \right) \leq \frac{1}{p}. \quad (13)$$

and for all $x_j \in X$ and $p > 0$.

$$\frac{1}{p} \sum_{j=1}^n \sum_{r=1}^p \frac{(t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j))}{\left(\begin{aligned} &([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2) + ([t_B^r(x_j)]^2 + [i_B^r(x_j)]^2 + [f_B^r(x_j)]^2) \\ &- [t_A^r(x_j)t_B^r(x_j) + i_A^r(x_j)i_B^r(x_j) + f_A^r(x_j)f_B^r(x_j)] \end{aligned} \right)} \leq n. \quad (14)$$

Similarly, Eq.(8) and Eq. (9) are true.

2. The proof is clear.

3. Let $A = B$. Then, $t_A^r(x_j) = t_B^r(x_j)$, $i_A^r(x_j) = i_B^r(x_j)$, and $f_A^r(x_j) = f_B^r(x_j)$ for all $x_j \in X$ and $r \in I_p$ and

$$\begin{aligned} (\tilde{A}, \tilde{B})_J &= \frac{1}{n} \sum_{j=1}^n \sum_{r=1}^p \frac{[t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2}{(2[t_A^r(x_j)]^2 + 2[i_A^r(x_j)]^2 + 2[f_A^r(x_j)]^2 - ([t_A^r(x_j)]^2 + [i_A^r(x_j)]^2 + [f_A^r(x_j)]^2))} \\ &= \frac{1}{p \cdot n} \sum_{j=1}^n \sum_{r=1}^p 1 = 1. \end{aligned}$$

The proofs for Dice and Cosine similarity measures can be made by similar way.

Each similarity measure between two SVN-R-sets $\tilde{A} = \{ \langle x, t_A^r(x), i_A^r(x), f_A^r(x) \rangle : x \in X, i \in I_p \}$ and $\tilde{B} = \{ \langle x, t_B^r(x), i_B^r(x), f_B^r(x) \rangle : x \in X, i \in I_p \}$ are undefined when $t_A^r(x) = i_A^r(x) = f_A^r(x) = 0$ and $t_B^r(x) = i_B^r(x) = f_B^r(x) = 0$ for all $x \in X$ and $i \in I_p$.

Proposition 2. *Let $\tilde{A}, \tilde{B} \in SVN-R_X^p$. Then, each weighted similarity measure $W(\tilde{A}, \tilde{B})_\Lambda (\Lambda = J, D, C)$ satisfies the following properties:*

1. $0 \leq W(\tilde{A}, \tilde{B})_\Lambda \leq 1$,
2. $W(\tilde{A}, \tilde{B})_\Lambda = W(\tilde{B}, \tilde{A})_\Lambda$,
3. $W(\tilde{A}, \tilde{B})_\Lambda = 1$ if $\tilde{B} = \tilde{A}$ i.e. $t_A^r(x_j) = t_B^r(x_j)$, $i_A^r(x_j) = i_B^r(x_j)$, and $f_A^r(x_j) = f_B^r(x_j)$ for every $x_j \in X$ and $r \in I_p$.

PROOF. The proofs can be made similar way to proof of Proposition 1.

Note that, if $w_j (j = 1, 2, \dots, n)$ values take as $\frac{1}{n}$, Eqs. (10), (11) and (12) are reduced Eqs. (4), (5) and (6), respectively.

Now similarity measures between two INR-sets will be defined as a extension of similarity measures between two IN-sets given in [34].

For convenience, $\ddot{A} = \left\{ \left\langle x, ([t_A^r(x), t_A^u(x)]), ([i_A^r(x), i_A^u(x)]), ([f_A^r(x), f_A^u(x)]) \right\rangle : x \in X \text{ and } r \in I_p \right\}$ will be meant by $\ddot{A} \in INR_X$

Definition 15. Let $\ddot{A}, \ddot{B} \in INR_X^p$. Then, the Jaccard similarity measure between INR-sets \ddot{A} and \ddot{B} is defined as follows:

$$(\ddot{A}, \ddot{B})_J = \frac{1}{p.n} \sum_{j=1}^n \sum_{r=1}^p \frac{\left(\begin{array}{c} (t^r_L(x_j)t^r_L(x_j) + t^r_U(x_j)t^r_U(x_j)) \\ + (i^r_L(x_j)i^r_L(x_j) + i^r_U(x_j)i^r_U(x_j)) \\ + (f^r_L(x_j)f^r_L(x_j) + f^r_U(x_j)f^r_U(x_j)) \end{array} \right)}{\left(\begin{array}{c} ([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2) \\ + ([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2) \\ - [t^r_L(x_j)t^r_L(x_j) + i^r_L(x_j)i^r_L(x_j) + f^r_L(x_j)f^r_L(x_j)] \\ - [t^r_U(x_j)t^r_U(x_j) + i^r_U(x_j)i^r_U(x_j) + f^r_U(x_j)f^r_U(x_j)] \end{array} \right)}. \quad (15)$$

Definition 16. Let $\ddot{A}, \ddot{B} \in INR_X^p$. Then, the Dice similarity measure between INR-sets \ddot{A} and \ddot{B} is defined as follows:

$$(\ddot{A}, \ddot{B})_D = \frac{1}{p.n} \sum_{j=1}^n \sum_{r=1}^p \frac{2 \left(\begin{array}{c} (t^r_L(x_j)t^r_L(x_j) + t^r_U(x_j)t^r_U(x_j)) \\ + (i^r_L(x_j)i^r_L(x_j) + i^r_U(x_j)i^r_U(x_j)) \\ + (f^r_L(x_j)f^r_L(x_j) + f^r_U(x_j)f^r_U(x_j)) \end{array} \right)}{\left(\begin{array}{c} ([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2) \\ + ([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2) \end{array} \right)}. \quad (16)$$

Definition 17. Let $\ddot{A}, \ddot{B} \in INR_X^p$. Then the cosine similarity measure between \ddot{A} and \ddot{B} is defined as follows:

$$(\ddot{A}, \ddot{B})_C = \frac{1}{p.n} \sum_{j=1}^n \sum_{r=1}^p \frac{\left(\begin{array}{c} (t^r_L(x_j)t^r_L(x_j) + t^r_U(x_j)t^r_U(x_j)) \\ + (i^r_L(x_j)i^r_L(x_j) + i^r_U(x_j)i^r_U(x_j)) \\ + (f^r_L(x_j)f^r_L(x_j) + f^r_U(x_j)f^r_U(x_j)) \end{array} \right)}{\left(\begin{array}{c} \sqrt{([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2)} \\ \sqrt{([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2)} \end{array} \right)}. \quad (17)$$

If $w_j \in [0, 1]$ be the weight of each element x_j for $j = 1, 2, \dots, n$ such that $\sum_{j=1}^n w_j = 1$, then the weighted Jaccard, Dice and Cosine similarity measures between INR-sets \ddot{A} and \ddot{B} is defined as follows:

$$W(\ddot{A}, \ddot{B})_J = \frac{1}{p} \sum_{j=1}^n \sum_{r=1}^p w_j \frac{\left(\begin{array}{c} (t^r_L(x_j)t^r_L(x_j) + t^r_U(x_j)t^r_U(x_j)) \\ + (i^r_L(x_j)i^r_L(x_j) + i^r_U(x_j)i^r_U(x_j)) \\ + (f^r_L(x_j)f^r_L(x_j) + f^r_U(x_j)f^r_U(x_j)) \end{array} \right)}{\left(\begin{array}{c} ([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2) \\ + ([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2) \\ - [t^r_L(x_j)t^r_L(x_j) + i^r_L(x_j)i^r_L(x_j) + f^r_L(x_j)f^r_L(x_j)] \\ - [t^r_U(x_j)t^r_U(x_j) + i^r_U(x_j)i^r_U(x_j) + f^r_U(x_j)f^r_U(x_j)] \end{array} \right)}, \quad (18)$$

$$W(\ddot{A}, \ddot{B})_D = \frac{1}{p} \sum_{j=1}^n \sum_{r=1}^p w_j \frac{2 \left(\begin{array}{c} (t^r_L(x_j)t^r_L(x_j) + t^r_U(x_j)t^r_U(x_j)) \\ + (i^r_L(x_j)i^r_L(x_j) + i^r_U(x_j)i^r_U(x_j)) \\ + (f^r_L(x_j)f^r_L(x_j) + f^r_U(x_j)f^r_U(x_j)) \end{array} \right)}{\left(\begin{array}{c} ([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2) \\ + ([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2) \end{array} \right)}. \quad (19)$$

and

$$W(\ddot{A}, \ddot{B})_C = \frac{1}{p} \sum_{j=1}^n \sum_{r=1}^p w_j \frac{\left(\begin{array}{c} (t^r_L(x_j)t^r_L(x_j) + t^r_U(x_j)t^r_U(x_j)) \\ + (i^r_L(x_j)i^r_L(x_j) + i^r_U(x_j)i^r_U(x_j)) \\ + (f^r_L(x_j)f^r_L(x_j) + f^r_U(x_j)f^r_U(x_j)) \end{array} \right)}{\left(\frac{\sqrt{([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2)}}{\sqrt{([t^r_L(x_j)]^2 + [i^r_L(x_j)]^2 + [f^r_L(x_j)]^2) + ([t^r_U(x_j)]^2 + [i^r_U(x_j)]^2 + [f^r_U(x_j)]^2)}} \right)}, \quad (20)$$

respectively.

Proposition 3. $\ddot{A}, \ddot{B} \in INR_X^p$. Then, each similarity measure $(\ddot{A}, \ddot{B})_\Lambda (\Lambda = J, D, C)$ satisfies the following properties:

1. $0 \leq (\ddot{A}, \ddot{B})_\Lambda \leq 1$
2. $(\ddot{A}, \ddot{B})_\Lambda = (\ddot{B}, \ddot{A})_\Lambda$;
3. $(\ddot{A}, \ddot{B})_\Lambda = 1$ if $\ddot{B} = \ddot{A}$ i.e. $[t^r_L(x_j), t^r_U(x_j)] = [t^r_L(x_j), t^r_U(x_j)]$, $[i^r_L(x_j), i^r_U(x_j)] = [i^r_L(x_j), i^r_U(x_j)]$, and $[f^r_L(x_j), f^r_U(x_j)] = [f^r_L(x_j), f^r_U(x_j)]$ for all $x_j \in X$ and $r \in I_p$.

PROOF. The proofs can be made similar way to proof of Proposition 1.

Proposition 4. $\ddot{A}, \ddot{B} \in INR_X^p$. Then, each weighted similarity measure $W(\ddot{A}, \ddot{B})_\Lambda (\Lambda = J, D, C)$ satisfies the following properties:

1. $0 \leq W(\ddot{A}, \ddot{B})_\Lambda \leq 1$
2. $W(\ddot{A}, \ddot{B})_\Lambda = W(\ddot{B}, \ddot{A})_\Lambda$;
3. $W(\ddot{A}, \ddot{B})_\Lambda = 1$ if $\ddot{B} = \ddot{A}$ i.e. $[t^r_L(x_j), t^r_U(x_j)] = [t^r_L(x_j), t^r_U(x_j)]$, $[i^r_L(x_j), i^r_U(x_j)] = [i^r_L(x_j), i^r_U(x_j)]$, and $[f^r_L(x_j), f^r_U(x_j)] = [f^r_L(x_j), f^r_U(x_j)]$ for every $x_j \in X$ and $r \in I_p$.

PROOF. The proofs can be made similar way to proof of Proposition 1.

Note that if $[t^r_L(x), t^r_U(x)] = [0, 0]$, $[i^r_L(x), i^r_U(x)] = [0, 0]$, $[f^r_L(x), f^r_U(x)] = [0, 0]$ and $[t^r_L(x), t^r_U(x)] = [0, 0]$, $[i^r_L(x), i^r_U(x)] = [0, 0]$, $[f^r_L(x), f^r_U(x)] = [0, 0]$, each similarity measure between two INR-sets \ddot{A} and \ddot{B} are undefined.

4. Similarity measure based multicriteria decision making under SVN- and INR-environment

In this section, applications of weighted similarity measures in multicriteria decision making problems under SVN- and INR-environment are given.

Let us consider a MCDM problem with k alternatives and r criteria. Let $A = \{A_1, A_2, \dots, A_k\}$ be a set of alternatives and $C = \{C_1, C_2, \dots, C_r\}$ be the set of criteria and $w = \{w_1, w_2, \dots, w_r\}$ be weights of the criteria $C_j (j = 1, 2, \dots, r)$ such that $w_j \geq 0 (j = 1, 2, \dots, r)$ and $\sum_{j=1}^r w_j = 1$.

4.1. Multi-criteria decision making under SVN-R-environment

Let $\{A_1, A_2, \dots, A_k\}$ be a set of alternatives and $\{C_1, C_2, \dots, C_s\}$ be a set of criterion. Alternatives $A_i (i = 1, 2, \dots, k)$ are characterized by SVN-R-values for each $C_j (j = 1, 2, \dots, s)$ as follows:

$$A_i = \left\{ \langle C_j, (t_{A_i}^1(C_j), \dots, t_{A_i}^p(C_j)), (i_{A_i}^1(C_j), \dots, i_{A_i}^p(C_j)), (f_{A_i}^1(C_j), \dots, f_{A_i}^p(C_j)) \rangle : C_j \in C \right\}.$$

For the sake of shortness, $(t_{A_i}^1(C_j), \dots, t_{A_i}^p(C_j))$, $(i_{A_i}^1(C_j), \dots, i_{A_i}^p(C_j))$ and $(f_{A_i}^1(C_j), \dots, f_{A_i}^p(C_j))$ are denoted by $(t_{ij}^1, \dots, t_{ij}^p)$, $(i_{ij}^1, \dots, i_{ij}^p)$ and $(f_{ij}^1, \dots, f_{ij}^p)$, respectively. Thus, the evaluation of the alternative A_i with respect to the criteria C_j made by expert or decision maker can be briefly written as $\gamma_{ij} = \langle (t_{ij}^1, \dots, t_{ij}^p), (i_{ij}^1, \dots, i_{ij}^p), (f_{ij}^1, \dots, f_{ij}^p) \rangle (i = 1, 2, \dots, k; j = 1, 2, \dots, r)$. Hence, SVN-R-decision matrix $D = [\gamma_{ij}]_{k \times s}$ can be constructed.

In MCDM environment, to characterize the best alternative properly in the decision set the notion of the ideal point is used. In order to evaluate the criteria, two type modifiers called benefit criteria (BC) and cost criteria (CC) are used generally.

In this study, for benefit criteria (BC) and cost criteria (CC) ideal SVN-R-values are defined, respectively, as follows:

- $\gamma_j^* = \langle (t_j^{1*}, \dots, t_j^{p*}), (i_j^{1*}, \dots, i_j^{p*}), (f_j^{1*}, \dots, f_j^{p*}) \rangle = \left\langle \begin{array}{c} (max_i(t_{ij}^1), \dots, max_i(t_{ij}^p)), (min_i(i_{ij}^1), \dots, min_i(i_{ij}^p)), \\ (min_i(f_{ij}^1), \dots, min_i(f_{ij}^p)) \end{array} \right\rangle$
- $\gamma_j^* = \langle (t_j^{1*}, \dots, t_j^{p*}), (i_j^{1*}, \dots, i_j^{p*}), (f_j^{1*}, \dots, f_j^{p*}) \rangle = \left\langle \begin{array}{c} (min_i(t_{ij}^1), \dots, min_i(t_{ij}^p)), (max_i(i_{ij}^1), \dots, max_i(i_{ij}^p)), \\ (max_i(f_{ij}^1), \dots, max_i(f_{ij}^p)) \end{array} \right\rangle,$

Here equations are called positive ideal solution and negative ideal solution, respectively. Also set of ideal SVN-R-values will be denoted by A^* .

Algorithm

Step 1: Determination of BC and CC criteria

Step 2: Determination of ideal SVN-R-values A^*

Step 3: Calculation of weighted similarity measures

In this step, using one of the Eq. (10), Eq. (11) or Eq.(12) weighted similarity measures between the ideal alternative A^* and $A_i (i = 1, 2, \dots, k)$ are calculated.

Step 4: Ranking of the alternative

Considering the values obtained using one of the Eq. (10), Eq. (11) or Eq.(12), the ranking order of all the alternatives can be easily determined.

Example 3. Let us consider the decision making problem given in [36]. We adapt this decision making problem to SVN-R-set. There is an investment company, which wants to invest a sum money in the best option. There is a list of four possible alternatives so as to invest the money: (1) A_1 is a car company; (2) A_2 is a food company; (3) A_3 is a computer company; (4) A_4 is an arms company. The investment company must take a decision according to the three criteria (1) C_1 is the risk; (2) C_2 is the growth; (3) C_3 is an environmental impact, The weights of criteria C_1, C_2 and C_3 are given by $w_1 = 0.35, w_2 = 0.25$ and $w_3 = 0.40$, respectively. The four alternatives

are evaluated under the criteria by SVNR-values provided by decision maker. These values are shown in SVNR-decision matrix as follows:

$$D = [\gamma_{ij}]_{k \times r} = \begin{pmatrix} \langle (.1, .2, .4), (.3, .3, .5), (.2, .4, .8) \rangle & \langle (.1, .4, .6), (.2, .3, .7), (.1, .3, .3) \rangle & \langle (.0, .3, .3), (.1, .5, .6), (.5, .6, .9) \rangle \\ \langle (.5, .6, .7), (.2, .4, .4), (.1, .6, .6) \rangle & \langle (.4, .6, .7), (.2, .5, .8), (.1, .5, .5) \rangle & \langle (.3, .3, .4), (.2, .6, .7), (.3, .4, .7) \rangle \\ \langle (.3, .3, .5), (.0, .1, .3), (.1, .4, .7) \rangle & \langle (.4, .5, .6), (.2, .3, .6), (.1, .3, .4) \rangle & \langle (.1, .3, .4), (.1, .4, .6), (.3, .3, .5) \rangle \\ \langle (.2, .4, .9), (.1, .5, .6), (.3, .5, .1) \rangle & \langle (.0, .2, .4), (.1, .5, .7), (.6, .7, .9) \rangle & \langle (.8, .8, .9), (.3, .4, .4), (.6, .6, .8) \rangle \end{pmatrix}$$

Step 1: Let us consider C_1 and C_2 as benefit criteria and C_3 as cost criterion.

Step 2: By using Eqs (4.1), (4.1) and SVNR-decision matrix, set of ideal SVNR-values A^* are obtained as follows:

$$A^* = \left\{ \langle (.5, .6, .9), (.0, .1, .3), (.1, .4, .6) \rangle, \langle (.4, .6, .7), (.1, .3, .6), (.1, .3, .3) \rangle, \langle (.0, .3, .3), (.3, .6, .7), (.6, .6, .9) \rangle \right\}.$$

Step 3: By using the Eqs. (7), (10), (8), (11), (9) and (12), for $\Lambda \in \{J, D, C\}$, similarity measures and weighted similarity measures are obtained as in Table 3

Table 3: Similarity measure values under SVNR-environment

Similarity measure	$i = 1$	$i=2$	$i=3$	$i=4$	Ranking order
$(A^*, A_i)_J$	0.74683	0.87717	0.84399	0.58717	$A_2 \succ A_3 \succ A_1 \succ A_4$
$(A^*, A_i)_D$	0.82330	0.92951	0.91115	0.71588	$A_2 \succ A_3 \succ A_1 \succ A_4$
$(A^*, A_i)_C$	0.84663	0.94562	0.96771	0.73688	$A_3 \succ A_2 \succ A_1 \succ A_4$
$W(A^*, A_i)_J$	0.74934	0.87590	0.85143	0.80483	$A_2 \succ A_3 \succ A_4 \succ A_1$
$W(A^*, A_i)_D$	0.82887	0.92437	0.90083	0.75112	$A_2 \succ A_3 \succ A_1 \succ A_4$
$W(A^*, A_i)_C$	0.85016	0.94076	0.96521	0.81158	$A_3 \succ A_2 \succ A_1 \succ A_4$

Step 4: Rankings of the alternatives are shown in last column of Table 3.

4.2. Comparative analysis of various similarity measures under SVNR-environment

In this section, proposed similarity measure methods in this paper are compared with existing similarity measures methods given in references [9], [10], [16] and [17]. Under SVNR-values given in Example 3, results of similarity measures and ordering of alternatives are shown in Table 4.

Table 4: Similarity measure values under SVNR-environment

Similarity measure	$i = 1$	$i=2$	$i=3$	$i=4$	Ranking order
$(A^*, A_i)_J$	0.74683	0.87717	0.84399	0.58717	$A_2 \succ A_3 \succ A_1 \succ A_4$
$(A^*, A_i)_D$	0.82330	0.92951	0.91115	0.71588	$A_2 \succ A_3 \succ A_1 \succ A_4$
$(A^*, A_i)_C$	0.84663	0.94562	0.96771	0.73688	$A_3 \succ A_2 \succ A_1 \succ A_4$
$C_{NRS}(A^*, A_i)$ [9]	0.96485	0.97138	0.96352	0.90499	$A_2 \succ A_1 \succ A_3 \succ A_4$
$S_{HNRS}^1(A^*, A_i)$ [10]	0.74444	0.78889	0.75556	0.50000	$A_2 \succ A_3 \succ A_1 \succ A_4$
$S_{HNRS}^2(A^*, A_i)$ [10]	0.64324	0.69892	0.65693	0.37754	$A_2 \succ A_3 \succ A_1 \succ A_4$
$S_{HNRS}^3(A^*, A_i)$ [10]	0.59292	0.65138	0.60714	0.33333	$A_2 \succ A_3 \succ A_1 \succ A_4$
$COT_{NRS}(A^*, A_i)$ [16]	0.80729	0.82576	0.79722	0.64835	$A_2 \succ A_1 \succ A_3 \succ A_4$
$T_{NRS}(A^*, A_i)$ [17]	0.88829	0.98252	0.96212	0.89749	$A_2 \succ A_3 \succ A_4 \succ A_1$

Note that, it is shown that similarity measure methods proposed in the paper is generally coherent by other similarity measure methods proposed before.

4.3. Multi-attribute decision making under INR-environment

Let alternatives $A_i (i = 1, 2, \dots, k)$ are characterized by INR-values for each criterion $C_j (j = 1, 2, \dots, s)$ as follows:

$$A_i = \left\{ \langle C_j, ([t_{A_i}^{1L}(C_j), t_{A_i}^{1U}(C_j)], \dots, [t_{A_i}^{pL}(C_j), t_{A_i}^{pU}(C_j)]), ([i_{A_i}^{1L}(C_j), i_{A_i}^{1U}(C_j)], \dots, [i_{A_i}^{pL}(C_j), i_{A_i}^{pU}(C_j)]), ([f_{A_i}^{1L}(C_j), f_{A_i}^{1U}(C_j)], \dots, [f_{A_i}^{pL}(C_j), f_{A_i}^{pU}(C_j)]) \rangle : C_j \in C \right\}.$$

For convenience, $([t_{A_i}^{1L}(C_j), t_{A_i}^{1U}(C_j)], \dots, [t_{A_i}^{pL}(C_j), t_{A_i}^{pU}(C_j)]), ([i_{A_i}^{1L}(C_j), i_{A_i}^{1U}(C_j)], \dots, [i_{A_i}^{pL}(C_j), i_{A_i}^{pU}(C_j)])$ and $([f_{A_i}^{1L}(C_j), f_{A_i}^{1U}(C_j)], \dots, [f_{A_i}^{pL}(C_j), f_{A_i}^{pU}(C_j)])$ are denoted by $([t_{ij}^{1L}], \dots, [t_{ij}^{pU}]), ([i_{ij}^{1L}], \dots, [i_{ij}^{pU}])$ and $([f_{ij}^{1L}], \dots, [f_{ij}^{pU}])$, respectively. So INR-value $\theta_{ij} = \langle ([t_{ij}^{1L}], \dots, [t_{ij}^{pU}]), ([i_{ij}^{1L}], \dots, [i_{ij}^{pU}]), ([f_{ij}^{1L}], \dots, [f_{ij}^{pU}]) \rangle (i = 1, 2, \dots, k; j = 1, 2, \dots, s)$ which is generally obtained from the evaluation of the alternative A_i with related to the criteria C_j by opinion of expert or decision maker. Thus, INR-decision matrix $D = [\theta_{ij}]_{k \times s}$ can be constructed.

Here, for benefit criteria (BC) and cost criteria (CC) ideal INR-values are defined, respectively, as follows:

$$\bullet \theta_j^* = \langle (t_j^{1*}, \dots, t_j^{p*}), (i_j^{1*}, \dots, i_j^{p*}), (f_j^{1*}, \dots, f_j^{p*}) \rangle = \left\langle \begin{array}{l} ([\max_i(t_{ij}^{1L}), \max_i(t_{ij}^{1U})], \dots, [\max_i(t_{ij}^{pL}), \max_i(t_{ij}^{pU})]), \\ ([\min_i(i_{ij}^{1L}), \min_i(i_{ij}^{1U})], \dots, [\min_i(i_{ij}^{pL}), \min_i(i_{ij}^{pU})]), \\ ([\min_i(f_{ij}^{1L}), \min_i(f_{ij}^{1U})], \dots, [\min_i(f_{ij}^{pL}), \min_i(f_{ij}^{pU})]) \end{array} \right\rangle$$

$$\bullet \theta_j^* = \langle (t_j^{1*}, \dots, t_j^{p*}), (i_j^{1*}, \dots, i_j^{p*}), (f_j^{1*}, \dots, f_j^{p*}) \rangle = \left\langle \begin{array}{l} ([\min_i(t_{ij}^{1L}), \min_i(t_{ij}^{1U})], \dots, [\min_i(t_{ij}^{pL}), \min_i(t_{ij}^{pU})]), \\ ([\max_i(i_{ij}^{1L}), \max_i(i_{ij}^{1U})], \dots, [\max_i(i_{ij}^{pL}), \max_i(i_{ij}^{pU})]), \\ ([\max_i(f_{ij}^{1L}), \max_i(f_{ij}^{1U})], \dots, [\max_i(f_{ij}^{pL}), \max_i(f_{ij}^{pU})]) \end{array} \right\rangle,$$

Here equations are called positive ideal solution and negative ideal solution, respectively. Also set of ideal INR-values will be denoted by \check{A}^* .

Algorithm

Step 1: Determination of BC and CC criteria

Step 2: Determination of ideal INR-values \check{A}^* solution

Step 3: Calculation of weighted similarity measures

In this step, using one of the Eq. (18), Eq. (19) or Eq.(20) weighted similarity measures between the ideal alternative \check{A}^* and INR-sets $A_i (i = 1, 2, \dots, k)$ are calculated.

Step 4: Ranking of the alternative

According to the values obtained using one of the Eq. (18), Eq. (19) or Eq.(20), the ranking order of all the alternatives can be easily determined.

Example 4. In this example, alternatives and criteria given in previous illustrative example will be considered under INR-environment. The four alternatives are to evaluated under the criteria by INR-values provided by decision maker. These values are shown in INR-decision matrix as follows:

$$D = [\theta_{ij}]_{k \times r} = \left(\begin{array}{ccc} \left\langle \begin{array}{l} ([.2, .3], [.2, .5], [.4, .7]), \\ ([.3, .4], [.3, .6], [.5, .9]), \\ ([.2, .5], [.4, .7], [.8, .8]) \end{array} \right\rangle & \left\langle \begin{array}{l} ([.1, .5], [.4, .5], [.6, 1]), \\ ([.2, .4], [.3, .7], [.7, .8]), \\ ([.1, .2], [.3, .8], [.3, .8]) \end{array} \right\rangle & \left\langle \begin{array}{l} ([.0, .3], [.3, .5], [.3, .9]), \\ ([.1, .2], [.5, .6], [.6, .6]), \\ ([.5, .5], [.6, .7], [.9, .9]) \end{array} \right\rangle \\ \left\langle \begin{array}{l} ([.1, .2], [.2, .8], [.4, .8]), \\ ([.4, .5], [.3, .6], [.5, .7]), \\ ([.1, .3], [.4, .5], [.8, .8]) \end{array} \right\rangle & \left\langle \begin{array}{l} ([.1, .4], [.4, .5], [.6, .6]), \\ ([.2, .3], [.3, .4], [.7, .8]), \\ ([.1, .5], [.3, .6], [.3, .7]) \end{array} \right\rangle & \left\langle \begin{array}{l} ([.0, .3], [.3, .4], [.3, .5]), \\ ([.1, .6], [.5, .6], [.6, .7]), \\ ([.5, .8], [.6, .8], [.9, 1]) \end{array} \right\rangle \\ \left\langle \begin{array}{l} ([.1, .4], [.2, .5], [.4, .6]), \\ ([.3, .4], [.3, .4], [.6, .7]), \\ ([.2, .3], [.4, .5], [.8, 1]) \end{array} \right\rangle & \left\langle \begin{array}{l} ([.2, .3], [.4, .5], [.6, .7]), \\ ([.2, .5], [.3, .6], [.7, .8]), \\ ([.1, .2], [.3, .4], [.4, .5]) \end{array} \right\rangle & \left\langle \begin{array}{l} ([.0, .1], [.3, .3], [.3, .4]), \\ ([.1, .2], [.5, .6], [.6, .7]), \\ ([.5, .6], [.6, .7], [.9, .9]) \end{array} \right\rangle \\ \left\langle \begin{array}{l} ([.1, .4], [.2, .4], [.4, .4]), \\ ([.3, .5], [.3, .6], [.5, .6]), \\ ([.2, .5], [.4, .6], [.8, .9]) \end{array} \right\rangle & \left\langle \begin{array}{l} ([.1, .5], [.4, .5], [.4, .6]), \\ ([.2, .4], [.3, .5], [.7, .9]), \\ ([.2, .2], [.3, .4], [.3, .4]) \end{array} \right\rangle & \left\langle \begin{array}{l} ([.0, .2], [.3, .4], [.3, .5]), \\ ([.1, .4], [.5, .6], [.6, .8]), \\ ([.5, .6], [.6, .7], [.9, 1]) \end{array} \right\rangle \end{array} \right)$$

Step 1: Let us consider C_1 and C_2 as benefit criteria and C_3 as cost criterion.

Step 2: From INR-decision matrix, ideal alternative \ddot{A}^* can be obtained as follows:

$$\ddot{A}^* = \left\{ \langle ([.2, .4], [.2, .8], [.4, .8]), ([.3, .4], [.3, .4], [.5, .6]), ([.1, .3], [.4, .5], [.8, .8]) \rangle, \right. \\ \langle ([.2, .5], [.4, .5], [.6, 1]), ([.2, .3], [.3, .4], [.7, .8]), ([.1, .2], [.3, .4], [.3, .4]) \rangle \\ \left. \langle ([.0, .1], [.3, .3], [.3, .4]), ([.1, .6], [.5, .6], [.6, .8]), ([.5, .8], [.6, .8], [.9, 1]) \rangle \right\}.$$

Step 3: By using the Eq. (18), Eq. (19) and Eq. (20) similarity measures and weighted similarity measures are obtained as in Table 5.

Step 4: Rankings of the alternatives are shown in last column of Table 5.

Similarity measure	$i = 1$	$i=2$	$i=3$	$i=4$	Ranking order
$(\ddot{A}^*, A_i)_J$	0.89674	0.94268	0.93521	0.93702	$A_2 \succ A_4 \succ A_3 \succ A_1$
$(\ddot{A}^*, A_i)_D$	0.94405	0.96944	0.96546	0.96689	$A_2 \succ A_4 \succ A_3 \succ A_1$
$(\ddot{A}^*, A_i)_C$	0.95629	0.97093	0.97344	0.97320	$A_2 \succ A_3 \succ A_4 \succ A_1$
$W(\ddot{A}^*, A_i)_J$	0.89348	0.94992	0.93656	0.93731	$A_2 \succ A_4 \succ A_3 \succ A_1$
$W(\ddot{A}^*, A_i)_D$	0.94212	0.97339	0.96612	0.96701	$A_2 \succ A_4 \succ A_3 \succ A_1$
$W(\ddot{A}^*, A_i)_C$	0.95421	0.97463	0.65693	0.97360	$A_3 \succ A_2 \succ A_4 \succ A_1$

5. Consistency analysis of similarity measures based INR-sets

In this section, so as to determine which similarity measure gives more consistent results, a methods is developed.

Let $A = \{A_1, A_2, \dots, A_k\}$ be a set of alternatives, $C = \{C_1, C_2, \dots, C_s\}$ be a set of criteria and \ddot{A}^* be set of ideal alternative values obtained from decision matrix defined in illustrative example of similarity measures based on INR-set. Then, consistencies of the similarity measures and weighted similarity measures based INR-values are defined, respectively, as follows:

$$Con(\Lambda) = \frac{1}{k} \sum_{i=1}^k |(\ddot{A}^{*L}, A_i^L)_\Lambda - (\ddot{A}^{*U}, A_i^U)_\Lambda|.$$

$$ConW(\Lambda) = \frac{1}{k} \sum_{i=1}^k |W(\ddot{A}^{*L}, A_i^L)_\Lambda - W(\ddot{A}^{*U}, A_i^U)_\Lambda|.$$

Here, \ddot{A}^{*L} and \ddot{A}^{*U} are determined with help of INR-decision matrix using formula of benefit criteria (BC) and cost criteria (CC) given as follows: For $\Delta \in \{L = lower, U = upper\}$

$$\begin{aligned} \bullet \delta_j^{\Delta} &= \langle (t_j^{1*\Delta}, \dots, t_j^{p*\Delta}), (i_j^{1*\Delta}, \dots, i_j^{p*\Delta}), (f_j^{1*\Delta}, \dots, f_j^{p*\Delta}) \rangle = \left\langle \begin{array}{l} (max_i(t_{ij}^{1\Delta}), \dots, max_i(t_{ij}^{p\Delta})), \\ (min_i(i_{ij}^{1\Delta}), \dots, min_i(i_{ij}^{p\Delta})), \\ (min_i(f_{ij}^{1\Delta}), \dots, min_i(f_{ij}^{p\Delta})) \end{array} \right\rangle \\ \bullet \delta_j^{\Delta} &= \langle (t_j^{1*\Delta}, \dots, t_j^{p*\Delta}), (i_j^{1*\Delta}, \dots, i_j^{p*\Delta}), (f_j^{1*\Delta}, \dots, f_j^{p*\Delta}) \rangle = \left\langle \begin{array}{l} (min_i(t_{ij}^{1\Delta}), \dots, min_i(t_{ij}^{p\Delta})), \\ (max_i(i_{ij}^{1\Delta}), \dots, max_i(i_{ij}^{p\Delta})), \\ (max_i(f_{ij}^{1\Delta}), \dots, max_i(f_{ij}^{p\Delta})) \end{array} \right\rangle, \end{aligned}$$

respectively.

$$\text{Also } \ddot{A}^{*L} = \left\{ \langle (t_j^{1*L}, \dots, t_j^{p*L}), (i_j^{1*L}, \dots, i_j^{p*L}), (f_j^{1*L}, \dots, f_j^{p*L}) : i \in I_p \right\} \text{ and}$$

$$\ddot{A}^{*U} = \left\{ \langle (t_j^{1*U}, \dots, t_j^{p*U}), (i_j^{1*U}, \dots, i_j^{p*U}), (f_j^{1*U}, \dots, f_j^{p*U}) : i \in I_p \right\}.$$

Example 5. Let us consider the Example 4. Then for all $\Lambda \in \{J, D, C\}$, results and orderings are obtained as in Table 6,

Note that, $Con(\ddot{A}^*, A_i)_J \geq Con(\ddot{A}^*, A_i)_D \geq Con(\ddot{A}^*, A_i)_C$. Since consistency degree of Jaccard similarity measure under INR environment is higher than consistency degrees of Dice and Cosine similarity measures, it is more convenient using the Jaccard similarity measure for discussed problem. If weighted similarity measures are considered, on the grounds that $ConW(\ddot{A}^*, A_i)_D \geq ConW(\ddot{A}^*, A_i)_C \geq ConW(\ddot{A}^*, A_i)_J$, it is more convenient using the Dice similarity measure for discussed problem.

6. Conclusion

In this paper, three similarity measure methods are developed for SVN-R-set and INR-sets based on Jaccard, Dice and Cosine similarity measures. Furthermore, applications of proposed similarity measure methods are given in multi-criteria decision making, and is given comparative analysis of various similarity measures under SVN-R-environment. Also a method is developed in order to compare similarity measures of INR-sets, and an application of this method is given. However, I hope that the main thrust of proposed formulas will be in the field of equipment evaluation, data mining and investment decision making. Also in future, similarity measure methods for INR-sets can be proposed based on the methods other than Jaccard, Dice and Cosine similarity measures.

References

- [1] K. Atanassov, Intuitionistic fuzzy sets, Fuzzy Set and Systems, 20, 87-96, (1986).
- [2] K. Atanassov and G. Gargov, Interval valued intuitionistic fuzzy sets, Fuzzy Set and Systems, 31(3), 343-349, (1989).

Table 6: Similarity measure values and ranking of alternatives under INR-environment

	$i = 1$	$i = 2$	$i = 3$	$i = 4$	$Con(\ddot{A}^*, A_i)_\Lambda$
$\{\ddot{A}^L, A_i\}_J, (\ddot{A}^U, A_i)_J\}$	{0.97917, 0.86550}	{0.97304, 0.93113}	{0.98305, 0.91904}	{0.95780, 0.92561}	0.06294
$\{\ddot{A}^L, A_i\}_D, (\ddot{A}^U, A_i)_D\}$	{0.98901, 0.92504}	{0.98565, 0.96309}	{0.99099, 0.95590}	{0.97707, 0.96035}	0.03459
$\{\ddot{A}^C, A_i\}_C, (\ddot{A}^U, A_i)_C\}$	{0.99165, 0.94804}	{0.98861, 0.96641}	{0.99120, 0.97189}	{0.97785, 0.97092}	0.02301
	$i = 1$	$i = 2$	$i = 3$	$i = 4$	$ConW(\ddot{A}^*, A_i)_\Lambda$
$\{W(\ddot{A}^L, A_i)_J, W(\ddot{A}^U, A_i)_J\}$	{0.90889, 0.88059}	{0.90245, 0.89762}	{0.89957, 0.89865}	{0.90062, 0.87023}	0.01611
$\{W(\ddot{A}^L, A_i)_D, W(\ddot{A}^U, A_i)_D\}$	{0.99068, 0.92172}	{0.98715, 0.96807}	{0.99071, 0.95672}	{0.98042, 0.95969}	0.03569
$\{W(\ddot{A}^L, A_i)_C, W(\ddot{A}^U, A_i)_C\}$	{0.99282, 0.94434}	{0.98962, 0.97080}	{0.99091, 0.97080}	{0.98101, 0.97093}	0.02345

- [3] A.Q. Ansari, R. Biswas, S. Aggarwal, Proposal for Applicability of Neutrosophic Set Theory in Medical AI, *International Journal of Computer Applications* 27(5), (2011).
- [4] A.Aydoğdu, On Similarity and Entropy of Single Valued Neutrosophic Sets, *General Mathematical Notes*, 29(1), (2015), 67-74.
- [5] L. Baowen, W. Peizhuang and L. Xihui, Fuzzy Bags and Relations with Set-Valued Statistic, *Computer Mathematics and Application*, 15(10), 811-818, (1988).
- [6] M. Bhowmik and M. Pal, Intuitionistic Neutrosophic Set Relations and Some of Its Properties, *Journal of Information and Computing Science*, 5(3) 183-192, (2010).
- [7] S. Broumi and F. Smarandache, Several similarity measures of neutrosophic sets, *Neutrosophic Sets and Systems* 1(1) (2013), 54-62.
- [8] S. Broumi, F. Smarandache, New Operations on Interval Neutrosophic Sets, *Journal of New Theory*, 1, 24-37, (2015).
- [9] S. Broumi, F. Smarandache, Neutrosophic Refined Similarity Measure Based on Cosine Function, *Neutrosophic Sets and Systems*, 6, (2014), 41-47.
- [10] S. Broumi, F. Smarandache, Extended Hausdorff Distance and Similarity Measures for Neutrosophic Refined Sets and Their Application In Medical Diagnosis, *Journal of New Theory*, 7, (2015), 64-78.
- [11] S. Broumi, F. Smarandache, New Distance and Similarity Measures of Interval Neutrosophic Sets, figshare. <http://dx.doi.org/10.6084/m9.figshare.1502585>
- [12] S. Broumi, I. Deli, F.Smarandache, N-Valued Interval Neutrosophic Sets and Their Application in Medical Diagnosis, *Critical Review. X*, (2015), 54-68.
- [13] I. Deli, S. Broumi and F. Smarandache, On neutrosophic refined sets and their applications in medical diagnosis, *Journal of New Theory*, 6, (2015), 88-98.
- [14] Y.-x. Ma, J.-q. Wang, J. Wang, X.-h Wu, An interval neutrosophic linguistic multi-criteria group decision-making method its application in selecting medical treatment options, *Neural Computing and Applications*, DOI:10.1007/s00521-016-2203-1, 2016.
- [15] P. Majumdar and S.K. Smanata, On Similarity and Entropy of Neutrosophic Sets, *Journal of Intelligent and Fuzzy Systems*, 26(3), (2014), 1245-1252.
- [16] K. Mondal and S. Pramanik, Neutrosophic refined similarity measure based on cotangent function and its application to multi-attribute decision making, *Global Journal of Advanced Research*, 2(2), (2015), 486-496.
- [17] K. Mondal and S. Pramanik, Neutrosophic refined similarity measure based on tangent function and its application to multi-attribute decision making, *Journal of New Theory*, 8, (2015), 41-50.
- [18] W. Peizhuang *Fuzzy Set and Fall-shadow of Random Sets* Publishing House of Beijing Normal University, Beijing, 42-57 (1984).

- [19] J.-j. Peng, J.-q. Wang, H.-y. Zhang, X.-h. Chen, An outranking approach for multi-criteria decision-making problems with simplified neutrosophic sets, *Applied Soft Computing*, 25, (2014), 336-346.
- [20] J.-j. Peng, J.-q. Wang, H.-y. Zhang, X.-h. Chen, Simplified neutrosophic sets and their applications in multi-criteria group decision-making problems, *International journal of Intelligent Systems Science*, DOI:10.1080/00207721.2014.994050
An outranking approach for multi-criteria decision-making problems with simplified neutrosophic sets, *Applied Soft Computing*, 25, (2014), 336-346.
- [21] A. Robinson, *Non-standard analysis*. North-Holland Publishing Co., Amsterdam 1966.
- [22] T. K. Shinoj, J.J. Sunil, *Intuitionistic Fuzzy Multisets And Its Application in Medical Diagnosis*, *World Academy of Science, Engineering and Technology*, 6, 01-28, (2012).
- [23] F. Smarandache, "Neutrosophy, Neutrosophic Probability, Set, and Logic, Amer. Res. Press, Rehoboth, USA, 105p., (1998).
- [24] F. Smarandache, Neutrosophic set - a generalization of the intuitionistic fuzzy set, *International Journal of Pure and Applied Mathematics*, 24/3, 287-297, (2005).
- [25] F. Smarandache, n-Valued Refined Neutrosophic Logic and Its Applications in Physics, *Progress in Physics*, 4, 143-146, (2013).
- [26] Z.-p Tian, H.-y. Zhang, J.-q. Wang, X.-h. Chen, Multi-criteria decision-making method based on a cross-entropy with interval neutrosophic sets, *International journal of Systems Science*, DOI:10.1080/00207721.2015.1102359.
- [27] H. Wang, *Interval Neutrosophic Sets and Logic: Theory and Applications in Computing*, Dissertation, Georgia State University, (2006).
- [28] H. Wang, F. Smarandache Y. Zhang, R. Sunderraman, Single Valued Neutrosophic Sets, Multispace and Multistructure, 4, 410-413, (2010).
- [29] X.-h. Wu, J.-j. Wang, J.-j. Peng, X.-h. Chen, Cross entropy and prioritized aggregation operator with simplified neutrosophic sets and their application in multi-criteria decision making problems, *International Journal of Fuzzy Systems*, DOI: 10.1007/s40815-016-0180-2, 2016.
- [30] L. A. Zadeh, Fuzzy sets, *Information and Control*, 8, 338-353, (1965).
- [31] H.Zhang, J. Wang, and X. Chen, Interval Neutrosophic Sets and Their Application in Multicriteria Decision Making Problems, *The Scientific World Journal* Volume 2014, Article ID 645953.
- [32] R.R. Yager On The Theory of Bags (Multi sets), *International Journal of General Systems*, 13(1), 23-37, (1986).
- [33] J. Ye and Q. Zhang, Single Valued Neutrosophic Similarity Measures for Multiple Attribute Decision-Making, *Neutrosophic Sets and Systems*, 2, (2014), 48-54.
- [34] J. Ye, Vector Similarity Measures of Simplified Neutrosophic Sets and Their Application in Multicriteria Decision Making, *International Journal of Fuzzy Systems*, 16(2), 204-211, (2014).

- [35] S.Ye and J. Ye, Dice Similarity Measure between Single Valued Neutrosophic Multisets and Its Application in Medical Diagnosis, *Neutrosophic Sets and Systems*, 6, 49-54, (2014).
- [36] J. Ye, Similarity measures between interval neutrosophic sets and their applications in multi-criteria decision-making, *Journal of Intelligent and Fuzzy Systems*, 26(1), (2014) 165-172,
- [37] J. Ye, A multicriteria decision-making method using aggregation operators for simplified neutrosophic sets, *J. Intell. Fuzzy Syst.* 26 (5) (2014) 24592466.
- [38] J.Ye, Clustering Methods Using Distance-Based Similarity Measures of Single-Valued Neutrosophic Sets, *Journal of Intelligent Systems*, 23(4), (2014), 48-54.
- [39] J. Ye and J. Fub, Multi-period medical diagnosis method using a single-valued neutrosophic similarity measure based on tangent function, *computer methods and programs in biomedicine* doi: 10.1016/j.cmpb.2015.10.002.
- [40] J. Ye, Single-valued neutrosophic similarity measures based on cotangent function and their application in the fault diagnosis of steam turbine, *Soft Computing*, DOI 10.1007/s00500-015-1818-y
- [41] J. Ye, Trapezoidal neutrosophic set and its application to multiple attribute decision-making, *Neural Computing and Application*, 26, 1157-1166, (2015).
- [42] H.-y. Zhang, P. Ji, J.-q. Wang, X.-h. Chen, A neutrosophic normal cloud and its application in decision-making, *Cognitive Computation*, DOI:10.1007/s12559-016-9394-8,2016.